

**Longitudinal observation of instrumental activities of daily living in Parkinson's disease patients**

Martina Kvapilova<sup>1</sup>, Josef Mana<sup>1</sup>, Daniel Janko<sup>1</sup>, Tereza Uhrova<sup>1</sup>, Evzen Ruzicka<sup>1</sup>, Filip Havlik<sup>1</sup>, Filip Ruzicka<sup>1</sup>, Dusan Urgosik<sup>2</sup>, Robert Jech<sup>1</sup>, and Ondrej Bezdicek<sup>1</sup>

<sup>1</sup>Department of Neurology and Centre of Clinical Neuroscience, First Faculty of Medicine and General University Hospital in Prague, Charles University, Czech Republic

<sup>2</sup>Department of stereotactic and radiation neurosurgery, Na Homolce Hospital, Prague, Czech Republic

### Author Note

Josef Mana  <https://orcid.org/0000-0002-7817-3978>

Filip Havlik  <https://orcid.org/0000-0003-4442-6743>

Robert Jech  <https://orcid.org/0000-0002-9732-8947>

Ondrej Bezdicek  <https://orcid.org/0000-0002-5108-0181>

Author roles were classified using the Contributor Role Taxonomy (CRediT; <https://credit.niso.org/>) as follows: Martina Kvapilova: Conceptualization, Investigation, Data curation, Writing - original draft; Josef Mana: Conceptualization, Data curation, Investigation, Formal analysis, Methodology, Software, Visualization, Writing - original draft; Daniel Janko: Conceptualization, Investigation, Methodology, Writing - original draft; Tereza Uhrova: Investigation; Evzen Ruzicka: Conceptualization, Investigation; Filip Havlik: Investigation; Filip Ruzicka: Investigation; Dusan Urgosik: Investigation; Robert Jech: Funding acquisition; Ondrej Bezdicek: Investigation, Data curation, Funding acquisition, Conceptualization, Project administration, Supervision, Writing - original draft

Correspondence concerning this article should be addressed to

## **Longitudinal observation of instrumental activities of daily living in Parkinson's disease patients**

### **Introduction**

Parkinson's disease (PD) is a progressive neurodegenerative disorder that impacts numerous aspects of day-to-day life. One of the areas of life that is affected early in the disease is the ability to perform activities of daily living (ADL) (Hariz & Forsgren, 2010). Worsening of motor and cognitive symptoms due to PD progression has gradual deteriorating effects on basic ADL (BADL) as well as instrumental activities of daily living (iADL) (Shulman et al., 2008). Given the different cognitive and social demands of BADL and iADL, it is important to distinguish between the two categories. BADL include bathing, toileting, eating, dressing, and personal hygiene (Lawton & Brody, 1969). On the other hand, iADL consist of more complex activities such as shopping, cooking, doing the laundry, or keeping financial records (Lawton & Brody, 1969). Generally, assessment of iADL across different populations has implications in characterizing aging or disease progression (Gold, 2012). Impairments in cognitive-driven iADL have also been shown to be one of potential predictors of severe worsening of cognitive impairment in PD patients (Becker et al., 2018a, 2021). Hence, studying the changes in iADL over time is crucial for a better understanding of disease progressions and treatment effects and better identification of individuals who are at risk of cognitive decline.

Correlates of iADL deterioration in PD have been studied and are persistent across different reports (Martin et al., 2013; Pirogovsky et al., 2013, 2014; Schmitter-Edgecombe et al., 2021). The most pronounced risk factors are PD related mild cognitive impairment (PD-MCI) and PD with dementia (PD-D) (Becker et al., 2022; Martin et al., 2013; Pirogovsky et al., 2013, 2014; Schmitter-Edgecombe et al., 2021). PD-MCI and PD-D are common symptoms of PD progression alongside worsening motor symptoms such as tremors, gait problems, and freezing. However, various therapeutical interventions are available that can alleviate some symptoms and slow down the progress of the disease (Chakraborty et al., 2020).

Subthalamic deep brain stimulation (STN DBS), an invasive neurosurgical treatment, is an option for patients with PD suffering from medication-refractory motor complications.

The motor effects of STN DBS have been studied extensively and are consistently positive (Deuschl et al., 2006). On the other hand, STN DBS has been shown to have little to no effect on cognitive performance and neuropsychiatric symptoms with the exception of specific domains (Accolla & Pollo, 2019; Alegret et al., 2001; Appleby et al., 2007; Bove et al., 2020; Drapier et al., 2006; Limousin & Foltynie, 2019; Mehanna et al., 2017; Parsons et al., 2006; Saint-Cyr, 2000). To this day, the non-motor effects of STN DBS are poorly understood and more research is needed to provide clarity on this topic.

While there have been a number of reports, both short and long-term, on the trajectories of BADL scores in the literature (Aviles-Olmos et al., 2014; Castrioto, 2011; Gervais-Bernard et al., 2009; Lezcano et al., 2016; Limousin & Foltynie, 2019; Piboolnurak et al., 2007; Rodriguez-Oroz et al., 2005; Schupbach, 2005; Zibetti et al., 2011), very few studies have examined the trends in iADL specific measures after STN DBS. To our knowledge, only two studies provide insight into the problematic of short-term trends in iADL (Bezdicek et al., 2022; Gorecka-Mazur et al., 2019). Even though these two studies reported improved iADL scores after STN DBS, we are still lacking a comprehensive understanding of short as well as long-term trends in iADL.

In this study, we try to address this issue by studying iADL with the Functional Activities Questionnaire (FAQ) (Pfeffer et al., 1982). FAQ is an indirect measure of iADL that can be divided into two subcategories - motor-driven and cognitive-driven iADL (Becker et al., 2018a). The two subcategories are distinguishable based on their correlations with cognitive (MoCA) and motor (UPDRS-III) batteries (Becker et al., 2018a). These features of FAQ allow us to examine each subcategory in time on its own and to also compare them to each other across different time points in PD patients undergoing DBS. This will help us draw better-informed conclusions about the overall effect of STN DBS on the patients' lives.

Among others, one of the treatment goals of STN DBS is a long-term improvement of iADL; thus, it is desirable to understand the effects STN DBS has on them. Given the lack of clarity on the post-DBS surgery trends in iADL, we sought to explore the following; (1) iADL scores at 1, 3, 5, 7, 11, and 13 years follow-ups, (2) differences in cognitive- and motor-driven iADL, and (3) the relationship between FAQ trends and trends in other neurological,

psychiatric and psychological measures (PDAQ, BDI, and UPDRS-III).

## Methods

### Participants

A total of ... patients (mean age ... pre-surgery, ...% males) were enrolled in the study. Patients were assessed at ... different time points: (a) before surgery (... months,  $n = \dots$ ), (b) 1 year after surgery (), (c) 3 years after surgery (), (d) 7 years after surgery (), (e) 11 years after surgery (), (f) ... ().

Patients with PD were recruited from the Movement Disorders Center, Department of Neurology, First Faculty of Medicine and General University hospital in Prague. All patients were examined by a trained neurologist specializing in movement disorders and met the UK PD Society Brain Bank criteria (Kubu, 2018). All of the participants were suffering from motor fluctuations and/or disabling dyskinesia and had been indicated for treatment with STN DBS. For clinical and demographic details see Table 1. Exclusion criteria included: PDD based on MDS criteria (), atypical or secondary parkinsonism, depression scores in the severe or moderate range according to Beck Depression Inventory (BDI-II) and psychiatric evaluation, florid psychotic manifestation (hallucination or delusions), anticholinergic medications and other medical or neurological conditions potentially causing cognitive impairment (e.g., epileptic seizures, stroke, head trauma, or tumor). All PD patients were undergoing dopaminergic therapy (i.e., levodopa or dopamine agonist, or a combination of the two), and their levodopa equivalent daily dose was calculated before and after surgery (Tomlinson et al., 2010). Bilateral STN DBS was performed as described in previous reports (JECH et al., 2006; Jech et al., 2012). All patients gave their written informed consent for participation. The study was approved by the Ethics Committee of the General Hospital in Prague, Czechia.

### Assessments

#### *Neuropsychological Assessment*

All patients underwent a comprehensive and clinically relevant pre-surgical evaluation (baseline) consisting of neuropsychological, psychiatric, and neurological examinations by trained movement disorders specialists in individual fields (). The patients were then followed

up 1 (% of the baseline population), 3 (), 5 (), 7 (), 9 (), 11 (), and 13 () years after the surgery. For details about the number of follow-up examinations undergone by each participant, see Fig. 2.

The neuropsychological assessment in pre-test-post-test followed the standard Movement Disorder Society neuropsychological battery at Level 1 for PD-MCI (Bezdzicek et al., 2016, 2017; Litvan et al., 2012) cognitive performance was assessed by Mattis Dementia Rating Scale, second edition (DRS-2) (Bezdzicek et al., 2015; Jurica et al., 2001). iADL and everyday functioning were measured by the PDAQ and FAQ self-reports. Lastly, depressive symptoms were assessed with the Beck Depression Scale (BDI-II) (Beck et al., 1996; Ciharova et al., 2020).

### ***Neurological Examination***

The neurological examination included medical history, medication status, and motor status assessed by the Movement Disorder Society Unified Parkinson's Disease Rating Scale, part three (MDS-UPDRS-III). Scores of the patients that were examined with the older version of the Unified Parkinson's Disease Rating Scale (UPDRS-III) were converted to the MDS-UPDRS III scale using the method described by (Hentz et al., 2015). All PD patients were treated with dopaminergic therapy, consisting of levodopa, dopamine agonists, or a combination of the two, and assessed during medication ON state. The LEDD was calculated at each assessment using the method from (Tomlinson et al., 2010).

A psychiatric evaluation was done before the surgery to exclude patients with surgery-contradicting symptoms (pre-psychotic or florid psychotic symptoms or mood disorders including suicidal thoughts or any other risky neuropsychiatric complications) (Foley et al., 2018).

### ***Statistical Analysis***

First, descriptive statistics (mean and standard deviation) were calculated at each time point.

Based on previously published data, we calculated the two subscores - cognitive-driven (FAQc) and motor-drive (FAQm) (Becker et al., 2018b).

**Results:****Discussion:****Conclusion:****Acknowledgements:**

Repovska Hana

**References:**

- Accolla, E. A., & Pollo, C. (2019). Mood effects after deep brain stimulation for parkinson's disease: An update. *Frontiers in Neurology*, 10. <https://doi.org/10.3389/fneur.2019.00617>
- Alegret, M., Junqué, C., Valldeoriola, F., Vendrell, P., Pilleri, M., Rumià, J., & Tolosa, E. (2001). Effects of Bilateral Subthalamic Stimulation on Cognitive Function in Parkinson Disease. *Archives of Neurology*, 58(8), 1223. <https://doi.org/10.1001/archneur.58.8.1223>
- Appleby, B. S., Duggan, P. S., Regenberg, A., & Rabins, P. V. (2007). Psychiatric and neuropsychiatric adverse events associated with deep brain stimulation: A meta-analysis of ten years' experience. *Movement Disorders*, 22(12), 1722–1728. <https://doi.org/10.1002/mds.21551>
- Aviles-Olmos, I., Kefalopoulou, Z., Tripoliti, E., Candelario, J., Akram, H., Martinez-Torres, I., Jahanshahi, M., Foltynie, T., Hariz, M., Zrinzo, L., & Limousin, P. (2014). Long-term outcome of subthalamic nucleus deep brain stimulation for Parkinson's disease using an MRI-guided and MRI-verified approach. *Journal of Neurology, Neurosurgery & Psychiatry*, 85(12), 1419–1425. <https://doi.org/10.1136/jnnp-2013-306907>
- Beck, A. T., Steer, R. A., & Brown, G. (1996). *Beck depression inventory–II*. American Psychological Association (APA). <https://doi.org/10.1037/t00742-000>
- Becker, S., Bäumer, A., Maetzler, W., Nussbaum, S., Timmers, M., Van Nueten, L., Salvatore, G., Zaunbrecher, D., Roeben, B., Brockmann, K., Streffer, J., Berg, D., & Liepelt-Scarfone, I. (2018a). Assessment of cognitive-driven activity of daily living impairment in non-demented Parkinson's patients. *Journal of Neuropsychology*, 14(1), 69–84. <https://doi.org/10.1111/jnp.12173>
- Becker, S., Bäumer, A., Maetzler, W., Nussbaum, S., Timmers, M., Van Nueten, L., Salvatore, G., Zaunbrecher, D., Roeben, B., Brockmann, K., Streffer, J., Berg, D., &

- Liepelt-Scarfone, I. (2018b). Assessment of cognitive-driven activity of daily living impairment in non-demented Parkinson's patients. *Journal of Neuropsychology*, 14(1), 69–84. <https://doi.org/10.1111/jnp.12173>
- Becker, S., Bode, M., Brockmann, K., Gasser, T., Michaelis, K., Solbrig, S., Nuerk, H.-C., Schulte, C., Maetzler, W., Zimmermann, M., Berg, D., & Liepelt-Scarfone, I. (2022). Cognitive-Driven Activities of Daily Living Impairment as a Predictor for Dementia in Parkinson Disease. *Neurology*, 99(23). <https://doi.org/10.1212/wnl.0000000000201201>
- Becker, S., Pauly, C., Lawton, M., Hipp, G., Bowring, F., Sulzer, P., Hu, M., Krüger, R., Gasser, T., & Liepelt-Scarfone, I. (2021). Quantifying activities of daily living impairment in Parkinson's disease using the Functional Activities Questionnaire. *Neurological Sciences*, 43(2), 1047–1054. <https://doi.org/10.1007/s10072-021-05365-1>
- Bezdicek, O., Mana, J., Růžicka, F., Havlik, F., Fečíková, A., Uhrová, T., Růžicka, E., Urgošík, D., & Jech, R. (2022). The instrumental activities of daily living in parkinson's disease patients treated by subthalamic deep brain stimulation. *Frontiers in Aging Neuroscience*, 14. <https://doi.org/10.3389/fnagi.2022.886491>
- Bezdicek, O., Michalec, J., Nikolai, T., Havránková, P., Roth, J., Jech, R., & Růžicka, E. (2015). Clinical Validity of the Mattis Dementia Rating Scale in Differentiating Mild Cognitive Impairment in Parkinson's Disease and Normative Data. *Dementia and Geriatric Cognitive Disorders*, 39(5-6), 303–311. <https://doi.org/10.1159/000375365>
- Bezdicek, O., Nikolai, T., Michalec, J., Růžicka, F., Havránková, P., Roth, J., Jech, R., & Růžicka, E. (2016). The Diagnostic Accuracy of Parkinson's Disease Mild Cognitive Impairment Battery Using the Movement Disorder Society Task Force Criteria. *Movement Disorders Clinical Practice*, 4(2), 237–244. <https://doi.org/10.1002/mdc3.12391>
- Bezdicek, O., Sulc, Z., Nikolai, T., Stepankova, H., Kopecek, M., Jech, R., & Růžicka, E. (2017). A parsimonious scoring and normative calculator for the Parkinson's disease mild cognitive impairment battery. *The Clinical Neuropsychologist*, 31(6-7), 1231–1247. <https://doi.org/10.1080/13854046.2017.1293161>
- Bove, F., Fraix, V., Cavallieri, F., Schmitt, E., Lhommée, E., Bichon, A., Meoni, S., Péliissier, P., Kistner, A., Chevrier, E., Ardouin, C., Limousin, P., Krack, P., Benabid, A. L.,



- Chabardès, S., Seigneuret, E., Castrioto, A., & Moro, E. (2020). Dementia and subthalamic deep brain stimulation in Parkinson disease. *Neurology*, 95(4).  
<https://doi.org/10.1212/wnl.00000000000009822>
- Castrioto, A. (2011). Ten-Year Outcome of Subthalamic Stimulation in Parkinson Disease. *Archives of Neurology*, 68(12), 1550. <https://doi.org/10.1001/archneurol.2011.182>
- Chakraborty, A., Brauer, S., & Diwan, A. (2020). Possible therapies of Parkinson's disease: A review. *Journal of Clinical Neuroscience*, 75, 1–4.  
<https://doi.org/10.1016/j.jocn.2020.03.024>
- Ciharova, M., Cígler, H., Dostálová, V., Šivicová, G., & Bezdicek, O. (2020). Beck depression inventory, second edition, Czech version: demographic correlates, factor structure and comparison with foreign data. *International Journal of Psychiatry in Clinical Practice*, 24(4), 371–379. <https://doi.org/10.1080/13651501.2020.1775854>
- Deuschl, G., Schade-Brittinger, C., Krack, P., Volkmann, J., Schäfer, H., Bötzel, K., Daniels, C., Deutschländer, A., Dillmann, U., Eisner, W., Gruber, D., Hamel, W., Herzog, J., Hilker, R., Klebe, S., Klob, M., Koy, J., Krause, M., Kupsch, A., ... Voges, J. (2006). A Randomized Trial of Deep-Brain Stimulation for Parkinson's Disease. *New England Journal of Medicine*, 355(9), 896–908. <https://doi.org/10.1056/nejmoa060281>
- Drapier, D., Drapier, S., Sauleau, P., Haegelen, C., Raoul, S., Biseul, I., Peron, J., Lallement, F., Rivier, I., Reymann, J. M., Edan, G., Verin, M., & Millet, B. (2006). Does subthalamic nucleus stimulation induce apathy in Parkinson's disease? *Journal of Neurology*, 253(8), 1083–1091. <https://doi.org/10.1007/s00415-006-0177-0>
- Foley, J. A., Foltynie, T., Limousin, P., & Cipolotti, L. (2018). Standardised Neuropsychological Assessment for the Selection of Patients Undergoing DBS for Parkinson's Disease. *Parkinson's Disease*, 2018, 1–13.  
<https://doi.org/10.1155/2018/4328371>
- Gervais-Bernard, H., Xie-Brustolin, J., Mertens, P., Polo, G., Klinger, H., Adamec, D., Broussolle, E., & Thobois, S. (2009). Bilateral subthalamic nucleus stimulation in advanced Parkinson's disease: Five year follow-up. *Journal of Neurology*, 256(2), 225–233. <https://doi.org/10.1007/s00415-009-0076-2>

- Gold, D. A. (2012). An examination of instrumental activities of daily living assessment in older adults and mild cognitive impairment. *Journal of Clinical and Experimental Neuropsychology*, 34(1), 11–34. <https://doi.org/10.1080/13803395.2011.614598>
- Gorecka-Mazur, A., Furgala, A., Krygowska-Wajs, A., Pietraszko, W., Kwinta, B., & Gil, K. (2019). Activities of Daily Living and Their Relationship to Health-Related Quality of Life in Patients with Parkinson Disease After Subthalamic Nucleus Deep Brain Stimulation. *World Neurosurgery*, 125, e552–e562. <https://doi.org/10.1016/j.wneu.2019.01.132>
- Hariz, G.-M., & Forsgren, L. (2010). Activities of daily living and quality of life in persons with newly diagnosed Parkinson's disease according to subtype of disease, and in comparison to healthy controls. *Acta Neurologica Scandinavica*, 123(1), 20–27. <https://doi.org/10.1111/j.1600-0404.2010.01344.x>
- Hentz, J. G., Mehta, S. H., Shill, H. A., Driver-Dunckley, E., Beach, T. G., & Adler, C. H. (2015). Simplified conversion method for unified Parkinson's disease rating scale motor examinations. *Movement Disorders*, 30(14), 1967–1970. <https://doi.org/10.1002/mds.26435>
- Jech, R., Mueller, K., Urgošik, D., Sieger, T., Holiga, Š., Růžicka, F., Dušek, P., Havránková, P., Vymazal, J., & Růžicka, E. (2012). The Subthalamic Microlesion Story in Parkinson's Disease: Electrode Insertion-Related Motor Improvement with Relative Cortico-Subcortical Hypoactivation in fMRI. *PLoS ONE*, 7(11), e49056. <https://doi.org/10.1371/journal.pone.0049056>
- JECH, R., RUZICKA, E., URGOSIK, D., SERRANOVA, T., VOLFOVA, M., NOVAKOVA, O., ROTH, J., DUSEK, P., & MECIR, P. (2006). Deep brain stimulation of the subthalamic nucleus affects resting EEG and visual evoked potentials in Parkinson's disease. *Clinical Neurophysiology*, 117(5), 1017–1028. <https://doi.org/10.1016/j.clinph.2006.01.009>
- Jurica, P. J., Leitten, C. L., & Mattis, S. (2001). *Dementia rating scale-2 (DRS-2) professional manual*. Psychological Assessment Resources.
- Kubu, C. S. (2018). The Role of a Neuropsychologist on a Movement Disorders Deep Brain Stimulation Team. *Archives of Clinical Neuropsychology*, 33(3), 365–374. <https://doi.org/10.1093/arclin/acx130>

- Lawton, M. P., & Brody, E. M. (1969). Assessment of Older People: Self-Maintaining and Instrumental Activities of Daily Living. *The Gerontologist*, 9(3 Part 1), 179–186.  
[https://doi.org/10.1093/geront/9.3\\_part\\_1.179](https://doi.org/10.1093/geront/9.3_part_1.179)
- Lezcano, E., Gómez-Esteban, J. C., Tijero, B., Bilbao, G., Lambarri, I., Rodriguez, O., Villoria, R., Dolado, A., Berganzo, K., Molano, A., Gopegui, E. R. de, Pomposo, I., Gabilondo, I., & Zarranz, J. J. (2016). Long-term impact on quality of life of subthalamic nucleus stimulation in Parkinson's disease. *Journal of Neurology*, 263(5), 895–905.  
<https://doi.org/10.1007/s00415-016-8077-4>
- Limousin, P., & Foltynie, T. (2019). Long-term outcomes of deep brain stimulation in Parkinson disease. *Nature Reviews Neurology*, 15(4), 234–242.  
<https://doi.org/10.1038/s41582-019-0145-9>
- Litvan, I., Goldman, J. G., Tröster, A. I., Schmand, B. A., Weintraub, D., Petersen, R. C., Mollenhauer, B., Adler, C. H., Marder, K., Williams-Gray, C. H., Aarsland, D., Kulisevsky, J., Rodriguez-Oroz, M. C., Burn, D. J., Barker, R. A., & Emre, M. (2012). Diagnostic criteria for mild cognitive impairment in Parkinson's disease: *Movement Disorder Society Task Force guidelines. Movement Disorders*, 27(3), 349–356.  
<https://doi.org/10.1002/mds.24893>
- Martin, R. C., Triebel, K. L., Kennedy, R. E., Nicholas, A. P., Watts, R. L., Stover, N. P., Brandon, M., & Marson, D. C. (2013). Impaired financial abilities in Parkinson's disease patients with mild cognitive impairment and dementia. *Parkinsonism & Related Disorders*, 19(11), 986–990. <https://doi.org/10.1016/j.parkreldis.2013.06.017>
- Mehanna, R., Bajwa, J. A., Fernandez, H., & Wagle Shukla, A. A. (2017). Cognitive Impact of Deep Brain Stimulation on Parkinson's Disease Patients. *Parkinson's Disease*, 2017, 1–15. <https://doi.org/10.1155/2017/3085140>
- Parsons, T. D., Rogers, S. A., Braaten, A. J., Woods, S. P., & Tröster, A. I. (2006). Cognitive sequelae of subthalamic nucleus deep brain stimulation in Parkinson's disease: a meta-analysis. *The Lancet Neurology*, 5(7), 578–588.  
[https://doi.org/10.1016/s1474-4422\(06\)70475-6](https://doi.org/10.1016/s1474-4422(06)70475-6)
- Pfeffer, R. I., Kurosaki, T. T., Harrah, C. H., Chance, J. M., & Filos, S. (1982). Measurement

- of Functional Activities in Older Adults in the Community. *Journal of Gerontology*, 37(3), 323–329. <https://doi.org/10.1093/geronj/37.3.323>
- Piboolnurak, P., Lang, A. E., Lozano, A. M., Miyasaki, J. M., Saint-Cyr, J. A., Poon, Y.-Y. W., Hutchison, W. D., Dostrovsky, J. O., & Moro, E. (2007). Levodopa response in long-term bilateral subthalamic stimulation for Parkinson's disease. *Movement Disorders*, 22(7), 990–997. <https://doi.org/10.1002/mds.21482>
- Pirogovsky, E., Martinez-Hannon, M., Schiehser, D. M., Lessig, S. L., Song, D. D., Litvan, I., & Filoteo, J. V. (2013). Predictors of performance-based measures of instrumental activities of daily living in nondemented patients with Parkinson's disease. *Journal of Clinical and Experimental Neuropsychology*, 35(9), 926–933. <https://doi.org/10.1080/13803395.2013.838940>
- Pirogovsky, E., Schiehser, D. M., Obtera, K. M., Burke, M. M., Lessig, S. L., Song, D. D., Litvan, I., & Filoteo, J. V. (2014). Instrumental activities of daily living are impaired in Parkinson's disease patients with mild cognitive impairment. *Neuropsychology*, 28(2), 229–237. <https://doi.org/10.1037/neu0000045>
- Rodriguez-Oroz, M. C., Obeso, J. A., Lang, A. E., Houeto, J.-L., Pollak, P., Rehncrona, S., Kulisevsky, J., Albanese, A., Volkmann, J., Hariz, M. I., Quinn, N. P., Speelman, J. D., Guridi, J., Zamarbide, I., Gironell, A., Molet, J., Pascual-Sedano, B., Pidoux, B., Bonnet, A. M., ... Van Blercom, N. (2005). Bilateral deep brain stimulation in Parkinson's disease: a multicentre study with 4 years follow-up. *Brain*, 128(10), 2240–2249. <https://doi.org/10.1093/brain/awh571>
- Saint-Cyr, J. A. (2000). Neuropsychological consequences of chronic bilateral stimulation of the subthalamic nucleus in parkinson's disease. *Brain*, 123(10), 2091–2108. <https://doi.org/10.1093/brain/123.10.2091>
- Schmitter-Edgecombe, M., McAlister, C., & Greeley, D. (2021). A Comparison of Functional Abilities in Individuals with Mild Cognitive Impairment and Parkinson's Disease with Mild Cognitive Impairment Using Multiple Assessment Methods. *Journal of the International Neuropsychological Society*, 28(8), 798–809. <https://doi.org/10.1017/s1355617721001077>

- Schupbach, W. M. M. (2005). Stimulation of the subthalamic nucleus in Parkinson's disease: a 5 year follow up. *Journal of Neurology, Neurosurgery & Psychiatry*, 76(12), 1640–1644. <https://doi.org/10.1136/jnnp.2005.063206>
- Shulman, L. M., Gruber-Baldini, A. L., Anderson, K. E., Vaughan, C. G., Reich, S. G., Fishman, P. S., & Weiner, W. J. (2008). The evolution of disability in Parkinson disease. *Movement Disorders*, 23(6), 790–796. <https://doi.org/10.1002/mds.21879>
- Tomlinson, C. L., Stowe, R., Patel, S., Rick, C., Gray, R., & Clarke, C. E. (2010). Systematic review of levodopa dose equivalency reporting in Parkinson's disease. *Movement Disorders*, 25(15), 2649–2653. <https://doi.org/10.1002/mds.23429>
- Zibetti, M., Merola, A., Rizzi, L., Ricchi, V., Angrisano, S., Azzaro, C., Artusi, C. A., Arduino, N., Marchisio, A., Lanotte, M., Rizzone, M., & Lopiano, L. (2011). Beyond nine years of continuous subthalamic nucleus deep brain stimulation in Parkinson's disease. *Movement Disorders*, 26(13), 2327–2334. <https://doi.org/10.1002/mds.23903>