



Contents lists available at ScienceDirect

The Journal of Arthroplasty

journal homepage: www.arthroplastyjournal.org

World Expert Meeting in Arthroplasty 2024

In Which Patients Should Cemented Femoral Components Be Used During Primary Total Hip Arthroplasty?



Zhaorui Wang, MD ^a, Abdelhak Adjel, MD ^b, Federico Burgo, MD, MBA ^c,
Muhammad Amin Chinoy, MD ^d, Dirk J.F. Moojen, MD, PhD ^e, Kenichi Oe, MD, PhD ^f,
Michael Reed, MD ^g, Marco Teloken, MD ^h, Michael M. Kheir, MD ^{a,*}

^a Department of Orthopaedic Surgery, University of Michigan, Ann Arbor, Michigan^b Department of Orthopedics, Hayet Clinic, Tiaret, Algeria^c Department of Orthopedics and Trauma, Hospital Universitario Austral, Buenos Aires, Argentina^d Department of Trauma & Orthopedics, Indus Hospital & Health Network, Karachi, Sindh, Pakistan^e Department of Orthopaedic and Trauma Surgery, OLVG Hospital, Amsterdam, The Netherlands^f Department of Orthopaedic Surgery, Kansai Medical University, Hirakata, Osaka, Japan^g Department of Trauma & Orthopaedics, Northumbria Healthcare NHS Foundation Trust and University of York, North Tyneside, Tyne and Wear, United Kingdom^h Orthopedic Department, Clinica Teloken, Porto Alegre, Rio Grande do Sul, Brazil

ARTICLE INFO

Article history:

Received 19 September 2024

Received in revised form

7 October 2024

Accepted 8 October 2024

Available online 16 October 2024

Keywords:

hip
arthroplasty
cemented
cementless
femoral
primary

In which patients should cemented femoral components be used during primary total hip arthroplasty?

Response/Recommendation: The literature supports the use of a cemented femoral component in women older than 70 years of age, in patients who have femoral neck fractures, in patients who have a Dorr type C femur, and in patients who have severe osteoporosis.

Level of Evidence: Moderate.

Expert Vote: 84.9% Agree, 11.4% Disagree, 3.7% Abstain.

Rationale

A large number of studies from multiple countries have been published on the topic of cemented femoral fixation during total hip arthroplasty, including large database and registry studies, retrospective reviews, meta-analyses, and some randomized controlled trials. Given the multitude of factors influencing the

decision-making for using cemented femoral components, each factor has been separately considered when reviewing the literature.

The most studied factors in the literature are age and sex, which consequently have the clearest evidence for cemented femoral fixation. Studies have found that cemented femoral fixation results in significantly decreased intraoperative periprosthetic fractures, postoperative periprosthetic fractures, and revisions in elderly women [1–9]. Studies consistently demonstrate that women appear to be an independent risk factor for periprosthetic fracture [3,10,11]. Compared to women, the risk reduction effects of cemented femoral components on periprosthetic fracture rates and revision rates appear to be less significant for men [1,2,6]. However, there are studies that demonstrate a reduction in periprosthetic fractures and revisions in all elderly patients regardless of sex

One or more of the authors of this paper have disclosed potential or pertinent conflicts of interest, which may include receipt of payment, either direct or indirect, institutional support, or association with an entity in the biomedical field which may be perceived to have potential conflict of interest with this work. For full disclosure statements refer to <https://doi.org/10.1016/j.arth.2024.10.034>.

* Address correspondence to: Michael M. Kheir, MD, University of Michigan, 1500 E. Medical Center Drive, Ann Arbor, Michigan 48109.

<https://doi.org/10.1016/j.arth.2024.10.034>

0883-5403/© 2024 Elsevier Inc. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

[1,12–20]. For patients aged less than 50 to 55 years, cementation does not appear to provide protection against periprosthetic fractures or revisions and might increase the risk of aseptic loosening and revisions [14,21,22]. It should also be noted that the definition of “elderly” varies significantly between studies, with common cutoffs between 65 and 75 years. Given the lack of consensus for age cutoff, considerations should be given for cementing the femoral stem in patients aged more than 75 years and perhaps aged more than 65 years in the presence of poor bone stock [9]. However, the benefits of reduced periprosthetic fractures need to be weighed against the potentially increased risk of aseptic loosening, pulmonary embolism, and infection [2,13,18,23,24].

In the challenging setting of hip arthroplasty in young patients who have hip dysplasia, some studies have demonstrated that cementless stems exhibited a higher ratio of intraoperative fracture and thinning of cortical bone, including stress shielding, medullary changes, stem alignment changes, and osteolysis, compared to cemented stems; furthermore, there appears to be no significant difference in survival at mean follow-up of 4 to 10 years [25–29]. This is not to say that cementless stems should not be used in these patients, but in the setting of challenging torsional deformities, cemented stems may be a reasonable alternative to diaphyseal-engaging stems to help correct these difficult deformities.

The literature overwhelmingly supports the use of cemented femoral components in elderly patients undergoing total hip arthroplasty for the treatment of a low-energy femoral neck fracture. Studies have shown that cementation of the femoral component in this setting significantly decreases the rates of periprosthetic fractures, complications, and revision rates and significantly increases patient-reported outcome measurements such as the Harris Hip Scores [30–37]. Notably, there were two randomized controlled trials evaluating cement usage in this population; early discontinuation of both trials was due to preliminary results demonstrating significantly increased complication rates (periprosthetic fractures, revisions, and dislocations) in the uncemented group [33,36].

Regarding patients who have radiographic osteoporosis or Dorr type C femoral anatomy, studies have shown an increased periprosthetic fracture rate [38,39] and cemented femoral fixation in these patients is correlated with better patient-reported outcomes and lower revision rates [10,40,41]. However, there is some disagreement in the literature, with one large database study showing similar rates of periprosthetic fractures and revisions in osteoporotic patients who have cemented versus cementless femoral components [42]. Overall, high-quality studies regarding cement usage for Dorr C/osteoporotic patients are lacking, which is likely in part due to patient selection bias given the general consensus of using cemented femoral components in these patients.

CRedit authorship contribution statement

Zhaorui Wang: Writing – original draft, Methodology, Investigation, Data curation. **Abdelhak Adjel:** Writing – review & editing. **Federico Burgo:** Writing – review & editing. **Muhammad Amin Chinoy:** Writing – review & editing. **Dirk J.F. Moojen:** Writing – review & editing. **Kenichi Oe:** Writing – review & editing. **Michael Reed:** Writing – review & editing. **Marco Teloken:** Writing – review & editing. **Michael M. Kheir:** Writing – review & editing. Writing – original draft, Supervision, Methodology, Investigation.

References

- [1] Tanzer M, Graves SE, Peng A, Shimmin AJ. Is cemented or cementless femoral stem fixation more durable in patients older than 75 years of age? A comparison of the best-performing stems. *Clin Orthop* 2018;476:1428–37. <https://doi.org/10.1097/01.blo.0000533621.57561.a4>.
- [2] Kelly MP, Chan PH, Prentice HA, Paxton EW, Hinman AD, Khatod M. Cause-specific stem revision risk in primary total hip arthroplasty using cemented vs cementless femoral stem fixation in a US cohort. *J Arthroplasty* 2022;37:89–96.e1. <https://doi.org/10.1016/j.arth.2021.09.020>.
- [3] Abdel MP, Watts CD, Houdek MT, Lewallen DG, Berry DJ. Epidemiology of periprosthetic fracture of the femur in 32 644 primary total hip arthroplasties: a 40-year experience. *Bone Jt J* 2016;98-B:461–7. <https://doi.org/10.1302/0301-620X.98B4.37201>.
- [4] Dale H, Børshiem S, Kristensen TB, Fenstad AM, Gjertsen JE, Hallan G, et al. Fixation, sex, and age: highest risk of revision for uncemented stems in elderly women - data from 66,995 primary total hip arthroplasties in the Norwegian Arthroplasty Register. *Acta Orthop* 2020;91:33–41. <https://doi.org/10.1080/17453674.2019.1682851>.
- [5] Omari A, Nielsen CS, Husted H, Otte KS, Troelsen A, Gromov K. Introduction of a new treatment algorithm reduces the number of periprosthetic femoral fractures after primary total hip arthroplasty in elderly females. *J Arthroplasty* 2020;35:3613–20. <https://doi.org/10.1016/j.arth.2020.06.077>.
- [6] Thien TM, Chatziagorou G, Garellick G, Furnes O, Havelin LI, Mäkelä K, et al. Periprosthetic femoral fracture within two years after total hip replacement: analysis of 437,629 operations in the nordic arthroplasty register association database. *J Bone Jt Surg Am* 2014;96:e167. <https://doi.org/10.2106/JBJS.M.00643>.
- [7] Edelstein AI, Hume EL, Pezzin LE, McGinley EL, Dillingham TR. The impact of femoral component cementation on fracture and mortality risk in elective total hip arthroplasty: analysis from a national medicare sample. *J Bone Jt Surg Am* 2022;104:523–9. <https://doi.org/10.2106/JBJS.21.00640>.
- [8] Gjertsen J-E, Nilsen D, Furnes O, Hallan G, Kroken G, Dyvik E, et al. Promoting cemented fixation of the femoral stem in elderly female hip arthroplasty patients and elderly hip fracture patients: a retrospective cohort study from the Norwegian Arthroplasty Register and the Norwegian Hip Fracture Register. *Acta Orthop* 2024;95:130–7. <https://doi.org/10.2340/17453674.2024.40073>.
- [9] Ladurner A, Zdravkovic V, Giesinger K. Sex-specific implant fixation can reduce revision rates in total hip arthroplasty: evidence from the Swiss national joint registry. *J Arthroplasty* 2024;39:421–6. <https://doi.org/10.1016/j.arth.2023.08.035>.
- [10] Lindberg-Larsen M, Jørgensen CC, Solgaard S, Kjersgaard AG, Kehlet H. Increased risk of intraoperative and early postoperative periprosthetic femoral fracture with uncemented stems. *Acta Orthop* 2017;88:390–4. <https://doi.org/10.1080/17453674.2017.1302908>.
- [11] Hopman SR, de Windt TS, van Erp JHJ, Bekkers JEJ, de Gast A. Uncemented total hip arthroplasty; increased risk of early periprosthetic fracture requiring revision surgery in elderly females. *J Orthop* 2021;25:40–4. <https://doi.org/10.1016/j.jor.2021.03.025>.
- [12] Kelly M, Chen AF, Ryan SP, Working ZM, Porter KR, De A, et al. Cemented femoral fixation in total hip arthroplasty reduces the risk of periprosthetic femur fracture in patients 65 Years and older: an analysis from the American joint replacement registry. *J Arthroplasty* 2023;38:S351–4. <https://doi.org/10.1016/j.arth.2023.04.039>.
- [13] Hameed D, McCormick BP, Sequeira SB, Dubin JA, Bains SS, Mont MA, et al. Cemented versus cementless femoral fixation for total hip arthroplasty following femoral neck fracture in patients aged 65 and older. *J Arthroplasty* 2024;39:1747–51. <https://doi.org/10.1016/j.arth.2024.01.034>.
- [14] Boyle AB, Zhu M, Frampton C, Poutawara V, Vane A. Comparing modern uncemented, hybrid and cemented implant combinations in older patients undergoing primary total hip arthroplasty, a New Zealand Joint Registry study. *Arch Orthop Trauma Surg* 2023;143:3597–604. <https://doi.org/10.1007/s00402-022-04610-2>.
- [15] Gray WK, Day J, Barker M, Briggs TWR. Fixation method and subsequent revision rates for elective primary hip arthroplasty in people aged 70 Years and older: analysis of national administrative data sets by the UK getting it right first time program. *J Arthroplasty* 2020;35:3631–7. <https://doi.org/10.1016/j.arth.2020.06.081>.
- [16] Jämsen E, Eskelinen A, Peltola M, Mäkelä K. High early failure rate after cementless hip replacement in the octogenarian. *Clin Orthop Relat Res* 2014;472:2779–89. <https://doi.org/10.1007/s11999-014-3641-7>.
- [17] Stea S, Comfort T, Sedrakyan A, Havelin L, Marinelli M, Barber T, et al. Multinational comprehensive evaluation of the fixation method used in hip replacement: interaction with age in context. *J Bone Jt Surg Am* 2014;96(Suppl 1):42–51. <https://doi.org/10.2106/JBJS.N.00463>.
- [18] Lindberg-Larsen M, Petersen PB, Jørgensen CC, Overgaard S, Kehlet H. Post-operative 30-day complications after cemented/hybrid versus cementless total hip arthroplasty in osteoarthritis patients > 70 years. *Acta Orthop* 2020;91:286–92. <https://doi.org/10.1080/17453674.2020.1745420>.
- [19] Gonzalez Della Valle A, Odum SM, De A, Barrington JW, Huddleston JI, Ilgen RL, et al. The effect of femoral fixation on revision and mortality following elective total hip arthroplasty in patients over the age of 65 years.

- An analysis of the American joint replacement registry. *J Arthroplasty* 2022;37:1105–10. <https://doi.org/10.1016/j.arth.2022.01.088>.
- [20] Bunyoz KI, Malchau E, Malchau H, Troelsen A. Has the use of fixation techniques in THA changed in this decade? The uncemented paradox revisited. *Clin Orthop* 2020;478:697–704. <https://doi.org/10.1097/CORR.0000000000001117>.
- [21] Pedersen AB, Mehnert F, Havelin LI, Furnes O, Herberts P, Kärrholm J, et al. Association between fixation technique and revision risk in total hip arthroplasty patients younger than 55 years of age. Results from the Nordic Arthroplasty Register Association. *Osteoarthritis Cartilage* 2014;22:659–67. <https://doi.org/10.1016/j.joca.2014.03.005>.
- [22] Kim YH, Park JW, Kim JS, Kim IW. Twenty-five- to twenty-seven-year results of a cemented vs a cementless stem in the same patients younger than 50 Years of age. *J Arthroplasty* 2016;31:662–7. <https://doi.org/10.1016/j.arth.2015.09.045>.
- [23] Moore MC, Dubin JA, Monárrez R, Bains SS, Hameed D, Nace J, et al. Cemented versus cementless femoral fixation for total hip arthroplasty following osteoarthritis. *J Arthroplasty* 2023;39:1545–9. <https://doi.org/10.1016/j.arth.2023.12.024>.
- [24] Yoon BH, Ha YC, Lee YK, Koo KH. Postoperative deep infection after cemented versus cementless total hip arthroplasty: a meta-analysis. *J Arthroplasty* 2015;30:1823–7. <https://doi.org/10.1016/j.arth.2015.04.041>.
- [25] Inoue D, Kabata T, Kajino Y, Ohmori T, Ueoka K, Tsuchiya H. Comparison of mid-term clinical results between cementless and cemented femoral stems in total hip arthroplasty with femoral shortening osteotomy for Crowe type IV hips. *Arch Orthop Trauma Surg* 2021;141:1057–64. <https://doi.org/10.1007/s00402-020-03749-0>.
- [26] Miyazaki T, Shimizu T, Ohura H, Katayama N, Iwasaki N, Takahashi D. Total hip arthroplasty with femoral shortening osteotomy using polished cemented stem vs. modular cementless stem in patients with Crowe type IV developmental dysplasia of the hip. *Arch Orthop Trauma Surg* 2023;143:3487–93. <https://doi.org/10.1007/s00402-022-04518-x>.
- [27] Huang Z, Ling J, Zeng Z, Di Z, Zhang J, Tao K. Mid-Term outcomes of cemented stem and subtrochanteric shortening derotational osteotomy in total hip arthroplasty for crowe IV developmental dysplasia. *Orthop Surg* 2022;14:3178–86. <https://doi.org/10.1111/os.13515>.
- [28] Kawai T, Tanaka C, Kanoe H. Total hip arthroplasty for Crowe IV hip without subtrochanteric shortening osteotomy -a long term follow up study. *BMC Musculoskel Disord* 2014;15:72. <https://doi.org/10.1186/1471-2474-15-72>.
- [29] Oe K, Iida H, Nakamura T, Okamoto N, Wada T. Subtrochanteric shortening osteotomy combined with cemented total hip arthroplasty for Crowe group IV hips. *Arch Orthop Trauma Surg* 2013;133:1763–70. <https://doi.org/10.1007/s00402-013-1869-4>.
- [30] Mao S, Chen B, Zhu Y, Qian L, Lin J, Zhang X, et al. Cemented versus uncemented total hip replacement for femoral neck fractures in elderly patients: a retrospective, multicentre study with a mean 5-year follow-up. *J Orthop Surg* 2020;15:447. <https://doi.org/10.1186/s13018-020-01980-4>.
- [31] Heckmann ND, Chen XT, Ballatori AM, Ton A, Shahrestani S, Chung BC, et al. Cemented vs cementless femoral fixation for total hip arthroplasty after displaced femoral neck fracture: a nationwide analysis of short-term complications and readmission rates. *J Arthroplasty* 2021;36:3667–3675.e4. <https://doi.org/10.1016/j.arth.2021.06.029>.
- [32] Raja BS, Gowda AKS, Singh S, Ansari S, Kalia RB, Paul S. Comparison of functional outcomes and complications of cemented vs uncemented total hip arthroplasty in the elderly neck of femur fracture patients: a systematic review and meta-analysis. *J Clin Orthop Trauma* 2022;29:101876. <https://doi.org/10.1016/j.jcot.2022.101876>.
- [33] Clement ND, van der Linden M, Keating JF. Higher rate of complications with uncemented compared to cemented total hip arthroplasty for displaced intracapsular hip fractures: a randomised controlled trial of 50 patients. *Eur J Orthop Surg Traumatol* 2021;31:587–94. <https://doi.org/10.1007/s00590-020-02808-x>.
- [34] Liu T, Hua X, Yu W, Lin J, Zhao M, Liu J, et al. Long-term follow-up outcomes for patients undergoing primary total hip arthroplasty with uncemented versus cemented femoral components: a retrospective observational study with a 5-year minimum follow-up. *J Orthop Surg* 2019;14:371. <https://doi.org/10.1186/s13018-019-1415-3>.
- [35] Andersen MF, Jakobsen T, Bensen AS, Krarup N. Lower reoperation rate for cemented femoral stem than for uncemented femoral stem in primary total hip arthroplasty following a displaced femoral neck fracture. *SICOT-J* 2015;1:26. <https://doi.org/10.1051/sicotj/2015028>.
- [36] Chammout G, Muren O, Laurencikas E, Bodén H, Kelly-Pettersson P, Sjöo H, et al. More complications with uncemented than cemented femoral stems in total hip replacement for displaced femoral neck fractures in the elderly. *Acta Orthop* 2017;88:145–51. <https://doi.org/10.1080/17453674.2016.1262687>.
- [37] Zhou X, Chen M, Yu W, Han G, Ye J, Zhuang J. Uncemented versus cemented total hip arthroplasty for displaced femoral neck fractures in elderly patients with osteoporosis: a retrospective analysis. *J Int Med Res* 2020;48:300060520944663. <https://doi.org/10.1177/0300060520944663>.
- [38] Kheir MM, Dilley JE, Speybroeck J, Kuyl EV, Ochenjele G, McLawhorn AS, et al. The influence of Dorr type and femoral fixation on outcomes following total hip arthroplasty for acute femoral neck fractures: a multicenter study. *J Arthroplasty* 2023;38:719–25. <https://doi.org/10.1016/j.arth.2022.10.028>.
- [39] Gkagkalis G, Goetti P, Mai S, Meinecke I, Helmy N, Bosson D, et al. Cementless short-stem total hip arthroplasty in the elderly patient - is it a safe option?: a prospective multicentre observational study. *BMC Geriatr* 2019;19:112. <https://doi.org/10.1186/s12877-019-1123-1>.
- [40] Yang C, Han X, Wang J, Yuan Z, Wang T, Zhao M, et al. Cemented versus uncemented femoral component total hip arthroplasty in elderly patients with primary osteoporosis: retrospective analysis with 5-year follow-up. *J Int Med Res* 2019;47:1610–9. <https://doi.org/10.1177/0300060518825428>.
- [41] Fiedler B, Patel V, Lygrisse KA, Kelly ME, Turcotte JJ, MacDonald J, et al. The effect of reduced bone mineral density on elective total hip arthroplasty outcomes. *Arch Orthop Trauma Surg* 2023;143:5993–9. <https://doi.org/10.1007/s00402-023-04830-0>.
- [42] Kuyl EV, Agarwal AR, Patel PK, Harris AB, Gu A, Rao S, et al. Osteoporotic patients undergoing total hip arthroplasty have a similar 5-year cumulative incidence rate of periprosthetic fracture regardless of cemented versus cementless femoral stem fixation. *J Arthroplasty* 2023;39:1285–1290.e1. <https://doi.org/10.1016/j.arth.2023.11.001>.