



Contents lists available at ScienceDirect

## The Journal of Arthroplasty

journal homepage: [www.arthroplastyjournal.org](http://www.arthroplastyjournal.org)

## World Expert Meeting in Arthroplasty 2024

## Is There a Limit to Lengthening in Patients Who Have Crowe IV Developmental Dysplasia of the Hip Undergoing Total Hip Arthroplasty?



Ibrahim El-Ganzoury, MD<sup>a</sup>, Mohammad T. Ghazavi, MD, FRCSC<sup>b</sup>,  
 Vahit Emre Özden, MD<sup>c,d,\*</sup>, Jesus Moreta, MD<sup>e</sup>, Oussama Chaar, MD<sup>f</sup>,  
 Vorawit Atipiboosin, MD<sup>g</sup>, Ömer F. Bilgen, MD<sup>h</sup>, Daisuke Inoue, MD, PhD<sup>i</sup>,  
 Peilai Liu, MD<sup>j</sup>, Yanguo Qin, MD<sup>k</sup>, Ahmed S. Younis, MD<sup>l</sup>

<sup>a</sup> Department of Orthopedic Surgery, Ain Shams University, Cairo, Egypt<sup>b</sup> Department of Orthopedic Surgery, Scarborough Health Network, Toronto, Ontario, Canada<sup>c</sup> Department of Orthopaedics and Traumatology, Acibadem Mehmet Ali Aydınlar University, İstanbul, Turkey<sup>d</sup> International Joint Center (IJC) Acibadem Maslak Hospital, İstanbul, Turkey<sup>e</sup> Department of Orthopedic Surgery, Hospital San Juan De Dios, Santurtzi, Bilbao, Spain<sup>f</sup> Burjeel Orthopaedic Institute, Abu Dhabi's Orthopaedic Institute, Abu Dhabi, United Arab Emirates<sup>g</sup> Faculty of Medicine, Department of Orthopedics, Khonkaen University, Khonkaen, Thailand<sup>h</sup> Department of Orthopaedics and Traumatology, Private Medicabil Hospital, Bursa, Turkey<sup>i</sup> Department of Orthopaedic Surgery, Graduate School of Medical Sciences, Kanazawa University, Kanazawa, Japan<sup>j</sup> Department of Orthopaedics Surgery, Qilu Hospital of Shandong University, Jinan, Shandong, China<sup>k</sup> Orthopaedic Medical Center, The Second Hospital of Jilin University, Changchun, Jilin, China<sup>l</sup> Faculty of Medicine, Ain Shams University, Cairo, Egypt

## ARTICLE INFO

## Article history:

Received 20 September 2024

Received in revised form

10 October 2024

Accepted 14 October 2024

Available online 22 October 2024

## Keywords:

type IV DDH

femoral shortening

high hip dislocation

hip

hip arthroplasty

## Is there a limit to lengthening in patients who have Crowe IV developmental dysplasia of the hip undergoing total hip arthroplasty?

**Response/Recommendation:** A review of the current literature does not identify a specific limit to acute limb lengthening in patients undergoing total hip arthroplasty. The traditional limits, such as 4 cm, are based on older studies with some methodological deficiencies. With careful intraoperative monitoring, restoration of length may be considered in selected cases.

**Level of Evidence:** Limited.

**Delegate Vote:** Agree: 83%, Disagree: 11%, and Abstain: 5% (Superior Majority, Strong Consensus).

## Rationale

A search was conducted for all studies investigating the outcomes of total hip arthroplasty (THA) for hip arthritis secondary to Crowe III to IV developmental dysplasia of the hip, aiming to define a safe acute lengthening limit. The search encompassed the following electronic databases: PubMed: 865 results on May 28, 2024; Central: 33 results on April 26, 2024; and Embase: 835 results on May 14, 2024. After the removal of duplicates and exclusion of noneligible studies following the screening of the title and abstract, 419 studies were eligible for full-text evaluation. Finally, 92 studies [1–92] were included in this review.

All included studies were cohort studies, whether prospective or retrospective. In all included cohorts, patients were treated with

Funding: There is no funding source.

One or more of the authors of this paper have disclosed potential or pertinent conflicts of interest, which may include receipt of payment, either direct or indirect, institutional support, or association with an entity in the biomedical field which may be perceived to have potential conflict of interest with this work. For full disclosure statements refer to <https://doi.org/10.1016/j.arth.2024.10.079>.

\* Address correspondence to: Vahit Emre Özden, MD, Department of Orthopaedics and Traumatology, Acibadem Mehmet Ali Aydınlar University, International Joint Centre (IJC), Acibadem Maslak Hospital, Büyükdere Cad No 40, Sarıyer, İstanbul 34457, Turkey.

<https://doi.org/10.1016/j.arth.2024.10.079>

0883-5403/© 2024 Elsevier Inc. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

THA and reportedly achieved good outcomes in terms of satisfaction, range of motion, restoration of limb length, and function. Debate regarding the extent of acute lengthening that is safe to undertake in patients who have dysplasia undergoing THA continues because of the established and well-recognized risk of neurological complications, specifically sciatic nerve palsy. Based on conventional teaching, acute lengthening of the limb > 4 cm and the resulting tension on the sciatic nerve puts the extremity at an unacceptable risk of neurological complications [49]. This accepted limit to acute lengthening means that some patients who have unilateral dysplasia and a short limb (> 4 cm) cannot completely restore their full length, and instead have a residual limb-length discrepancy. The belief in the stated limit leads many surgeons to resort to performing a femoral shortening osteotomy if lengthening is likely to be > 4 cm.

Our review of the contemporary literature did not reveal a specific limit to lengthening during arthroplasty for this condition. In fact, there are reports in the literature suggesting that the limit for acute lengthening should be determined using femoral length as a reference [44]. In a study by Kabata et al., the risk for neurological complications appeared to increase when lengthening exceeded 8.7% of the femoral length [44]. There were two comparative studies by Fujishiro et al. that highlighted the influence of factors other than lengthening, such as hip flexion contracture, regarding neurological complications in patients undergoing THA [9,60]. Another study proposed 50 mm as a limit for acute lengthening [92]. Lengthening up to 51 mm was seen in some of the 58 studies that used femoral shortening osteotomy. There were 13 studies included in cohorts of patients who had Crowe IV dysplasia when femoral osteotomy was not performed during THA. The mean limb lengthening in these patients ranged from 21 to 48 mm. In nine studies, patients were subjected to staged procedures whereby lengthening occurred during a preliminary procedure. If neurological complications did not occur, definitive THA was performed during a second surgery. These studies reported a mean lengthening ranging from 36 to 100 mm, with the first-stage operations involving a variety of procedures. The initial lengthening might be achieved with a distraction nail, ilio-femoral fixator, or even initial hip periarticular extensive soft-tissue releases and traction.

Some techniques, distinct from femoral osteotomy and staged procedures, have been described with the aim of achieving reduction of the femoral head into the true acetabulum with minimal complications. Yan et al. described maintaining the hip and knee in a flexed position following surgery and then allowing for gradual knee extension until the patient notes the onset of numbness. They achieved an average lengthening of 40 mm (range, 29 to 62) using this technique [12]. Shemesh et al. reported using an intraoperative nerve stimulator to assess neurological insults during THA procedures [81]. They stated that intraoperative nerve monitoring mitigated the risk of nerve palsy and reduced the need for femoral shortening osteotomies. These findings were corroborated by Kong et al. in their comparative retrospective cohort, demonstrating a lower incidence of nerve palsy when intraoperative nerve monitoring was used during complex THA [73].

In conclusion, there is only limited and conflicting evidence about the potential limit of acute lengthening in patients undergoing THA, after which the risk for neurological complications increases. In the absence of any definitive limit, surgeons may decide to lengthen the extremity more than 4 cm to achieve equal limb length and avoid leaving these young patients who have a possibly major residual limb-length discrepancy. With careful intraoperative monitoring and maintaining the knee in a flexed position intraoperatively and postoperatively, restoration of length may be considered in selected cases.

## Availability of Data and Material (Data Transparency)

All data and material are available when required.

## Code Availability (Software Application or Custom Code)

All software application and custom codes are available when needed.

## Consent for Publication

Consent for publication was obtained.

## CRediT authorship contribution statement

**Ibrahim El-Ganzoury:** Writing – review & editing, Writing – original draft, Visualization, Methodology, Formal analysis, Data curation. **Mohammad T. Ghazavi:** Writing – review & editing, Writing – original draft. **Vahit Emre Özden:** Writing – review & editing, Supervision, Methodology, Data curation, Conceptualization. **Jesus Moreta:** Writing – review & editing, Methodology. **Oussama Chaar:** Writing – review & editing. **Vorawit Atipiboosin:** Writing – review & editing. **Ömer F. Bilgen:** Writing – review & editing. **Daisuke Inoue:** Writing – review & editing. **Peilai Liu:** Writing – review & editing. **Yanguo Qin:** Writing – review & editing. **Ahmed S. Younis:** Writing – review & editing, Writing – original draft, Methodology, Investigation, Data curation, Conceptualization.

## References

- [1] Oinuma K, Shiratsuchi H, Nakakita Y, Miura Y, Ninomiya T. A new anterior total hip arthroplasty reduction technique for patients who require leg lengthening. *Hip Int* 2020;30:587–91. <https://doi.org/10.1177/1120700020937248>.
- [2] Li H, Yuan Y, Xu J, Chang Y, Dai K, Zhu Z. Direct leverage for reducing the femoral head in total hip arthroplasty without femoral shortening osteotomy for Crowe type 3 to 4 dysplasia of the hip. *J Arthroplast* 2018;33:794–9. <https://doi.org/10.1016/j.arth.2017.09.011>.
- [3] Nishino T, Mishima H, Kawamura H, Yoshizawa T, Miyakawa S, Yamazaki M. Ten-year results of 55 dysplasia hips of hip offset and leg length reconstruction in total hip arthroplasty with cementless tapered stems having a high offset option designed for dysplastic femur. *J Orthop Surg* 2020;28:2309499020909499. <https://doi.org/10.1177/2309499020909499>.
- [4] Imarisio D, Trecci A, Sabatini L, Uslenghi M, Leone C, Scagnelli R. Cementless total hip replacement for severe developmental dysplasia of the hip: our experience in Crowe's group IV. *Musculoskelet Surg* 2013;97:25–30. <https://doi.org/10.1007/s12306-012-0227-y>.
- [5] Altay M, Demirkale I, Çatma MF, Karaduman M. Results of Crowe type IV developmental dysplasia of hip treated by subtrochantric osteotomy and total hip arthroplasty. *Indian J Orthop* 2018;52:374–9. [https://doi.org/10.4103/ortho.IJOrtho.445\\_16](https://doi.org/10.4103/ortho.IJOrtho.445_16).
- [6] Huang Y, Zhou Y, Shao H, Chu Y, Gu J, Li H. Total hip arthroplasties for har-tofilakidis type C1 and C2 high hip dislocations demonstrate similar survivorship and clinical function at minimum 10-year follow-up with cementless implants. *J Arthroplast* 2022;37:2374–80. <https://doi.org/10.1016/j.arth.2022.06.005>.
- [7] Holinka J, Pfeiffer M, Hofstaetter JG, Lass R, Kotz RI, Giurea A. Total hip replacement in congenital high hip dislocation following iliofemoral monoto-bute distraction. *Int Orthop* 2011;35:639–45. <https://doi.org/10.1007/s00264-010-1001-0>.
- [8] Ozden VE, Dikmen G, Beksac B, Tozun IR. Total hip arthroplasty with step-cut subtrochanteric femoral shortening osteotomy in high riding hip dislocated patients with previous femoral osteotomy. *J Orthop Sci* 2017;22:517–23. <https://doi.org/10.1016/j.jos.2017.01.017>.
- [9] Fujishiro T, Nishiyama T, Hayashi S, Hashimoto S, Kurosaka M, Kanno T, et al. Predicting leg-length change after total hip arthroplasty by measuring pre-operative hip flexion under general anaesthesia. *J Orthop Surg* 2012;20:327–30. <https://doi.org/10.1177/230949901202000312>.
- [10] Akman YE, Yavuz U, Çetinkaya E, Gür V, Gül M, Demir B. Cementless total hip arthroplasty for severely dislocated hips previously treated with Schanz osteotomy of the proximal femur. *Arch Orthop Trauma Surg* 2018;138:427–34. <https://doi.org/10.1007/s00402-018-2879-z>.

- [11] Chu YM, Zhou YX, Han N, Yang DJ. Two different total hip arthroplasties for hartofilakidis type C1 developmental dysplasia of hip in adults. *Chin Med J* 2016;129:289–94. <https://doi.org/10.4103/0366-6999.174507>.
- [12] Yan F, Chen G, Yang L, He R, Gu L, Wang F. A reduction technique of arthroplasty without subtrochanteric femoral shortening osteotomy for the treatment of developmental high dislocation of hip: a case series of 28 hips. *J Arthroplast* 2014;29:2289–93. <https://doi.org/10.1016/j.arth.2013.11.016>.
- [13] Liu RY, Bai CY, Song QC, Dang XQ, Wu YJ, Wang KZ. Partial greater trochanter osteotomy for hip reduction in total hip arthroplasty for high dislocated hip: a preliminary report. *BMC Musculoskelet Disord* 2014;15:293. <https://doi.org/10.1186/1471-2474-15-293>.
- [14] Li L, Yu M, Yang C, Gu G. Total hip arthroplasty (S-ROM stem) and subtrochanteric osteotomy for Crowe type IV developmental dysplasia of the hip. *Indian J Orthop* 2016;50:195–200. <https://doi.org/10.4103/0019-5413.177575>.
- [15] Hasegawa Y, Iwase T, Kanoh T, Seki T, Matsuoka A. Total hip arthroplasty for Crowe type IV developmental dysplasia. *J Arthroplast* 2012;27:1629–35. <https://doi.org/10.1016/j.arth.2012.02.026>.
- [16] Zhong C, Cai XZ, Yan SG, He RX. S-ROM modular arthroplasty combined with transverse subtrochanteric shortening for Crowe type IV congenital dislocation of hip. *Chin Med J* 2011;124:3891–5.
- [17] Binazzi R. Two-stage progressive femoral lowering followed by cementless total hip arthroplasty for treating Crowe IV-hartofilakidis type 3 developmental dysplasia of the hip. *J Arthroplast* 2015;30:790–6. <https://doi.org/10.1016/j.arth.2014.12.019>.
- [18] Masonis JL, Patel JV, Miu A, Bourne RB, McCalden R, Macdonald SJ, et al. Subtrochanteric shortening and derotational osteotomy in primary total hip arthroplasty for patients with severe hip dysplasia: 5-year follow-up. *J Arthroplast* 2003;18:68–73. <https://doi.org/10.1054/arth.2003.50104>.
- [19] Imbuldeniya AM, Walter WL, Zicat BA, Walter WK. Cementless total hip replacement without femoral osteotomy in patients with severe developmental dysplasia of the hip: minimum 15-year clinical and radiological results. *Bone Jt J* 2014;96-b:1449–54. <https://doi.org/10.1302/0301-620x.96b11.33698>.
- [20] Zhao E, Liu Z, Ding Z, Luo Z, Li H, Zhou Z. A propensity score-matched analysis between patients with high hip dislocation after childhood pyogenic infection and Crowe IV developmental dysplasia of the hip in total hip arthroplasty with subtrochanteric shortening osteotomy. *J Orthop Surg Res* 2020;15:418. <https://doi.org/10.1186/s13018-020-01947-5>.
- [21] Mu W, Yang D, Xu B, Mamtimin A, Guo W, Cao L. Midterm outcome of cementless total hip arthroplasty in Crowe IV-hartofilakidis type III developmental dysplasia of the hip. *J Arthroplast* 2016;31:668–75. <https://doi.org/10.1016/j.arth.2015.10.011>.
- [22] Tahta M, Isik C, Uluyardimci E, Cepni S, Oltulu I. Total hip arthroplasty without subtrochanteric femoral osteotomy is possible in patients with Crowe III/IV developmental dysplasia: total hip arthroplasty without femoral osteotomy. *Arch Orthop Trauma Surg* 2020;140:409–13. <https://doi.org/10.1007/s00402-019-03320-6>.
- [23] Krych AJ, Howard JL, Trousdale RT, Cabanela ME, Berry DJ. Total hip arthroplasty with shortening subtrochanteric osteotomy in Crowe type-IV developmental dysplasia. *J Bone Jt Surg Am* 2009;91:2213–21. <https://doi.org/10.2106/jbjs.H.01024>.
- [24] Ding ZC, Zeng WN, Mou P, Liang ZM, Wang D, Zhou ZK. Risk of dislocation after total hip arthroplasty in patients with Crowe type IV developmental dysplasia of the hip. *Orthop Surg* 2020;12:589–600. <https://doi.org/10.1111/os.12665>.
- [25] Makita H, Inaba Y, Hirakawa K, Saito T. Results on total hip arthroplasties with femoral shortening for Crowe's group IV dislocated hips. *J Arthroplast* 2007;22:32–8. <https://doi.org/10.1016/j.arth.2006.02.157>.
- [26] Oinuma K, Tamaki T, Miura Y, Kaneyama R, Shiratsuchi H. Total hip arthroplasty with subtrochanteric shortening osteotomy for Crowe grade 4 dysplasia using the direct anterior approach. *J Arthroplast* 2014;29:626–9. <https://doi.org/10.1016/j.arth.2013.07.038>.
- [27] Kiliçoğlu ÖI, Türker M, Akgül T, Yazicioğlu O. Cementless total hip arthroplasty with modified oblique femoral shortening osteotomy in Crowe type IV congenital hip dislocation. *J Arthroplast* 2013;28:117–25. <https://doi.org/10.1016/j.arth.2012.06.014>.
- [28] Kawai T, Tanaka C, Ikenaga M, Kanoe H. Cemented total hip arthroplasty with transverse subtrochanteric shortening osteotomy for Crowe group IV dislocated hip. *J Arthroplast* 2011;26:229–35. <https://doi.org/10.1016/j.arth.2010.03.029>.
- [29] Ors C, Caylak R, Togrul E. Total hip arthroplasty with the wagner cone femoral stem in patients with Crowe IV developmental dysplasia of the hip: a retrospective study. *J Arthroplast* 2022;37:103–9. <https://doi.org/10.1016/j.arth.2021.09.007>.
- [30] Yang TC, Chen CF, Tsai SW, Chen WM, Chang MC. Does restoration of hip center with subtrochanteric osteotomy provide preferable outcome for Crowe type III-IV irreducible development dysplasia of the hip? *J Chin Med Assoc* 2017;80:803–7. <https://doi.org/10.1016/j.jcma.2017.06.006>.
- [31] Ahmed E, Ibrahim el G, Ayman B. Total hip arthroplasty with subtrochanteric osteotomy in neglected dysplastic hip. *Int Orthop* 2015;39:27–33. <https://doi.org/10.1007/s00264-014-2554-0>.
- [32] Wu K, Zhang X, Chen M, Shang X. Restoration of proximal femoral anatomy during total hip arthroplasty for high developmental dysplasia of the hip: an original technique. *Orthop Surg* 2020;12:343–50. <https://doi.org/10.1111/os.12614>.
- [33] Nagoya S, Kaya M, Sasaki M, Tateda K, Kosukegawa I, Yamashita T. Cementless total hip replacement with subtrochanteric femoral shortening for severe developmental dysplasia of the hip. *J Bone Jt Surg Br* 2009;91:1142–7. <https://doi.org/10.1302/0301-620x.91b9.21736>.
- [34] Lu Z, Chen Q, Lan Y, Xie S, Lin F, Feng E. Subtrochanteric osteotomy in direct anterior approach total hip arthroplasty for Crowe IV dysplasia-surgical technique and literature review. *Orthop Surg* 2024;16:766–74. <https://doi.org/10.1111/os.13996>.
- [35] Lee SJ, Yoon KS. Two-stage total hip arthroplasty following skeletal traction after extensive soft tissue release for severe limb-length discrepancy. *Hip Int* 2021;31:223–30. <https://doi.org/10.1177/1120700019865742>.
- [36] Lai KA, Shen WJ, Huang LW, Chen MY. Cementless total hip arthroplasty and limb-length equalization in patients with unilateral Crowe type-IV hip dislocation. *J Bone Jt Surg Am* 2005;87:339–45. <https://doi.org/10.2106/jbjs.D.02097>.
- [37] Peng J, Liu Z, Ding Z, Qian Q, Wu Y. Clinical efficacy of greater trochanter osteotomy with tension wire fixation in total hip arthroplasty for Crowe type IV developmental dysplasia of the hip. *J Orthop Surg Res* 2024;19:12. <https://doi.org/10.1186/s13018-023-04344-w>.
- [38] Kawai T, Tanaka C, Kanoe H. Total hip arthroplasty for Crowe IV hip without subtrochanteric shortening osteotomy –a long term follow up study. *BMC Musculoskelet Disord* 2014;15:72. <https://doi.org/10.1186/1471-2474-15-72>.
- [39] Zhu J, Shen C, Chen X, Cui Y, Peng J, Cai G. Total hip arthroplasty with a non-modular conical stem and transverse subtrochanteric osteotomy in treatment of high dislocated hips. *J Arthroplast* 2015;30:611–4. <https://doi.org/10.1016/j.arth.2014.11.002>.
- [40] Faldini C, Tassinari L, Pederiva D, Rossomando V, Brunello M, Pilla F, et al. Direct anterior approach in total hip arthroplasty for severe Crowe IV dysplasia: retrospective clinical and radiological study. *Med* 2024;60:114. <https://doi.org/10.3390/medicina60010114>.
- [41] Li X, Lu Y, Sun J, Lin X, Tang T. Treatment of Crowe type-IV hip dysplasia using cementless total hip arthroplasty and double chevron subtrochanteric shortening osteotomy: a 5- to 10-year follow-up study. *J Arthroplast* 2017;32:475–9. <https://doi.org/10.1016/j.arth.2016.07.050>.
- [42] Baumgart R, Dahl M, Horn S, Plötz W, Baumgart T, Morrison S. Reduction of high hip dislocation with a distraction nail and arthroplasty (preliminary results of a novel technique). *Arthroplast Today* 2023;24:101257. <https://doi.org/10.1016/j.artd.2023.101257>.
- [43] Sun JY, Ma HY, Shen JM, Du YQ, Dong Y, Zhang YC, et al. Factors influencing knee valgus alignment in Crowe type IV hip dysplasia after total hip arthroplasty. *J Orthop Traumatol* 2021;22:41. <https://doi.org/10.1186/s10195-021-00601-y>.
- [44] Kabata T, Kajino Y, Inoue D, Ohmori T, Yoshitani J, Ueno T, et al. Safety range for acute limb lengthening in primary total hip arthroplasty. *Int Orthop* 2019;43:2047–56. <https://doi.org/10.1007/s00264-018-4158-6>.
- [45] Karaismailoglu B, Karaismailoglu TN. Comparison of trochanteric slide and subtrochanteric shortening osteotomy in the treatment of severe hip dysplasia: mid-term clinical outcomes of cementless total hip arthroplasty. *J Arthroplast* 2020;35:2529–36. <https://doi.org/10.1016/j.arth.2020.04.039>.
- [46] Inoue D, Kabata T, Kajino Y, Ohmori T, Ueoka K, Tsuchiya H. Comparison of mid-term clinical results between cementless and cemented femoral stems in total hip arthroplasty with femoral shortening osteotomy for Crowe type IV hips. *Arch Orthop Trauma Surg* 2021;141:1057–64. <https://doi.org/10.1007/s00402-020-03749-0>.
- [47] Sofu H, Kockara N, Gursu S, Issin A, Oner A, Sahin V. Transverse subtrochanteric shortening osteotomy during cementless total hip arthroplasty in Crowe type-III or IV developmental dysplasia. *J Arthroplast* 2015;30:1019–23. <https://doi.org/10.1016/j.arth.2015.01.045>.
- [48] Demirkale İ, Yaradılmış YU, Ateş A, Altay M. Total hip arthroplasty for high-riding hips: a retrospective analysis of 79 cases and proposal of a new classification. *J Orthop Surg* 2020;28:2309499020924164. <https://doi.org/10.1177/2309499020924164>.
- [49] Lee SJ, Yoo JJ, Kim HJ. Cementless total hip arthroplasty involving trochanteric osteotomy without subtrochanteric shortening for high hip dislocation. *Clin Orthop Surg* 2017;9:19–28. <https://doi.org/10.4055/cios.2017.9.1.19>.
- [50] Bilgen ÖF, Yaray O, Mutlu M, Aksakal AM. Intraoperative two-stage evaluation of muscle contractures in Crowe type IV hips in total hip arthroplasty (a new surgical technique). *Hip Int* 2022;32:391–400. <https://doi.org/10.1177/1120700020959784>.
- [51] Wu X, Li SH, Lou LM, Cai ZD. The techniques of soft tissue release and true socket reconstruction in total hip arthroplasty for patients with severe developmental dysplasia of the hip. *Int Orthop* 2012;36:1795–801. <https://doi.org/10.1007/s00264-012-1622-6>.
- [52] Kayaalp ME, Can A, Erdogan F, Özşahin MK, Aydingöz O, Kaynak G. Clinical and radiological results of Crowe type 3 or 4 dysplasia patients operated on with total hip arthroplasty using a cementless rectangular femoral component without fixating or grafting the transverse osteotomy site. *J Arthroplast* 2020;35:2537–42. <https://doi.org/10.1016/j.arth.2020.04.045>.
- [53] Biçici V, Bingöl İ, Sazak T. Mid-term results of total hip arthroplasty with subtrochanteric Z-osteotomy in Crowe type 3-4 developmental hip dysplasia. *Turk J Med Sci* 2021;51:1976–83. <https://doi.org/10.3906/sag-2102-217>.
- [54] Bao N, Meng J, Zhou L, Guo T, Zeng X, Zhao J. Lesser trochanteric osteotomy in total hip arthroplasty for treating CROWE type IV developmental dysplasia of hip. *Int Orthop* 2013;37:385–90. <https://doi.org/10.1007/s00264-012-1758-4>.



- [55] McHale KA. CORR Insights®: assessing leg length discrepancy is necessary before arthroplasty in patients with unilateral Crowe type IV hip dislocation. *Clin Orthop Relat Res* 2023;481:1790–1. <https://doi.org/10.1097/corr.0000000000002754>.
- [56] Xu Q, Wang Q, Zhu J, Lin J, Lu Z, Wang T, et al. Clinical outcomes of proximal femoral reconstruction technique combined with THA in the treatment of high dislocation secondary to septic arthritis: a retrospective single-center study. *BMC Musculoskelet Disord* 2023;24:732. <https://doi.org/10.1186/s12891-023-06818-8>.
- [57] Higuchi Y, Hasegawa Y, Ishiguro N. Leg lengthening of more than 5 cm is a risk factor for sciatic nerve injury after total hip arthroplasty for adult hip dislocation. *Nagoya J Med Sci* 2015;77:455–63.
- [58] Sun C, Zhang Y, Li LT, Ding H, Guo T, Zhao JN. Long-Term outcomes of total hip arthroplasty with transverse subtrochanteric shortening osteotomy and modular stem in Crowe IV developmental dysplasia. *J Arthroplast* 2021;36:630–5. <https://doi.org/10.1016/j.arth.2020.08.031>.
- [59] Zeng WN, Liu JL, Wang FY, Zhang X, Fan HQ, Chen GX, et al. Total hip arthroplasty for patients with Crowe type IV developmental dysplasia of the hip: ten years results. *Int J Surg* 2017;42:17–21. <https://doi.org/10.1016/j.jisu.2017.04.029>.
- [60] Fujishiro T, Nishiyama T, Hayashi S, Kurosaka M, Kanno T, Masuda T. Leg length change in total hip arthroplasty with subtrochanteric femoral shortening osteotomy for Crowe type IV developmental hip dysplasia. *J Arthroplast* 2012;27:1019–22. <https://doi.org/10.1016/j.arth.2012.01.032>.
- [61] Oe K, Iida H, Nakamura T, Okamoto N, Wada T. Subtrochanteric shortening osteotomy combined with cemented total hip arthroplasty for Crowe group IV hips. *Arch Orthop Trauma Surg* 2013;133:1763–70. <https://doi.org/10.1007/s00402-013-1869-4>.
- [62] Hameed D, Dubin JA, Bains SS, Moore MC, Standard SC, Mont MA. Innovative hip distraction approaches for total hip arthroplasty in Crowe IV developmental dysplasia or similar high-riding patients. *J Arthroplast* 2024;39:5124–30. <https://doi.org/10.1016/j.arth.2024.04.019>.
- [63] Wang D, Li DH, Li Q, Wang HY, Luo ZY, Yang Y, et al. Subtrochanteric shortening osteotomy during cementless total hip arthroplasty in young patients with severe developmental dysplasia of the hip. *BMC Musculoskelet Disord* 2017;18:491. <https://doi.org/10.1186/s12891-017-1857-x>.
- [64] Li Y, Zhang X, Wang Q, Peng X, Wang Q, Jiang Y, et al. Equalisation of leg lengths in total hip arthroplasty for patients with Crowe type-IV developmental dysplasia of the hip: classification and management. *Bone Jt J* 2017;99-b:872–9. <https://doi.org/10.1302/0301-620x.99b7.Bjj-2016-1328.R1>.
- [65] Zhao X, Zhu ZA, Xie YZ, Yu B, Yu DG. Total hip replacement for high dislocated hips without femoral shortening osteotomy. *J Bone Jt Surg Br* 2011;93:1189–93. <https://doi.org/10.1302/0301-620x.93b9.25914>.
- [66] Erdem Y, Bek D, Atbasi Z, Neyisci C, Yildiz C, Basbozkurt M. Total hip arthroplasty with rectangular stems and subtrochanteric transverse shortening osteotomy in Crowe type IV hips: a retrospective study. *Arthroplast Today* 2019;5:234–42. <https://doi.org/10.1016/j.arth.2019.03.002>.
- [67] Ibrahim FMF, Gameel S. Total cementless unconstrained hip arthroplasty in Crowe type IV hip dysplasia with subtrochanteric derotation osteotomy. *Acta Orthop Belg* 2022;88:517–24. <https://doi.org/10.52628/88.3.7416>.
- [68] Sukur E, Senel A, Ozdemir U, Akman YE, Azboy İ, Ozturkmen Y. Cementless modular total hip arthroplasty with subtrochanteric transverse shortening osteotomy for high hip dislocations. *J Orthop Surg Res* 2022;17:139. <https://doi.org/10.1186/s13018-022-03025-4>.
- [69] Lai KA, Liu J, Liu TK. Use of iliofemoral distraction in reducing high congenital dislocation of the hip before total hip arthroplasty. *J Arthroplast* 1996;11:588–93. [https://doi.org/10.1016/s0883-5403\(96\)80114-8](https://doi.org/10.1016/s0883-5403(96)80114-8).
- [70] Tu Q, Ding HW, Chen H, Shen JJ, Miao QJ, Liu B, et al. Preliminary application of 3D-printed individualised guiding templates for total hip arthroplasty in Crowe type IV developmental dysplasia of the hip. *Hip Int* 2022;32:334–44. <https://doi.org/10.1177/1120700020948006>.
- [71] Akinci O, Turgut A. Long-Term results of total hip arthroplasty with step-cut osteotomy in Crowe type IV dysplastic hips. *Indian J Orthop* 2022;56:672–9. <https://doi.org/10.1007/s43465-021-00567-3>.
- [72] Luo Z, Chen M, Hu F, Ni Z, Ji X, Zhang X, et al. Cementless total hip arthroplasty with extended sliding trochanteric osteotomy for high congenital hip dislocation: a retrospective study. *Med* 2017;96:e6581. <https://doi.org/10.1097/md.0000000000006581>.
- [73] Kong X, Chai W, Chen J, Yan C, Shi L, Wang Y. Intraoperative monitoring of the femoral and sciatic nerves in total hip arthroplasty with high-riding developmental dysplasia. *Bone Jt J* 2019;101-b:1438–46. <https://doi.org/10.1302/0301-620x.101b11.Bjj-2019-0341.R2>.
- [74] Imam MA, Fathalla I, Holton J, Nabil M, Kashif F. Cementless total hip replacement for the management of severe developmental dysplasia of the hip in the middle eastern population: a prospective analysis. *Front Surg* 2016;3:31. <https://doi.org/10.3389/fsurg.2016.00031>.
- [75] Liu T, Wang S, Huang G, Wang W. Treatment of Crowe IV developmental dysplasia of the hip with cementless total hip arthroplasty and shortening subtrochanteric osteotomy. *J Int Med Res* 2019;47:3223–33. <https://doi.org/10.1177/0300060519853383>.
- [76] Akiyama H, Kawanabe K, Yamamoto K, Kuroda Y, So K, Goto K, et al. Cemented total hip arthroplasty with subtrochanteric femoral shortening transverse osteotomy for severely dislocated hips: outcome with a 3- to 10-year follow-up period. *J Orthop Sci* 2011;16:270–7. <https://doi.org/10.1007/s00776-011-0049-z>.
- [77] Zhao HY, Kang PD, Shi XJ, Zhou ZK, Yang J, Shen B, et al. Effects of total hip arthroplasty on axial alignment of the lower limb in patients with unilateral developmental hip dysplasia (Crowe type IV). *J Arthroplast* 2019;34:2406–14. <https://doi.org/10.1016/j.arth.2019.04.028>.
- [78] Chareancholvanich K, Becker DA, Gustilo RB. Treatment of congenital dislocated hip by arthroplasty with femoral shortening. *Clin Orthop Relat Res* 1999;127–35. <https://doi.org/10.1097/00003086-199903000-00016>.
- [79] Rollo G, Solarino G, Vicenti G, Picca G, Carrozzo M, Moretti B. Subtrochanteric femoral shortening osteotomy combined with cementless total hip replacement for Crowe type IV developmental dysplasia: a retrospective study. *J Orthop Traumatol* 2017;18:407–13. <https://doi.org/10.1007/s10195-017-0466-7>.
- [80] Ueoka K, Kabata T, Kajino Y, Inoue D, Ohmori T, Ueno T, et al. Patient-reported outcomes following primary total hip arthroplasty in Crowe type III or IV developmental dysplasia are comparable to those in Crowe type I: a case-control study of 96 hips with intermediate-term follow-up. *BMC Musculoskelet Disord* 2020;21:344. <https://doi.org/10.1186/s12891-020-03371-6>.
- [81] Shemesh SS, Robinson J, Overley S, Bronson MJ, Moucha CS, Chen D. Novel technique for intraoperative sciatic nerve assessment in complex primary total hip arthroplasty: a pilot study. *Hip Int* 2018;28:210–7. <https://doi.org/10.5301/hipint.5000553>.
- [82] Kerboul M, Hamadouche M, Kerboul L. Total hip arthroplasty for Crowe type IV developmental hip dysplasia: a long-term follow-up study. *J Arthroplast* 2001;16:170–6. <https://doi.org/10.1054/arth.2001.28368>.
- [83] Caylak R, Ors C, Togrul E. Minimum 10-year results of cementless ceramic-on-ceramic total hip arthroplasty performed with transverse subtrochanteric osteotomy in Crowe type IV hips. *J Arthroplast* 2021;36:3519–26. <https://doi.org/10.1016/j.arth.2021.05.035>.
- [84] Chen M, Gittings DJ, Yang S, Liu X. Total hip arthroplasty for Crowe type IV developmental dysplasia of the hip using a titanium mesh cup and subtrochanteric femoral osteotomy. *Iowa Orthop J* 2018;38:191–5.
- [85] Huang ZY, Ling J, Zeng ZM, Di ZL, Zhang JH, Tao K. Mid-Term outcomes of cemented stem and subtrochanteric shortening derotational osteotomy in total hip arthroplasty for Crowe IV developmental dysplasia. *Orthop Surg* 2022;14:3178–86. <https://doi.org/10.1111/os.13515>.
- [86] Wang D, Li LL, Wang HY, Pei FX, Zhou ZK. Long-Term results of cementless total hip arthroplasty with subtrochanteric shortening osteotomy in Crowe type IV developmental dysplasia. *J Arthroplast* 2017;32:1211–9. <https://doi.org/10.1016/j.arth.2016.11.005>.
- [87] Jin JY, Yoon TR, Park KS, Jin SY, Jung DM, Li QS. Mid-term results of total hip arthroplasty with modified trochanteric osteotomy in Crowe type IV developmental dysplasia of the hip. *BMC Surg* 2021;21:9. <https://doi.org/10.1186/s12893-020-01002-4>.
- [88] Lerch M, Thorey F, von Lewinski G, Klages P, Wirth CJ, Windhagen H. An alternative treatment method to restore limb-length discrepancy in osteoarthritis with high congenital hip dislocation. *Arch Orthop Trauma Surg* 2009;129:1593–9. <https://doi.org/10.1007/s00402-009-0846-4>.
- [89] Park CV, Lim SJ, Cha YT, Park YS. Total hip arthroplasty with subtrochanteric shortening osteotomy in patients with high hip dislocation secondary to childhood septic arthritis: a matched comparative study with Crowe IV developmental dysplasia. *J Arthroplast* 2020;35:204–11. <https://doi.org/10.1016/j.arth.2019.08.034>.
- [90] Sonohata M, Kitajima M, Kawano S, Mawatari M. Nerve palsy after total hip arthroplasty without subtrochanteric femoral shortening osteotomy for a completely dislocated hip joint. *Open Orthop J* 2016;10:785–92. <https://doi.org/10.2174/1874325001610010785>.
- [91] Wang S, Zeng M, Zhang X, Liu T. Total hip arthroplasty with soft tissue release for patients with severe development dysplasia of the hip. *Zhong Nan Da Xue Xue Bao Yi Xue Ban* 2018;43:54–7. <https://doi.org/10.11817/j.issn.1672-7347.2018.01.009>.
- [92] Eggi S, Hankemayer S, Müller ME. Nerve palsy after leg lengthening in total replacement arthroplasty for developmental dysplasia of the hip. *J Bone Joint Surg Br* 1999;81:843–5.