



ELSEVIER

Contents lists available at ScienceDirect

The Journal of Arthroplasty

journal homepage: www.arthroplastyjournal.org

World Expert Meeting in Arthroplasty 2024

Is There Evidence Supporting the Use of All-Polyethylene Tibial Components in Primary Total Knee Arthroplasty?



Hamidreza Yazdi, MD ^a, Mohammadamin Haghbin, MD ^{b,*},
 Mohammadmahdi Omidian, MD ^{b,c}, Aidin Eslam Pour, MD, MS ^d,
 Victor H. Hernandez, MD, MS ^e, Fouad Zamel Sadek, MD ^f, Masood Umer, MD ^g,
 Antonio Fraguas Castany, MD ^h

^a Department of Knee Surgery, Firoozgar Hospital, Iran University of Medical Sciences, Tehran, Iran^b Bone and Joint Reconstruction Research Center, Department of Orthopedics, School Of Medicine, Iran University of Medical Sciences, Tehran, Iran^c Department of Orthopedic Surgery and Traumatology, Imam Hossein Medical Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran^d Department of Orthopaedic Surgery, Yale University, New Haven, Connecticut^e Division of Arthroplasty & Adult Joint Reconstruction, Orthopaedic Department, Miller School of Medicine, University of Miami, MIA^f Pelvic and Arthroplasty Unit, Cairo University Hospitals, Gize, Egypt^g Department of Orthopedic Surgery, Aga Khan University Karachi, Karachi, Pakistan^h Knee and Hip Surgery Centro Medico, Teknon Barcelon, Barcelona, Spain

ARTICLE INFO

Article history:

Received 16 September 2024

Received in revised form

10 October 2024

Accepted 14 October 2024

Available online 22 October 2024

Keywords:

all-polyethylene

metal-backed

tibial component

total knee arthroplasty

outcomes

Question: Is There Evidence Supporting the use of All-Polyethylene Tibial Components in Primary Total Knee Arthroplasty?**Response/Recommendation:** Excellent performance across various patient demographics, including age, body mass index (BMI), and activity level.

More than two years' follow-up studies indicate that all-polyethylene tibial (APT) components offer major cost savings with excellent performance across various patient demographics, including age, body mass index (BMI), and activity level.

Level of evidences: Moderate.**Expert vote:** Agree: 81.2%. Disagree: 12.6%. Abstain: 6.3%.

The survival and revision rates, complications, and functional scores of APT components are comparable to those of metal-backed tibial (MBT) components.

Level of Evidence: Moderate.**Expert vote:** Agree: 61.5%. Disagree: 24%. Abstain: 14.5%.

Rationale

Basic Science

Early designs of total knee arthroplasty (TKA) primarily utilized all-polyethylene tibial (APT) components, with studies from that

No author associated with this paper has disclosed any potential or pertinent conflicts which may be perceived to have impending conflict with this work. For full disclosure statements refer to <https://doi.org/10.1016/j.arth.2024.10.073>.

* Address correspondence to: Mohammadamin Haghbin, MD, Shafa Orthopedic Hospital, Baharestan Square, Tehran, Iran 1157637131.

<https://doi.org/10.1016/j.arth.2024.10.073>

0883-5403/© 2024 Elsevier Inc. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

era reporting survival rates exceeding 90% at more than 10 years of follow-up for both cruciate-retaining and posterior-stabilized designs with APT components [1–7]. Early aseptic loosening, often due to poor surgical techniques or design flaws [8–13], led to the introduction of mobile-bearing tibial (MBT) components in the 1970s, providing theoretical advantages such as modularity, which allows for intraoperative flexibility, and isolated polyethylene liner exchanges without disturbing tibial fixation [14,15]. However, designs, components, and surgical techniques when utilizing APT have improved dramatically over the years, leading to better clinical outcomes and increased use among orthopaedic surgeons. To evaluate the outcomes of APT components, studies with more than

two years of follow-up, including meta-analyses, randomized controlled trials, and retrospective and prospective cohort studies published in peer-reviewed journals, were analyzed.

The APT components demonstrate good to excellent survival with revision rates comparable to MBT components [15–46]. These results were consistent across different age groups [24,32,37,39–41,47–51], with reports from the Total Joint Replacement Registry and other large-scale studies confirming satisfactory outcomes and a low risk of revision in patients below 65 years [37,40]. The APT technique also demonstrated satisfactory outcomes among active patients [26,37,39–41,51]. Most studies evaluating the effect of BMI on outcomes reported that both APT and MBT components performed similarly [37,43,52–55]. Notably, some studies indicated that APT components had better results in morbidly obese patients (BMI > 40) [24,29]. The APT components also exhibit favorable functional scores [22,23,25,27–29,31,33,34,36,39,41,42,45,46,50,51]. Their use in TKA can reduce hospital charges and save costs [8,27,32,34,36,40,45,46,56–62]. The complication rates associated with APT components in primary TKA, such as infections, osteolysis, periprosthetic fractures, component migration, and wound complications, are acceptable and comparable to MBT [20,21,24,25,27,29–31,33,35,37,39,40,42–46,63]. The clinical performance of APT components in TKA has demonstrated consistent benefits, particularly with respect to more than 10 years implant survival and patient satisfaction, but their success is closely linked to both their design and surgical technique [2–6,8,9,12,13,20,23,30,37,42,49,60,61,63–98]. Many randomized controlled trials also confirm these results [26,33,60,66,75]. Nevertheless, patient-specific factors and other potential limitations must be carefully considered when selecting APT components. Notable limitations include the inability to fine-tune knee balancing postimplantation, the lack of stemmed options, and the challenges of undertaking debridement, antibiotics and implant retention in early infections [20,99,100]. However, these concerns are mitigated by the lower infection rates reported with APT constructs [20,21,24,25,27,29–31,33,35,37,39,40,42–46,60,63,65,66].

CRedit authorship contribution statement

Hamidreza Yazdi: Supervision, Conceptualization. **Mohammadamin Haghighi:** Methodology, Investigation. **Mohammadmahdi Omidian:** Methodology, Investigation. **Aidin Eslam Pour:** Methodology, Conceptualization. **Victor H. Hernandez:** Conceptualization. **Fouad Zamel Sadek:** Conceptualization. **Masood Umer:** Conceptualization. **Antonio Fraguas Castany:** Conceptualization.

References

- [1] Font-Rodriguez DE, Scuderi GR, Insall JN. Survivorship of cemented total knee arthroplasty. *Clin Orthop Relat Res* 1997;345:79–86. <https://doi.org/10.1097/00003086-199712000-00012>.
- [2] Gill GS, Joshi AB, Mills DM. Total condylar knee arthroplasty. 16- to 21-year results. *Clin Orthop Relat Res* 1999;367:210–5. <https://doi.org/10.1097/00003086-199910000-00026>.
- [3] Ranawat CS, Boachie-Adjei O. Survivorship analysis and results of total condylar knee arthroplasty. Eight- to 11-year follow-up period. *Clin Orthop Relat Res* 1988;226:6–13. <https://doi.org/10.1097/00003086-198810000-00003>.
- [4] Ranawat CS, Flynn Jr WF, Saddler S, Hansraj KK, Maynard MJ. Long-term results of the total condylar knee arthroplasty. A 15-year survivorship study. *Clin Orthop Relat Res* 1993;286:94–102. <https://doi.org/10.1097/00003086-199301000-00015>.
- [5] Rodriguez JA, Bhende H, Ranawat CS. Total condylar knee replacement: a 20-year followup study. *Clin Orthop Relat Res* 2001;388:10–7. <https://doi.org/10.1097/00003086-200107000-00004>.
- [6] Scuderi GR, Insall JN, Windsor RE, Moran MC. Survivorship of cemented knee replacements. *J Bone Joint Surg Br* 1989;71:798–803. <https://doi.org/10.1302/0301-620X.71B5.2584250>.
- [7] Vince KG, Insall JN. Long-term results of cemented total knee arthroplasty. *Orthop Clin North Am* 1988;19:575–80. [https://doi.org/10.1016/s0030-5898\(20\)32262-8](https://doi.org/10.1016/s0030-5898(20)32262-8).
- [8] Apel DM, Tozzi JM, Dorr LD. Clinical comparison of all-polyethylene and metal-backed tibial components in total knee arthroplasty. *Clin Orthop Relat Res* 1991;273:243–52. <https://doi.org/10.1097/00003086-199112000-00035>.
- [9] Ducheyne P, Kagan A, Lacey JA. Failure of total knee arthroplasty due to loosening and deformation of the tibial component. *J Bone Joint Surg* 1978;60:384–91. <https://doi.org/10.2106/00004623-197860030-00021>.
- [10] Evanski PM, Waugh TR, Orofino CF, Anzel SH. UCI knee replacement. *Clin Orthop Relat Res* 1976;33–8. <https://doi.org/10.1097/00003086-197610000-00007>.
- [11] Faris PM, Ritter MA, Keating EM, Meding JB, Harty LD. The agc all-polyethylene tibial component. *J Bone Joint Surg Am* 2003;85:489–93. <https://doi.org/10.2106/00004623-200303000-00014>.
- [12] Hyldahl H, Regnér L, Carlsson L, Kärrholm J, Weidenhielm L. All-polyethylene vs. metal-backed tibial component in total knee arthroplasty—a randomized RSA study comparing early fixation of horizontally and completely cemented tibial components: Part 1. Horizontally cemented components: AP better fixated than MB. *Acta Orthop* 2005;76:769–77. <https://doi.org/10.1080/17453670510045354>.
- [13] Ritter MA. The cemented all-poly tibia. *Orthopedics* 1994;17:841. <https://doi.org/10.3928/0147-7447-19940901-32>.
- [14] AbuMoussa S, White CCT, Eichinger JK, Friedman RJ. All-polyethylene versus metal-backed tibial components in total knee arthroplasty. *J Knee Surg* 2019;32:714–8. <https://doi.org/10.1055/s-0039-1683979>.
- [15] Elmhireh A, Abuodeh Y, Alzobi O, Zikria B, Alkhayarin M, Morrey BF. All-polyethylene versus metal-backed tibial components in total knee arthroplasty: a meta-analysis of randomized controlled trials. *Eur J Orthop Surg Traumatol* 2023;33:3611–22. <https://doi.org/10.1007/s00590-023-03594-y>.
- [16] Forster M. Survival analysis of primary cemented total knee arthroplasty Which designs last? *J Arthroplasty* 2003;18:265–70. <https://doi.org/10.1054/arth.2003.50051>.
- [17] Voss B, El-Othmani MM, Schnur A-K, Botchway A, Mihalko WM, Saleh KJ. A meta-analysis comparing all-polyethylene tibial component to metal-backed tibial component in total knee arthroplasty: assessing survivorship and functional outcomes. *J Arthroplasty* 2016;31:2628–36. <https://doi.org/10.1016/j.arth.2015.08.035>.
- [18] Longo UG, Ciuffreda M, D'Andrea V, Mannering N, Locher J, Denaro V. All-polyethylene versus metal-backed tibial component in total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc* 2017;25:3620–36. <https://doi.org/10.1007/s00167-016-4168-0>.
- [19] Bonanzinga T, Gambaro FM, Iacono F, Leogrande F, Di Lascio L, Marcacci M. All-polyethylene versus metal-backed tibial component in total knee arthroplasty: a meta-analysis of randomized controlled trials. *J Arthroplasty* 2023;39:1353–60. <https://doi.org/10.1016/j.arth.2023.10.058>.
- [20] Gudnason A, Hailer NP, A WD, Sundberg M, Robertsson O. All-polyethylene versus metal-backed tibial components—an analysis of 27,733 cruciate-retaining total knee replacements from the Swedish knee arthroplasty register. *J Bone Joint Surg Am* 2014;96:994–9. <https://doi.org/10.2106/JBJS.M.00373>.
- [21] Kelley B, Mullen K, De A, Sassoon A. Modular metal-backed tibial components provide minimal mid-term survivorship benefits despite increased cost and frequency of use: a retrospective review of the American joint replacement registry database. *J Arthroplasty* 2022;37:1570–1574.e1. <https://doi.org/10.1016/j.arth.2022.01.031>.
- [22] Nachtnabl L, Apostolopoulos V, Mahdal M, Pazourek L, Brancik P, Valousek T, et al. Implant preference and clinical outcomes of patients with staged bilateral total knee arthroplasty: all-polyethylene and contralateral metal-backed tibial components. *J Clin Med* 2023;12:7438. <https://doi.org/10.3390/jcm12237438>.
- [23] Van Hamersveld KT, Marang-Van De Mheen PJ, Nelissen R, Toksvig-Larsen S. Migration of all-polyethylene compared with metal-backed tibial components in cemented total knee arthroplasty. *Acta Orthop* 2018;89:412–7. <https://doi.org/10.1080/17453674.2018.1464317>.
- [24] Houdek MT, Watts CD, Martin JR, Trousdale RT, Taunton MJ. Metal or modularity: why do metal-backed tibias have inferior outcomes to all-polyethylene tibial components in patients with osteoarthritis. *J Arthroplasty* 2017;32:836–42. <https://doi.org/10.1016/j.arth.2016.09.036>.
- [25] Apostolopoulos V, Nachtnabl L, Mahdal M, Pazourek L, Bohac P, Janicek P, et al. Clinical outcomes and survival comparison between NexGen all-poly and its metal-backed equivalent in total knee arthroplasty. *Int Orthop* 2023;47:2207–13. <https://doi.org/10.1007/s00264-023-05772-3>.
- [26] Bettinson KA, Pinder IM, Moran CG, Weir DJ, Lingard EA. All-polyethylene compared with metal-backed tibial components in total knee arthroplasty at ten years. A prospective, randomized controlled trial. *J Bone Joint Surg Am* 2009;91:1587–94. <https://doi.org/10.2106/JBJS.G.01427>.
- [27] Dalury DF, Pomeroy DL, Gonzales RA, Gruen TA, Adams MJ, Empson JA. Midterm results of all-polyethylene tibial components in primary total knee arthroplasty. *J Arthroplasty* 2009;24:620–4. <https://doi.org/10.1016/j.arth.2008.03.004>.
- [28] Dalury DF, Tucker KK, Kelley TC. All-polyethylene tibial components in obese patients are associated with low failure at midterm followup. *Clin Orthop Relat Res* 2012;470:117–24. <https://doi.org/10.1007/s11999-011-1964-1>.

- [29] Diaz Dileria F, Blanchard T, Vasarhelyi E, Lanting B, Howard J. Minimum 2-year outcomes of all-polyethylene tibial components in patients who have body mass index of 35 or greater. *J Arthroplasty* 2023;38(6S):S157–63. <https://doi.org/10.1016/j.arth.2023.03.078>.
- [30] Faris PM, Ritter MA, Keating EM, Meding JB, Harty LD. The AGC all-polyethylene tibial component: a ten-year clinical evaluation. *J Bone Joint Surg Am* 2003;85:489–93. <https://doi.org/10.2106/00004623-200303000-00014>.
- [31] Garabano G, Rodriguez J, Perez Alamino L, Pesciallo CA, Del Sel H, Lopreite F. Stress shielding in total knee replacements: comparative analysis between titanium and all-polyethylene bases at 10 years follow-up. *J Orthop* 2022;34:276–81. <https://doi.org/10.1016/j.jor.2022.09.007>.
- [32] Gioe TJ, Sinner P, Mehle S, Ma W, Killeen KK. Excellent survival of all-polyethylene tibial components in a community joint registry. *Clin Orthop Relat Res* 2007;464:88–92. <https://doi.org/10.1097/BLO.0b013e31812f7879>.
- [33] Gioe TJ, Stroemer ES, Santos ER. All-polyethylene and metal-backed tibias have similar outcomes at 10 years: a randomized level I [corrected] evidence study. *Clin Orthop Relat Res* 2007;455:212–8. <https://doi.org/10.1097/01.blo.0000238863.69486.97>.
- [34] Gustke KA, Gelbke MK. All-polyethylene tibial component use for elderly, low-demand total knee arthroplasty patients. *J Arthroplasty* 2017;32:2421–6. <https://doi.org/10.1016/j.arth.2017.02.077>.
- [35] Houdek MT, Wagner ER, Wyles CC, Watts CD, Cass JR, Trousdale RT. All-polyethylene tibial components: an analysis of long-term outcomes and infection. *J Arthroplasty* 2016;31:1476–82. <https://doi.org/10.1016/j.arth.2015.12.048>.
- [36] Jabbal M, Clement N, Walmsley PJ. All-polyethylene tibia components have the same functional outcomes and survival, and are more cost-effective than metal-backed components in patients 70 years and older undergoing total knee arthroplasty: propensity match study with a minimum five-year follow-up. *Bone Jt Open* 2022;3:969–76. <https://doi.org/10.1302/2633-1462.3.12.BJO-2022-0063.R1>.
- [37] Kremers HM, Sierra RJ, Schleck CD, Berry DJ, Cabanela ME, Hanssen AD, et al. Comparative survivorship of different tibial designs in primary total knee arthroplasty. *J Bone Joint Surg Am* 2014;96:e121. <https://doi.org/10.2106/JBJS.M.00820>.
- [38] Maradit Kremers H, Kremers WK, Sierra RJ, Lewallen DG, Berry DJ. Competing risk of death when comparing tibial implant types in total knee arthroplasty. *J Bone Joint Surg Am* 2016;98:591–6. <https://doi.org/10.2106/JBJS.15.00488>.
- [39] Meftah M, Ranawat AS, Sood AB, Rodriguez JA, Ranawat CS. All-polyethylene tibial implant in young, active patients a concise follow-up, 10 to 18 years. *J Arthroplasty* 2012;27:10–4. <https://doi.org/10.1016/j.arth.2011.05.013>.
- [40] Mohan V, Inacio MC, Namba RS, Sheth D, Paxton EW. Monoblock all-polyethylene tibial components have a lower risk of early revision than metal-backed modular components. *Acta Orthop* 2013;84:530–6. <https://doi.org/10.3109/17453674.2013.862459>.
- [41] Ranawat AS, Mohanty SS, Goldsmith SE, Rasquinha VJ, Rodriguez JA, Ranawat CS. Experience with all-polyethylene total knee arthroplasty in younger, active patients with follow-up from 2 to 11 years. *J Arthroplasty* 2005;20(7 Suppl 3):7–11. <https://doi.org/10.1016/j.arth.2005.04.027>.
- [42] Rodriguez JA, Baez N, Rasquinha V, Ranawat CS. Metal-backed and all-polyethylene tibial components in total knee replacement. *Clin Orthop Relat Res* 2001;392:174–83. <https://doi.org/10.1097/00003086-200111000-00021>.
- [43] Toman J, Iorio R, Healy WL. All-polyethylene and metal-backed tibial components are equivalent with BMI of less than 37.5. *Clin Orthop Relat Res* 2012;470:108–16. <https://doi.org/10.1007/s11999-011-2124-3>.
- [44] Trousdale RT. The cemented all-polyethylene tibia: solution for all seasons – affirms. *Orthop Proc* 2018;100-B(SUPP_10):99. <https://doi.org/10.1302/1358-992x.2018.10.099>.
- [45] Udomkiat P, Dorr LD, Long W. Matched-pair analysis of all-polyethylene versus metal-backed tibial components. *J Arthroplasty* 2001;16:689–96. <https://doi.org/10.1054/arth.2001.25500>.
- [46] Yassin M, Garti A, Weissbrot M, Ashkenazi U, Khatib M, Robinson D. All-polyethylene tibial components are not inferior to metal-backed tibial components in long-term follow-up of knee arthroplasties. *Eur J Orthop Surg Traumatol* 2015;25:1087–91. <https://doi.org/10.1007/s00590-015-1645-y>.
- [47] Rand JA, Trousdale RT, Ilstrup DM, Harmsen WS. Factors affecting the durability of primary total knee prostheses. *J Bone Joint Surg Am* 2003;85:259–65. <https://doi.org/10.2106/00004623-200302000-00012>.
- [48] L'Insalata JL, Stern SH, Insall JN. Total knee arthroplasty in elderly patients. Comparison of tibial component designs. *J Arthroplasty* 1992;7:261–6. [https://doi.org/10.1016/0883-5403\(92\)90046-s](https://doi.org/10.1016/0883-5403(92)90046-s).
- [49] Pagnano MW, Levy BA, Berry DJ. Cemented all polyethylene tibial components in patients age 75 years and older. *Clin Orthop Relat Res* 1999;73–80.
- [50] Najibi S, Iorio R, Surdam JW, Whang W, Appleby D, Healy WL. All-polyethylene and metal-backed tibial components in total knee arthroplasty: a matched pair analysis of functional outcome. *J Arthroplasty* 2003;18(7 Suppl 1):9–15. [https://doi.org/10.1016/s0883-5403\(03\)00304-8](https://doi.org/10.1016/s0883-5403(03)00304-8).
- [51] Singh A, Singh KK. Clinical evaluation of efficacy and performance of all-poly tibial freedom(R) total knee system for treating osteoarthritis patients: three-year follow up study. *J Clin Diagn Res* 2017;11:RC01–5. <https://doi.org/10.7860/JCDR/2017/26418.10671>.
- [52] Winiarsky R, Barth P, Lotke P. Total knee arthroplasty in morbidly obese patients. *JBJS* 1998;80:1770–4.
- [53] Jämsen E, Nevalainen P, Eskelinen A, Huotari K, Kalliovalkama J, Moilanen T. Obesity, diabetes, and preoperative hyperglycemia as predictors of peri-prosthetic joint infection: a single-center analysis of 7181 primary hip and knee replacements for osteoarthritis. *JBJS* 2012;94:e101. <https://doi.org/10.2106/jbjs.j.01935>.
- [54] Friedman RJ, Hess S, Berkowitz SD, Homering M. Complication rates after hip or knee arthroplasty in morbidly obese patients. *Clin Orthop Relat Res* 2013;471:3358–66. <https://doi.org/10.1007/s11999-013-3049-9>.
- [55] Tsao AK, Jones LC, Lewallen DG. What patient and surgical factors contribute to implant wear and osteolysis in total joint arthroplasty? *JAAOS-Journal of the American Academy of Orthopaedic Surgeons* 2008;16:S7–13. <https://doi.org/10.5435/00124635-200800001-00004>.
- [56] Mendenhall S. Where are implant prices going. *Orthop Network News* 1996;7:2–6.
- [57] Gioe TJ, Stroemer ES, Santos ER. All-polyethylene and metal-backed tibias have similar outcomes at 10 Years: a randomized level I evidence study. *Clin Orthop Relat Res* 2007;455:212–8.
- [58] Healy WL, Iorio R, Ko J, Appleby D, Lemos DW. Impact of cost reduction programs on short-term patient outcome and hospital cost of total knee arthroplasty. *J Bone Joint Surg Am* 2002;84:348–53. <https://doi.org/10.2106/00004623-200203000-00003>.
- [59] Pomeroy DL, Schaper LA, Badenhansen WE, Suthers KE, Smith MW, Empson JA, et al. Results of all-polyethylene tibial components as a cost-saving technique. *Clin Orthop Relat Res* 2000;380:140–3. <https://doi.org/10.1097/00003086-200011000-00018>.
- [60] Gioe TJ, Glynn J, Sembrano J, Suthers K, Santos ER, Singh J. Mobile and fixed-bearing (all-polyethylene tibial component) total knee arthroplasty designs. A prospective randomized trial. *J Bone Joint Surg Am* 2009;91:2104–12. <https://doi.org/10.2106/JBJS.H.01442>.
- [61] Muller SD, Deehan DJ, Holland JP, Outterside SE, Kirk LM, Gregg PJ, et al. Should we reconsider all-polyethylene tibial implants in total knee replacement? *J Bone Joint Surg Br* 2006;88:1596–602. <https://doi.org/10.1302/0301-620X.88B12.17695>.
- [62] Sabeh K, Alam M, Rosas S, Hussain S, Schneiderbauer M. Cost analysis of all-polyethylene compared to metal-backed implants in total knee arthroplasty. *Surg Technol Int* 2018;32:249–55.
- [63] Robinson RP, Green TM. Eleven-year implant survival rates of the all-polyethylene and metal-backed modular Optetrak posterior stabilized knee in bilateral simultaneous cases. *J Arthroplasty* 2011;26:1165–9. <https://doi.org/10.1016/j.arth.2011.01.009>.
- [64] Rand JA. Comparison of metal-backed and all-polyethylene tibial components in cruciate condylar total knee arthroplasty. *J Arthroplasty* 1993;8:307–13. [https://doi.org/10.1016/s0883-5403\(06\)80094-x](https://doi.org/10.1016/s0883-5403(06)80094-x).
- [65] Selvan DR, Santini AJA, Davidson JS, Pope JA. The medium-term survival analysis of an all-polyethylene tibia in a single-series cohort of over 1000 knees. *J Arthroplasty* 2020;35:2837–42. <https://doi.org/10.1016/j.arth.2020.05.017>.
- [66] Adalberth G, Nilsson KG, Byström S, Kolstad K, Milbrink J. Low-conforming all-polyethylene tibial component not inferior to metal-backed component in cemented total knee arthroplasty: prospective, randomized radio-stereometric analysis study of the AGC total knee prosthesis. *J Arthroplasty* 2000;15:783–92. <https://doi.org/10.1054/arth.2000.8101>.
- [67] Dorr LD, Boiaro RA. Technical considerations in total knee arthroplasty. *Clin Orthop Relat Res* 1986;5–11.
- [68] Weightman B, Isherwood DP, Swanson SA. The fracture of ultrahigh molecular weight polyethylene in the human body. *J Biomed Mater Res* 1979;13:669–72. <https://doi.org/10.1002/jbm.820130412>.
- [69] Ryd L, Lindstrand A, Stenstrom A, Selvik G. Cold flow reduced by metal backing. An in vivo roentgen stereophotogrammetric analysis of unicompartmental tibial components. *Acta Orthop Scand* 1990;61:21–5. <https://doi.org/10.3109/17453679008993058>.
- [70] Bargren JH, Day WH, Freeman MA, Swanson SA. Mechanical tests on the tibial components of non-hinged knee prostheses. *J Bone Joint Surg Br* 1978;60-B:256–61. <https://doi.org/10.1302/0301-620X.60B2.659476>.
- [71] Bartel DL, Bicknell VL, Wright TM. The effect of conformity, thickness, and material on stresses in ultra-high molecular weight components for total joint replacement. *J Bone Joint Surg Am* 1986;68:1041–51. <https://doi.org/10.2106/00004623-198668070-00010>.
- [72] Goodfellow J, O'Connor J. The mechanics of the knee and prosthesis design. *J Bone Joint Surg Br* 1978;60-B:358–69. <https://doi.org/10.1302/0301-620X.60B3.581081>.
- [73] Ryd L, Albrektsson BE, Herberts P, Lindstrand A, Selvik G. Micromotion of noncemented Freeman-Samuelson knee prostheses in gonarthrosis. A roentgen-stereophotogrammetric analysis of eight successful cases. *Clin Orthop Relat Res* 1988;229:205–12. <https://doi.org/10.1097/00003086-198804000-00028>.
- [74] Ryd L, Lindstrand A, Rosenquist R, Selvik G. Micromotion of conventionally cemented all-polyethylene tibial components in total knee replacements. A roentgen stereophotogrammetric analysis of migration and inducible displacement. *Arch Orthop Trauma Surg* 1978;106:82–8. <https://doi.org/10.1007/BF00435419>.
- [75] Adalberth G, Nilsson KG, Byström S, Kolstad K, Milbrink J. All-polyethylene versus metal-backed and stemmed tibial components in cemented total

- knee arthroplasty. A prospective, randomised RSA study. *J Bone Joint Surg Br* 2001;83:825–31. <https://doi.org/10.1302/0301-620x.83b6.11430>.
- [76] Bartel DL, Burstein AH, Santavica EA, Insall JN. Performance of the tibial component in total knee replacement. *J Bone Joint Surg Am* 1982;64:1026–33.
- [77] Lewis JL, Askew M, Jaycox D. A comparative evaluation of tibial component designs of total knee prostheses. *JBJS* 1982;64:129–35.
- [78] Reilly D, Walker P, Ben-Dov M, Ewald F. Effects of tibial components on load transfer in the upper tibia. *Clin Orthop Relat Res* 1982;165:273–82.
- [79] Emerson Jr RH, Higgins LL, Head WC. The AGC total knee prosthesis at average 11 years. *J Arthroplasty* 2000;15:418–23. <https://doi.org/10.1054/arth.2000.4337>.
- [80] Parks NL, Engh GA, Topoleski LD, Emperado J. The Coventry Award. Modular tibial insert micromotion. A concern with contemporary knee implants. *Clin Orthop Relat Res* 1998;356:10–5. <https://doi.org/10.1097/00003086-199811000-00003>.
- [81] Parks NL, Engh GA, Topoleski LT, Emperado J. Modular tibial insert micromotion: a concern with contemporary knee implants. *Clin Orthop Relat Res* 1998;356:10–5.
- [82] Wasielewski RC, Parks N, Williams I, Surprenant H, Collier JP, Engh G. Tibial insert undersurface as a contributing source of polyethylene wear debris. *Clin Orthop Relat Res* 1997;345:53–9.
- [83] Davis PF, Bocell JR Jr, Tullos HS. Dissociation of the tibial component in total knee replacements. *Clin Orthop Relat Res* 1991;272:199–204.
- [84] Stern S, Insall J. Posterior stabilized prosthesis. Results after follow-up of nine to twelve years. *JBJS* 1992;74:980–6.
- [85] Ritter MA, Berend ME, Meding JB, Keating EM, Faris PM, Crites BM. Long-term followup of anatomic graduated components posterior cruciate-retaining total knee replacement. *Clin Orthop Relat Res* 2001;388:51–7. <https://doi.org/10.1097/00003086-200107000-00009>.
- [86] Herschmiller T, Bradley KE, Wellman SS, Attarian DE. Early to midterm clinical and radiographic survivorship of the all-polyethylene versus modular metal-backed tibia component in primary total knee replacement. *J Surg Orthop Adv* 2019;28:108–14.
- [87] Rao AR, Engh GA, Collier MB, Lounici S. Tibial interface wear in retrieved total knee components and correlations with modular insert motion. *JBJS* 2002;84:1849–55. <https://doi.org/10.2106/00004623-200210000-00017>.
- [88] de Andrade MA, Dos Santos JR, Gonzaga LG, Silva GM. A functional and roentgenographic preliminary comparative study using metal-backed and all-polyethylene tibial components in total knee arthroplasty. *Rev Bras Ortop* 2010;45:295–301. [https://doi.org/10.1016/S2255-4971\(15\)30372-4](https://doi.org/10.1016/S2255-4971(15)30372-4).
- [89] Robinson RP. The early innovators of today's resurfacing condylar knees. *J Arthroplasty* 2005;20(1 Suppl 1):2–26. <https://doi.org/10.1016/j.arth.2004.11.002>.
- [90] Ritter MA, Keating EM, Faris PM, Brugo G. Metal-backed acetabular cups in total hip arthroplasty. *J Bone Joint Surg Am* 1990;72:672–7.
- [91] Brihault J, Navacchia A, Pianigiani S, Labey L, De Corte R, Pascale V, et al. All-polyethylene tibial components generate higher stress and micromotions than metal-backed tibial components in total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc* 2016;24:2550–9. <https://doi.org/10.1007/s00167-015-3630-8>.
- [92] Taylor M, Tanner K, Freeman M. Finite element analysis of the implanted proximal tibia: a relationship between the initial cancellous bone stresses and implant migration. *J Biomech* 1998;31:303–10. [https://doi.org/10.1016/S0021-9290\(98\)00022-0](https://doi.org/10.1016/S0021-9290(98)00022-0).
- [93] Karrholm J, Jonsson H, Nilsson KG, Soderqvist I. Kinematics of successful knee prostheses during weight-bearing: three-dimensional movements and positions of screw axes in the Tricon-M and Miller-Galante designs. *Knee Surg Sports Traumatol Arthrosc* 1994;2:50–9. <https://doi.org/10.1007/BF01552655>.
- [94] Nilsson KG, Karrholm J, Gadegaard P. Abnormal kinematics of the artificial knee. Roentgen stereophotogrammetric analysis of 10 Miller-Galante and five New Jersey LCS knees. *Acta Orthop Scand* 1991;62:440–6. <https://doi.org/10.3109/17453679108996640>.
- [95] Engh GA, Dwyer KA, Hanes CK. Polyethylene wear of metal-backed tibial components in total and unicompartmental knee prostheses. *J Bone Joint Surg Br* 1992;74:9–17. <https://doi.org/10.1302/0301-620X.74B1.1732274>.
- [96] Feng EL, Stulberg SD, Wixson RL. Progressive subluxation and polyethylene wear in total knee replacements with flat articular surfaces. *Clin Orthop Relat Res* 1994;299:60–71 (1976–2007).
- [97] Stiehl JB, Komistek RD, Dennis DA. Detrimental kinematics of a flat on flat total condylar knee arthroplasty. *Clin Orthop Relat Res* 1999;365:139–48. <https://doi.org/10.1097/00003086-199908000-00019>.
- [98] Insall JN, Hood RW, Flawn LB, Sullivan DJ. The total condylar knee prosthesis in gonarthrosis. A five to nine-year follow-up of the first one hundred consecutive replacements. *J Bone Joint Surg Am* 1983;65:619–28. <https://doi.org/10.2106/00004623-198365050-00006>.
- [99] Naylor PT, Myrvik QN, Gristina A. Antibiotic resistance of biomaterial-adherent coagulase-negative and coagulase-positive staphylococci. *Clin Orthop Relat Res* 1990;261:126–33.
- [100] Mihalko WM, Manaswi A, Cui Q, Parvizi J, Schmalzried TP, Saleh KJ. Diagnosis and treatment of the infected primary total knee arthroplasty. *Instr Course Lect* 2008;57:327–39.