



Contents lists available at ScienceDirect

The Journal of Arthroplasty

journal homepage: www.arthroplastyjournal.org

World Expert Meeting in Arthroplasty 2024

Do Functional Outcomes Differ Among Total Knee Arthroplasty Approaches at six, 12, and Beyond 18 Months of Follow-Up?



Seyed Mohammad Javad Mortazavi, MD ^{a,*}, Ali Soltani Farsani ^a, George Babis, MD ^b, Julio Cesar Palacio, MD ^c, David Mateu-Vicent, MD ^d, Joao Mauricio Barretto, MD ^e, Mohammad Razi, MD ^f, Parag Sancheti, FRCS (Ed), MS (Orth), PhD (Dundee, UK) ^g, Mohammad Saeed, MD ^h, Eleftherios Tsiridis, MD ⁱ, Seyed Hadi Kalantar, MD ^j

^a Joint Reconstruction Research Center, Tehran University of Medical Sciences, Tehran, Iran^b National & Kapodistrian University of Athens Medical School, Athens, Greece^c Ortopedia y Traumatología, Cirugía de Cadera y Rodilla, Imbanaco Medical Center, Cali, Colombia^d Universitat Autònoma de Barcelona, Barcelona, Spain^e Orthopedic Surgery Department, Universidade do Estado do Rio de Janeiro (UERJ), Rio de Janeiro, Brazil^f Iran University of Medical Sciences, Tehran, Iran^g Sancheti Institute For Orthopaedics & Rehabilitation & PG College, Senate Member- Maharashtra University Health Sciences (MUHS), Nashik, India^h Medical City Complex, Baghdad, Iraqⁱ President European Hip Society, President Hellenic Association of Orthopaedics & Trauma, PGH Hospital, Aristotle University Medical School, Thessaloniki, Greece^j Department of Orthopedic Surgery, Imam Khomeini Hospital Complex, Tehran University of Medical Sciences, Tehran, Iran

ARTICLE INFO

Article history:

Received 20 September 2024

Received in revised form

8 October 2024

Accepted 11 October 2024

Available online 23 November 2024

Keywords:

total knee arthroplasty

surgical approaches

functional outcomes

midterm outcome

parapatellar

subvastus

Do Functional Outcomes Differ Among Total Knee Arthroplasty Approaches at six, 12, and Beyond 18 Months of follow-up?

Response/Recommendation: Current literature indicates no clinically relevant or statistical differences in early functional outcomes (within 6 months post-surgery) when comparing various surgical approaches to the medial parapatellar approach. However, in more extended follow-up (beyond 18 months), both the quadriceps-sparing and mid-vastus approaches showed clinically relevant and statistically significant improvements in functional outcomes compared to the medial parapatellar approach, with the mid-vastus approach exhibiting superior results at beyond 18 months of follow-up.

Level of Evidence: high**Expert Vote:** Agree 57.2%, Disagree 32.8%, and Abstain 9.8%.

Rationale

Various total knee arthroplasty (TKA) approaches, including the medial parapatellar (MP), mini-MP, sub-vastus (SV), mini-SV, mid-vastus (MV), mini-MV, and quadriceps-sparing (QS), exhibit

One or more of the authors of this paper have disclosed potential or pertinent conflicts of interest, which may include receipt of payment, either direct or indirect, institutional support, or association with an entity in the biomedical field which may be perceived to have potential conflict of interest with this work. For full disclosure statements refer to <https://doi.org/10.1016/j.arth.2024.10.046>.

* Address correspondence to: Seyed Mohammad Javad Mortazavi, MD, Joint Reconstruction Research Center, Tehran University of Medical Sciences, End of Keshavarz Blvd, Tehran 1419733141, Iran.

differences in incision configuration and intraoperative factors such as blood loss and tourniquet time. Due to similarities in incision techniques, the mini-SV approach was grouped with the QS approach [1]. Likewise, the mini-MV and MV approaches were combined due to their similar techniques and the limited number of studies reporting outcomes specifically for the MV approach. The MP approach was selected as the reference group, as it represents the traditional and most commonly used technique in most randomized controlled trials (RCTs) [2,3]. This methodology allowed for a comprehensive comparison of four different approaches (SV, MV, QS, and mini-MP) against the MP approach.

<https://doi.org/10.1016/j.arth.2024.10.046>

0883-5403/© 2024 Elsevier Inc. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

Table 1
Comparing Different TKA Approaches With Medial Parapatellar Approach.

PROMs	6 months	1 year	>1.5 years
KSS knee	No differences	QS > MP (MD = 3.94, CI: 95% [0.68 to 7.20], $P = 0.01$)	No differences
KSS function	No differences	MV > MP (MD = 3.60, CI: 95% [1.67 to 5.54], $P = 0.00$) QS > MP (MD = 2.32, CI: 95% [0.61 to 4.02], $P = 0.00$)	MV > MP (MD = 8.28, CI: 95% [3.37 to 13.19], $P = 0.00$) QS > MP (MD = 5.13, CI: 95% [1.88 to 8.38], $P = 0.00$) mini-MP > MP (MD = 6.21, CI: 95% [4.42 to 8.01], $P = 0.00$) MV > MP (MD = 5.46, CI: 95% [3.08 to 7.83], $P = 0.00$) QS > MP (MD = 4.78, CI: 95% [2.71 to 6.85], $P = 0.00$)
ROM	No differences	No differences	No differences
OKS	No differences	MV > MP (MD = 4.40, CI to 95% [0.61; 8.19], $P = 0.02$)	No differences
WOMAC	No differences	No differences	Not enough data

KSS, Knee Society Score; OKS, Oxford Knee Score; ROM, range of motion; WOMAC, Western Ontario and McMaster Universities Osteoarthritis Index; MP, medial parapatellar; MD, mean differences; CI, confidence interval; SV, sub-vastus; MV, mid-vastus; QS, quadriceps-sparing.

The minimum clinically important difference was set at 5 to 6 for the Knee Society Score (KSS), 4 to 5 for the Oxford Knee Score (OKS), and more than five degrees for the range of motion (ROM) [4,5]. Previous studies were often limited by their focus on endpoint scores without accounting for baseline variations, potentially obscuring treatment effects. In this analysis, we addressed these limitations by comparing the improvement (delta) in functional outcomes between approaches at the endpoint, providing a more accurate assessment of treatment effects.

A comprehensive network meta-analysis (NMA) was conducted, encompassing all RCTs from 2000 onwards that reported patient-reported outcomes (PROMs) at least six months after surgery. The analysis included 51 RCTs [1–3],[6–40],[41–53] that reported on KSS, Western Ontario and McMaster Universities Osteoarthritis Index, OKS, and ROM. The findings indicate that there were no significant variations in PROMs between different TKA approaches at the 6-month mark compared to the MP approach (Table 1). However, at the 1-year follow-up, both the QS and MV approaches showed superior PROMs compared to the MP approach. The QS approach led to greater improvement in the KSS knee, while both QS and MV improved regarding KSS function. However, these differences did not exceed the minimum clinically important difference of approximately 5 points. The MV approach also demonstrated meaningful clinical improvement in the OKS compared to MP. In the final analysis of data beyond 18 months, the QS, MV, and mini-MP approaches showed superior outcomes over MP. Both QS and MV achieved significantly and clinically better KSS function results. Also, mini-MP, MV, and QS provided greater ROM enhancement, with clinically meaningful differences observed for mini-MP and MV.

An evaluation of a recent 5-year meta-analysis of RCTs was performed. Yang et al. reported no significant differences in the PROMs at 6-month and 1-year follow-ups between the MP and MV approaches; however, the PROMs were superior in the MV group at the early follow-up (three months) [54]. Bouché et al., in their NMA, found that all approaches exhibited comparable PROMs during early follow-up (six months), except for ROM, where the SV approach demonstrated superior results [55]. They further reported no significant differences between approaches at one and two years postoperatively for PROMs, except for KSS, where the mini-MP approach showed superior outcomes [55]. An NMA conducted by Zhang et al. also found no significant differences in KSS among different approaches during the short-term follow-up [56]. Additionally, a meta-analysis by Yuan et al. indicated that the QS approach led to a more significant improvement in KSS compared to the MP approach at mean follow-up of two years [57]. Furthermore, Berstock et al., in their meta-analysis evaluating KSS at six weeks and 12 months of follow-up, found no significant differences between the MP, mini-SV, and mini-SV approaches [58]. However, statistically significant advantages were observed only in the mini-MP group for functional KSS compared to the conventional MP approach.

A possible explanation for the similarity in early outcomes and the divergence in extended follow-ups may be attributable to the initial postoperative recovery phase. During this period, improvements in pain relief and joint function are generally consistent across different surgical techniques due to factors such as pain management, soft tissue healing, and standardized rehabilitation protocols. In contrast, extended follow-ups highlight the advantages of less invasive approaches, such as QS, MV, and mini-MP techniques, which demonstrate superior functional outcomes, particularly in KSS function and ROM. The prolonged period permits the manifestation of the mid-term benefits of these approaches, as they may better preserve muscle function, minimize soft tissue trauma, and promote more natural joint mechanics, leading to enhanced functional recovery over time.

CRediT authorship contribution statement

Seyed Mohammad Javad Mortazavi: Writing – review & editing, Writing – original draft, Project administration, Investigation. **Ali Soltani Farsani:** Writing – original draft, Formal analysis, Data curation, Conceptualization. **George Babis:** Writing – review & editing, Supervision. **Julio Cesar Palacio:** Validation, Supervision. **David Mateu-Vicent:** Writing – review & editing, Validation, Supervision. **Joao Mauricio Barretto:** Writing – review & editing, Supervision. **Mohammad Razi:** Supervision. **Parag Sancheti:** Writing – review & editing, Validation, Supervision. **Mohammad Saeed:** Investigation. **Eleftherios Tsiridis:** Validation, Supervision. **Seyed Hadi Kalantar:** Writing – original draft, Validation, Investigation.

References

- [1] Mehta N, Bhat MS, Goyal A, Mishra P, Joshi D, Chaudhary D. Quadriceps sparing (subvastus/midvastus) approach versus the conventional medial parapatellar approach in primary knee arthroplasty. *J Arthrosc Joint Surg* 2017;4:15–20. <https://doi.org/10.1016/j.jajs.2017.02.004>.
- [2] Cho KY, Kim KI, Umran S, Kim SH. Better quadriceps recovery after minimally invasive total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc* 2014;22:1759–64. <https://doi.org/10.1007/s00167-013-2556-2>.
- [3] Zhu M, Ang CL, Yeo SJ, Lo NN, Chia SL, Chong HC. Minimally invasive computer-assisted total knee arthroplasty compared with conventional total knee arthroplasty: a prospective 9-year follow-up. *J Arthroplasty* 2016;31:1000–4. <https://doi.org/10.1016/j.arth.2015.11.023>.
- [4] Lee WC, Kwan YH, Chong HC, Yeo SJ. The minimal clinically important difference for knee society clinical rating system after total knee arthroplasty for primary osteoarthritis. *Knee Surg Sports Traumatol Arthrosc* 2017;25:3354–9. <https://doi.org/10.1007/s00167-016-4208-9>.
- [5] Murray DW, Fitzpatrick R, Rogers K, Pandit H, Beard DJ, Carr AJ, et al. The use of the oxford hip and knee scores. *J Bone Joint Surg Br* 2007;89:1010–4. <https://doi.org/10.1302/0301-620x.89b8.19424>.
- [6] Aglietti P, Baldini A, Sensi L. Quadriceps-sparing versus mini-subvastus approach in total knee arthroplasty. *Clin Orthop Relat Res* 2006;452:106–11. <https://doi.org/10.1097/01.blo.0000238789.51972.16>.
- [7] Kelly MJ, Rumi MN, Kothari M, Parentis MA, Bailey KJ, Parrish WM, et al. Comparison of the vastus-splitting and median parapatellar approaches for primary total knee arthroplasty: a prospective, randomized study. *J Bone Jt Surg Am Vol* 2006;88A:715–20. <https://doi.org/10.2106/JBJS.E.00107>.

- [8] Kim YH, Kim JS, Kim DY. Clinical outcome and rate of complications after primary total knee replacement performed with quadriceps-sparing or standard arthroscopy. *J Bone J Surg Br* 2007;89:467–70. <https://doi.org/10.1302/0301-620X.89B4.18663>.
- [9] Seon JK, Song EK, Yoon TR, Park SJ, Bae BH, Cho SG. Comparison of functional results with navigation-assisted minimally invasive and conventional techniques in bilateral total knee arthroplasty. *Comput Aided Surg* 2007;12: 189–93. <https://doi.org/10.3109/10929080701311861>.
- [10] Tashiro Y, Miura H, Matsuda S, Okazaki K, Iwamoto Y. Minimally invasive versus standard approach in total knee arthroplasty. *Clin Orthop Relat Res* 2007;463:144–50.
- [11] Karachalios T, Giotikas D, Roidis N, Poultides L, Bargiotas K, Malizos KN. Total knee replacement performed with either a mini-midvastus or a standard approach - a prospective randomised clinical and radiological trial. *J Bone Joint Surg Br* 2008;90B:584–91. <https://doi.org/10.1302/0301-620X.90B5.20122>.
- [12] Kashyap SN, van Ommeren JW. Clinical experience with less invasive surgery techniques in total knee arthroplasty: a comparative study. *Knee Surg Sports Traumatol Arthrosc* 2008;16:544–8. <https://doi.org/10.1007/s00167-008-0523-0>.
- [13] Bridgman SA, Walley G, MacKenzie G, Clement D, Griffiths D, Maffulli N. Subvastus approach is more effective than a medial parapatellar approach in primary total knee arthroplasty: a randomized controlled trial. *Knee* 2009;16: 216–22. <https://doi.org/10.1016/j.knee.2008.11.012>.
- [14] Jung YB, Lee YS, Lee EY, Jung HJ, Nam CH. Comparison of the modified subvastus and medial parapatellar approaches in total knee arthroplasty. *Int Orthop* 2009;33:419–23. <https://doi.org/10.1007/s00264-007-0510-y>.
- [15] Karpman RR, Smith HL. Comparison of the early results of minimally invasive vs standard approaches to total knee arthroplasty: a prospective, randomized study. *J Arthroplasty* 2009;24:681–8. <https://doi.org/10.1016/j.arth.2008.03.011>.
- [16] Watanabe T, Muneta T, Ishizuki M. Is a minimally invasive approach superior to a conventional approach for total knee arthroplasty? Early outcome and 2- to 4-year follow-up. *J Orthop Sci* 2009;14:589–95. <https://doi.org/10.1007/s00776-009-1383-2>.
- [17] Bonutti PM, Zywiell MG, Ulrich SD, Stroh DA, Seyler TM, Mont MA. A comparison of subvastus and midvastus approaches in minimally invasive total knee arthroplasty. *J Bone Joint Surg* 2010;92:575–82. <https://doi.org/10.2106/JBJS.I.00268>.
- [18] Hay GC, Kampshoff J, Kuster MS. Lateral subvastus approach with osteotomy of the tibial tubercle for total knee replacement a two-year prospective, randomised, blinded controlled trial. *J Bone Joint Surg Br* 2010;92B:862–6. <https://doi.org/10.1302/0301-620X.92B6.24027>.
- [19] Hernandez-Vaquero D, Noriega-Fernandez A, Suarez-Vazquez A. Total knee arthroplasties performed with a mini-incision or a standard incision. Similar results at six months follow-up. *BMC Musculoskelet Disord* 2010;11:27. <https://doi.org/10.1186/1471-2474-11-27>.
- [20] Pan WM, Li XG, Tang TS, Qian ZL, Zhang Q, Zhang CM. Mini-subvastus versus a standard approach in total knee arthroplasty: a prospective, randomized, controlled study. *J Int Med Res* 2010;38:890–900. <https://doi.org/10.1177/147323001003800315>.
- [21] Varela-Egocheaga JR, Suárez-Suárez MA, Fernández-Villán M, González-Sastre V, Varela-Gómez JR, Rodríguez-Merchán C. Minimally invasive subvastus approach: improving the results of total knee arthroplasty: a prospective, randomized trial. *Clin Orthop Relat Res* 2010;468:1200–8. <https://doi.org/10.1007/s11999-009-1160-8>.
- [22] Dutka J, Skowronek M, Sosin P, Skowronek P. Subvastus and medial parapatellar approaches in tka: comparison of functional results. *Orthopedics* 2011;34:6. <https://doi.org/10.3928/01477447-20110427-05>.
- [23] Kim JG, Lee SW, Ha JK, Choi HJ, Yang SJ, Lee MY. The effectiveness of minimally invasive total knee arthroplasty to preserve quadriceps strength: a randomized controlled trial. *Knee* 2011;18:443–7. <https://doi.org/10.1016/j.knee.2010.08.008>.
- [24] Lee DH, Choi J, Nha KW, Kim HJ, Han SB. No difference in early functional outcomes for mini-midvastus and limited medial parapatellar approaches in navigation-assisted total knee arthroplasty: a prospective randomized clinical trial. *Knee Surg Sports Traumatol Arthrosc* 2011;19:66–73. <https://doi.org/10.1007/s00167-010-1130-4>.
- [25] Bourke MG, Jull GA, Buttrum PJ, FitzPatrick PL, Dalton PA, Russell TG. Comparing outcomes of medial parapatellar and subvastus approaches in total knee arthroplasty. A randomized controlled trial. *J Arthroplasty* 2012;27: 347–353.e1. <https://doi.org/10.1016/j.arth.2011.06.005>.
- [26] Chiang H, Lee CC, Lin WP, Jiang CC. Comparison of quadriceps-sparing minimally invasive and medial parapatellar total knee arthroplasty: a 2-year follow-up study. *J Formos Med Assoc* 2012;111:698–704. <https://doi.org/10.1016/j.jfma.2011.11.025>.
- [27] Essving P, Axelsson K, Otterborg L, Spännar H, Gupta A, Magnuson A, et al. Minimally invasive surgery did not improve outcome compared to conventional surgery following unicompartmental knee arthroplasty using local infiltration analgesia: a randomized controlled trial with 40 patients. *Acta Orthop* 2012;83:634–41. <https://doi.org/10.3109/17453674.2012.736169>.
- [28] Guy SP, Farndon MA, Conroy JL, Bennett C, Grainger AJ, London NJ. A prospective randomised study of minimally invasive midvastus total knee arthroplasty compared with standard total knee arthroplasty. *Knee* 2012;19: 866–71. <https://doi.org/10.1016/j.knee.2012.04.009>.
- [29] Masjudin T, Kamari ZH. A comparison between subvastus and midvastus approaches for staged bilateral total knee arthroplasty: a prospective, randomised study. *Malaysian Orthop J* 2012;6:31–6.
- [30] Avci CC, Güllabi D, Erdem M, Kurnaz R, Güneş T, Bostan B. Minimal invasive midvastus versus standard parapatellar approach in total knee arthroplasty. *Acta Orthop Traumatol Turc* 2013;47:1–7. <https://doi.org/10.3944/aott.2013.2801>.
- [31] Lin SY, Chen CH, Fu YC, Huang PJ, Lu CC, Su JY, et al. Comparison of the clinical and radiological outcomes of three minimally invasive techniques for total knee replacement at two years. *Bone Joint J* 2013;95-b:906–10. <https://doi.org/10.1302/0301-620X.95b7.29694>.
- [32] Pongcharoen B, Yakampor T, Charoencholvani K. Patellar tracking and anterior knee pain are similar after medial parapatellar and midvastus approaches in minimally invasive tka knee. *Clin Orthop Relat Res* 2013;471: 1654–60. <https://doi.org/10.1007/s11999-012-2778-5>.
- [33] Thienpont E. Faster recovery after minimally invasive surgery in total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc* 2013;21:2412–7. <https://doi.org/10.1007/s00167-012-1978-6>.
- [34] Zhang Z, Zhu W, Gu B, Zhu L, Chen C. Mini-midvastus versus mini-medial parapatellar approach in total knee arthroplasty: a prospective, randomized study. *Arch Orthop Trauma Surg* 2013;133:389–95. <https://doi.org/10.1007/s00402-012-1645-x>.
- [35] Heekin RD, Fokin AA. Mini-midvastus versus mini-medial parapatellar approach for minimally invasive total knee arthroplasty: outcomes pendulum is at equilibrium. *J Arthroplasty* 2014;29:339–42. <https://doi.org/10.1016/j.arth.2013.05.016>.
- [36] Nutton RW, Wade FA, Coutts FJ, Van Der Linden ML. Short term recovery of function following total knee arthroplasty: a randomised study of the medial parapatellar and midvastus approaches. *Arthritis* 2014;2014:1737857. <https://doi.org/10.1155/2014/173857>.
- [37] Tasker A, Hassaballa M, Murray J, Lancaster S, Artz N, Harries W, et al. Minimally invasive total knee arthroplasty: a pragmatic randomised controlled trial reporting outcomes up to 2-year follow up. *Knee* 2014;21:189–93. <https://doi.org/10.1016/j.knee.2013.07.010>.
- [38] Zhou C, Peng W, Si ZP. Effect of total knee arthroplasty via different approaches on extensor mechanism. *Chinese J Tissue Eng Res* 2014;18:1337–42. <https://doi.org/10.3969/j.issn.2095-4344.2014.09.005>.
- [39] Huang AB, Wang HJ, Yu JK, Yang B, Ma D, Zhang JY. Are there any clinical and radiographic differences between quadriceps-sparing and mini-medial parapatellar approaches in total knee arthroplasty after a minimum 5 years of follow-up? *Chin Med J (Engl)* 2015;128:1898–904. <https://doi.org/10.4103/0366-6999.160521>.
- [40] Liu H, Mei X, Zhang Z, Sun J. Mini-midvastus versus mini-medial parapatellar approach in simultaneous bilateral total knee arthroplasty with 24-month follow-up. *Acta Orthop Traumatol Turcica* 2015;49:586–92. <https://doi.org/10.3944/AOTT.2015.15.0078>.
- [41] Feczko P, Engelmann L, Arts JJ, Campbell D. Computer-assisted total knee arthroplasty using mini midvastus or medial parapatellar approach technique: a prospective, randomized, international multicentre trial. *BMC Musculoskelet Disord* 2016;17:19. <https://doi.org/10.1186/s12891-016-0872-7>.
- [42] Huang AB, Wang HJ, Yu JK, Yang B, Ma D, Zhang JY. Optimal patellar alignment with minimally invasive approaches in total knee arthroplasty after a minimum five year follow-up. *Int Orthop* 2016;40:487–92. <https://doi.org/10.1007/s00264-015-2896-2>.
- [43] Koh IJ, Kim MW, Kim MS, Jang SW, Park DC, In Y. The patient's perception does not differ following subvastus and medial parapatellar approaches in total knee arthroplasty: a simultaneous bilateral randomized study. *J Arthroplasty* 2016;31:112–7. <https://doi.org/10.1016/j.arth.2015.08.004>.
- [44] Verburg H, Mathijssen NMC, Niesten DD, Verhaar JAN, Pilot P. Comparison of mini-midvastus and conventional total knee arthroplasty with clinical and radiographic evaluation: a prospective randomized clinical trial with 5-year follow-up. *J Bone Joint Surg Am* 2016;98:1014–22. <https://doi.org/10.2106/JBJS.15.00654>.
- [45] Aslam MA, Sabir AB, Tiwari V, Abbas S, Tiwari A, Singh P. Approach to total knee replacement: a randomized double blind study between medial parapatellar and midvastus approach in the early postoperative period in asian population. *J Knee Surg* 2017;30:793–7. <https://doi.org/10.1055/s-0036-1597978>.
- [46] Shukla R, Mahajan P, Singh M, Jain RK, Kumar R. Outcome of total knee replacement via two approaches in indian scenario. *J Knee Surg* 2017;30: 174–8. <https://doi.org/10.1055/s-0036-1584192>.
- [47] Unwin O, Hassaballa M, Murray J, Harries W, Porteous A. Minimally invasive surgery (mis) for total knee replacement; medium term results with minimum five year follow-up. *Knee* 2017;24:454–9. <https://doi.org/10.1016/j.knee.2017.01.010>.
- [48] Li Z, Cheng W, Sun L, Yao Y, Cao Q, Ye S, et al. Mini-subvastus versus medial parapatellar approach for total knee arthroplasty: a prospective randomized controlled study. *Int Orthop* 2018;42:543–9. <https://doi.org/10.1007/s00264-017-3703-z>.
- [49] Jhurani A, Agarwal P, Aswal M, Rasquinha C, Srivastava M. Subvastus exposure compared to parapatellar approach in navigated sequential bilateral total knee arthroplasty (sbtk): a prospective randomized study. *J Knee Surg* 2021;34:635–43. <https://doi.org/10.1055/s-0039-1700496>.
- [50] Madadi F, Osias E, Washington III ER, Shamie AN, Yazdanasheh H. Comparison of effects of medial parapatellar and subvastus bilateral simultaneous

- approaches in total knee arthroplasty. *J Arthrosc Joint Surg* 2021;8:350–3. <https://doi.org/10.1016/j.jaj.2021.06.001>.
- [51] Sogbein OA, Zomar BO, Bryant DM, Howard JL, Marsh JD, Lanting BA. Effects of surgical approach and tourniquet use on patient-reported outcomes following total knee arthroplasty: a pilot randomized clinical trial. *Orthop Res Rev* 2022;14:407–17. <https://doi.org/10.2147/ORR.S381894>.
- [52] Phruetthiphat OA, Mokmongkolkul K, Apinyankul R. Total knee arthroplasty with lateral parapatellar approach results in less early postoperative skin numbness than medial approach, but no difference at mid-term follow-up: a randomized control trial. *Arthroplast Today* 2024;27:101365. <https://doi.org/10.1016/j.artd.2024.101365>.
- [53] Berth A, Urbach D, Neumann W, Awiszus F. Strength and voluntary activation of quadriceps femoris muscle in total knee arthroplasty with midvastus and subvastus approaches. *J Arthroplasty* 2007;22:83–8. <https://doi.org/10.1016/j.arth.2006.02.161>.
- [54] Yang X, Cheng QH, Yang YZ, Zhang AR, Fan H, Guo HZ. Minimally invasive medial femoral approach to total knee arthroplasty improves short-term outcomes compared to the standard medial parapatellar approach: a systematic review and meta-analysis. *J Orthop Surg Res* 2023;18:657. <https://doi.org/10.1186/s13018-023-04136-2>.
- [55] Bouché PA, Corsia S, Nizard R, Resche-Rigon M. Comparative efficacy of the different surgical approaches in total knee arthroplasty: a systematic-review and network meta-analysis. *J Arthroplasty* 2021;36:1187–1189.e1. <https://doi.org/10.1016/j.arth.2020.09.052>.
- [56] Zhang L, Li X, Rüwald JM, Welle K, Schildberg FA, Kabir K. Comparison of minimally invasive approaches and standard median parapatellar approach for total knee arthroplasty: a systematic review and network meta-analysis of randomized controlled trials. *Technol Health Care* 2021;29:557–74. <https://doi.org/10.3233/thc-192078>.
- [57] Yuan FZ, Zhang JY, Jiang D, Yu JK. Quadriceps-sparing versus traditional medial parapatellar approaches for total knee arthroplasty: a meta-analysis. *BMC Musculoskelet Disord* 2019;20:117. <https://doi.org/10.1186/s12891-019-2482-7>.
- [58] Berstock JR, Murray JR, Whitehouse MR, Blom AW, Beswick AD. Medial subvastus versus the medial parapatellar approach for total knee replacement: a systematic review and meta-analysis of randomized controlled trials. *EFORT Open Rev* 2018;3:78–84. <https://doi.org/10.1302/2058-5241.3.170030>.