



World Expert Meeting in Arthroplasty 2024

Should Routine Radiographic Screening for Spino-pelvic Relationship Be Performed in Patients Undergoing Primary Total Hip Arthroplasty?



Seyed Mohammad Javad Mortazavi, MD^{*}, Mohammad Poursalehian, MD, Marcus Crestani, MD, Alberto Di Martino, MD, Valentin Antoci, MD, Valeriy Murylev, MD, Alberto Delgado-Martinez, MD, Mahdi Sahebi

Joint Reconstruction Research Center, Tehran University of Medical Sciences, Tehran, Iran

ARTICLE INFO

Article history:

Received 20 September 2024

Received in revised form

14 October 2024

Accepted 15 October 2024

Available online 21 October 2024

Keywords:

spinopelvic relationship
primary total hip arthroplasty
radiographic screening
revision
dislocation

Should routine radiographic screening for spino-pelvic relationships be performed in patients undergoing primary total hip arthroplasty?

Response/Recommendation: There is no need for routine radiographic screening for the spino-pelvic relationship, and this should be reserved for patients who have spinal disease and risk factors for instability.

Level of Evidence: Low.

Expert Vote: Agree 82.0%, Disagree 15.2%, and Abstain 2.8%

Rationale

The spino-pelvic relationship plays a critical role in the biomechanics and functional outcomes of patients undergoing primary total hip arthroplasty (THA). Accurate preoperative assessment of this dynamic relationship is crucial to optimize implant positioning and reduce complications such as dislocation, impingement, aseptic loosening, and revision surgery. While some surgeons advocate for routine radiographic screening of the spino-pelvic relationship in all patient candidates for THA, others argue that it should be reserved for select cases only. The existing literature was reviewed to identify factors associated with impaired spino-pelvic mobility.

There are multiple studies that reported impaired spino-pelvic motion in patients who have surgical or nonsurgical spinal fusion

[1–3]. The impaired spino-pelvic mobility in these patients could cause hip instability after THA [3]. Therefore, spino-pelvic assessment in this subset of patients is recommended.

Older age is associated with impaired spino-pelvic mobility [4–7]. Innmann et al. proposed a cutoff point of 65 years of age suggesting the performance of a radiological screening for spino-pelvic mobility [6].

Patients who have spinal deformity, including scoliosis, flat back, and hyperlordosis, are at increased risk of spino-pelvic mobility impairment [8–10]. Patients who have clinical spinal deformity are recommended to undergo further assessments. A study by Innmann et al. stated that patients who have lumbar lordosis lower than 45 degrees are at higher risk of spino-pelvic mobility impairment [6]. Further cutoff points for flat back and scoliosis are yet to be determined in the literature. Vigdorich et al. reported that patients who have severe sagittal spinal deformity (lumbar lordosis minus pelvic incidence mismatch of greater than 20 degrees) are at increased risk of spino-pelvic mobility impairment [11].

Moreover, patients who have lumbar degenerative disk disorders are reported to have impaired spino-pelvic mobility [12–15].

No author associated with this paper has disclosed any potential or pertinent conflicts which may be perceived to have impending conflict with this work. For full disclosure statements refer to <https://doi.org/10.1016/j.arth.2024.10.066>.

^{*} Address correspondence to: Seyed Mohammad Javad Mortazavi, MD, Joint Reconstruction Research Center, Tehran University of Medical Sciences, Tehran 1419733141, Iran.

However, the cost-effectiveness of lumbar degenerative disk disorder evaluation in patients undergoing THA has not been studied in literature.

Patients who have limited hip range of motion are reported to have impaired spino-pelvic mobility [16,17]. A study by Innmann et al. reported a cutoff point of 88 degrees for hip flexion for screening of spino-pelvic assessment [16].

Contralateral hip degenerative joint disease has been reported to have effects on spino-pelvic mobility [8,17,18]. Moreover, contralateral THA may have an effect on spino-pelvic mobility [18]. Therefore, spino-pelvic mobility evaluation in bilateral hip osteoarthritis might be necessary.

Multiple studies have reported that increased pelvic tilt on standing X-rays is associated with impaired spino-pelvic mobility [11,16,19]. Multiple cutoff points of 10, 13, and 19 degrees have been reported in the literature [11,16,19]. Patients who have increased pelvic tilt should undergo a thoughtful spino-pelvic evaluation; however, the exact cutoff point is yet to be determined. Pelvic tilt on lateral radiographs is correlated with pubic symphysis to sacro-coccygeal junction distance on supine antero-posterior (AP) radiographs [20]. Carender et al. reported that patients who have overlap of the sacro-coccygeal junction/pubis symphysis in AP radiographs have 10 times greater risk of impaired spino-pelvic mobility [21]. Rainer et al. reported that patients who have an overlap of the sacro-coccygeal junction/pubis symphysis in AP radiographs have nine times greater risk of dislocation [20].

Sacral slope on standing radiographs is correlated with spino-pelvic mobility [7,8,16]. Innmann et al. reported that patients who have sacral slopes greater than 42 degrees have spino-pelvic hypermobility [16]. Therefore, patients who have impaired standing sacral slopes should undergo further spino-pelvic evaluation.

In summary, the above-mentioned factors can be categorized into two groups.

1. Factors assessable through physical examination and radiographic studies routinely performed for primary THA:
 - a. Spinal fusion (both surgical and nonsurgical) [1–3,21].
 - b. Older age ≥ 65 years old [4–7].
 - c. Clinically evident spinal column deformity (flat back, hyperlordosis, and scoliosis) [8–10].
 - d. Limited range of motion of the hip (flexion $< 88^\circ$) [16,17].
 - e. Contralateral hip degenerative joint disease or contralateral THA [8,17,18].
 - f. Lumbar degenerative disc disease [12–15].
 - g. Overlap of the sacro-coccygeal junction/pubis symphysis in AP radiograph [20,21].
2. Factors that could be assessed on standing lateral lumbo-sacral radiographs:
 - a. Pelvic tilt $> 19^\circ$, 13° , 10° [11,16,19].
 - b. Sacral slope $> 42^\circ$ [7,8,16].
 - c. Pelvic incidence-lumbar lordosis PI-LL mismatch $> 20^\circ$ [11].
 - d. Lumbar lordosis $< 45^\circ$ [6].

No clear screening method for spino-pelvic mobility is defined in the literature. The outlined factors above represent a collation of available studies, as none of the articles evaluate all the predictors in combination or their efficacy in determining outcomes. Additionally, the cost-effectiveness of using an additional standing lateral pelvic X-ray for screening purposes has yet to be determined.

References

- [1] Steff M, Lundergan W, Heckmann N, McKnight B, Ike H, Murgai R, et al. Spinopelvic mobility and acetabular component position for total hip arthroplasty. *Bone Joint J* 2017;99-B(1 Suppl A):37–45. <https://doi.org/10.1302/0301-620x.99b1.Bjj-2016-0415.R1>.
- [2] Salib CG, Reina N, Perry KI, Taunton MJ, Berry DJ, Abdel MP. Lumbar fusion involving the sacrum increases dislocation risk in primary total hip arthroplasty. *Bone Joint J* 2019;101-B:198–206. <https://doi.org/10.1302/0301-620x.101b2.Bjj-2018-0754.R1>.
- [3] Yang DS, McDonald CL, DiSilvestro KJ, Patel SA, Li NY, Cohen EM, et al. Risk of dislocation and revision following primary total hip arthroplasty in patients with prior lumbar fusion with spinopelvic fixation. *J Arthroplasty* 2023;38:700–705.e1. <https://doi.org/10.1016/j.arth.2022.03.061>.
- [4] Heckmann N, McKnight B, Steff M, Trasolini NA, Ike H, Dorr LD. Late dislocation following total hip arthroplasty: spinopelvic imbalance as a causative factor. *J Bone Joint Surg Am* 2018;100:1845–53. <https://doi.org/10.2106/jbjs.18.00078>.
- [5] Haffer H, Hu Z, Wang Z, Müllner M, Hardt S, Pumberger M. Association of age and spinopelvic function in patients receiving a total hip arthroplasty. *Sci Rep* 2023;13:2589. <https://doi.org/10.1038/s41598-023-29545-5>.
- [6] Innmann M, Verhaegen J, Renkawitz T, Merle C, Grammatopoulos G. How to screen for lumbar spine stiffness in patients awaiting total hip arthroplasty. *J Arthroplasty* 2024;39:124–31. <https://doi.org/10.1016/j.arth.2023.08.006>.
- [7] Qoreishy M, Movahedinia M, Keyhani S, Salehi B, Goodarzi M, Naghibi B. Evaluation of changes in radiographic spinopelvic parameters after primary total hip arthroplasty and its relationship with short-term patient function. *Eur J Orthop Surg Traumatol* 2024;34:225–30. <https://doi.org/10.1007/s00590-023-03639-2>.
- [8] Homma Y, Ishii S, Yanagisawa N, Ochi H, Baba T, Nojiri H, et al. Pelvic mobility before and after total hip arthroplasty. *Int Orthop* 2020;44:2267–74. <https://doi.org/10.1007/s00264-020-04688-6>.
- [9] Buckland AJ, Fernandez L, Shimmmin AJ, Bare JV, McMahon SJ, Vigdorchik JM. Effects of sagittal spinal alignment on postural pelvic mobility in total hip arthroplasty candidates. *J Arthroplasty* 2019;34:2663–8. <https://doi.org/10.1016/j.arth.2019.06.036>.
- [10] Kleeman-Försthuber L, Vigdorchik JM, Pierrepont JW, Dennis DA. Pelvic incidence significance relative to spinopelvic risk factors for total hip arthroplasty instability. *Bone Joint J* 2022;104-b:352–8. <https://doi.org/10.1302/0301-620x.104b3.Bjj-2021-0894.R1>.
- [11] Vigdorchik JM, Sharma AK, Madurawe CS, Pierrepont JW, Dennis DA, Shimmmin AJ. Prevalence of risk factors for adverse spinopelvic mobility among patients undergoing total hip arthroplasty. *J Arthroplasty* 2021;36:2371–8. <https://doi.org/10.1016/j.arth.2020.12.029>.
- [12] Berliner JL, Esposito CI, Miller TT, Padgett DE, Mayman DJ, Jerabek SA. What preoperative factors predict postoperative sitting pelvic position one year following total hip arthroplasty? *Bone Joint J* 2018;100-b:1289–96. <https://doi.org/10.1302/0301-620x.100b10.Bjj-2017-1336.R2>.
- [13] Limmahakhun S, Box HN, Arazu P, Hennessy DW, Klemm C, Kwon YM. In vivo analysis of spinopelvic kinematics and peak head-cup contact in total hip arthroplasty patients with lumbar degenerative disc disease. *J Orthop Res* 2019;37:674–80. <https://doi.org/10.1002/jor.24234>.
- [14] Klemm C, Limmahakhun S, Bounajem G, Xiong L, Yeo I, Kwon YM. Effect of postural changes on in vivo pelvic tilt and functional component anteversion in total hip arthroplasty patients with lumbar disc degenerations. *Bone Joint J* 2020;102-b:1505–10. <https://doi.org/10.1302/0301-620x.102b11.Bjj-2020-0777.R1>.
- [15] Vigdorchik JM, Sharma AK, Dennis DA, Walter LR, Pierrepont JW, Shimmmin AJ. The majority of total hip arthroplasty patients with a stiff spine do not have an instrumented fusion. *J Arthroplasty* 2020;35:S252–4. <https://doi.org/10.1016/j.arth.2020.01.031>.
- [16] Innmann MM, Merle C, Gotterbarm T, Ewerbeck V, Beaulé PE, Grammatopoulos G. Can spinopelvic mobility be predicted in patients awaiting total hip arthroplasty? A prospective, diagnostic study of patients with end-stage hip osteoarthritis. *Bone Joint J* 2019;101-b:902–9. <https://doi.org/10.1302/0301-620x.101b8.Bjj-2019-0106.R1>.
- [17] Ouchida J, Nakashima H, Kanemura T, Satake K, Ando K, Ito K, et al. Impact of the hip joint mobility on whole-body sagittal alignment: prospective analysis in case with hip arthroplasty. *Eur Spine J* 2022;31:2399–407. <https://doi.org/10.1007/s00586-022-07251-6>.
- [18] Muellner M, Becker L, Wang Z, Hu Z, Hardt S, Pumberger M, et al. Spinopelvic mobility is influenced by pre-existing contralateral hip arthroplasty: a matched-pair analysis in patients undergoing hip replacement. *J Orthop Surg Res* 2022;17:64. <https://doi.org/10.1186/s13018-022-02945-5>.
- [19] Innmann MM, Merle C, Phan P, Beaulé PE, Grammatopoulos G. How can patients with mobile hips and stiff lumbar spines be identified prior to total hip arthroplasty? A prospective, diagnostic cohort study. *J Arthroplasty* 2020;35:S255–61. <https://doi.org/10.1016/j.arth.2020.02.029>.
- [20] Rainer WG, Abdel MP, Freedman BA, Berry DJ, Taunton MJ. Pelvic tilt and the pubic symphysis to sacro-coccygeal junction distance: risk factors for hip dislocation observed on anteroposterior pelvis radiographs. *J Arthroplasty* 2021;36:S367–73. <https://doi.org/10.1016/j.arth.2021.02.079>.
- [21] Carender CN, Feuchtenberger BW, DeMik DE, An Q, Brown TS, Bedard NA. Can abnormal spinopelvic relationships be identified by anteroposterior pelvic radiographs? *J Arthroplasty* 2022;37:507–12. <https://doi.org/10.1016/j.arth.2021.11.032>.