



## World Expert Meeting in Arthroplasty 2024

## What Are the Indications for Hinged Implants in Revision Total Knee Arthroplasty?



Ashraf T. Hantouly, MD, MSc <sup>a,\*</sup>, Sathish Muthu, MSc, PhD <sup>b, c, d</sup>,  
 Mahmood Shahab, MD <sup>e</sup>, Maritin Sarungi, MD, PhD <sup>f</sup>, Aasis Unnanuntana, MD, MSc <sup>g</sup>,  
 Brian de Beaubien, MD <sup>h</sup>, Jacobus D. Jordaan, MD <sup>i</sup>, Thorsten Gehrke, MD <sup>j</sup>,  
 Javad Parvizi, MD <sup>k</sup>, Mustafa Citak, MD, MBA <sup>j</sup>

<sup>a</sup> Aspetar Orthopaedic and Sports Medicine Hospital, Doha, Qatar<sup>b</sup> Department of Orthopaedics, Government Medical College, Karur, Tamil Nadu, India<sup>c</sup> Faculty of Engineering, Department of Biotechnology, Karpagam Academy of Higher Education, Coimbatore, Tamil Nadu, India<sup>d</sup> Orthopedic Research Group, Coimbatore, Tamil Nadu, India<sup>e</sup> Orthopedic Surgery Department, Medical City Complex, Baghdad, Iraq<sup>f</sup> Department of Orthopaedics, Golden Jubilee National Hospital, Glasgow, Scotland, United Kingdom<sup>g</sup> Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok, Thailand<sup>h</sup> Department of Orthopedics, Covenant Medical Center, Saginaw, Michigan<sup>i</sup> Faculty of medicine and Health Sciences, Division of orthopaedic Surgery, University of Stellenbosch, South Africa<sup>j</sup> Department of Orthopaedic Surgery, Helios ENDO-Clinic Hamburg, Hamburg, Germany<sup>k</sup> Orthopedic Surgery, Acibadem University, Istanbul, Turkey

## ARTICLE INFO

## Article history:

Received 21 September 2024

Received in revised form

23 October 2024

Accepted 24 October 2024

Available online 5 November 2024

## Keywords:

arthroplasty

knee

revision

hinged

total knee arthroplasty

## What are the indications for a hinged implants in revision total knee arthroplasty?

**Response/Recommendation:** The three most commonly reported indications for using hinged-design implants in revision total knee arthroplasty (rTKA) are infection, instability, and aseptic loosening. However, these conditions rarely present in isolation and are often accompanied by additional factors such as pain, stiffness/arthrofibrosis, periprosthetic fracture, dislocation/subluxation, malalignment/malposition, mechanical failure, bone loss, patellar complications, and the need for revision of a previously implanted hinged prosthesis. Therefore, hinged implants should be considered in cases where major bone loss or compromised soft tissue and ligamentous integrity renders semiconstrained devices prone to failure.

## Strength of Recommendation: Moderate.

Delegate vote: agree: 69%, disagree: 21%, abstain: 10%

## Rationale

A systematic review was conducted to analyze the indications for hinged knee implants in revision total knee arthroplasty (rTKA).

One or more of the authors of this paper have disclosed potential or pertinent conflicts of interest, which may include receipt of payment, either direct or indirect, institutional support, or association with an entity in the biomedical field which may be perceived to have potential conflict of interest with this work. For full disclosure statements refer to <https://doi.org/10.1016/j.arth.2024.10.126>.

\* Address correspondence to: Ashraf T. Hantouly, MD, MSc, Department of Orthopaedic Surgery, Helios ENDO-Clinic Hamburg, Holstenstraße, Hamburg 222767, Germany.

<https://doi.org/10.1016/j.arth.2024.10.126>

0883-5403/© 2024 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

PubMed, Web of Science, and Scopus were searched from inception till March 27, 2024, for original articles on hinged implant indications in rTKA. Exclusion criteria were non-English language, case reports, sample size < 10, review articles, registry-based studies, studies with oncology cases, distal femur replacements, technique articles, nonhuman studies, and studies not reporting implant use indications. The Preferred Reporting Items for Systematic Review and Meta-Analysis guidelines were strictly followed [1]. Of the 2,349 articles identified, 58 met the inclusion criteria [2–59], representing 2,803 revision TKAs. Most studies were retrospective, providing level IV evidence. Table 1 summarizes the indications for hinged designs in rTKA. Table 1 shows the

**Table 1**  
Indications of Hinged Implant Designs in rTKA.

Indication	n	%
Infection	781	27.86
Instability	671	23.94
Aseptic loosening	665	23.72
Stiffness/Arthrofibrosis	228	8.13
Multiple reasons	128	4.57
Periprosthetic fracture	110	3.92
Dislocation/Subluxation	41	1.46
Malalignment/Malposition	40	1.43
Mechanical Failure	32	1.14
Bone loss	27	0.96
Patellar complications	24	0.86
Pain	17	0.61
Others	17	0.61
Revision of hinged as only indication	15	0.54
Not specified	7	0.25

rTKA, revision total knee arthroplasty.

list of indications with their share for the use of hinged designs in rTKA.

Hinged implants are increasingly used in complex rTKA cases involving substantial instability and bone loss, which compromise the performance and longevity of standard implant designs [60]. The three most commonly reported indications for considering hinged-implants in rTKA are infection, instability, and aseptic loosening. Although rTKA for infection could be done in a single-stage or two-stage revision, hinged design can be considered to compensate for the lack of stability after extensive debridement of bone and soft tissue. Apart from infection, instability—whether from direct ligamentous insufficiency or as a result of revision procedures that cannot be managed by less-constrained implants—is the next common indication, followed by aseptic loosening. From a clinical perspective, the above-mentioned indications are not usually as distinctive; rather, they present in a combined fashion, thereby making a case for the use of hinged knee designs. These high-risk scenarios are associated with an increased risk of reoperations and revisions. The common reasons for reoperation and revision following rTKA with hinged knee prosthesis are listed below in Table 2. The overall reported reoperation and revision rate in the included studies was 28.1%.

With the increase in modularity and next-generation rotating designs, hinged implants are now becoming an essential component in the arsenal of not only rTKA, but also in complex primary scenarios.

**Table 2**  
The Common Reasons for Reoperation and Revision Following rTKA With hinged Knee Prosthesis.

Revision/Reoperation Causes	n	%
Infection	188	34.4
Aseptic Loosening	85	15.5
Patellar Complications	52	9.5
Wound complication	48	8.8
Arthrofibrosis	40	7.3
Implant Failure	24	4.4
Periprosthetic Fracture	50	9.1
Chronic Pain	12	2.2
Dislocation	15	2.7
Extensor Mechanism Failure	14	2.6
Instability	8	1.5
Polyethylene wear	7	1.3
Others	4	0.7
Total	547	100.00

rTKA, revision total knee arthroplasty.

Like all systematic reviews, this study had potential limitations. Significant heterogeneity and selection bias were present, and only English-language articles were included. Additionally, the inconsistency in reporting outcomes impeded a comprehensive quantitative analysis, and some studies lacked detailed patient demographics and comorbidity data, key factors influencing outcomes. Additionally, variations in the follow-up durations across studies may have affected long-term assessment of outcomes and mortality rates. Moreover, the diversity of implant designs and indications likely influenced the observed outcomes, reoperation/revision rates, and complication rates. Also, inconsistent differentiation between revisions and reoperations prevented definitive conclusions on these rates.

To conclude, in rTKA, the main reasons reported in the literature for considering hinged-design implants are infection, instability, and aseptic loosening. These indications, however, are rarely encountered in isolation. More often, they are combined or accompanied by additional factors such as pain, stiffness/arthrofibrosis, periprosthetic fracture, dislocation/subluxation, malalignment/malposition, mechanical failure, bone loss, patellar issues, and the need to revise an existing hinged prosthesis. Therefore, hinged implants should be considered in cases where major bone loss or compromised soft tissue and ligamentous integrity renders semiconstrained devices prone to failure.

## CRediT authorship contribution statement

**Ashraf T. Hantouly:** Writing – review & editing, Writing – original draft, Supervision, Methodology, Investigation, Data curation, Conceptualization. **Sathish Muthu:** Writing – review & editing, Writing – original draft, Methodology, Formal analysis, Conceptualization. **Mahmood Shahab:** Writing – review & editing, Writing – original draft, Supervision, Conceptualization. **Maritin Sarungi:** Writing – review & editing, Supervision, Conceptualization. **Aasis Unnanuntana:** Writing – review & editing, Conceptualization. **Brian Debeaubien:** Writing – review & editing, Writing – original draft, Supervision, Conceptualization. **Jacobus D. Jordaan:** Writing – review & editing, Writing – original draft, Supervision, Methodology, Conceptualization. **Thorsten Gehrke:** Writing – review & editing, Writing – original draft, Supervision, Methodology, Conceptualization. **Javad Parvizi:** Writing – review & editing, Writing – original draft, Supervision, Methodology, Conceptualization. **Mustafa Citak:** Writing – review & editing, Writing – original draft, Supervision, Methodology, Conceptualization.

## References

- [1] Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. PLoS Med 2009;6:e1000097. <https://doi.org/10.1371/journal.pmed.1000097>.
- [2] Minamoto STN, Mozella AP, Cossich VRA, Gavilão UF, Machado HS, Barreto JM. Comparative functional and isokinetic analysis between implants with posterior stabilization and rotating hinge total knee arthroplasty. Rev Bras Ortop (Sao Paulo) 2024;59:e68–75. <https://doi.org/10.1055/s-0044-1779685>.
- [3] Dzidzishvili L, Sáez D, Calvo E. Metaphyseal cones combined with diaphyseal impaction grafting provide good outcomes and survival in a complex revision total knee arthroplasty: a matched comparative analysis. Eur J Orthop Surg Traumatol 2024;34:577–83. <https://doi.org/10.1007/s00590-023-03659-y>.
- [4] Schlechter M, Theil C, Gosheger G, Moellenbeck B, Schwarze J, Puetzler J, et al. Good mid-term implant survival of a novel single-design rotating-hinge total knee arthroplasty. J Clin Med 2023;12:6113. <https://doi.org/10.3390/jcm12196113>.
- [5] Koch KA, Spranz DM, Westhauser F, Bruckner T, Lehner B, Alvand A, et al. Impact of comorbidities and previous surgery on mid-term results of revision total knee arthroplasty for periprosthetic joint infection. J Clin Med 2023;12:5542. <https://doi.org/10.3390/jcm12175542>.
- [6] Hecker A, Pütz HA, Wangler S, Eberlein SC, Klenke FM. Indications, clinical outcome and survival of rotating hinge total knee arthroplasty in a

- retrospective study of 63 primary and revision cases. *Eur J Orthop Surg Traumatol* 2023;33:1885–94. <https://doi.org/10.1007/s00590-022-03349-1>.
- [7] Van Laarhoven SN, Te Molder MEM, Van Hellemont GG, Heesterbeek PJC. Acceptable migration of a fully cemented rotating hinge-type knee revision system measured in 20 patients with model-based RSA with a 2-year follow-up. *Acta Orthop* 2023;94:185–90. <https://doi.org/10.2340/17453674.2023.12305>.
- [8] Matar HE, Bloch BV, James PJ. Satisfactory medium- to long-term outcomes of cemented rotating hinge prosthesis in revision total knee arthroplasty. *Bone Jt Open* 2023;4:776–81. <https://doi.org/10.1302/2633-1462.410BJO-2022-0032.R1>.
- [9] Miralles-Muñoz FA, Pineda-Salazar M, Rubio-Morales M, González-Parrón S, Ruiz-Lozano M, Lizarra-Utrilla A. Similar outcomes of constrained condylar knee and rotating hinge prosthesis in revision surgery for extension instability after primary total knee arthroplasty. *Orthop Traumatol Surg Res* 2022;108:103265. <https://doi.org/10.1016/j.otsr.2022.103265>.
- [10] Özdemir M, Yaradılmış YU, Ateş A, Evren AT, Okkaoglu MC, Altay M. Hinged prosthesis in the knee revision surgery: is there a great need? *Ann Clin Anal Med* 2021;13:1061–5. <https://doi.org/10.4328/ACAM.20990>.
- [11] Sanz-Ruiz P, León-Román VE, Matas-Díez JA, Villanueva-Martínez M, Vaquero J. Long-term outcomes of one single-design varus valgus constrained versus one single-design rotating hinge in revision knee arthroplasty after over 10-year follow-up. *J Orthop Surg Res* 2022;17:135. <https://doi.org/10.1186/s13018-022-03026-3>.
- [12] Neumann DR, Hofstaedter T, Dorn U. Follow-up of a modular rotating hinge knee system in salvage revision total knee arthroplasty. *J Arthroplasty* 2012;27:814–9. <https://doi.org/10.1016/j.arth.2011.08.015>.
- [13] Schnetz M, Hofmann L, Ewald L, Klug A, Hoffmann R, Gramlich Y. Midterm results of modular hinge total knee arthroplasty using cementless osseointegrating stems: low fixation associated complications and good functional outcome in primary and revision knee arthroplasty. *Arch Orthop Trauma Surg* 2024;144:831–45. <https://doi.org/10.1007/s00402-023-05148-7>.
- [14] Vertesich K, Staats K, Böhler C, Koza R, Lass R, Giurea A. Long term results of a rotating hinge total knee prosthesis with carbon-fiber reinforced poly-ether-ether-ketone (CFR-PEEK) as bearing material. *Front Bioeng Biotechnol* 2022;4:845859. <https://doi.org/10.3389/fbioe.2022.845859>.
- [15] Rajgopal A, Agrawal U. Long term results of rotating hinge total knee arthroplasty in complex primary and revision cases. *Acta Orthop Belg* 2020;86:114–23.
- [16] Memori N, Iqbal F, Noor SS, Najjad KR, Sozera MF, Abro A, et al. Mid-term results and survival rates following a single-design rotating hinge knee arthroplasty in non-tumor conditions in a Pakistani population. *Knee Surg Relat Res* 2021;33:15. <https://doi.org/10.1186/s43019-021-00102-6>.
- [17] Bourbotte-Salmon F, Ferry T, Cardinale M, Servien E, Rongieras F, Fessy MH, et al., Lyon Bone and Joint Infections Study Group. Rotating hinge knee arthroplasty for revision prosthetic-knee infection: good functional outcomes but a crucial need for superinfection prevention. *Front Surg* 2021;8:551814. <https://doi.org/10.3389/fsurg.2021.551814>.
- [18] Wignadasan W, Chang JS, Kayani B, Kontoghiorghe C, Haddad FS. Long-term results of revision total knee arthroplasty using a rotating hinge implant. *Knee* 2021;28:72–80. <https://doi.org/10.1016/j.knee.2020.11.009>.
- [19] Bingham JS, Bukowski BR, Wyles CC, Pareek A, Berry DJ, Abdel MP. Rotating-Hinge revision total knee arthroplasty for treatment of severe arthrofibrosis. *J Arthroplasty* 2019;34(7S):S271–6. <https://doi.org/10.1016/j.arth.2019.01.072>.
- [20] Hintze J, Niemeläinen M, Sintonen H, Nieminen J, Eskelinen A. Outcomes of the rotating hinge knee in revision total knee arthroplasty with a median follow-up of 6.2 years. *BMC Musculoskelet Disord* 2021;22:336. <https://doi.org/10.1186/s12891-021-04205-9>.
- [21] Arnholdt J, Boelch SP, Dogan F, Hoberg M, Holzapfel BM, Rudert M. Revision arthroplasty with rotating hinge systems for total knee arthroplasty instability. *Oper Orthop Traumatol* 2020;32:298–308. <https://doi.org/10.1007/s00064-020-00663-x>. English.
- [22] Brown LR, Clement ND, MacDonald DJ, Breusch SJ. The survivorship of the link endo-rotational hinge total knee arthroplasty: 5–12-year follow-up of 100 patients. *Arch Orthop Trauma Surg* 2019;139:107–12. <https://doi.org/10.1007/s00402-018-3064-0>.
- [23] van Rensch PJH, Heesterbeek PJC, Hannink G, van Hellemont GG, Wymenga AB. Improved clinical outcomes after revision arthroplasty with a hinged implant for severely stiff total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc* 2019;27:1043–8. <https://doi.org/10.1007/s00167-018-5235-5>.
- [24] Lim JBT, Pang HN, Tay KJD, Chia SL, Lo NN, Yeo SJ. Increased constraint of rotating hinge knee prosthesis is associated with poorer clinical outcomes as compared to constrained condylar knee prosthesis in total knee arthroplasty. *Eur J Orthop Surg Traumatol* 2020;30:529–35. <https://doi.org/10.1007/s00590-019-02598-x>.
- [25] Böhler C, Kolbitsch P, Schuh R, Lass R, Kubista B, Giurea A. Midterm results of a new rotating hinge knee implant: a 5-year follow-up. *BioMed Res Int* 2017;2017:7532745. <https://doi.org/10.1155/2017/7532745>.
- [26] Abdelaziz H, Biewald P, Anastasiadis Z, Haasper C, Gehrke T, Hawi N, et al. Midterm results after tantalum cones in 1-stage knee exchange for periprosthetic joint infection: a single-center study. *J Arthroplasty* 2020;35:1084–9. <https://doi.org/10.1016/j.arth.2019.11.016>.
- [27] Kearns SM, Culp BM, Bohl DD, Sporer SM, Della Valle CJ, Levine BR. Rotating hinge implants for complex primary and revision total knee arthroplasty. *J Arthroplasty* 2018;33:766–70. <https://doi.org/10.1016/j.arth.2017.10.009>.
- [28] Chen MJ, Hung JF, Chang CH, Lee SH, Shih HN, Chang YH. Periprosthetic knee infection reconstruction with a hinged prosthesis: implant survival and risk factors for treatment failure. *Knee* 2020;27:1035–42. <https://doi.org/10.1016/j.knee.2020.03.004>.
- [29] Rouquette L, Batailler C, Muller B, Neyret P, Servien E, Lustig S. Early complications and causes of revision after rotating-hinge TKA. *Arch Orthop Trauma Surg* 2020;140:109–19. <https://doi.org/10.1007/s00402-019-03290-9>.
- [30] Hermans K, Vandenneucker H, Truijen J, Oosterbosch J, Bellermans J. Hinged versus CCK revision arthroplasty for the stiff total knee. *Knee* 2019;26:222–7. <https://doi.org/10.1016/j.knee.2018.10.012>.
- [31] van Laarhoven SN, van Eerd AHJ, van Hellemont GG, Schreurs BW, Wymenga AB, Heesterbeek PJC. Superior survival of fully cemented fixation compared to hybrid fixation in a single design rotating hinge knee implant. *J Arthroplasty* 2022;37:482–7. <https://doi.org/10.1016/j.arth.2021.11.037>.
- [32] León-Román VE, García-Mato D, López-Torres II, Vaquero-Martín J, Calvo-Haro JA, Pascau J, et al. Is a greater degree of constraint really harmful? Clinical biomechanical comparative study between condylar constrained knee and rotating hinge prosthesis. *Clin Biomech* 2024;111:106149. <https://doi.org/10.1016/j.clinbiomech.2023.106149>.
- [33] Schneider AM, Rice SJ, Lancaster N, McGraw M, Farid Y, Finn HA. Low-dose irradiation and rotating-hinge revision for the treatment of severe idiopathic arthrofibrosis following total knee arthroplasty: a review of 60 patients with a mean 6-year follow-up. *J Arthroplasty* 2024;39:1075–82. <https://doi.org/10.1016/j.arth.2023.10.021>.
- [34] Pradhan NR, Bale L, Kay P, Porter ML. Salvage revision total knee replacement using the Endo-Model rotating hinge prosthesis. *Knee* 2004;11:469–73. <https://doi.org/10.1016/j.knee.2004.03.001>.
- [35] Hernández-Vaquero D, Sandoval-García MA. Hinged total knee arthroplasty in the presence of ligamentous deficiency. *Clin Orthop Relat Res* 2010;468:1248–53. <https://doi.org/10.1007/s11999-009-1226-7>.
- [36] Gudnason A, Milbrink J, Hailer NP. Implant survival and outcome after rotating-hinge total knee revision arthroplasty: a minimum 6-year follow-up. *Arch Orthop Trauma Surg* 2011;131:1601–7. <https://doi.org/10.1007/s00402-011-1330-5>.
- [37] Jones RE, Skedros JG, Chan AJ, Beauchamp DH, Harkins PC. Total knee arthroplasty using the S-ROM mobile-bearing hinge prosthesis. *J Arthroplasty* 2001;16:279–87. <https://doi.org/10.1054/arth.2001.21498>.
- [38] Ochs BG, Schreiner AJ, de Zwart PM, Stöckle U, Gonser CE. Computer-assisted navigation is beneficial both in primary and revision surgery with modular rotating-hinge knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc* 2016;24:64–73. <https://doi.org/10.1007/s00167-014-3316-7>.
- [39] Shen C, Lichstein PM, Austin MS, Sharkey PF, Parvizi J. Revision knee arthroplasty for bone loss: choosing the right degree of constraint. *J Arthroplasty* 2014;29:127–31. <https://doi.org/10.1016/j.arth.2013.04.042>.
- [40] Baier C, Lüring C, Schaumburger J, Köck F, Beckmann J, Tingart M, et al. Assessing patient-oriented results after revision total knee arthroplasty. *J Orthop Sci* 2013 Nov;18:955–61. <https://doi.org/10.1007/s00776-013-0467-1>.
- [41] Utting MR, Newman JH. Customised hinged knee replacements as a salvage procedure for failed total knee arthroplasty. *Knee* 2004;11:475–9. <https://doi.org/10.1016/j.knee.2003.12.007>.
- [42] Back DL, David L, Hilton A, Blunn G, Briggs TW, Cannon SR. The SMILES prosthesis in salvage revision knee surgery. *Knee* 2008;15:40–4. <https://doi.org/10.1016/j.knee.2007.09.002>.
- [43] Neumann DR, Hofstaedter T, Dorn U. Follow-up of a modular rotating hinge knee system in salvage revision total knee arthroplasty. *J Arthroplasty* 2012;27:814–9. <https://doi.org/10.1016/j.arth.2011.08.015>.
- [44] Farid YR, Thakral R, Finn HA. Intermediate-term results of 142 single-design, rotating-hinge implants: frequent complications may not preclude salvage of severely affected knees. *J Arthroplasty* 2015;30:2173–80. <https://doi.org/10.1016/j.arth.2015.06.033>.
- [45] Barrack RL, Lyons TR, Ingraham RQ, Johnson JC. The use of a modular rotating hinge component in salvage revision total knee arthroplasty. *J Arthroplasty* 2000;15:858–66. <https://doi.org/10.1054/arth.2000.9056>.
- [46] Zahar A, Kendoff DO, Klatte TO, Gehrke TA. Can good infection control be obtained in one-stage exchange of the infected TKA to a rotating hinge design? 10-year results. *Clin Orthop Relat Res* 2016;474:81–7. <https://doi.org/10.1007/s11999-015-4408-5>.
- [47] Rodríguez-Merchán EC, Gómez-Cardero P, Martínez-Lloreda Á. Revision knee arthroplasty with a rotating-hinge design in elderly patients with instability following total knee arthroplasty. *J Clin Orthop Trauma* 2015;6:19–23. <https://doi.org/10.1016/j.jcot.2014.11.001>.

- [48] Singer J, Merz A, Frommelt L, Fink B. High rate of infection control with one-stage revision of septic knee prostheses excluding MRSA and MRSE. *Clin Orthop Relat Res* 2012;470:1461–71. <https://doi.org/10.1007/s11999-011-2174-6>.
- [49] Inglis AE, Walker PS. Revision of failed knee replacements using fixed-axis hinges. *J Bone Joint Surg Br* 1991;73:757–61. <https://doi.org/10.1302/0301-620X.73B5.1894661>.
- [50] Efe T, Roessler PP, Heyse TJ, Hauk C, Pahrmann C, Getgood A, et al. Mid-term results after implantation of rotating-hinge knee prostheses: primary versus revision. *Orthop Rev* 2012;4:e35. <https://doi.org/10.4081/or.2012.e35>.
- [51] Bistolfi A, Massazza G, Rosso F, Crova M. Rotating-hinge total knee for revision total knee arthroplasty. *Orthopedics* 2012;35:e325–30. <https://doi.org/10.3928/01477447-20120222-34>.
- [52] Streitbuerger A, Hardes J, Gosheger G, Dieckmann R, Hoell S. Knee salvage in revision arthroplasty after massive bone loss of the femur condyles ( $\geq$ Engh III) with a single-modular-hinged knee revision implant. *Arch Orthop Trauma Surg* 2016;136:1077–83. <https://doi.org/10.1007/s00402-016-2491-z>.
- [53] Joshi N, Navarro-Quilis A. Is there a place for rotating-hinge arthroplasty in knee revision surgery for aseptic loosening? *J Arthroplasty* 2008;23:1204–11. <https://doi.org/10.1016/j.arth.2007.10.016>.
- [54] Lutjeboer JS, Bénard MR, Defoort KC, van Hellemont GG, Wymenga AB. Revision total knee arthroplasty for instability—outcome for different types of instability and implants. *J Arthroplasty* 2016;31:2672–6. <https://doi.org/10.1016/j.arth.2016.06.062>.
- [55] Angelini FJ, Helito CP, Veronesi BA, Guimarães TM, Pécora JR, Demange MK. Knee arthroplasty revision with a constrained implant using hinge and rotating tibial basis. *Acta Ortop Bras* 2016;24:22–6. <https://doi.org/10.1590/1413-785220162401153984>.
- [56] Sanguineti F, Mangano T, Formica M, Franchin F. Total knee arthroplasty with rotating-hinge Endo-Model prosthesis: clinical results in complex primary and revision surgery. *Arch Orthop Trauma Surg* 2014;134:1601–7. <https://doi.org/10.1007/s00402-014-2061-1>.
- [57] Hwang SC, Kong JY, Nam DC, Kim DH, Park HB, Jeong ST, et al. Revision total knee arthroplasty with a cemented posterior stabilized, condylar constrained or fully constrained prosthesis: a minimum 2-year follow-up analysis. *Clin Orthop Surg* 2010;2:112–20. <https://doi.org/10.4055/cios.2010.2.2.112>.
- [58] Steens W, Loehr JF, Wodtke J, Katzer A. Morselized bone grafting in revision arthroplasty of the knee: a retrospective analysis of 34 reconstructions after 2–9 years. *Acta Orthop* 2008;79:683–8. <https://doi.org/10.1080/17453670810016713>.
- [59] Fuchs S, Sandmann C, Gerdemann G, Skwara A, Tibesku CO, Bottner F. Quality of life and clinical outcome in salvage revision total knee replacement: hinged vs total condylar design. *Knee Surg Sports Traumatol Arthrosc* 2004;12:140–3. <https://doi.org/10.1007/s00167-003-0401-8>.
- [60] Salari P, Baldini A. Revision knee surgery: the practical approach. *EJORT Open Rev* 2021;6:495–500. <https://doi.org/10.1302/2058-5241.6.210018>.