



Contents lists available at ScienceDirect

The Journal of Arthroplasty

journal homepage: www.arthroplastyjournal.org

World Expert Meeting in Arthroplasty 2024

Is There a Difference Between Modular Versus Monoblock Femoral Stems Used During Revision Total Hip Arthroplasty?



Alisina Shahi, MD, PhD ^{a,*}, Hari Bezwada, MD ^b, Daniel Kendoff, MD ^c,
Fouad Sadek, MD ^d, Margarita Veloso Duran, MD ^e, Heinz Winkler, MD ^f,
Weijun Wang, MD ^g, Xiaogang Zhang, MD ^h

^a Orthopaedic Department, University of Texas Health Science Center at Houston, Houston, Texas^b Hospital of the University of Pennsylvania, Philadelphia, Pennsylvania^c HELIOS Kliniken Berlin-Buch, Berlin, Germany^d Department of Orthopaedics and Traumatology, Cairo University, Cairo, Egypt^e Hospital Universitari Mútua Terrassa, Barcelona, Spain^f Osteitis Center Döbling, Vienna, Austria^g Orthopaedic Department, Nanjing Medical University, Nanjing, China^h Department of Orthopedics, Xinjiang Medical University, Xinjiang, China

ARTICLE INFO

Article history:

Received 19 September 2024

Received in revised form

19 October 2024

Accepted 23 October 2024

Available online 13 November 2024

Keywords:

revision total hip arthroplasty

modular stem

monoblock stem

modular femoral stem

monoblock femoral stem

femoral stem design

Is there a difference between modular versus monoblock femoral stems used during revision total hip arthroplasty?

Response/Recommendation: The literature delineates distinct differences between modular and monoblock femoral stems in revision total hip arthroplasty each presenting unique advantages and limitations. Modular stems offer intraoperative flexibility, enabling precise restoration of leg length and offset. However, they are associated with complications such as junctional fatigue, corrosion, and implant breakage, and they incur much higher costs compared to monoblock stems. In contrast, monoblock stems provide enhanced mechanical stability and no risk of modular junction complications, though their implantation is technically more challenging. Both modular and monoblock femoral stem designs perform well in revision total hip arthroplasty, exhibiting no significant differences in long-term (3 to 5 years) clinical outcomes or subsidence rates. The choice between modular and monoblock stems should be tailored to individual patient needs and specific clinical scenarios.

Level of Evidence: Moderate.**Delegates Vote:** 92.7% Agree, 5.7% Disagree, 1.7% Abstain.

Rationale

A comprehensive literature search was conducted using PubMed, Embase, and Cochrane Library databases. The search terms included “Revision Arthroplasty, Hip,” “Femoral Stem,”

“Modular Femoral Stem,” “Monoblock Femoral Stem,” “Complications,” “Outcomes,” and “Prosthesis Design.” Articles were limited to English-language publications from the past 15 years. Inclusion criteria were studies involving adult patients undergoing revision total hip arthroplasty (THA) with modular or monoblock femoral stems, with outcomes related to stability, mechanical complications, and clinical performance. Exclusion criteria included case reports, nonrevision THA studies, and studies not comparing modular and monoblock stems. Data extracted included study design, population characteristics, outcomes, and complications. A total of 789 articles were identified, with 25 meeting the inclusion criteria after title and abstract screening and full-text review.

One or more of the authors of this paper have disclosed potential or pertinent conflicts of interest, which may include receipt of payment, either direct or indirect, institutional support, or association with an entity in the biomedical field which may be perceived to have potential conflict of interest with this work. For full disclosure statements refer to <https://doi.org/10.1016/j.arth.2024.10.123>.

* Address correspondence to: Alisina Shahi, MD, PhD, Orthopaedic Department, University of Texas Health Science Center at Houston, Houston, TX.

<https://doi.org/10.1016/j.arth.2024.10.123>

0883-5403/© 2024 Elsevier Inc. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

The literature suggests no significant differences between modular and monoblock femoral stems in revision THA. Modular femoral stems offer intraoperative flexibility, allowing surgeons to customize components for optimal fit and alignment. This flexibility is particularly advantageous in complex revisions where restoring leg length, offset, and achieving stable fixation is critical. However, modular stems are associated with specific complications such as junctional fatigue, corrosion, and implant breakage, which can lead to mechanical failure over time. Despite these concerns, modern modular implants have been designed to mitigate these issues, with improved materials and designs reducing the incidence of such complications [1–5].

In contrast, monoblock femoral stems, constructed from a single piece of material, eliminate the risk of modular junction complications. These stems provide increased mechanical stability and durability, which can be particularly beneficial in patients who have poor bone quality or extensive bone loss. However, the lack of intraoperative modularity can limit the ability to make precise adjustments during surgery, and their implantation can be technically more challenging [6–9].

Modular stems are generally more costly than monoblock stems, which is an important consideration in their routine use. The cost implications must be weighed against the potential benefits of modular stems in achieving better biomechanical restoration and patient-specific customization [10,11].

Clinical outcomes between the two types of stems also vary. Studies have shown that modular stems can achieve better restoration of biomechanics and improved functional outcomes in the short term (1 to 3 years). In contrast, monoblock stems have demonstrated lower revision rates and fewer mechanical failures in the long term (3 to 5 years) [12,13]. Additionally, no significant differences were found between modular and monoblock tapered stems regarding postoperative hip function, re-revision rates, and complications [7,9,14].

Overall, the choice between modular and monoblock femoral stems should be individualized based on patient-specific factors, including bone quality, the extent of bone loss, and the need for precise biomechanical restoration. Surgeons must weigh the benefits of intraoperative flexibility against the potential risks of mechanical complications associated with modular stems.

CRediT authorship contribution statement

Alisina Shahi: Writing – original draft, Investigation. **Hari Bezwada:** Investigation. **Daniel Kendoff:** Investigation. **Fouad**

Sadek: Investigation. **Margarita Veloso Duran:** Investigation. **Heinz Winkler:** Investigation. **Weijun Wang:** Investigation. **Xiaogang Zhang:** Investigation.

References

- [1] Restrepo C, Mashadi M, Parvizi J, Austin MS, Hozack WJ. Modular femoral stems for revision total hip arthroplasty. *Clin Orthop Relat Res* 2011;469:476–82. <https://doi.org/10.1007/s11999-010-1561-8>.
- [2] Snijders TE, van Erp JHJ, de Gast A. Restoring femoral offset and leg length; the potential of a short curved stem in total hip arthroplasty. *J Orthop* 2019;16:396–9. <https://doi.org/10.1016/j.jor.2019.04.013>.
- [3] Dou Y, Zhou Y, Tang Q, Yang D, Liu J. Leg-length discrepancy after revision hip arthroplasty: are modular stems superior? *J Arthroplasty* 2013;28:676–9. <https://doi.org/10.1016/j.arth.2012.08.009>.
- [4] Rodriguez JA, Fada R, Murphy SB, Rasquinha VJ, Ranawat CS. Two-year to five-year follow-up of femoral defects in femoral revision treated with the link MP modular stem. *J Arthroplasty* 2009;24:751–8. <https://doi.org/10.1016/j.arth.2008.09.011>.
- [5] Fink B. What can the surgeon do to reduce the risk of junction breakage in modular revision stems? *Arthroplast Today* 2018;4:306–9. <https://doi.org/10.1016/j.artd.2018.03.002>.
- [6] Hozack WJ. Monoblock OR modular? Making the right stem choice. *Orthop Proc* 2016;98-B:54. https://doi.org/10.1302/1358-992X.98BSUPP_22.CCJR2015-054.
- [7] Siljander BR, Chandi SK, Cox FR, Nguyen JT, Sculco PK, Chalmers BP, et al. A consecutive series of vancouver B2 periprosthetic femur fractures treated with contemporary monoblock versus modular revision stems: clinical and radiographic outcomes. *J Arthroplasty* 2024;39:S213. <https://doi.org/10.1016/j.arth.2024.03.046>.
- [8] Wang D, Li H, Zhang W, Li H, Xu C, Liu W, et al. Efficacy and safety of modular versus monoblock stems in revision total hip arthroplasty: a systematic review and meta-analysis. *J Orthop Traumatol* 2023;24:50. <https://doi.org/10.1186/s10195-023-00731-5>.
- [9] Yacovelli S, Ottaway J, Banerjee S, Courtney PM. Modern revision femoral stem designs have No difference in rates of subsidence. *J Arthroplasty* 2021;36:268–73. <https://doi.org/10.1016/j.arth.2020.07.078>.
- [10] Srinivasan A, Jung E, Levine BR. Modularity of the femoral component in total hip arthroplasty. *J Am Acad Orthop Surg* 2012;20:214. <https://doi.org/10.5435/JAAOS-20-04-214>.
- [11] Schwartz AJ, Spanghel MJ, Beauchamp CP. Is modularity necessary when using a tapered stem in revision total hip arthroplasty? *Curr Orthop Pract* 2014;25:143. <https://doi.org/10.1097/BCO.000000000000082>.
- [12] Confalonieri N, Manzotti A, Montironi F, Pullen C. Leg length discrepancy, dislocation rate, and offset in total hip replacement using a short modular stem: navigation vs conventional freehand. *Orthopedics* 2008;31:orthosupersite.com/view.asp?rID=35541.
- [13] Garbuz DS, Toms A, Masri BA, Duncan CP. Improved outcome in femoral revision arthroplasty with tapered fluted modular titanium stems. *Clin Orthop Relat Res* 2006;453:199. <https://doi.org/10.1097/01.blo.0000238875.86519.cf>.
- [14] Pomeroy E, Flynn SO, Grigoras M, Murphy TP, Stavrakis AI, Rowan FE. Subsidence of monoblock and modular titanium fluted tapered stems in revision hip arthroplasty: a retrospective multicentre comparison study. *J Clin Orthop Trauma* 2022;34:102021. <https://doi.org/10.1016/j.jcot.2022.102021>.