



Contents lists available at ScienceDirect

The Journal of Arthroplasty

journal homepage: www.arthroplastyjournal.org

World Expert Meeting in Arthroplasty 2024

Is There a Difference in Outcomes Between Hemiarthroplasty and Total Hip Arthroplasty for Patients Who Have Displaced Femoral Neck Fractures?



Geert Meermans, MD ^a, Darko Talevski, MD, MSc ^b, Pēteris Studers, MD, PhD ^c,
Emre Togrul, MD ^d, Ruslan G. Nazaruddin, MD ^e, Kamolsak Sukhonthamarn, MD ^f,
Mark Phillips, PhD ^g, Seper Ekhtiari, MD, MSc ^{g,*}

^a Bravis Hospital, Bergen op Zoom and Roosendaal, Netherlands^b General City Hospital, Skopje, North Macedonia^c Department of Orthopaedics, Riga Stradiņš University Hospital of Traumatology and Orthopaedics, Riga, Latvia^d Acibadem Adana Hospital, Adana, Turkey^e ALTY Orthopaedic Hospital, Kuala Lumpur, Malaysia^f Department of Orthopaedics, Faculty of Medicine, Khon Kaen University, Khon Kaen, Thailand^g Department of Surgery, McMaster University, Hamilton, Canada

ARTICLE INFO

Article history:

Received 20 September 2024

Received in revised form

16 October 2024

Accepted 17 October 2024

Available online 22 October 2024

Keywords:

femoral neck fracture

hip fracture

arthroplasty

total hip arthroplasty

hemiarthroplasty

bipolar

Is there a difference in outcomes between hemiarthroplasty and total hip arthroplasty for patients who have displaced femoral neck fractures?

Response/Recommendation: The current evidence, with at up to 5 years follow-up, does not show a clinically meaningful difference between hemiarthroplasty and total hip arthroplasty performed for elderly patients who have displaced femoral neck fractures. However, based on expert opinion, the use of total hip arthroplasty may be considered for younger and more active patients.

Voting Results (N = 186): Agree: 93% (N = 173), Disagree: 6% (N = 11), and Abstain: 1% (n = 2).

Level of Evidence: High.

Rationale

Given the preponderance of randomized controlled trial (RCT) studies available, this analysis was limited to published RCTs on the topic, with up to 5 years of follow-up. There were 19 RCTs that were included with a total sample size of 3,414 patients who had a mean

age of 78 years [1–18]. The outcomes of interest included all important clinical outcomes that were reported in a sufficient number of studies to allow for quantitative meta-analysis. These outcomes included mortality, revision surgery, periprosthetic fracture, complications, dislocations, operative time, estimated blood loss, function, and health-related quality of life (HRQoL).

In summary, based on quantitative meta-analyses, there was no significant difference between hemiarthroplasty (HA) and total hip arthroplasty (THA) in terms of mortalities, revision surgeries, periprosthetic fractures, total complications, or dislocations. The most robust evidence exists for revision and dislocation, each of which was reported in 14 studies [1–3,5,8,10–15,17,18]. It should be noted that dislocations, which have previously been reported as being more likely with THA, were not significantly different between the groups in this analysis. While the absolute dislocation rate was

One or more of the authors of this paper have disclosed potential or pertinent conflicts of interest, which may include receipt of payment, either direct or indirect, institutional support, or association with an entity in the biomedical field which may be perceived to have potential conflict of interest with this work. For full disclosure statements refer to <https://doi.org/10.1016/j.arth.2024.10.085>.

* Address correspondence to: Seper Ekhtiari, MD, MSc, Sinai Health, McMaster University, 476B-600 University Ave, Toronto, ON, Canada M5G 1X5.

E-mail address: seper.ekhtiari@medportal.ca

<https://doi.org/10.1016/j.arth.2024.10.085>

0883-5403/© 2024 Published by Elsevier Inc.

indeed higher in THA patients (4.6%, 59 of 1,270) compared to HA (2.9%, 38 of 1,325), this difference was not significant (odds ratio: 1.48, 95% confidence interval [CI]: 0.60 to 3.65). Mortality rates (11 RCTs) were nearly identical between groups, at 15.3% for THA patients versus 15.8% for HA patients (odds ratio: 0.98, 95% CI: 0.79 to 1.22) [2,3,8–14,18,19].

An option that may be considered to help reduce the risk of dislocation is dual mobility THA. While beyond the scope of this question, a systematic review mostly comprised of nonrandomized studies did find a significantly lower dislocation rate with dual mobility THA compared to HA in patients who have femoral neck fractures [20]. The only RCT included in the present review ($n = 60$) demonstrated a significant reduction in dislocation rates in the dual mobility THA group (0%) compared to the bipolar HA group (16.6%) [19]. However, the role of dual mobility THA remains unclear due to a lack of strong RCT evidence, and in particular, it is unclear if there is a sufficient reduction in dislocation rates to justify the added cost and unique complications associated with dual mobility implants.

There were small and statistically significant differences in favor of THA in terms of function and HRQoL. Based on nine RCTs, and with scores converted to the most common instrument (Harris Hip Score, range 0 to 100, higher scores = better), THA patients had a significantly higher mean score than HA patients (mean difference [MD]: 4.59, 95% CI: 1.65 to 7.53) [2,4–6,11–13,15,17]. However, this is smaller than the previously established minimal clinically important difference for the Harris Hip Score following arthroplasty (eight points) [21]. Similarly, based on five studies, there was a significantly higher HRQoL score based on the EuroQol-5 Dimensions for THA versus HA patients (MD: 0.05, 95% CI: 0.02 to 0.08) [2,11–13,17]. The minimal clinically important difference estimate for EuroQol-5 Dimensions has been determined to be 0.145 based on the best available literature [21], exceeding the mean difference.

Patients undergoing HA had significantly lower estimated blood loss (MD: 133.04 mL, 95% CI: 96.39 to 169.69) [3–7,9,15,16] and significantly shorter operative times (MD: 29.7 minutes, 95% CI: 17.95 to 41.49) [2–7,15,17,19]. These findings are not surprising given that acetabular work is not required in HA, whereas it is required in THA. The clinical importance of these findings is difficult to estimate; on the one hand, it is well established that longer operative times increase the risk of infection. However, the weighted mean operative time for the THA patients was only 96 minutes; thus, operative times in all patients were relatively short. As well, total complications (which included infection in some studies) did not differ between groups. In terms of estimated blood loss, the difference is statistically significant, but less than 150 mL, which may not have clinical significance. Transfusion rates, which were not reported in enough studies to allow for quantitative analysis, would be a more clinically important outcome to compare between groups.

Some interesting secondary analyses have been performed using data from the largest RCT on this topic, the Hip Fracture Evaluation with Alternatives of Total Hip Arthroplasty versus Hemiarthroplasty trial. Patients who underwent HA or THA by nonarthroplasty-trained surgeons had significantly higher rates of periprosthetic joint infection and discharge to a facility other than home. There was no significant difference in the reoperation rate [22]. In terms of the functional and HRQoL differences, THA compared to monopolar arthroplasty was associated with a small but clinically unimportant benefit, whereas THA versus bipolar arthroplasty was not. Higher American Society of Anesthesiologists scores and preoperative use of a mobility aid were associated with lower functional and HRQoL scores regardless of treatment [23]. Also, a secondary analysis of the Hip Fracture Evaluation with Alternatives of Total Hip Arthroplasty versus Hemiarthroplasty trial

identified a range of factors associated with the risk of revision surgery, including age, body mass index, comorbidities, and length of operation [24].

While not possible to include in our meta-analysis, registry data do provide additional support for the recommendation. Based on data from the Australian Orthopaedic Association National Joint Replacement Registry, there was no significant difference in revision rates at a maximum of 19.7 years follow-up between bipolar HA and THA, although monopolar HA did demonstrate a significantly higher revision rate [25].

Overall, there is no evidence to support recommending THA or HA universally in patients who have femoral neck fractures. Given high mortality rates following hip fractures, and the reduced costs, operative time, and blood loss, HA remains an excellent option in treating patients who have a displaced femoral neck fracture. Given that the arguments regarding THA in this population commonly revolve around potentially improved function, longevity/long-term survival of the implant, and the small, but statistically significant benefits found in these data, THA can be considered in younger, healthier, and more active patients; however, this recommendation cannot be said to be evidence-based as the studies analyzed did not differentiate among patients on these criteria.

CRediT authorship contribution statement


Geert Meermans: Writing – review & editing, Writing – original draft, Methodology, Data curation, Conceptualization. **Darko Talevski:** Writing – review & editing, Writing – original draft, Methodology, Data curation, Conceptualization. **Peteris Studers:** Writing – review & editing, Writing – original draft, Methodology, Data curation. **Emre Togrul:** Writing – review & editing, Writing – original draft, Methodology, Data curation. **G. Ruslan Nazaruddin Simanjuntak:** Writing – review & editing, Writing – original draft, Methodology, Data curation. **Kamolsak Sukhonthamarn:** Writing – review & editing, Writing – original draft, Methodology, Data curation. **Mark Phillips:** Writing – review & editing, Writing – original draft, Methodology, Data curation, Conceptualization. **Seper Ekhtiari:** Writing – review & editing, Writing – original draft, Visualization, Validation, Supervision, Software, Methodology, Formal analysis, Data curation, Conceptualization.

References


- [1] Dorr LD, Glousman R, Hoy AL, Vanis R, Chandler R. Treatment of femoral neck fractures with total hip replacement versus cemented and noncemented hemiarthroplasty. *J Arthroplasty* 1986;1:21–8.
- [2] Macaulay W, Nellans KW, Garvin KL, Iorio R, Healy WL, Rosenwasser MP. Prospective randomized clinical trial comparing hemiarthroplasty to total hip arthroplasty in the treatment of displaced femoral neck fractures. Winner of the dorr award. *J Arthroplasty* 2008;23(6 Suppl 1):2–8.
- [3] Parker MJ, Cawley S. A long term follow-up for a randomised trial of total hip arthroplasty versus hemiarthroplasty for displaced intracapsular fractures. *Injury* 2023;54:110925.
- [4] Li X, Zhao L, Chen R, Cao H, Wei Y, Wu X, et al. Effects of total hip arthroplasty and hemiarthroplasty on hip function in patients with traumatic femoral neck fracture. *Arch Orthop Trauma Surg* 2023;143:873–8.
- [5] Uikey K, Kumar Bakode H, Singh Rahangdale P, Bhalavi M. Comparative study of total hip arthroplasty versus bipolar hemiarthroplasty in treating neck of femur fractures in geriatric patients with osteoporosis. *Int J Pharmaceut Clin Res Orig Res Artic Orthop Spec* 2024;16:101–5.
- [6] Chatterji G, Shukla S, Singhania S, Singh MP, Mohanty SS, Jaiswal A, et al. A prospective study comparing the functional outcome of bipolar hemiarthroplasty versus total hip replacement in elderly patients with fracture of the neck of femur. *Cureus* 2022;14:e29529.
- [7] Ren C, Guo J, Gao Y. Comparison of total hip arthroplasty and hemiarthroplasty in elderly patients with femoral neck fracture. *Biomed Res* 2017;28:7127–30.
- [8] Baker RP, Squires B, Gargan MF, Bannister GC. Total hip arthroplasty and hemiarthroplasty in mobile, independent patients with a displaced intracapsular fracture of the femoral neck: a randomized, controlled trial. *J Bone Joint Surg Am* 2006;88:2583–9.

- [9] Blomfeldt R, Törnkvist H, Eriksson K, Söderqvist A, Ponzer S, Tidermark J. A randomised controlled trial comparing bipolar hemiarthroplasty with total hip replacement for displaced intracapsular fractures of the femoral neck in elderly patients. *J Bone Joint Surg Br* 2007;89:160–5.
- [10] Cadossi M, Chiarello E, Savarino L, Tedesco G, Baldini N, Faldini C, et al. A comparison of hemiarthroplasty with a novel polycarbonate-urethane acetabular component for displaced intracapsular fractures of the femoral neck: a randomised controlled trial in elderly patients. *Bone Joint J* 2013;95:609–15.
- [11] Chammout G, Kelly-Pettersson P, Hedbeck CJ, Stark A, Mukka S, Sköldenberg O. HOPE-trial: hemiarthroplasty compared with total hip arthroplasty for displaced femoral neck fractures in octogenarians. *JB JS Open Access* 2019;4:e0059.
- [12] The HEALTH Investigators. Total hip arthroplasty or hemiarthroplasty for hip fracture. *N Engl J Med* 2019;381:2199–208.
- [13] Mouzopoulos G, Stamatakis M, Arabatzi H, Vasiliadis G, Batanis G, Tsembeli A, et al. The four-year functional result after a displaced subcapital hip fracture treated with three different surgical options. *Int Orthop* 2008;32:367–73.
- [14] Skinner P, Riley D, Ellery J, Beaumont A, Coumine R, Shafaghian B. Displaced subcapital fractures of the femur: a prospective randomized comparison of internal fixation, hemiarthroplasty and total hip replacement. *Injury* 1989;20:291–3.
- [15] Sonaje JC, Meena PK, Bansawal RC, Bobade SS. Comparison of functional outcome of bipolar hip arthroplasty and total hip replacement in displaced femoral neck fractures in elderly in a developing country: a 2-year prospective study. *Eur J Orthop Surg Traumatol* 2018;28:493–8.
- [16] Sharma V, Awasthi B, Kumar K, Kohli N, Katoch P. Outcome analysis of hemiarthroplasty vs. Total hip replacement in displaced femoral neck fractures in the elderly. *J Clin Diagn Res* 2016;10:RC11–3.
- [17] Keating JF, Grant A, Masson M, Scott NW, Forbes JF. Randomized comparison of reduction and fixation, bipolar hemiarthroplasty, and total hip arthroplasty: treatment of displaced intracapsular hip fractures in healthy older patients. *J Bone Joint Surg Am* 2006;88:249–60.
- [18] Van Den Bekerom MPJ, Hilverdink EF, Sierevelt IN, Reuling EMBP, Schnater JM, Bonke H, et al. A comparison of hemiarthroplasty with total hip replacement for displaced intracapsular fracture of the femoral neck: a randomised controlled multicentre trial in patients aged 70 years and over. *J Bone Joint Surg Br* 2010;92:1422–8.
- [19] Iorio R, Iannotti F, Mazza D, Speranza A, Massafra C, Guzzini M, et al. Is dual cup mobility better than hemiarthroplasty in patients with dementia and femoral neck fracture? A randomized controlled trial. *SICOT J* 2019;5:38.
- [20] Cha YH, Yoo JI, Kim JT, Park CH, Ahn YS, Choy WS, et al. Dual mobility total hip arthroplasty in the treatment of femoral neck fractures: systematic review and meta-analysis. *Bone Joint J* 2020;102-B:1457–66.
- [21] Ekhtiari S, Gormley J, Axelrod DE, Devji T, Bhandari M, Guyatt GH. Total hip arthroplasty versus hemiarthroplasty for displaced femoral neck fracture: a systematic review and meta-analysis of randomized controlled trials. *J Bone Joint Surg Am* 2020;102:1638–45.
- [22] DeAngelis RD, Minutillo GT, Stein MK, Schemitsch EH, Bzovsky S, Sprague S, et al. Who did the arthroplasty? Hip fracture surgery reoperation rates are not affected by type of training-an analysis of the HEALTH database. *J Orthop Trauma* 2020;34:S64.
- [23] Axelrod D, Comeau-Gauthier M, Bzovsky S, Schemitsch EH, Poolman RW, Frihagen F, et al. What predicts health-related quality of life for patients with displaced femoral neck fractures managed with arthroplasty? A secondary analysis of the HEALTH trial. *J Orthop Trauma* 2020;34:S29–36.
- [24] Blankstein M, Schemitsch EH, Bzovsky S, Poolman RW, Frihagen F, Axelrod D, et al. What factors increase revision surgery risk when treating displaced femoral neck fractures with arthroplasty: a secondary analysis of the HEALTH trial. *J Orthop Trauma* 2020;34:S49.
- [25] Farey JE, Cuthbert AR, Adie S, Harris IA. Bipolar hemiarthroplasty does not result in a higher risk of revision compared with total hip arthroplasty for displaced femoral neck fractures. *J Bone Joint Surg Am* 2022;104:919–27.


Appendix–Forest Plots




Appendix Figure 1. Revision surgery.




Appendix Figure 2. Health-related quality of life.




Appendix Figure 3. Dislocation rates.




Appendix Figure 4. Mortality rates.




Appendix Figure 5. Periprosthetic fracture.



Appendix Figure 6. Function.



Appendix Figure 7. Operative time.



Appendix Figure 8. Estimated blood loss.