

Reverse Survival x5

Jose H C Monteiro da Silva, Helena C Castanheira

2021-06-02

Reverse Survival x5 version

In this short note we present the differences between the original version and the x5 version of the Reverse Survival method implemented in the package *fertestr*.

The original version (let's call it x1) reconstructs the TFRs and birth counts (we call it B^t) for 15 points in time preceding the date of inquiry each one corresponding to each single age of children's population count input (for ages 0 to 14). In the x5 version, we reconstruct the TFRs and B^t for each five year period before the inquiry based on population counts of children reported in five-year age groups (0-4, 5-9, 10-14). Therefore, the main difference of these two approaches lies in the reconstruction of B^t through reverse survival.

In the x1 approach, the inputs for computing B^t are: 1) population of children from 0-14 in single age groups, 2) a 'standard' survival function in single ages from ages 0 to 15 (l_x), 3) estimates of child mortality probability for each three five year period preceding the inquiry (${}_5q_0^{t-2.5}$, ${}_5q_0^{t-7.5}$, ${}_5q_0^{t-12.5}$). The original version of the x1 approach (Moultrie et al. 2013) uses the logit transformation of the standard survival function and the mortality probabilities of prior periods to retrieve survival functions and from them compute the "proportion of births occurring x to $x + 1$ years earlier that survive to the time of the inquiry, c_x^L ." Details of this approach can be viewed in the spreadsheet attached to the method's explanation in <http://demographicestimation.iussp.org/content/estimation-fertility-reverse-survival>.

For the x5 alternative version, the inputs for computing B^t are similar: 1) population counts of children reported in five-year age groups, 2) a 'standard' survival function for ages 5, 10 and 15 (l_5, l_{10}, l_{15} - l_x radix l_0 set to 1), 3) estimates of child mortality probability for each three five year period preceding the inquiry (${}_5q_0^{t-2.5}$, ${}_5q_0^{t-7.5}$, ${}_5q_0^{t-12.5}$). The estimation procedure of c_x^L (proportion of births occurring x to $x + 5$ years earlier that survive to the time of the inquiry) for periods $x = 2.5$ (0-4 years earlier), $x = 7.5$ (5-9 years earlier) and $x = 10-14$ (10-14 years earlier) follow the same strategy of the the original x1 version of the method, and we will describe it below.

Step 1: From ${}_5q_0$, compute α^t

We compute the intercept α^t of the linear relation between logits of observed (${}_5q_0$) and standard survival function (l_5) for each period preceding the inquiry. As Moultrie et al. (2013), we set the slope values as $\beta = 1$. Therefore,

$$\begin{aligned} Y_{obs}_x^t &= \alpha^t + \beta * Y_{std}_x \\ \alpha^t &= Y_{obs}_5^t - Y_{std}_5. \end{aligned}$$

Step 2: from standard l_x and α , compute l_x^t

Then, by using the estimated α values, we can compute a set of survival functions l_5, l_{10} and l_{15} for each period ealier than the date of inquiry:

$$l_x^t = \frac{1}{1 + \exp[2(\alpha^t + Y std_x)]}.$$

Step 3: compute ${}_nL_x$

Using estimated values for l_5^t, l_{10}^t and l_{15}^t and $l_0^t = 1$, we compute the person-years lived between ages x and $x + 5$, for $x = 0, 5$ and 10 :

$${}_nL_x^t = 5 * l_{x+5}^t + {}_n a_x * (l_x^t - l_{x+5}^t),$$

where ${}_5a_0 = 0.50$, ${}_5a_5 = 2$ and ${}_5a_{10} = 2.5$ by default (these values can be modified in the function).

Step 4: compute survivorship ratios P_x^t

$$P_0^t = \frac{{}_5L_0^t}{{}_5l_0^t}$$

$$P_5^t = \frac{{}_5L_5^t}{{}_5L_0^t}$$

$$P_{10}^t = \frac{{}_5L_{10}^t}{{}_5L_5^t}$$

Step 5: define S_x^t and compute cL_x

We compute S_x^t as the survivorship ratio between ages x and $x + 5$ in periods t equal 0-4, 5-9 and 10-14 before the inquiry.

Then,

$$S_0^{0-4} = P_0^{0-4}$$

$$S_0^{5-9} = P_0^{5-9}$$

$$S_0^{10-14} = P_0^{10-14}$$

$$S_5^{0-4} = P_5^{0-4}$$

$$S_5^{5-9} = P_5^{5-9}$$

$$S_{10}^{0-4} = P_{10}^{0-4},$$

then:

$${}^cL_x = S_0^x S_5^{x-5} \dots S_x^0,$$

and

$${}^cL_0 = S_0^{0-4} = P_0^{0-4}$$

$${}^cL_5 = S_5^{0-4} S_0^{5-9} = P_5^{0-4} P_0^{5-9}$$

$${}^cL_{10} = S_{10}^{0-4} S_5^{5-9} S_0^{10-14} = P_{10}^{0-4} P_5^{5-9} P_0^{10-14}.$$

Step 6: reconstruct B^t

$$B^{x-2.5} = \frac{P_{0-4}}{5^c L_0}$$

$$B^{x-7.5} = \frac{P_{5-9}}{5^c L_5}$$

$$B^{x-12.5} = \frac{P_{10-14}}{5^c L_{10}}$$

References

Moultrie, Tom, Rob Dorrington, Allan Hill, Kenneth Hill, Ian Timæus, and Basia Zaba. 2013. *Tools for Demographic Estimation*. Paris: International Union for the Scientific Study of Population (IUSSP).