

# Germline melanoma susceptibility and prognostic genes: A review of the literature

Katherine A. Ward, MD,<sup>a</sup> DeAnn Lazovich, PhD,<sup>b</sup> and Maria K. Hordinsky, MD<sup>c</sup>  
Minneapolis, Minnesota

In recent years, there have been increasing efforts to identify germline genetic variants that may alter melanoma susceptibility and prognosis. The findings of these studies have indicated the presence of rare, high-penetrance alleles with large effects, such as *CDKN2A* and *CDK4*, more common, moderately penetrant genes like *MC1R*, and very common, low-penetrance polymorphisms with small effects that are related to pigmentation, nevus count, immune responses, DNA repair, metabolism, and the vitamin D receptor. The study of these low-penetrance single nucleotide polymorphisms is relatively new; thus many of them are termed 'candidate melanoma susceptibility or prognostic genes.' This review summarizes the research on germline polymorphisms that have been implicated in melanoma susceptibility and prognosis in order to provide a framework for additional studies to meet the ultimate goal of predicting a patient's risk of, and prognosis in, cutaneous malignant melanoma. (J Am Acad Dermatol 2012;67:1055-67.)

**Key words:** cutaneous malignant melanoma; DNA repair; genetic susceptibility; germline; immune responses; pigmentation; prognosis; metabolism; nevus counts; vitamin D receptor.

## INTRODUCTION

It has long been known that mutations in the *CDKN2A* and *CDK4* genes play a role in familial melanoma, which comprises approximately 10% of all cutaneous malignant melanoma (CMM) cases.<sup>1-4</sup> Recently, other CMM risk alleles have been found in genes related to pigmentation,<sup>5</sup> nevus count,<sup>6</sup> immune responses,<sup>7</sup> DNA repair,<sup>8</sup> metabolism,<sup>9</sup> and the vitamin D receptor (VDR).<sup>10</sup>

The penetrance of these genes in CMM is a spectrum, with high-risk alleles being *CDKN2A* and *CDK4*, moderate-risk genes including *MC1R*, and low-risk alleles such as *TYR*, *ASIP*, *TYRP1*, *OCA2*, *SLC45A2*, *GSTM1*, *CYP2D6*, *VDR*, *IL-9* [interleukin 9], *IL-10*, *TNF*, *HLA-DQB0301*, *XPC*, *PLA2G6*, and numerous others.<sup>1,11-16</sup> The lower risk alleles are much more prevalent in the population, but their effects are less profound, whereas the higher risk alleles, although rare, when present almost always confer a phenotype of CMM.<sup>16</sup>

### Abbreviations used:

CDK:	cyclin-dependent kinase
CMM:	cutaneous malignant melanoma
<i>CYP2D6</i> :	cytochrome P450-debrisoquine hydroxylase locus
GSTs:	glutathione S-transferases
HLA:	human leukocyte antigen
ICAM-1:	intracellular adhesion molecule-1
IL:	interleukin
<i>MC1R</i> :	melanocortin-1 receptor
MSH:	$\alpha$ -melanocyte-stimulating hormone
<i>OCA2</i> :	oculocutaneous albinism-related gene
SNP:	single nucleotide polymorphism
VDR:	vitamin D receptor

The purpose of this review is to provide a synopsis of what is currently known about the relationship between the various high, moderate, and low-risk germline polymorphisms, as well as other candidate single nucleotide polymorphisms (SNPs), and susceptibility and prognosis in CMM in order to provide a guide for future larger and more comprehensive research undertakings.

From the University of Minnesota Medical School,<sup>a</sup> Division of Epidemiology and Community Health, University of Minnesota,<sup>b</sup> and the Department of Dermatology, University of Minnesota.<sup>c</sup>

Funding sources: None.

Conflicts of interest: Drs Lazovich and Ward have no conflicts of interest to declare. Dr Hordinsky has declared financial relationships with Johnson & Johnson, Novartis, Medics, and Allergan.

Accepted for publication February 29, 2012.

Reprints not available from the authors.

Correspondence to: DeAnn Lazovich, PhD, Division of Epidemiology and Community Health, University of Minnesota, West Bank Office Building (WBOB), 1300 S. Second St, Suite 300, Minneapolis, MN 55454-1015. E-mail: lazov001@umn.edu.

Published online May 14, 2012.

0190-9622/\$36.00

© 2012 by the American Academy of Dermatology, Inc.

doi:10.1016/j.jaad.2012.02.042

**HIGH-PENETRANCE GENES*****CDKN2A***

The best-established high-risk locus for melanoma susceptibility is the cyclin-dependent kinase (CDK) inhibitor 2A gene (*CDKN2A*), which is located on chromosome 9p21.<sup>17,18</sup> Germline mutations in this locus were initially reported in 1995 among multiple melanoma-predisposed families.<sup>19-25</sup> Approximately 40% of familial melanoma cases carry *CDKN2A* mutations.<sup>14</sup> In addition to melanoma, the 9p21 locus has also been implicated in certain other cancers, including pancreatic cancer, highlighting its importance in tumor suppression.<sup>3,26-28</sup>

The *CDKN2A* locus encodes for 2 different proteins via alternative splicing of four exons.<sup>16</sup> The two proteins, p16/Ink4a and p14/Arf, are both extremely important tumor suppressors that regulate the progression of the cell cycle from G<sub>1</sub> to S phase and apoptosis, respectively.<sup>16,29</sup> The main function of the p16/Ink4a protein is to bind *CDK4*, which thereby inhibits this protein kinase from phosphorylating the retinoblastoma (Rb) tumor suppressor, such that the E2F restriction at G<sub>1</sub> is not released and progression to the S phase is halted.<sup>30,31</sup> This yields an overall effect of blocking cell division and proliferation. Whereas the p16/Ink4a gene protein works through the Rb tumor suppressor pathway, the p14/Arf gene product is directly involved in p53 regulation and apoptosis.<sup>16-32</sup> p14/Arf exerts its effects by binding to human double minute-2 through the N terminal domain, which sequesters human double minute-2 in the nucleolus so it cannot bind and down-regulate p53.<sup>16</sup> Thus p53 is stabilized as human double minute-2 is depleted.<sup>33-35</sup> Therefore mutation in the 9p21 locus of Ink4a and Arf simultaneously impairs two of the most important tumor suppressor pathways, Rb and p53, greatly increasing the likelihood of malignant transformation.

***CDK4***

A second high-penetrance melanoma susceptibility gene, *CDK4*, has also been established for familial melanoma.<sup>18</sup> This gene is located on chromosome 12q13 and encodes the kinase that is the target of p16/Ink4a in the Rb pathway of tumor suppression.<sup>17</sup> Any mutation in *CDK4* that inhibits the binding of p16/Ink4a rendering *CDK4* resistant to

inactivation is, therefore, oncogenic. Only a single activating mutation in the *CDK4* germline is necessary for tumorigenesis.<sup>19</sup>

Two *CDK4* germline mutations have been found in patients with melanoma, namely *Arg24His* and *Arg24Cys*.<sup>36</sup> Coincidentally, these two mutations are also the most common somatic *CDK4* mutations

found in melanoma tumors.<sup>36</sup> These germline mutations in *CDK4* are much rarer compared with *CDKN2A* mutations in familial melanoma. Fewer than 15 families with these mutations have been reported worldwide and only 2% of families in the most extensive study of familial melanoma conducted by the Melanoma Genetics Consortium (GenoMEL), exhibited the mutations.<sup>2,3,37-40</sup>

Therefore it can be con-

cluded that *CDKN2A* germline mutations are much more prevalent among inherited melanoma kindreds than *CDK4* mutations, but larger studies are warranted to attempt to decipher the role that *CDK4* mutations play in melanoma susceptibility.

**MODERATE PENETRANCE GENES*****MC1R***

The melanocortin-1 receptor (*MC1R*) is considered a moderate-risk gene for melanoma susceptibility and is a key regulator of skin pigmentation.<sup>17</sup> The *MC1R* is a 7-transmembrane G-protein coupled receptor that acts on adenylate cyclase in order to increase cAMP levels in response to interaction with its ligand,  $\alpha$ -melanocyte-stimulating hormone (MSH).<sup>41,42</sup> The binding of MSH affects the activity of various enzymes and proteins involved in the production of melanin, with the ultimate effect of switching production from red/yellow pheomelanins (photosensitive and potentially mutagenic because of its production of free radicals in response to UV irradiation) to brown/black eumelanins (photoprotective).<sup>43-45</sup> This activation of *MC1R* on the surface of melanocyte cell membranes is an integral part of the tanning response following UV irradiation.<sup>46,47</sup>

The *MC1R* gene locus is highly polymorphic; it is also a major cause of the various pigmentation phenotypes and skin phototypes in humans.<sup>47,48</sup> *MC1R* variant alleles, with amino acid substitutions within the coding region, have been shown to reduce receptor function; as a result, they show increased pheomelanin to eumelanin ratios within cells.<sup>41,49-52</sup>

**CAPSULE SUMMARY**

- *CDKN2A* and *CDK4* are high-penetrance melanoma susceptibility alleles.
- *MC1R* is a moderate-penetrance melanoma risk gene.
- Many low-penetrance genes also likely play a role in melanoma susceptibility and prognosis; these relate to pigmentation, nevus count, immune responses, DNA repair, metabolism, and the vitamin D receptor (VDR).

The higher pheomelanin levels associated with these *MC1R* variant alleles cause the red hair and fair skin phenotype (red hair color [RHC]), also known as the *MC1R* null phenotype.<sup>53</sup> This subpopulation is known to be more sensitive to the effects of UV irradiation, demonstrated by a lack of tanning ability, and is also at increased risk of developing CMM, by mechanisms yet to be fully understood.

Melanoma patients are significantly more likely to carry *MC1R* variants than are healthy control subjects. In fact, having a *MC1R* variant carries a 2.2- to 3.9-fold risk of developing melanoma, and the effects of having multiple variants are additive, as carriers of two variants have a 4.1- to 4.8 fold risk.<sup>1,19,54</sup> In a recent meta-analysis, 7 *MC1R* variants (*Asp84Glu*, *Arg142His*, *Arg151Cys*, *Ile155Thr*, *Arg160Trp*, *Arg163Gln*, and *Asp294His*) were all significantly associated with CMM development, with odds ratios (ORs) ranging from 1.42 (95% confidence interval [CI], 1.09-1.85) for *Arg163Gln* to 2.45 (95% CI, 1.32-4.55) for *Ile155Thr*, and most were associated with red hair and fair skin or red hair only.<sup>55</sup> Another recent study implicated the *MC1R* variant, R151C, in CMM development in the Ashkenazi Jewish population with an OR of 2.6 (95% CI, 1.3-5.3).<sup>56</sup> A previous study also showed a significant association between R151C, in addition to R160W, and D294H *MC1R* variants (RHC phenotype), carrying a more than two-fold increased risk of melanoma.<sup>57</sup> Mössner et al<sup>58</sup> confirmed the significant risk associated with R151C with an OR of 1.69 (95% CI, 1.12-2.55), but also added another *MC1R* variant to the list of susceptibility alleles, D84E, with an OR of 4.96 (95% CI, 1.06-23.13). In a very recent, comprehensive field synopsis and systematic meta-analysis, variant *MC1R* alleles were correlated with increased risk of CMM with an OR of 1.83 (95% CI, 1.56-2.15) as compared with controls.<sup>59</sup>

Certain *MC1R* variants not only act as independent risk factors for melanoma development, but they also serve as modifier alleles, exerting effects on the *CDKN2A* gene.<sup>60,61</sup> *MC1R* variants have been shown to increase the penetrance of *CDKN2A* mutations from 50% to 84%, in addition to lowering the mean age of onset by 20 years.<sup>1,19,60</sup>

## LOW-PENETRANCE GENES

### Pigmentation/nevus count-related genes

Individuals with fair skin, blond or red hair, who never tan and always burn (Fitzpatrick phototypes I and II) are at the highest risk for CMM development.<sup>1,62</sup> As a result of this correlation, there has been a search in recent years to identify pigmentation genes in addition to *MC1R* that may have an impact on CMM susceptibility.

In a large, recent study of populations of European descent, a significant association was found between *ASIP*, which encodes agouti signaling protein, the antagonist of MSH binding to *MC1R*, on chromosome 20q11.22 and CMM, with an OR of 1.45 (95% CI, 1.29-1.64).<sup>63</sup> A previous study also demonstrated a significant association between *ASIP* SNPs, rs4911414[T] and rs1015362[G], and CMM among European populations, again with an OR of 1.45.<sup>64</sup> Another study found a significant association between these same two *ASIP* SNPs and CMM, but with an OR of 1.68.<sup>65</sup> An Australian genome-wide association study also indicated the presence of a melanoma susceptibility locus on chromosome 20q11.22, with an OR of 1.72 for *ASIP* SNPs, rs910873 and rs1885120.<sup>66</sup> However, a study with 423 melanoma cases and 147 controls showed no association between *ASIP* polymorphisms and CMM, possibly because of the low number of control subjects.<sup>67</sup>

The *TYR* gene on chromosome 5q34, which encodes for the tyrosinase protein that affects eye color and tanning response, has also been implicated in CMM susceptibility.<sup>36,68</sup> One study found a significant association between the *Arg402Gln* variant and CMM, with an OR of 1.21.<sup>12</sup> Bishop et al<sup>5</sup> found an association between a coding variant in *TYR* and CMM, with an OR of 1.27. A very recent study found a significant association between the *TYR* variant, rs1126809, and CMM risk with an OR of 1.22.<sup>59</sup> Two studies have described an interesting inverse relationship between the *TYR Arg402Gln* polymorphism and generalized vitiligo and CMM. This SNP confers protection from generalized vitiligo, while increasing susceptibility to CMM among European-derived white individuals.<sup>5,63</sup>

Another candidate melanoma susceptibility gene is *TYRP1* on chromosome 9p23, encoding for tyrosine-related protein 1, which stabilizes tyrosinase.<sup>12</sup> One study found that the *TYRP1* variant, rs1408799, actually decreased the risk of CMM by an OR of 0.77 (95% CI, 0.60-0.98).<sup>65</sup> Chatzinasiou et al<sup>59</sup> demonstrated a significant association between the *TYRP1* variant, rs1408799, and protection from CMM (OR, 0.86 [95% CI, 0.80-0.93]).

*OCA2* (human type II oculocutaneous albinism-related gene) is another candidate melanoma susceptibility gene located on chromosome 15q11.2-12.<sup>12</sup> The protein encoded by this gene is a melanosomal transmembrane protein that modifies human eye color and pigmentation.<sup>41,69</sup> Fernandez et al<sup>62</sup> found a variant allele of this gene, R419Q (rs1800407), to significantly increase the risk of CMM by an OR of 1.55 (95% CI, 1.04-3.31). An earlier study also found an association between CMM and

**Table I.** Summary of low-penetrance candidate melanoma susceptibility and prognostic genes

Pigmentation/nevus count genes	Immune genes	DNA-repair genes	Metabolism genes	Vitamin D receptor polymorphisms
<i>ASIP</i>	<i>IL-10</i>	<i>XPB/ERCC2</i>	<i>CYP2D6</i>	<i>FokI</i>
<i>TYR</i>	<i>IL-1<math>\beta</math></i>	<i>ERCC1</i>	<i>GSTM1</i>	<i>BsmI</i>
<i>TYRP1</i>	<i>TNF-<math>\alpha</math></i>	<i>XPB</i>	<i>GSTT1</i>	<i>TaqI</i>
<i>OCA2</i>	<i>LT-<math>\alpha</math></i>	<i>XRCC3</i>	<i>GSTP1</i>	<i>Apal</i>
<i>SLC45A2 (MATP)</i>	<i>IL-6R</i>	<i>MGMT</i>		<i>A-1012G</i>
<i>MYO7A</i>	<i>IFN-<math>\gamma</math></i>	<i>XRCC1</i>		
<i>NID1</i>	<i>HLA class II allele DQB1*0301</i>	<i>MDM2</i>		
<i>KIT</i>	<i>ICAM-1</i>	<i>APEX1</i>		
<i>KITLG</i>		<i>TERT1</i>		
<i>IRF</i>		<i>TRF1</i>		
<i>HERC2</i>		<i>TERT-CLPTMIL</i>		
<i>PAX3</i>				
<i>EDNRB</i>				
<i>ADTB3A</i>				
<i>CHS1</i>				
<i>MLANA</i>				
<i>ATRNL</i>				
<i>SOX10</i>				
<i>HPS</i>				
<i>MGRN1</i>				
<i>MYO5A</i>				
<i>SLC24A4</i>				
<i>PLA2G6</i>				

the *OCA2* locus, with a *P* value of .030.<sup>5</sup> Another study implicated *OCA2* polymorphisms and mole count, which is positively correlated with CMM.<sup>41</sup>

*SLC45A2* (*MATP*) is a gene on chromosome 5p13.3 that is involved in normal human pigmentation and has recently been investigated as a CMM susceptibility candidate gene.<sup>12,70</sup> A Spanish population study found that the *SLC45A2* variant allele, rs16891982 (p.Phe374Leu), was associated with protection from CMM development, with an OR of 0.41 (95% CI, 0.24-0.70).<sup>70</sup> A recent French study exhibited similar findings, as the *SLC45A2* variant, p.Phe374Leu, was again found to be significantly and strongly protective for CMM, with an OR of 0.32 (95% CI, 0.34-0.43).<sup>71</sup> Nan et al<sup>65</sup> found a different *SLC45A2* variant, 1721 C>G, to be significantly associated with a reduction in risk for CMM, with an OR of 0.75 (95% CI, 0.60-0.95). A recent study found the *SLC45A2* variant, 16891982, to be associated with a decreased CMM risk with an OR of 0.40.<sup>59</sup> Finally, a fifth study found an association with CMM and the *SLC45A2* variants, rs28777, rs35391, and rs16891982.<sup>72</sup>

A variant in the *MYO7A* gene was recently described in a single study as being associated with CMM susceptibility. The variant, S1666C (rs2276288), had an increased risk of CMM development, with an OR of 1.35 (95% CI, 1.04-1.76), although its

association appeared to be stronger among certain phenotypic groups.<sup>62</sup>

Increased nevus counts occurring on sun-exposed sites is a well-known risk factor for CMM development.<sup>6,73</sup> Both benign and dysplastic nevi are presumed to be precursors for CMM, but dysplastic melanocytic nevi are believed to be associated with a higher risk of CMM transformation.<sup>74</sup> Two genetic variants, at 9p21 and 22q13, have recently been identified by genome-wide association studies (GWAS) to be associated with melanocytic nevi development.<sup>75</sup>

A recent GWAS identified a novel susceptibility locus for nevus count and CMM risk.<sup>76</sup> This gene, known as nidogen 1 (*NID1*) on 1q42, produces a member of the nidogen family of basement membrane proteins involved in basement membrane assembly and is a biologically plausible locus for neovogenesis and melanoma development.<sup>76</sup> Results of the study indicated that increased expression of nidogen in one variant *NID1* SNP (rs10754833 T allele) was significantly associated with decreased CMM risk (OR, 0.86) (Table I).<sup>76</sup>

A recent study reported that the SNPs of 5 genes mentioned above (*MC1R*, *SLC45A2*, *OCA2*, *TYR*, and *ASIP*) explain approximately one third to one half of the difference in risk of CMM due to pigmentation phenotype.<sup>72</sup> However, other pigmentation-related

CMM candidate susceptibility genes have begun to be researched and include *KIT*, *KITLG*, *IRF*, *HERC2*, *PAX3*, *EDNRB*, *ADTB3A*, *CHS1*, *MLANA*, *ATRN*, *SOX10*, *HPS*, *MGRN1*, *MYO5A*, *SLC24A4*, and *PLA2G6* (see Table I). The study of these genes is relatively new, but most are related to the production or transport of melanin, hair color, tanning ability, nevus counts, or melanocyte proliferation, differentiation, and survival.<sup>14,62,77-79</sup> To date, results on these candidate genes and relationship to CMM development have been unrevealing, but more studies are warranted to assess the effects of these and other pigmentation genes on CMM susceptibility.

### Immune-related genes

There is much evidence that indicates CMM patients develop an immune-mediated response to their tumors.<sup>80,81</sup> However, in most cases the immune response is insufficient to halt tumor growth. This is especially evident in immunosuppressed transplant patients, as they exhibit higher than normal rates of CMM development.<sup>82,83</sup> It is possible, then, that allelic variations in the genes controlling the immune response may be one avenue to which differences in CMM susceptibility and prognosis may be attributed.<sup>80</sup> A number of immune-modulating genes have been examined in relation to CMM in the past two decades (see Table I).

Differences in ability to produce IL-10 have been widely studied and may be relevant in the development and course of CMM, most likely due to its immunosuppressor and anti-angiogenic properties.<sup>84</sup> The major IL-10 promoter haplotypes are *GCC*, *ACC*, and *ATA* formed by SNPs *IL-10-1087AG*, *IL-10-824CT*, and *IL-10-597AC*, with *GCC/GCC* (*GG*) being high in vitro IL-10 producers, *GCC/ACC* and *GCC/ATA* (*AG*) as medium producers, and *ACC/ACC*, *ACC/ATA*, and *ATA/ATA* (*AA*) defined as low producers.<sup>11,84-86</sup>

Recent findings have implicated low producing IL-10 haplotypes in CMM susceptibility and poor prognosis. Alonso et al<sup>84</sup> reported that the low producer *-1082AA* genotype was significantly associated with decreased survival in CMM patients with advanced disease (Table II). A recent case-control study found that the IL-10 higher producing haplotype *ITAGC* was found to be significantly associated with a reduced risk of CMM development ( $P = 0.02$ ).<sup>11</sup> A previous study did not find a correlation of the above-mentioned SNPs with CMM susceptibility, but did find an association with prognosis; the IL-10 low-producing haplotypes were significantly linked to the poor prognostic indicator of increased tumor thickness and decreased survival time (see Table II).<sup>87</sup> Two other studies reported

significant associations between the IL-10 low-producing haplotypes and poor disease outcome and/or susceptibility in CMM (see Table II).<sup>7,88</sup>

However, Dummer et al<sup>89</sup> reported results in opposition to these findings, with increased IL-10 levels in CMM patients with metastatic disease. Another study identified IL-10 as a growth factor of melanoma cells in vitro.<sup>90</sup> One study also reported that an IL-10 low-producing haplotype was associated with prolonged survival in CMM patients with metastases, but a more recent study found that the low-producing *AA* genotypes were associated with more rapid progression (see Table II).<sup>91,92</sup>

Because of the immunogenicity of melanoma tumors, other cytokines have also been researched in relation to susceptibility and prognosis (see Table I). One study investigated SNPs associated with IL-1 $\beta$ , IL-2, IL-4, IL-6, IL-8, and interferon gamma (IFN- $\gamma$ ) and susceptibility and prognosis in CMM and found that none of the cytokines had any significant association with susceptibility, and only the *IL-1 $\beta$ -511 TT* genotype had an association with prognosis, as it was correlated with thinner tumors (see Table II).<sup>80</sup> Another study researched the influence of tumor necrosis factor alpha (TNF- $\alpha$ ) and lymphotoxin-alpha (LT- $\alpha$ ) on CMM and reported that the TNF- $\alpha$  -238 *GG* genotype was significantly associated with increased susceptibility and the *GA* genotype showed a significant association with decreased susceptibility, whereas LT- $\alpha$  +252 genotype may have had a prognostic correlation with increased mitotic count in vertical growth phase tumors (see Table II).<sup>93</sup> Another study did not find any susceptibility or prognostic associations between IL-6 or IFN- $\gamma$  and CMM.<sup>87</sup> However, Gu et al<sup>82</sup> evaluated the associations between *IL-4*, *IL-6*, and *IL-10* and the IL receptor genes, *IL-4R*, *IL-6R*, and *IL-10R*, and CMM susceptibility; they reported that 4 SNPs in the *IL-6R* gene were associated with an increased risk of CMM, but none of the other genes exhibited any other associations.<sup>83</sup> Finally, another study found that the *interferon- $\gamma$  1874A*  $\rightarrow$  *T* gene polymorphism independently predicted overall survival in CMM patients (see Table II).<sup>91</sup>

The association between human leukocyte antigen (HLA) class II loci and CMM is a topic of widespread debate as it has been postulated that this relationship might be the immunogenetic basis for melanoma susceptibility.<sup>94,95</sup> In an early study of 45 Caucasian patients with CMM, Lee et al reported that 56% carried the HLA class II allele, *DQB1\*0301*, versus only 27% of Caucasian controls ( $P = .003$ ); additionally, the allele was associated with advanced disease at diagnosis, with 44% of the *DQB1\*0301*-positive patients presenting with metastases compared with only 5% of *DQB1\*0301*-negative patients



**Table II.** Summary of studies evaluating candidate melanoma prognostic genes

Authors	Year	Gene(s)	Genotype	No. of patients	No. of controls	Study design	Statistical significance	Outcome
Alonso et al <sup>84</sup>	2005	<i>IL-10</i>	−1082AA	98	100	Case-control	$P < .05$	Decreased survival time and mean age at diagnosis
Martinez-Escribano et al <sup>87</sup>	2002	<i>IL-10</i>	−1082AA, −819CT, −592CA	42	48	Case-control	$P = .002$ ; $P = .050$	Decreased survival time; increased Breslow thickness
Howell et al <sup>7</sup>	2001	<i>IL-10</i>	−1082AA; 1082GA	165	158	Case-control	$P = .005$ ; $P = .009$	Increased Breslow thickness; increased Breslow thickness
Liu et al <sup>91</sup>	2005	<i>IL-10</i>	−1082AA;	90	N/A	Prospective clinical trial	$P = 0.065$ ; $P = 0.33$	Overall survival; Response to therapy
		<i>Interferon-γ</i>	+874A → T				$P < 0.001$ ; $P = 0.001$	Overall survival; Response to therapy
		<i>ERCC1</i>	Codon 118 (TT/CT)				$P = 0.045$ ; $P = 0.03$	Overall survival; Response to therapy in stage IV melanoma patients treated with biochemotherapy
von Euw et al <sup>92</sup>	2008	<i>IL-10</i>	−1082AA	16	N/A	Phase I clinical trial	$P = .04$	Accelerated disease progression after lymph node surgery and vaccine
Howell et al <sup>80</sup>	2003	<i>IL-1β</i>	−511TT	169	261	Case-control	$P = .03$	Decreased Breslow thickness
Howell et al <sup>93</sup>	2002	<i>Lymphotoxin-α</i>	+255AA	146	220	Case-control	$P = .02$	Higher mitotic count in vertical growth-phase tumors
Lee et al <sup>96</sup>	1994	<i>HLA-DQB1*0301</i>	Present	45	200	Case-control	$P = .02$ ; $P = .003$	Increased Breslow thickness; metastases on presentation
Lee et al <sup>97</sup>	1996	<i>HLA-DQB1*0301</i>	Present	259	N/A	Cohort	$P = .0002$	Decreased disease-free survival
Figl et al <sup>110</sup>	2009	<i>XRCC1</i>	R399Q*AA	400 + 529	N/A	Cohort	$P = .0006$ ; $P = .04$	Increased metastasis-free survival.
Bu et al <sup>120</sup>	2007	<i>GSTM1</i>	−77T>C Null	11	4	Case-control	$P = .03$	Increased survival following first metastasis
Hutchinson et al <sup>10</sup>	2000	<i>VDR (TaqI+FokI)</i>	TTff	316	108	Case-control	$P = .001$	Breslow thickness >2.5 mm
Santonocito et al <sup>126</sup>	2007	<i>VDR (BsmI)</i>	bb	101	101	Case-control	$P = .001$	Breslow thickness ≥ 3.5 mm Increased Breslow thickness

N/A, Not applicable.

( $P = 0.003$ ) (see Table II).<sup>96</sup> Another study by these same authors found that stage I or II CMM patients who carried the HLA-*DQB1\*0301* allele were at an increased risk of developing recurrent disease compared with stage-matched patients who were *DQB1\*0301*-negative (see Table II).<sup>97</sup>

A study among the Spanish population indicated that the contribution of HLA class II alleles to CMM susceptibility was not significant, but did report a statistically significant increase in HLA-DQA1 homozygosity among CMM patients versus controls and a significant association between HLA-*DQB1\*0301* and red or fair-haired persons (relative risk, 5.65).<sup>94</sup> A variety of other studies have reported even more conflicting results with HLA-*DQB1\*0301* among CMM populations. One study reported no association between *DQB1\*0301* and risk of CMM,<sup>98</sup> whereas another found a slight, but not significant, positive association with susceptibility,<sup>99</sup> and yet another indicated a negative association with risk of CMM.<sup>100</sup> The variability of these results could, in part, be due to sample size, patient series heterogeneity, control populations, or technical differences in detecting HLA genotype.<sup>94</sup>

Other studies have shown associations between CMM and additional HLA alleles. Negative associations were found between *DQB1\*0302/0303* and CMM, whereas a positive association was found for *DQB1\*0501* in two Italian studies.<sup>99,100</sup> In Japanese patients with CMM, a positive association was found with *DQB1\*0302*, whereas negative associations were reported for *DRB1\*0802* and *DQA1\*0101/0104/0401*.<sup>101</sup> Finally, in a British population, an increase in the frequency of *DQB1\*0303* was found among CMM patients.<sup>102</sup>

Another candidate CMM susceptibility gene that has recently been investigated is the intracellular adhesion molecule-1 (*ICAM-1*). *ICAM-1* appears to play a role in the inflammatory and immune response through its interaction with the leukocyte integrins leukocyte function-associated antigen-1 (LFA-1) and Mac-1.<sup>103,104</sup> Higher levels of *ICAM-1* may therefore interfere with lymphocyte recognition of tumor cells and lymphocyte localization.<sup>104-106</sup> In one study, individuals carrying at least one variant *ICAM-1* SNP *R241* allele (conferring increased *ICAM-1* expression) versus wild-type *GG* genotype had a 4.3 relative risk of melanoma ( $P = .022$ ).<sup>103</sup> However, in this study, the CMM patient and control sample sizes each only numbered 59, so larger studies are needed to confirm this preliminary finding.

### DNA repair-related genes

UV irradiation is the major environmental risk factor for melanoma.<sup>36</sup> The body's main defense

against UV-induced DNA damage (cyclobutane pyrimidine dimers and 6-4 photoproducts) is the nucleotide excision repair pathway.<sup>107</sup> Many genes implicated in this pathway have been examined in relation to CMM (see Table I).

A recent meta-analysis found a significant association between the *XPD/ERCC2* SNP rs13181 variant C allele and CMM susceptibility, with an OR of 1.12 (95% CI, 1.03-1.21).<sup>8</sup> Povey et al<sup>107</sup> analyzed multiple genes, including *ERCC1*, *XPD*, *XPF*, *XPG*, *XRCC1*, *OGG1*, *XRCC3*, *GSTT1*, and *p53*, but only found significant associations between *ERCC1* and *XPF* and CMM, especially in CMM patients who are 50 years of age and younger (*ERCC1* OR = 1.59 and *XPF* OR = 1.69).<sup>107</sup> Another recent study found only one significant association out of 13 different polymorphisms in 8 DNA repair genes; in *XRCC3*; carriers of variant alleles had a decreased risk of CMM (OR 0.83, 95% CI, 0.79-0.98).<sup>108</sup> An additional study reported a significant association between *MGMT* haplotypes and CMM risk, with a greater risk observed among 84Phe or 143Val carriers, who have a lower alkylation-damage repair capacity due to the variant alleles.<sup>109</sup>

One study focused on the effects of two DNA repair genes (*XRCC1* and *APEX1*) and prognosis in CMM; a significant association was reported between the AA genotype of the R399Q *XRCC1* polymorphism and CMM, which showed a median overall survival of 24.4 years compared with 11.5 years for two other genotypes (hazard ratio of 0.40) and a median metastasis-free survival of 20.9 years versus 5.3 for two other genotypes (hazard ratio of 0.32) (see Table II).<sup>110</sup> This same study also found a significant association for -77 T>C *XRCC1* and survival following the first metastasis, with a hazard ratio of 1.73 (see Table II). An additional study found an association between *ERCC1* polymorphisms at codon 118 (TT and CT genotypes) and increased response to chemotherapy and overall survival in CMM (see Table II).<sup>91</sup>

Another study evaluated *MDM2* SNP309 (implicated in the p53 tumor suppressor pathway) and its role in CMM and reported that at ages younger than 50 years, women with a *GG* genotype had a 3.89 times greater chance of being diagnosed with CMM than women with *TG* or *TT* genotypes ( $P = .01$ ).<sup>111</sup> Additionally, one study reviewed the role of *BRCA1* and *BRCA2* in CMM patients and found that no mutation was present in any of the 92 familial melanoma cases.<sup>112</sup> Li et al<sup>113</sup> reported a significant association between *APEX1* and CMM, with an OR of 0.59 (95% CI, 0.42-0.83), but another study by the same authors<sup>114</sup> failed to find an association for *XPC*.

An additional study analyzed the effects of various genes involved in regulation of telomerase activity, which is widely known to cause elongation of telomeres in tumor cells. This study found a significant association between SNPs in the *TERT* and *TRF1* genes and increased CMM risk, with the OR ranging from 1.43 to 1.87.<sup>115</sup> This study<sup>115</sup> also found a decreased risk of CMM in the *TERT-CLPTMIL* locus, with an OR of 0.73 (see Table I).

### Metabolism-related genes

Epidemiological studies over the past few decades have indicated that CMM is related to both UV exposure and host factors. Since UV irradiation generates reactive oxygen species that are damaging to DNA, the extent of damage done and potential for carcinogenesis is dependent on how the body metabolizes and detoxifies these oxygen radicals.<sup>9</sup> A variety of host metabolic factors have been investigated as possible CMM susceptibility genes for this reason (see Table I).

Three studies have reviewed inactivating polymorphisms in the cytochrome P450-debrisoquine hydroxylase locus (*CYP2D6*) in relation to melanoma, with homozygotes for mutant alleles being termed poor metabolizers.<sup>1</sup> The first study found no significant difference in the frequency of *CYP2D6* poor metabolizers between 127 CMM cases and 720 controls, but did find a significant increase in the number of mutant alleles in the CMM cases ( $P = .02$ ).<sup>116</sup> A second study found no significant associations between the number of *CYP2D6* mutant alleles among cases and controls in the Slovene population.<sup>117</sup> Strange et al<sup>118</sup> reported a significant increase in frequency of homozygosity for the mutant *CYP2D6* alleles in the 333 CMM cases compared with the 467 controls (OR, 2.2; 95% CI, 1.2-3.9) among Northern European Caucasians. Larger studies are warranted to confirm a true increase in CMM susceptibility with mutant *CYP2D6* alleles.<sup>119</sup>

The glutathione S-transferases (GSTs), involved in detoxification of drugs and potential carcinogens, have also been examined in CMM.<sup>120-122</sup> Homozygosity of null alleles in *GSTM1* and *GSTT1* genes result in decreased production of the corresponding enzymes and increased cancer susceptibility.<sup>120,123</sup> One study in the Swedish population examined the *GSTM1 null*, *GSTT1 null*, and *GSTP1 GG* genotypes, and no significant differences were seen in frequency of these genotypes between CMM patients and controls.<sup>120</sup> However, there was a significant association between the *GSTM1 null* genotype and tumor Breslow thickness greater than 2.5 mm (poor prognostic indicator) as well as a particular type of CMM, nodular (see Table II).<sup>120</sup> In

contrast, an earlier study did report a significantly higher portion of controls ( $n = 47$ ) with measurable *GSTM1* levels than CMM patients ( $n = 197$ ), with 51% versus 42%, respectively ( $P = .002$ ).<sup>9</sup> Kanetsky et al<sup>124</sup> did not find an association between *GSTM1 null* or *GSTT1 null* genotypes and CMM patients, but did report that CMM patients with red or blond hair were twice as likely to carry *GSTM1 null* and nearly 10-fold more likely to carry both *GSTM1 null* and *GSTT1 null* genotypes compared with controls without CMM. A more recent study did not find any statistically significant difference between *GSTM1*, *GSTT1*, or *GSTP1* genotypes in CMM patients and healthy controls in the Slovenian population.<sup>125</sup>

### Vitamin D receptor polymorphisms

There is evidence to suggest that vitamin D exerts antiproliferative, prodifferentiation, proapoptotic, and antiangiogenic effects on cells, including melanoma cells.<sup>126-129</sup> The vitamin D receptor (VDR) is an intracellular hormone receptor that binds the biologically active form of vitamin D, 1,25-dihydroxyvitamin D or calcitriol, to mediate its effects by interacting with response elements of target genes.<sup>128</sup> The *VDR* gene is located on chromosome 12p12-q14 and has been investigated as a CMM susceptibility gene, with the idea that certain SNPs, including *FokI*, *BsmI*, *TaqI*, *Apal*, and *A-1012G* may alter activity of the VDR and, in effect, act similarly to a systemic deficiency of vitamin D, which has been associated with an increased risk of certain cancers (including CMM) in many epidemiological studies (see Table I).<sup>10,130,131</sup>

In an early study, the *FokI*, but not *TaqI* polymorphism was associated with an altered susceptibility to CMM, with the wild-type *FF* genotype associated with a risk reduction of 23.7%.<sup>9</sup> Another relatively early study reported a novel association between the variant *BsmI* genotype *bb* and increased CMM susceptibility ( $P = .02$ ), but no association for *FokI* or *A-1012G*.<sup>126</sup> However, Halsall et al<sup>132</sup> found that the *A* allele of *A-1012G* was more frequent in CMM patients than in controls ( $P = .011$ ). A case-control study investigated the combined effects of *TaqI*, *BsmI*, and *FokI* polymorphisms on CMM risk and reported that the *tBf* and *tBf* haplotypes were both significantly associated with a reduced melanoma risk (OR of 0.52 and 0.51, respectively) when the putative risk alleles, *T*, *b*, and *f* were used as a referent.<sup>133</sup> An earlier study exhibited similar results when the *TaqI* and *FokI* polymorphisms were examined, with a reduced risk of CMM with the *Tt* and *tt* genotypes (OR of 0.70) compared with the *TT* genotype, and an increased risk with the *Ff* genotype (OR of 1.32) compared with the *FF* genotype.<sup>131</sup> In a



meta-analysis of 10 studies, the summary relative risks (SRRs) for the *FokI* polymorphisms *Ff* and *ff* versus wild-type were 1.21 and 1.21, respectively, whereas the SRR for the *BsmI* polymorphisms *Bb* and *BB* versus wild-type were 0.78 and 0.75, respectively.<sup>134</sup>

In addition to their role in CMM susceptibility, VDR polymorphisms have also been implicated in CMM prognosis. Hutchinson et al<sup>10</sup> reported that homozygosity for both *FokI* and *TaqI* variant alleles was significantly associated with thicker tumors ( $\geq 3.5$  mm;  $P = .001$ ; OR of 31.5), with Breslow thickness being the most important prognostic indicator in CMM (see Table II). Another study reported a significant association between *BsmI* polymorphisms and Breslow thickness, with the variant *bb* alleles being associated with tumors of the greatest mean Breslow thickness and the wild-type *BB* alleles correlating with the lowest mean Breslow thickness tumors (see Table II).<sup>126</sup> In a recent cohort study, Newton-Bishop et al<sup>135</sup> reported that higher 25-hydroxyvitamin D<sub>3</sub> levels were associated with lower Breslow thickness at diagnosis ( $P = .002$ ) and were independently protective of relapse and death.

## CONCLUSIONS

While two of the rare, high-penetrance genetic variants (*CDKN2A* and *CDK4*) responsible for some of the familial melanomas have been identified, we are far from understanding all of the genetics behind the majority of CMMs. It is likely that a large number of SNPs, each with a small effect and low penetrance, in addition to the small number of large effect, high-penetrance SNPs, are responsible for CMM risk. Unfortunately, it is likely to be challenging and technically difficult to identify the remaining polymorphisms that have an effect on CMM susceptibility and prognosis. In addition, the published research to date has various flaws that make it difficult to draw accurate conclusions about the effects of these low-penetrance SNPs, such as small sample size, hospital-based controls, failure to record a major confounding variable, for example, UV irradiation exposure, a small number of studies on various individual SNPs, as well as others. In addition, it should be noted that even GWAS have significant limitations, including the fact that many of the platforms used in the studies do not always cover the exact genes mentioned. Rather, the SNPs are found near the gene loci and thus remain speculative.

This review was intended to provide a summary of the research on the known germline SNPs that have been implicated in CMM susceptibility and prognosis. The ultimate goal of this research is to pinpoint patients who are known to have a genetic

predisposition to CMM so that steps can be taken to ensure early detection or more intense treatment in individuals with poor prognostic indicators so as to minimize morbidity and mortality from this deadly skin cancer.

However, to date, science is far from achieving this, as both the American Society of Clinical Oncology and the Melanoma Genetics Consortium (GenoMEL) do not advocate the use of routine clinical genetic testing until there is further evidence to indicate clinical utility.<sup>136,137</sup> With no widely accepted guidelines in place, it remains difficult for clinicians to discuss the risks and benefits of genetic testing with patients.<sup>138</sup> Reasons for recommending against routine genetic testing are as follows: (1) most families with hereditary CMM have no detected mutations; (2) our understanding of CMM risk to carriers is limited; (3) other factors (ie, UV irradiation) appear to affect penetrance; and (4) negative testing may provide false reassurance, as up to 9% of noncarriers in *CDKN2A*-positive families have been reported to develop CMM.<sup>136</sup> So, where does this leave clinicians? Should anyone be tested? What about genetic testing for prognosis? These are important questions and, unfortunately, clinicians are without definitive answers at this time. Additional studies are needed and warranted to further evaluate the role of the above-mentioned and other yet-to-be-discovered SNPs, in order to make genetic testing for CMM possible.

The authors thank Dr Erin Warshaw for her guidance in the submission process.

## REFERENCES

1. Hayward NK. Genetics of melanoma predisposition. *Oncogene* 2003;22:3053-62.
2. Zuo L, Weger J, Yang Q, Goldstein AM, Tucker MA, Walker GJ, et al. Germline mutations in the p16INK42 binding domain of CDK4 in familial melanoma. *Nat Genet* 1996;12:97-9.
3. Soufir N, Avril MF, Chompret A, Demeais F, Bombléd J, Spatz A, et al. Prevalence of p16 and CDK4 germline mutations in 48 melanoma-prone families in France. The French Familial Melanoma Study Group. *Hum Mol Genet* 1998;7:209-16.
4. Hayward N. New developments in melanoma genetics. *Curr Oncol Rep* 2000;2:300-6.
5. Bishop DT, Demeais F, Iles MM, Harland M, Taylor JC, Corda E, et al. Genome-wide association study identifies three loci associated with melanoma risk. *Nat Genet* 2009;41:920-8.
6. Gandini S, Sera F, Cattaruzza MS, Pasquini P, Abeni D, Boyle P, et al. Meta-analysis of risk factors for cutaneous melanoma: I. Common and atypical naevi. *Eur J Cancer* 2005;41:28-44.
7. Howell WM, Turner SJ, Bateman AC, Theaker JM. IL-10 promoter polymorphisms influence tumour development in cutaneous malignant melanoma. *Genes Immun* 2001;2:25-31.
8. Mocellin S, Verdi D, Nitti D. DNA repair gene polymorphisms and risk of cutaneous melanoma: a systematic review and meta-analysis. *Carcinogenesis* 2009;30:1735-43.
9. Lafuente A, Molina R, Palou J, Castel T, Moral A, Trias M. Phenotype of glutathione S-transferase Mu (GSTM1) and susceptibility to malignant melanoma. *Br J Cancer* 1995;72:324-6.

10. Hutchinson PE, Osborne JE, Lear JT, Smith AG, Bowers PW, Morris PN, et al. Vitamin D receptor polymorphisms are associated with altered prognosis in patients with malignant melanoma. *Clin Cancer Res* 2000;6:498-504.
11. Schoof N, von Bonin F, König IR, Mössner R, Krüger U, Reich K, et al. Distal and proximal interleukin (IL)-10 promoter polymorphisms associated with risk of cutaneous melanoma development: a case-control study. *Genes Immun* 2009;10:586-90.
12. Pho LN, Leachman SA. Genetics of pigmentation and melanoma predisposition. *G Ital Dermatol Venereol* 2010;145:37-45.
13. Sondak VK, Chang AE. Melanoma and human leukocyte antigen status: the missing link? *Ann Surg Oncol* 2002;9:723-4.
14. Yeh I, Bastian BC. Genome-wide associations: studies for melanoma and nevi. *Pigment Cell Melanoma Res* 2009;22:527-8.
15. Yang XR, Pfeiffer RM, Wheeler W, Yeager M, Chanock S, Turner MA, et al. Identification of modifier genes for cutaneous malignant melanoma in melanoma-prone families with and without CDKN2A mutations. *Int J Cancer* 2009;125:2912-7.
16. Udayakumar D, Tsao H. Melanoma genetics: an update on risk-associated genes. *Hematol Oncol Clin North Am* 2009;23:415-29.
17. Nelson AA, Tsao H. Melanoma and genetics. *Clin Dermatol* 2009;27:46-52.
18. Bishop JN, Harland M, Randerson-Moor J, Bishop DT. Management of familial melanoma. *Lancet Oncol* 2007;8:46-54.
19. Stahl S, Bar-Meir E, Friedman E, Regev E, Orenstein A, Winkler E. Genetics in melanoma. *Isr Med Assoc J* 2004;6:774-7.
20. Sharpless NE, Chin L. The INK4a/ARF locus and melanoma. *Oncogene* 2003;22:3092-8.
21. Hussussian CJ, Struewing JP, Goldstein AM, Higgins PA, Ally DS, Sheahan MD, et al. Germline p16 mutations in familial melanoma. *Nat Genet* 1994;8:15-21.
22. Kamb A, Shattuck-Eidens D, Eeles R, Liu Q, Gruis NA, Ding W, et al. Analysis of the p16 gene (CDKN2) as a candidate for the chromosome 9p melanoma susceptibility locus. *Nat Genet* 1994;8:23-6.
23. Holland EA, Beaton SC, Becker TM, Grulet OM, Peters BA, Rizos H, et al. Analysis of the p16 gene, CDKN2, in 17 Australian melanoma kindreds. *Oncogene* 1995;11:2289-94.
24. Liu L, Lassam NJ, Slingerland JM, Bailey D, Cole D, Jenkins R, et al. Germline p16INK4A mutation and protein dysfunction in a family with inherited melanoma. *Oncogene* 1995;11:405-12.
25. Walker GJ, Hussussian CJ, Flores JF, Glendening JM, Haluska FG, Dracopoli NC, et al. Mutations of the CDKN2/p16INK4 gene in Australian melanoma kindreds. *Hum Mol Genet* 1995;4:1845-52.
26. Kamb A, Gruis NA, Weaver-Feldhaus J, Liu Q, Harshman K, Tavitgian SV, et al. A cell cycle regulator potentially involved in genesis of many tumor types. *Science* 1994;264:436-40.
27. Whelan AJ, Bartsch D, Goodfellow PJ. Brief Report: a familial syndrome of pancreatic cancer and melanoma with a mutation in the CDKN2 tumor-suppressor gene. *N Engl J Med* 1995;333:975-7.
28. Goldstein AM, Fraser MC, Struewing JP, Hussussian CJ, Ranade K, Zimetkin DP, et al. Increased risk of pancreatic cancer in melanoma-prone kindreds with p16INK4 mutations. *N Engl J Med* 1995;333:970-1.
29. Sharpless NE, Chin L. The INK4a/ARF locus and melanoma. *Oncogene* 2003;22:3092-8.
30. Tsoa H, Benoit E, Sober AJ, Thiele C, Haluska FG. Novel mutations in the p16/CDKN2A binding region of the cyclin-dependent kinase-4 gene. *Cancer Res* 1998;58:109-13.
31. Wagner SN, Wagner C, Briedigkeit L, Goos M. Homozygous deletion of the p16INK4a and the p15INK4b tumour suppressor genes in a subset of human sporadic cutaneous malignant melanoma. *Br J Dermatol* 1998;138:13-21.
32. Pomerantz J, Schreiber-Agus N, Liegeois NJ, Silverman A, Alland L, Chin L, et al. The Ink4a tumor suppressor gene product, p19Arf, interacts with MDM2 and neutralizes MDM2's inhibition of p53. *Cell* 1998;92:713-23.
33. Kamijo T, Weber JD, Zambetti G, Zindy F, Roussel MF, Sherr CJ. Functional and physical interactions of the ARF tumor suppressor with p53 and Mdm2. *Proc Natl Acad Sci U S A* 1998;95:8292-7.
34. Stott FJ, Bates S, James MC, McConnell BB, Starborg M, Brookes S, et al. The alternative product from the human CDKN2A locus, p14(ARF), participates in a regulatory feedback loop with p53 and MDM2. *EMBO J* 1998;17:5001-14.
35. Zhang Y, Xiong Y, Yarbrough WG. ARF promotes MDM2 degradation and stabilizes p53: ARF-INK4a locus deletion impairs both the RB and p53 tumor suppression pathways. *Cell* 1998;92:725-34.
36. Meyle KD, Gulberg P. Genetic risk factors for melanoma. *Hum Genet* 2009;126:499-510.
37. Goldstein AM, Chan M, Harland M, Gillanders EM, Hayward NK, Avril MF, et al. High-risk melanoma susceptibility genes and pancreatic cancer, neural system tumors, and uveal melanoma across GenoMEL. *Cancer Res* 2006;66:9818-28.
38. Helsing P, Nymoen DA, Ariansen S, Steine SJ, Maehle L, Aamdal S, et al. Population-based prevalence of CDKN2A and CDK4 mutations in patients with multiple primary melanomas. *Genes Chromosomes Cancer* 2008;47:175-84.
39. Molven A, Grimstvedt MB, Steine SJ, Harland M, Avril MF, Hayward NK, et al. A large Norwegian family with inherited malignant melanoma, multiple atypical nevi, and CDK4 mutations. *Genes Chromosomes Cancer* 2005;44:10-8.
40. Pjanova D, Molven A, Akslen LA, Engele L, Streinert B, Azarjana K, et al. Identification of a CDK4 R24H mutation-positive melanoma family by analysis of early-onset melanoma patients in Latvia. *Melanoma Res* 2009;19:119-22.
41. Duffy DL, Box NF, Chen W, Palmer JS, Montgomery GW, James MR, et al. Interactive effects of MC1R and OCA2 on melanoma risk phenotypes. *Hum Mol Genet* 2004;13:447-61.
42. Busca R, Ballotti R. Cyclic AMP: a key messenger in the regulation of skin pigmentation. *Pigment Cell Res* 2000;13:60-9.
43. Valverde P, Healy E, Sikkink S, Haldane F, Thody AJ, Carothers A, et al. The Asp84Glu variant of the melanocortin 1 receptor (MC1R) is associated with melanoma. *Hum Mol Genet* 1996;5:1663-6.
44. Robbins LS, Nadeau JH, Johnson KR, Kelly MA, Roselli-Reh fuss L, Baack E, et al. Pigmentation phenotypes of variant extension locus alleles result from point mutations that alter MSH receptor function. *Cell* 1993;72:827-34.
45. Jackson IJ. Molecular and developmental genetics of mouse coat color. *Ann Rev Genet* 1994;28:189-217.
46. Sturm RA, Duffy DL, Box NF, Chen W, Smit DJ, Brown DL, et al. The role of melanocortin-1 receptor polymorphism in skin cancer risk phenotypes. *Pigment Cell Res* 2003;16:266-72.
47. Suzuki I, Im S, Tada A, Scott C, Akcali C, Davis MB, et al. Participation of the melanocortin-1 receptor in the UV control of pigmentation. *J Invest Dermatol Symp Proc* 1999;4:29-34.

48. Sturm RA, Teasdale RD, Box NF. Human pigmentation genes: identification, structure and consequences of polymorphic variation. *Gene* 2001;277:49-62.
49. Frandberg PA, Doufexis M, Kapas S, Chhājlani V. Human pigmentation phenotype: a point mutation generates non-functional MSH receptor. *Biochem Biophys Res Commun* 1998;245:490-2.
50. Schioth HB, Sr Phillips, Rudzish R, Birch-Machin MA, Wikberg JE, Rees JL. Loss of function mutations of the human melanocortin 1 receptor are common and are associated with red hair. *Biochem Biophys Res Commun* 1999;260:488-91.
51. Thody AJ, Higgins EM, Wakamatsu K, Ito S, Burchill SA, Marks JM. Pheomelanin as well as eumelanin is present in human epidermis. *J Invest Dermatol* 1991;97:340-4.
52. Napolitano A, Vincensi MR, Di Donato P, Monfrecola G, Prota G. Microanalysis of melanins in mammalian hair by alkaline hydrogen peroxide degradation: identification of a new structural marker of pheomelanins. *J Invest Dermatol* 2000;114:1141-7.
53. Robinson S, Dixon S, Augus S, Diffey B, Wakamatsu K, Ito S, et al. Protection against UVR involves MC1R-mediated non-pigmentary and pigmentary mechanisms in vivo. *J Invest Dermatol* 2010;130:1904-13.
54. Valverde P, Healy E, Jackson I, Rees JL, Thody AJ. Variants of the melanocyte-stimulating hormone receptor gene are associated with red hair and fair skin in humans. *Nat Genet* 1995;11:328-30.
55. Raimondi S, Sera F, Gandini S, Iodice S, Caini S, Maisonneuve P, et al. MC1R variants, melanoma and red hair color phenotype: a meta-analysis. *Int J Cancer* 2008;122:2753-60.
56. Galore-Haskel G, Azizi E, Mohamdi H, Scope A, Chaudru V, Laitman Y, et al. MC1R variant alleles and malignant melanoma risk in Israel. *Eur J Cancer* 2009;45:2015-22.
57. Palmer JS, Duffy DL, Box NF, Aitken JF, O'Gorman LE, Green AC, et al. Melanocortin-1 receptor polymorphisms and risk of melanoma: is the association explained solely by pigmentation phenotype? *Am J Hum Genet* 2000;66:176-86.
58. Mössner R, Anders N, König I, Krüger U, Schmidt D, Berking C, et al. Variations of the melanocortin-1 receptor and the glutathione-S transferase T1 and M1 genes in cutaneous malignant melanoma. *Arch Dermatol Res* 2007;298:371-9.
59. Chatzinasiou F, Lill CM, Kypreou K, Stefanaki I, Nicolaou V, Spyrou G, et al. Comprehensive field synopsis and systematic meta-analyses of genetic association studies in cutaneous melanoma. *J Natl Cancer Inst* 2011;103:1227-35.
60. Box NF, Duffy DL, Chen W, Stark M, Martin NG, Sturm RA, et al. MC1R genotype modifies risk of melanoma in families segregating CDKN2A mutations. *Am J Hum Genet* 2001;69:765-73.
61. van der Velden PA, Sandkuijl LA, Bergman W, Pavel S, van Mourik L, Frants RR, et al. Melanocortin-1 receptor variant R151C modifies melanoma risk in Dutch families with melanoma. *Am J Hum Genet* 2001;69:774-9.
62. Fernandez LP, Milne RL, Pita G, Floristan U, Sendagorta E, Feito M, et al. Pigmentation-related genes and their implication in malignant melanoma susceptibility. *Exp Dermatol* 2009;18:634-42.
63. Gudbjartsson DF, Sulem P, Stacey SN, Goldstein AM, Rafnar T, Sigurgeirsson B, et al. ASIP and TYR pigmentation variants associate with cutaneous melanoma and basal cell carcinoma. *Nat Genet* 2008;48:886-91.
64. Sulem P, Gudbjartsson DF, Stacey SN, Helgason A, Rafnar T, Jakobsdottir M, et al. Two newly identified genetic determinants of pigmentation in Europeans. *Nat Genet* 2008;40:835-7.
65. Nan H, Kraft P, Hunter DJ, Han J. Genetic variants in pigmentation genes, pigmentary phenotypes, and risk of skin cancer in Caucasians. *Int J Cancer* 2009;125:909-17.
66. Brown KM, Macgregor S, Montgomery GW, Craig DW, Zhao ZZ, Iyadurai K, et al. Common sequence variants on 20q11.22 confer melanoma susceptibility. *Nat Genet* 2008;40:838-40.
67. Kanetsky PA, Swoyer J, Panossian S, Holmes R, Guerry D, Rebbeck TR. A polymorphism in the agouti signaling protein gene is associated with human pigmentation. *Am J Hum Genet* 2002;70:770-5.
68. Berson JF, Frank DW, Calvo PA, Bieler BM, Marks MS. A common temperature-sensitive allelic form of human tyrosinase is retained in the endoplasmic reticulum at the nonpermissive temperature. *J Biol Chem* 2000;275:12281-9.
69. Jannot AS, Meziani R, Bertrand G, Gérard B, Descamps V, Archimbaud A, et al. Allele variations in the OCA2 gene (pink-eyed-dilution locus) are associated with genetic susceptibility to melanoma. *Eur J Hum Genet* 2005;13:913-20.
70. Fernandez LP, Milne RL, Pita G, Avilés JA, Lázaro P, Benítez J, et al. SLC45A2: a novel malignant melanoma-associated gene. *Hum Mutat* 2008;29:1161-7.
71. Guedj M, Bourillon A, Combadières C, Rodero M, Dieudé P, Descamps V, et al. Variants of the MATP/SLC45A2 gene are protective for melanoma in the French population. *Hum Mutat* 2008;29:1154-60.
72. Duffy DL, Zhao ZZ, Sturm RA, Hayward NK, Martin NG, Montgomery GW. Multiple pigmentation gene polymorphisms account for a substantial proportion of risk of cutaneous malignant melanoma. *J Invest Dermatol* 2010;130:520-8.
73. Whiteman DC, Whiteman CA, Green AC. Childhood sun exposure as a risk factor for melanoma: a systematic review of epidemiological studies. *Cancer Causes Control* 2001;12:69-82.
74. Greene MH, Clark WH Jr, Tucker MA, Kraemer KH, Elder DE, Fraser MC. High risk of malignant melanoma in melanoma-prone families with dysplastic nevi. *Ann Intern Med* 1985;102:458-65.
75. Falchi M, Bataille V, Hayward NK, Duffy DL, Bishop JA, Pastinen T, et al. Genome-wide association study identifies variants at 9p21 and 22q13 associated with development of cutaneous nevi. *Nat Genet* 2009;41:915-9.
76. Nan H, Xu M, Zhang J, Zhang M, Kraft P, Qureshi AA, et al. Genome-wide associated study identifies nidogen 1 (NID1) as a susceptibility locus to cutaneous nevi and melanoma risk. *Hum Mol Genet* 2011;20:2673-9.
77. Han J, Kraft P, Nan H, Guo Q, Chen C, Qureshi A, et al. A genome-wide association study identifies novel alleles associated with hair color and skin pigmentation. *PLoS Genet* 2008;4:1-11.
78. Sulem P, Gudbjartsson DF, Stacey S, Helgason A, Rafnar T, Magnusson KP, et al. Genetic determinants of hair, eye and skin pigmentation in Europeans. *Nat Genet* 2007;39:1443-52.
79. Wang ZQ, Si L, Tang Q, Lin D, Fu Z, Zhang J, et al. Gain-of-function mutation of KIT ligand on melanin synthesis causes familial progressive hyperpigmentation. *Am J Hum Genet* 2009;84:672-7.
80. Howell WM, Turner SJ, Theaker JM, Bateman AC. Cytokine gene single nucleotide polymorphisms and susceptibility to and prognosis in cutaneous malignant melanoma. *Eur J Immunogenet* 2003;30:409-14.
81. Wolfel T, Hauer M, Klehmann E, Brichard V, Ackermann B, Knuth A, et al. Analysis of antigens recognised on human melanoma cells by A2-restricted cytolytic T lymphocytes (CTL). *Int J Cancer* 1993;55:237.

82. Gu F, Qureshi AA, Niu T, Kraft P, Guo Q, Hunter DJ, et al. Interleukin and interleukin receptor gene polymorphisms and susceptibility to melanoma. *Melanoma Res* 2008;18:330-5.
83. Penn I. Depressed immunity and skin cancer. *Immunol Today* 1984;5:291-3.
84. Alonso R, Suarez A, Castro P, Lacave AJ, Gutierrez C. Influence of interleukin-10 genetic polymorphism on survival rates in melanoma patients with advanced disease. *Melanoma Res* 2005;15:53-60.
85. Turner DM, Williams DM, Sankaran D, Lazarus M, Sinnott PJ, Hutchinson IV. An investigation of polymorphisms in the interleukin-10 gene promoter. *Eur J Immunogenet* 1997;24:1-8.
86. Hajeer AH, Lazarus M, Turner D, Mageed RA, Vencovsky J, Sinnott P, et al. IL-10 gene promoter polymorphisms in rheumatoid arthritis. *Scand J Rheumatol* 1998;27:142-5.
87. Martinez-Escribano JA, Moya-Quiles MR, Muro M, Montes-Ares O, Hernández-Caselles T, Frías JF, et al. Interleukin-10, interleukin-6 and interferon-gamma gene polymorphisms in melanoma patients. *Melanoma Res* 2002;12:465-9.
88. Nikolova PN, Pawelec GP, Mihailova SM, Ivanova MI, Myhalova AP, Baltadjieva DN, et al. Association of cytokine gene polymorphisms with malignant melanoma in Caucasian population. *Cancer Immunol Immunother* 2007;56:371-9.
89. Dummer W, Becker JC, Schwaaf A, Leverkus M, Moll T, Bröcker EB. Elevated serum levels of interleukin-10 in patients with metastatic malignant melanoma. *Melanoma Res* 1995;5:67-8.
90. Yue FY, Dummer R, Geersten R, Hofbauer G, Laine E, Manolio S, et al. Interleukin-10 is a growth factor for human melanoma cells and down-regulates HLA class-I, HLA class-II and ICAM-1 molecules. *Int J Cancer* 1997;71:630-7.
91. Liu D, O'Day SJ, Yang D, Boasberg P, Milford R, Kristedja T, et al. Impact of gene polymorphisms on clinical outcome for stage IV melanoma patients treated with biochemotherapy: an exploratory study. *Clin Cancer Res* 2005;11:1237-46.
92. von Euw EM, Barrio MM, Furman D, Levy EM, Bianchini M, Pequillet I, et al. A phase I clinical study of vaccination of melanoma patients with dendritic cells loaded with allogeneic apoptotic/necrotic melanoma cells. Analysis of toxicity and immune response to the vaccine and of IL-10 -1082 promoter genotype as predictor of disease progression. *J Transl Med* 2008;6:1-14.
93. Howell WM, Turner SJ, Collins A, Bateman AC, Theaker JM. Influence of TNF $\alpha$  and LT $\alpha$  single nucleotide polymorphisms on susceptibility to and prognosis in cutaneous malignant melanoma in the British population. *Eur J Immunogenet* 2002;29:17-23.
94. Planelles D, Nagore E, Moret A, Botella-Estrada R, Villa E, Guillén C, et al. HLA class II polymorphisms in Spanish melanoma patients: homozygosity for HLA-DQA1 locus can be a potential melanoma risk factor. *Br J Dermatol* 2006;154:261-6.
95. Lee JE, Loflin PT, Laud PR, Lu M, Reveille JD, Lawlor DA. The human leukocyte antigen TAP2 gene defines the centromeric limit of melanoma susceptibility on chromosome 6p. *Tissue Antigens* 1996;47:117-21.
96. Lee JE, Reveille JD, Ross MI, Platsoucas CD. HLA-DQB1\*0301 association with increased cutaneous melanoma risk. *Int J Cancer* 1994;59:510-3.
97. Lee JE, Lu M, Mansfield PF, Platsoucas CD, Reveille JD, Ross MI. Malignant melanoma: relationship of the human leukocyte antigen class II gene DQB1\*0301 to disease recurrence in American Joint Committee on Cancer stage I or II. *Cancer* 1996;78:758-63.
98. Marincola FM, Shamamian P, Rivoltini L, Salgaller M, Cormier J, Restifo NP, et al. HLA associations in the antitumor response against malignant melanoma. *J Immunother* 1995;18:242-52.
99. Lombardi ML, Mercurio O, Pirozzi G, Ionna F, Lombardi V, Mozzillo N, et al. Molecular analysis of HLA DRB1 and DQB1 polymorphism in Italian melanoma patients. *J Immunother* 1998;21:435-9.
100. Lulli P, Grammatico P, Brioli G, Catricalá C, Morellini M, Roccella M, et al. HLA-DR and -DQ alleles in Italian patients with melanoma. *Tissue Antigens* 1998;51:276-80.
101. Kageshita T, Naruse T, Hirai S, Ono T, Horikoshi T, Nakagawa H, et al. Molecular genetic analysis of HLA class II alleles in Japanese patients with melanoma. *Tissue Antigens* 1997;49:466-70.
102. Bateman AC, Turner SJ, Theaker JM, Howell WM. HLA-DQB1\*0303 and \*0301 alleles influence susceptibility to and prognosis in cutaneous malignant melanoma in the British Caucasian population. *Tissue Antigens* 1998;52:67-73.
103. Vinceti M, Pellacani G, Casali B, Malagoli C, Nicoli D, Farnetti E, et al. High risk of cutaneous melanoma amongst carriers of the intercellular adhesion molecule-1 R241 allele. *Melanoma Res* 2006;16:93-6.
104. van de Stolpe A, van der Saag PT. Intercellular adhesion molecule-1. *J Mol Med* 1996;74:13-33.
105. Viganò P, Infantino M, Lattuada D, Lauletta R, Ponti E, Somigliana E, et al. Intercellular adhesion molecule-1 (ICAM-1) gene polymorphisms in endometriosis. *Mol Hum Reprod* 2003;9:47-52.
106. Diamond MS, Staunton DE, Marlin SD, Springer TA. Binding of the integrin Mac-1 (CD11b/CD18) to the third immunoglobulin-like domain of ICAM-1 (CD54) and its regulation by glycosylation. *Cell* 1991;65:961-71.
107. Povey JE, Darakhshan F, Robertson K, Bisset Y, Mekky M, Rees J, et al. DNA repair gene polymorphisms and genetic predisposition to cutaneous melanoma. *Carcinogenesis* 2007;28:1087-93.
108. Figl A, Scherer D, Nagore E, Bermejo JL, Botella-Estrada R, Gast A, et al. Single-nucleotide polymorphisms in DNA-repair genes and cutaneous melanoma. *Mutat Res* 2010;702(1):8-16.
109. Gu F, Qureshi AA, Kraft P, Guo Q, Hunter DJ, Han J. Polymorphisms in genes involved in DNA repair, cell growth, oxidative stress, and inflammatory response and melanoma risk. *Br J Dermatol* 2009;161:209-12.
110. Figl A, Scherer D, Nagore E, Bermejo JL, Dickes E, Thirumaran RK, et al. Single nucleotide polymorphisms in DNA repair genes XRCC1 and APEX1 in progression and survival of primary cutaneous melanoma patients. *Mutat Res* 2009;661:78-84.
111. Firoz EF, Warycha M, Zakrzewski J, Pollens D, Wang G, Shapiro R, et al. Association of MDM2 SNP309, age of onset, and gender in cutaneous melanoma. *Clin Cancer Res* 2009;15:2573-80.
112. Kadouri L, Temper M, Grenader T, Abeliovich D, Hamburger T, Peretz T, et al. Absence of founder BRCA1 and BRCA2 mutations in cutaneous malignant melanoma patients of Ashkenazi origin. *Fam Cancer* 2009;8:29-32.
113. Li C, Liu Z, Wang LE, Strom SS, Lee JE, Gershenwald JE, et al. Genetic variants of the ADPRT, XRCC1 and APE1 genes and risk of cutaneous melanoma. *Carcinogenesis* 2006;27:1894-901.
114. Li C, Hu Z, Liu Z, Strom SS, Gershenwald JE, Lee JE, et al. Polymorphisms in the DNA repair genes XPC, XPD, and XPG and risk of cutaneous melanoma: a case-control analysis. *Cancer Epidemiol Biomarkers Prev* 2006;15:2526-32.

115. Nan H, Qureshi AA, Prescott J, De Vivo I, Han J. Genetic variants in telomere-maintaining genes and skin cancer risk. *Hum Genet* 2011;129:247-53.
116. Wolf CR, Smith CA, Gough AC, Moss JE, Vallis KA, Howard G, et al. Relationship between the debrisoquine hydroxylase polymorphism and cancer susceptibility. *Carcinogenesis* 1992;13:1035-8.
117. Dolzan V, Rudolf Z, Breskvar K. Human CYP2D6 gene polymorphism in Slovene cancer patients and healthy controls. *Carcinogenesis* 1995;16:2675-8.
118. Strange RC, Ellison T, Ichii-Jones F, Bath J, Hoban P, Lear JT, et al. Cytochrome P450 CYP2D6 genotypes: association with hair colour, Breslow thickness and melanocyte stimulating hormone receptor alleles in patients with malignant melanoma. *Pharmacogenetics* 1999;9:269-76.
119. Agúndez J. Cytochrome P450 gene polymorphism and cancer. *Current Drug Metab* 2004;5:211-24.
120. Bu H, Rosdahl I, Holmdahl-Källen K, Sun XF, Zhang H. Significance of glutathione S-transferases M1, T1, and P1 polymorphisms in Swedish melanoma patients. *Oncol Rep* 2007;17:859-64.
121. Lusini L, Tripodi SA, Rossi R, Giannerini F, Giustarini D, del Vecchio MT, et al. Altered glutathione antioxidant metabolism during tumor progression in human renal-cell carcinoma. *Int J Cancer* 2001;91:55-9.
122. Baars AJ, Breimer DD. The glutathione S-transferases: their role in detoxification and toxification of xenobiotics. *Ann Biol Clin* 1980;38:49-56.
123. Rebbeck TR. Molecular epidemiology of the human glutathione S-transferase genotypes GSTM1 and GSTT1 in cancer susceptibility. *Cancer Epidemiol Biomarkers Prev* 1997;6:733-74.
124. Kanetsky PA, Holmes R, Walker A, Najarian D, Swoyer J, Guerry D, et al. Interaction of glutathione S-transferase M1 and T1 genotypes and malignant melanoma. *Cancer Epidemiol Biomarkers Prev* 2001;10:509-13.
125. Dolzan V, Rudolf Z, Breskvar K. Genetic susceptibility to environmental carcinogenesis in Slovenian melanoma patients. *Acta Dermatovenol Alp Panonica Adriat* 2006;15:69-78.
126. Santonocito C, Capizzi R, Concolino MM, Lavieri MM, Paradisi A, Gentileschi S, et al. Association between cutaneous melanoma, Breslow thickness and vitamin D receptor Bsm1 polymorphism. *Br J Dermatol* 2007;156:277-82.
127. Seifert M, Rech M, Meineke V, Tilgen W, Reichrath J. Differential biological effects of 1,25-dihydroxyvitamin D3 on melanoma cell lines in vitro. *J Steroid Biochem Mol Biol* 2004;89-90:375-9.
128. Calvo MS, Whiting SJ, Barton CN. Vitamin D intake: a global perspective of current status. *J Nutr* 2005;135:310-6.
129. Raimondi S, Johansson H, Maisonneuve P, Gandini S. Review and meta-analysis on vitamin D receptor polymorphisms and cancer risk. *Carcinogenesis* 2009;30:1170-80.
130. Baker AR, McDonnell DP, Hughes M, Crisp TM, Mangelsdorf DJ, Haussler MR, et al. Cloning and expression of full-length cDNA encoding human vitamin D receptor. *Proc Natl Acad Sci U S A* 1988;85:3294-8.
131. Li C, Liu Z, Zhang Z, Strom SS, Gershenwald JE, Prieto VG, et al. Genetic variants of the vitamin D receptor gene alter risk of cutaneous melanoma. *J Invest Dermatol* 2007;127:276-80.
132. Halsall JA, Osborne JE, Epstein MP, Pringle JH, Hutchinson PE. The unfavorable effect of the A allele of the vitamin D receptor promoter polymorphism A-1012G has different mechanisms related to susceptibility and outcome of malignant melanoma. *Dermatoendocrinol* 2009;1:54-7.
133. Li C, Liu Z, Wang LE, Gershenwald JE, Lee JE, Prieto VG, et al. Haplotype and genotypes of the VDR gene and cutaneous melanoma risk in non-Hispanic whites in Texas: a case-control study. *Int J Cancer* 2008;122:2077-84.
134. Gandini S, Raimondi S, Gnagnarella P, Doré JF, Maisonneuve P, Testori A. Vitamin D and skin cancer: a meta-analysis. *Eur J Cancer* 2009;45:634-41.
135. Newton-Bishop JA, Beswick S, Randerson-Moor J, Chang YM, Affleck P, Elliott F, et al. Serum 25-hydroxyvitamin D3 levels are associated with Breslow thickness at presentation and survival from melanoma. *J Clin Oncol* 2009;27:5439-44.
136. Kefford R, Bishop JN, Tucker M, Bressac-de Paillerets B, Bianchi-Scarrá G, Bergman W, et al. Genetic testing for melanoma. *Lancet Oncol* 2002;3:653-4.
137. Kefford RF, Newton Bishop JA, Bergman W, Tucker MA. Counseling and DNA testing for individuals perceived to be genetically predisposed to melanoma: a consensus statement of the Melanoma Genetics Consortium. *J Clin Oncol* 1999;17:3245-51.
138. Lin J, Hocker TL, Singh M, Tsao H. Genetics of melanoma predisposition. *Br J Dermatol* 2008;159:286-91.