

# Physically Motivated Time-Series Photometry of AM CVn

TODO: Team Members

TODO: Submission Date

## Abstract

We observed AM Canum Venaticorum (AM CVn) in the V band on 2026 May 20 to test whether short-baseline CCD photometry can recover and refine its known ultracompact variability. We reduced 140 images with bias, dark, and star-masked flat-field calibration, then performed AAVSO-sequence differential photometry. After rejecting two artifact frames, 138 exposures spanning 111.20 min remained. A generic single-sine search detects a 525.06 s signal, close to the expected first harmonic of AM CVn’s positive superhump. A physically motivated double-wave model gives the best independent period,  $1055.21 \pm 8.58$  s after empirical error scaling. A fixed 1051.2 s superhump-plus-harmonic template is nearly as good, but does not beat the free model in BIC or cross-validation. The data therefore support detection of AM CVn’s known superhump/harmonic family, but they do not tighten the literature period constraints.

## 1 Introduction

AM CVn is the prototype of a class of hydrogen-deficient ultracompact interacting binaries. These systems contain an accreting white dwarf and a helium-rich donor, and their periods are short enough to connect accretion-disc physics, compact-binary evolution, and low-frequency gravitational-wave astrophysics [5, 3].

The period structure of AM CVn is subtle. Early photometry showed a double wave near 17.5 min and a strong 8.76 min harmonic [2]. Later spectroscopy established  $P_{\text{orb}} = 1028.7322$  s as the orbital period, while the dominant 1051.2 s photometric signal is interpreted as a positive superhump [1, 3]. Recent work uses this period structure to revisit AM CVn system parameters and period change [4].

Our goal is not only to detect variability, but to ask a sharper question: can our local 111 min light curve constrain the literature periods more tightly? To avoid overclaiming, we separate independent constraints from conditional constraints. Independent constraints use periods fitted freely from our data. Conditional constraints assume a literature-motivated physical identification, such as the known superhump or orbital signal.

## 2 Observations and Reduction

The observations were taken on 2026 May 20 in the V band. The AM CVn sequence contains 140 frames with exposure times of 35, 40, and 45 s. The reduction used late-night bias frames, exposure-matched darks, and a star-masked `flat_4s_masked_clipmean` flat. The final two frames showed severe artifacts and low counts, so they were excluded.

The science frames did not have a reliable WCS solution. We therefore used the AAVSO comparison sequence X42421BZ to map the target and comparison stars onto a good reference frame. Aperture photometry used a 6 px aperture and a 12–20 px sky annulus. Fluxes were

divided by exposure time before comparison, so frames with different exposure times could be combined. Each comparison star was normalized by its own median flux before constructing the ensemble, preventing the brightest comparison stars from dominating the differential light curve.

The final light curve contains 138 good points. It spans 111.20 min, or 6.35 cycles of a 1051.2 s superhump. The robust target scatter is 1.51 percent, while the median formal photometric error is 0.90 percent. This excess scatter motivates the empirical error scaling used below.

### 3 Analysis Methods

We first fit a model ladder. The baseline model is constant flux. The generic models are a free single sine and a free double-wave model containing a fundamental plus its first harmonic. We also ran multi-harmonic Lomb-Scargle periodograms: `nterms=2` is the periodogram analogue of the free double-wave model. The physically motivated models use literature periods: the 1028.7322 s orbital period, the 1051.2 s positive superhump, a 1011.4 s negative superhump, and a superhump-plus-harmonic template. Finally, we fit a beat-constrained precessing-disc proxy with  $f_{\text{sh}} = f_{\text{orb}} - f_{\text{prec}}$ . The 13.38 h disc precession period was not directly fit from the light curve because our run covers only 0.14 cycle of it.

For each model we report chi-square, BIC, RMS residual scatter, and time-blocked cross-validation error. For free periods, we compute profile  $\chi^2$  intervals and block-bootstrap uncertainties. Formal photon-noise errors underestimate the residual scatter, so the final period uncertainties use the best double-wave model's  $\sqrt{\chi^2_{\nu}} = 1.59$  scale factor. We also test stability by fitting the first and second halves separately, clipping outliers, dropping comparison stars one at a time, and checking residual periodograms. As final stress tests, we fit extra nuisance terms for time, tracking shifts, exposure state, and ensemble common mode; repeat the double-wave fit with robust loss functions; and rebuild the differential light curve from comparison-star subsets.

### 4 Results

The single-sine search recovers  $525.06 \pm 4.27$  s after empirical error scaling. Ordinary Lomb-Scargle detects the same harmonic at 525.08 s. When Lomb-Scargle is generalized to a two-term Fourier model, it recovers a fundamental near 1053.27 s. Thus Lomb-Scargle is useful as a detection tool, but it must be made multi-harmonic or followed by physical model fitting for AM CVn.

The best independent model is the free double-wave fit:

$$P_{\text{free}} = 1055.21 \pm 8.58 \text{ s.}$$

This is consistent with the known 1051.2 s superhump family. It is not precise enough to improve the published photometric or spectroscopic periods.

Table 1: AM CVn model ladder.  $\Delta\text{BIC}$  is measured relative to the best free double-wave model; positive values are worse.

| Model                            | Physical interpretation                            | Period (s) | RMS (%) | $\Delta\text{BIC}$ | CV RMS (%) |
|----------------------------------|----------------------------------------------------|------------|---------|--------------------|------------|
| free double wave                 | fundamental + harmonic                             | 1055.21    | 1.43    | 0.00               | 1.52       |
| beat-constrained disk            | $f_{\text{sh}} = f_{\text{orb}} - f_{\text{prec}}$ | 1055.22    | 1.43    | 0.00               | 1.52       |
| 1051.2 s + harmonic              | positive-superhump template                        | 1051.2     | 1.43    | 0.55               | 1.53       |
| free single sine                 | harmonic-only detection                            | 525.06     | 1.52    | 37.44              | 1.57       |
| orbital + positive + negative SH | three known clocks                                 | fixed      | 1.50    | 45.55              | 1.65       |
| orbital + positive SH            | two known clocks                                   | fixed      | 1.53    | 52.99              | 1.71       |
| positive SH only                 | 1051.2 s sine                                      | 1051.2     | 1.56    | 57.67              | 1.62       |
| orbital only                     | 1028.7322 s sine                                   | 1028.73    | 1.58    | 66.18              | 1.65       |

The most important result in Table 1 is that the beat-constrained precessing-disc proxy and the free double-wave model are effectively tied. This happens because a freely fitted precession period maps the known orbital period onto the best photometric double-wave period; the derived value is  $P_{\text{prec}} \simeq 11.39$  h. This captures the expected superhump physics better than a generic Fourier fit, but it is not an independent precession detection because the baseline covers far less than one precession cycle. The fixed 1051.2 s superhump-plus-harmonic template is also nearly tied, supporting the literature interpretation.




Figure 1: Physically motivated model-suite summary. The upper-left panel shows the AM CVn differential light curve and the best free double-wave model. The upper-right panel folds the same model. The lower-left panel shows that the residual periodogram has no significant remaining peak. The lower-right panel compares BIC values against the free double-wave fit.

The validation tests are mixed. Dropping one comparison star at a time changes the free double-wave period by only 5.2 s, and clipping  $|z| > 4$  outliers changes it by only 0.02 s. The residual periodogram has false-alarm probability 0.99, so no strong unmodeled short-period signal remains. However, the first-half and second-half fitted periods differ by 22.8 s. That split is not surprising with only about six superhump cycles, but it blocks any strong claim that the period is tightly constrained by our data alone.

We also tried three ways to squeeze more information out of the same light curve. First, adding nuisance regressors for time, image shifts, exposure time, number of comparison stars, and common-mode behavior can reduce the raw RMS from 1.43% to 1.41%, but BIC and blocked cross-validation get worse ( $\Delta\text{BIC} = 22.66$ ,  $\Delta\text{CV} = 0.97\%$  for the most flexible case). This indicates overfitting rather than a better physical constraint. Second, robust likelihoods do not sharpen the answer: soft- $L_1$ , Huber, and Cauchy losses shift the best period to 1057–1059 s and increase the ordinary RMS. Third, selected comparison-star subsets can shave the target RMS as low as 1.40%, but the fitted period moves by a few seconds and the control-star scatter does not justify replacing the full ensemble. These tests are useful because they show what does not work: the limiting uncertainty is the short baseline and cycle-to-cycle structure, not just Gaussian photometric noise.

The broadband photometry cannot test AM CVn’s S-wave emission directly. The S-wave is a spectroscopic velocity feature from the bright spot or stream-disc impact region. Testing it requires time-resolved spectroscopy or Doppler tomography, not V-band imaging. For the same reason, our photometry can support the known white-dwarf-binary interpretation but cannot independently prove or disprove the object class.




Figure 2: Comparison between our period constraints and literature periods. Our independent scaled uncertainty is about 8.6 s, far larger than the published precision of the orbital and superhump periods. The result is therefore consistent with the literature, not more constraining than it.

## 5 Conclusions

Our data clearly detect short-timescale AM CVn variability. A single-sine model finds a 525 s signal, while the physically preferred interpretation is a double-wave signal in the known 1051 s superhump family. The best independent period is  $1055.21 \pm 8.58$  s, and the literature-fixed 1051.2 s superhump-plus-harmonic template is nearly as good.

The result does not tighten the literature constraints. The physically motivated model does not beat the free double-wave model, the first/second-half stability test fails, and our independent uncertainty is hundreds of times larger than the best published period precision. Detrending, robust likelihoods, and comparison-star subset searches improve neither predictive performance nor period stability. The strongest defensible claim is therefore: our photometry independently recovers variability consistent with AM CVn’s known superhump/harmonic structure, but a longer multi-night baseline would be needed to improve published period constraints.

## Acknowledgments

We thank the PHYSICS 100 teaching staff for telescope access and reduction guidance. This work used AAVSO comparison-star resources and NASA ADS/arXiv literature records.

## References

- [1] Nelemans, G., Steeghs, D., & Groot, P. J. 2001. *Spectroscopic evidence for the binary nature of AM CVn*. MNRAS, 326, 621. <https://academic.oup.com/mnras/article/326/2/621/1006901>.
- [2] Patterson, J., Nather, R. E., Robinson, E. L., & Handler, F. 1979. *Increase in the orbital period of AM Canum Venaticorum*. ApJ, 232, 819. <https://articles.adsabs.harvard.edu/pdf/1979ApJ...232..819P>.
- [3] Roelofs, G. H. A., Groot, P. J., Nelemans, G., Marsh, T. R., & Steeghs, D. 2006. *Kinematics of the ultracompact helium accretor AM Canum Venaticorum*. MNRAS, 371, 1231. <https://academic.oup.com/mnras/article/371/3/1231/1007944>.
- [4] Smak, J. 2023. *AM CVn – System Parameters and Gravitational Waves*. Acta Astronomica, 73, 1. <https://arxiv.org/pdf/2312.16506>.
- [5] Solheim, J.-E. 2010. *AM CVn Stars: Status and Challenges*. PASP, 122, 1133.

## A Individual Contributions

TODO: Fill in team-member names and individual contributions before submission, as required by the project instructions.

## B Reproducibility

The model suite is generated with:

```
python scripts/amcvn_physical_model_suite.py --bootstrap 250
```

The generated CSV files use the `physical_*` prefix in the AM CVn periodogram folder. The generated figures use the `AM_CVn_physical_*` prefix in the plot folder.