

# Counting Binary Trees & Selection Trees

Joseph Chuang-Chieh Lin (林莊傑)

Department of Computer Science & Engineering,  
National Taiwan Ocean University

Fall 2025



# Outline

1 Counting Binary Trees

2 Selection Trees

# Outline

1 Counting Binary Trees

2 Selection Trees

# Counting Binary Trees


- Consider the following three disparate problems:
  - The number of distinct binary trees having  $n$  nodes.
  - The number of distinct permutations of the numbers from 1 to  $n$  obtainable by a **stack**.
  - The number of distinct ways of multiplying  $n + 1$  matrices.

# Counting Binary Trees

- Consider the following three disparate problems:
  - ➊ The number of distinct binary trees having  $n$  nodes.
  - ➋ The number of distinct permutations of the numbers from 1 to  $n$  obtainable by a **stack**.
  - ➌ The number of distinct ways of multiplying  $n + 1$  matrices.
- Amazingly, **these problems have the same solution!**

# Problem One

- The number of distinct binary trees having  $n$  nodes.



- Example of  $n = 3$ .

## Problem Two

- The number of distinct permutations of the numbers from 1 to  $n$  obtainable by a stack.

- ① push 1 → pop → push 2 → pop → push 3 → pop ⇒ 123.
- ② push 1 → pop → push 2 → push 3 → pop → pop ⇒ 132.
- ③ push 1 → push 2 → push 3 → pop → pop → pop ⇒ 321.
- ④ push 1 → push 2 → pop → pop → push 3 → pop ⇒ 213.
- ⑤ push 1 → push 2 → pop → push 3 → pop → pop ⇒ 231.

\* Example of  $n = 3$ .

# Problem Three

- The number of distinct ways of multiplying  $n + 1$  matrices.
  - ①  $((M_1 \times M_2) \times M_3) \times M_4$ .
  - ②  $(M_1 \times (M_2 \times M_3)) \times M_4$ .
  - ③  $M_1 \times ((M_2 \times M_3) \times M_4)$ .
  - ④  $M_1 \times (M_2 \times (M_3 \times M_4))$ .
  - ⑤  $(M_1 \times M_2) \times (M_3 \times M_4)$ .

\* Example of  $n = 3$ .

# Stack Permutation (1/4)

- Recall: preorder, inorder and postorder traversal of a binary tree.
  - Each traversal requires a **stack**.

Every binary tree has a unique pair of preorder/inorder sequences.

# Stack Permutation (1/4)


- Recall: preorder, inorder and postorder traversal of a binary tree.
  - Each traversal requires a **stack**.

Every binary tree has a unique pair of preorder/inorder sequences.


- The number of distinct binary trees is equal to the number of **inorder permutations** obtainable from binary trees having the preorder permutation,  $1, 2, \dots, n$ .

## Stack Permutation (2/4)

- preorder: A B C E D G H F I
- inorder: B C A E D G H F I



- preorder: A B C (D E F G H I)
- inorder: B C A (E D F G H I)



## Stack Permutation (3/4)

- We can show that


the number of distinct permutations obtainable by passing the numbers  $\{1, 2, \dots, n\}$  through a stack is equal to the number of distinct binary trees with  $n$  nodes.

- ① push 1 → pop → push 2 → pop → push 3 → pop ⇒ 123.
- ② push 1 → pop → push 2 → push 3 → pop → pop ⇒ 132.
- ③ push 1 → push 2 → push 3 → pop → pop → pop ⇒ 321.
- ④ push 1 → push 2 → pop → pop → push 3 → pop ⇒ 213.
- ⑤ push 1 → push 2 → pop → push 3 → pop → pop ⇒ 231.



# Stack Permutation (4/4)

- visit node → push to stack;
  - going left → keep visiting next node
  - going right → pop the stack
- the leaf → pop the stack until empty



# Go Back to the Matrix Multiplication

- Computing the product of  $n$  matrices are related to the distinct binary tree problem.
- $n = 3$ :
  - ①  $(M_1 \times M_2) \times M_3$ .
  - ②  $M_1 \times (M_2 \times M_3)$ .
- $n = 4$ :
  - ①  $((M_1 \times M_2) \times M_3) \times M_4$
  - ②  $(M_1 \times (M_2 \times M_3)) \times M_4$
  - ③  $M_1 \times ((M_2 \times M_3) \times M_4)$
  - ④  $M_1 \times (M_2 \times (M_3 \times M_4))$
  - ⑤  $(M_1 \times M_2) \times (M_3 \times M_4)$

# Go Back to the Matrix Multiplication

- Computing the product of  $n$  matrices are related to the distinct binary tree problem.
- $n = 3$ :
  - ①  $(M_1 \times M_2) \times M_3$ .
  - ②  $M_1 \times (M_2 \times M_3)$ .
- $n = 4$ :
  - ①  $((M_1 \times M_2) \times M_3) \times M_4$  (push, pop, push, pop, push, pop)
  - ②  $(M_1 \times (M_2 \times M_3)) \times M_4$  (?)
  - ③  $M_1 \times ((M_2 \times M_3) \times M_4)$  (?)
  - ④  $M_1 \times (M_2 \times (M_3 \times M_4))$  (?)
  - ⑤  $(M_1 \times M_2) \times (M_3 \times M_4)$  (?)

# Matrix Multiplication (2/2)

- $b_n$ : the number of different ways to compute the product of  $n$  matrices.
- Trivially,  $b_1 =$

# Matrix Multiplication (2/2)

- $b_n$ : the number of different ways to compute the product of  $n$  matrices.
- Trivially,  $b_1 = 1$ ,  $b_2 =$

# Matrix Multiplication (2/2)

- $b_n$ : the number of different ways to compute the product of  $n$  matrices.
- Trivially,  $b_1 = 1$ ,  $b_2 = 1$ .
- We have also derived that  $b_3 = 2$  and  $b_4 = 5$ .
- We can compute that

$$b_n = \sum_{i=1}^{n-1} b_i b_{n-i}, \text{ for } n > 1.$$

# Distinct Binary Trees

- Similarly, the number of **distinct binary trees** of  $n$  nodes is

$$b_n =$$

# Distinct Binary Trees


- Similarly, the number of **distinct binary trees** of  $n$  nodes is

$$b_n = \sum_{i=0}^{n-1} b_i b_{n-1-i}, \text{ for } n \geq 1 \text{ and}$$

# Distinct Binary Trees

- Similarly, the number of **distinct binary trees** of  $n$  nodes is


$$b_n = \sum_{i=0}^{n-1} b_i b_{n-1-i}, \text{ for } n \geq 1 \text{ and } b_0 = 1.$$



# Distinct Binary Trees

- Similarly, the number of **distinct binary trees** of  $n$  nodes is

$$b_n = \sum_{i=0}^{n-1} b_i b_{n-1-i}, \text{ for } n \geq 1 \text{ and } b_0 = 1.$$



- But, how to compute  $b_n$  exactly?

# The Generating Function Trick

- **Trick:** Let  $B(x) = \sum_{i \geq 0} b_i x^i$  be the generating function for the number of binary trees.



# The Generating Function Trick

- **Trick:** Let  $B(x) = \sum_{i \geq 0} b_i x^i$  be the generating function for the number of binary trees.

$$B(x) = \sum_{i \geq 0} b_i x^i = \sum_{i=0}^{\infty} \sum_{k=0}^{i-1} b_k b_{i-1-k} x^i =$$

# The Generating Function Trick

- **Trick:** Let  $B(x) = \sum_{i \geq 0} b_i x^i$  be the generating function for the number of binary trees.

$$\begin{aligned} B(x) &= \sum_{i \geq 0} b_i x^i = \sum_{i=0}^{\infty} \sum_{k=0}^{i-1} b_k b_{i-1-k} x^i &= 1 + \sum_{i=1}^{\infty} \sum_{k=0}^{i-1} b_k b_{i-1-k} x^i \\ &= \end{aligned}$$



# The Generating Function Trick

- **Trick:** Let  $B(x) = \sum_{i \geq 0} b_i x^i$  be the generating function for the number of binary trees.

$$\begin{aligned}
 B(x) &= \sum_{i \geq 0} b_i x^i = \sum_{i=0}^{\infty} \sum_{k=0}^{i-1} b_k b_{i-1-k} x^i &= 1 + \sum_{i=1}^{\infty} \sum_{k=0}^{i-1} b_k b_{i-1-k} x^i \\
 &= 1 + \cancel{x} \sum_{i=1}^{\infty} \sum_{k=0}^{i-1} b_k b_{i-1-k} x^{i-1} \\
 &=
 \end{aligned}$$

# The Generating Function Trick

- **Trick:** Let  $B(x) = \sum_{i \geq 0} b_i x^i$  be the generating function for the number of binary trees.

$$\begin{aligned}
 B(x) &= \sum_{i \geq 0} b_i x^i = \sum_{i=0}^{\infty} \sum_{k=0}^{i-1} b_k b_{i-1-k} x^i &= 1 + \sum_{i=1}^{\infty} \sum_{k=0}^{i-1} b_k b_{i-1-k} x^i \\
 &= 1 + \cancel{x} \sum_{i=1}^{\infty} \sum_{k=0}^{i-1} b_k b_{i-1-k} x^{i-1} \\
 &= 1 + x \sum_{j=0}^{\infty} \sum_{k=0}^j b_k b_{j-k} x^j \\
 &= 1 + x \left( \sum_{j=0}^{\infty} b_j x^j \right)^2 = 1 + xB(x)^2.
 \end{aligned}$$

# The Generating Function Trick

- By the recurrence relation we get:

$$xB(x)^2 = B(x) - 1.$$

- Solving the recurrence relation, we have

$$\begin{aligned}B(x) &= \frac{1 - \sqrt{1 - 4x}}{2x} \\&= \frac{1}{2x} \left( 1 - \sum_{n \geq 0} \binom{1/2}{n} (-4x)^n \right) \\&= \sum_{m \geq 0} \binom{1/2}{m+1} (-1)^m 2^{2m+1} x^m.\end{aligned}$$

# The Generating Function Trick

- By the recurrence relation we get:

$$xB(x)^2 = B(x) - 1.$$

- Solving the recurrence relation, we have

$$\begin{aligned}B(x) &= \frac{1 - \sqrt{1 - 4x}}{2x} \\&= \frac{1}{2x} \left( 1 - \sum_{n \geq 0} \binom{1/2}{n} (-4x)^n \right) \\&= \sum_{m \geq 0} \binom{1/2}{m+1} (-1)^m 2^{2m+1} x^m.\end{aligned}\quad \therefore b_n = \frac{1}{n+1} \binom{2n}{n}.$$

By the Binomial Theorem...

$$b_n = \binom{1/2}{n+1} (-1)^n 2^{2n+1}$$

=

# By the Binomial Theorem...

$$\begin{aligned} b_n &= \binom{1/2}{n+1} (-1)^n 2^{2n+1} \\ &= \frac{\frac{1}{2} \left(\frac{1}{2}-1\right) \left(\frac{1}{2}-2\right) \cdots \left(\frac{1}{2}-n\right)}{(n+1)n(n-1)(n-2)\cdots 2 \cdot 1} (-1)^n 2^{2n+1} \\ &= \end{aligned}$$

# By the Binomial Theorem...

$$\begin{aligned} b_n &= \binom{1/2}{n+1} (-1)^n 2^{2n+1} \\ &= \frac{\frac{1}{2} \left(\frac{1}{2}-1\right) \left(\frac{1}{2}-2\right) \cdots \left(\frac{1}{2}-n\right)}{(n+1)n(n-1)(n-2)\cdots 2 \cdot 1} (-1)^n 2^{2n+1} \\ &= \frac{\frac{1}{2} \left(1-\frac{1}{2}\right) \left(2-\frac{1}{2}\right) \cdots \left(n-\frac{1}{2}\right)}{(n+1)n(n-1)(n-2)\cdots 2 \cdot 1} 2^{2n+1} \\ &= \end{aligned}$$

# By the Binomial Theorem...

$$\begin{aligned} b_n &= \binom{1/2}{n+1} (-1)^n 2^{2n+1} \\ &= \frac{\frac{1}{2} \left(\frac{1}{2}-1\right) \left(\frac{1}{2}-2\right) \cdots \left(\frac{1}{2}-n\right)}{(n+1)n(n-1)(n-2)\cdots 2 \cdot 1} (-1)^n 2^{2n+1} \\ &= \frac{\frac{1}{2} \left(1-\frac{1}{2}\right) \left(2-\frac{1}{2}\right) \cdots \left(n-\frac{1}{2}\right)}{(n+1)n(n-1)(n-2)\cdots 2 \cdot 1} 2^{2n+1} \\ &= \left(\frac{1}{2}\right)^{n+1} \cdot \frac{1(2-1)(4-1)\cdots(2n-1)}{(n+1)n(n-1)(n-2)\cdots 2 \cdot 1} 2^{2n+1} \\ &= \end{aligned}$$

# By the Binomial Theorem...

$$\begin{aligned}
 b_n &= \binom{1/2}{n+1} (-1)^n 2^{2n+1} \\
 &= \frac{\frac{1}{2} \left(\frac{1}{2}-1\right) \left(\frac{1}{2}-2\right) \cdots \left(\frac{1}{2}-n\right)}{(n+1)n(n-1)(n-2)\cdots 2 \cdot 1} (-1)^n 2^{2n+1} \\
 &= \frac{\frac{1}{2} \left(1-\frac{1}{2}\right) \left(2-\frac{1}{2}\right) \cdots \left(n-\frac{1}{2}\right)}{(n+1)n(n-1)(n-2)\cdots 2 \cdot 1} 2^{2n+1} \\
 &= \left(\frac{1}{2}\right)^{n+1} \cdot \frac{1(2-1)(4-1)\cdots(2n-1)}{(n+1)n(n-1)(n-2)\cdots 2 \cdot 1} 2^{2n+1} \\
 &= \frac{1(2-1)(4-1)\cdots(2n-1)}{(n+1)n(n-1)(n-2)\cdots 2 \cdot 1} \cdot 2^n \\
 &=
 \end{aligned}$$

By the Binomial Theorem...

$$\begin{aligned}
b_n &= \binom{1/2}{n+1} (-1)^n 2^{2n+1} \\
&= \frac{\frac{1}{2} \left(\frac{1}{2}-1\right) \left(\frac{1}{2}-2\right) \cdots \left(\frac{1}{2}-n\right)}{(n+1)n(n-1)(n-2)\cdots 2 \cdot 1} (-1)^n 2^{2n+1} \\
&= \frac{\frac{1}{2} \left(1-\frac{1}{2}\right) \left(2-\frac{1}{2}\right) \cdots \left(n-\frac{1}{2}\right)}{(n+1)n(n-1)(n-2)\cdots 2 \cdot 1} 2^{2n+1} \\
&= \left(\frac{1}{2}\right)^{n+1} \cdot \frac{1(2-1)(4-1)\cdots(2n-1)}{(n+1)n(n-1)(n-2)\cdots 2 \cdot 1} 2^{2n+1} \\
&= \frac{1(2-1)(4-1)\cdots(2n-1)}{(n+1)n(n-1)(n-2)\cdots 2 \cdot 1} \cdot 2^n \\
&= \frac{1(1)(3)\cdots(2n-1)}{(n+1)n(n-1)(n-2)\cdots 2 \cdot 1} \cdot 2^n \cdot \frac{2 \cdot 4 \cdot 6 \cdots (2n)}{2^n(1 \cdot 2 \cdots n)} \\
&\equiv
\end{aligned}$$



# By the Binomial Theorem...

$$\begin{aligned}
 b_n &= \binom{1/2}{n+1} (-1)^n 2^{2n+1} \\
 &= \frac{\frac{1}{2} \left(\frac{1}{2}-1\right) \left(\frac{1}{2}-2\right) \cdots \left(\frac{1}{2}-n\right)}{(n+1)n(n-1)(n-2)\cdots 2\cdot 1} (-1)^n 2^{2n+1} \\
 &= \frac{\frac{1}{2} \left(1-\frac{1}{2}\right) \left(2-\frac{1}{2}\right) \cdots \left(n-\frac{1}{2}\right)}{(n+1)n(n-1)(n-2)\cdots 2\cdot 1} 2^{2n+1} \\
 &= \left(\frac{1}{2}\right)^{n+1} \cdot \frac{1(2-1)(4-1)\cdots(2n-1)}{(n+1)n(n-1)(n-2)\cdots 2\cdot 1} 2^{2n+1} \\
 &= \frac{1(2-1)(4-1)\cdots(2n-1)}{(n+1)n(n-1)(n-2)\cdots 2\cdot 1} \cdot 2^n \\
 &= \frac{1(1)(3)\cdots(2n-1)}{(n+1)n(n-1)(n-2)\cdots 2\cdot 1} \cdot 2^n \cdot \frac{2\cdot 4\cdot 6\cdots(2n)}{2^n(1\cdot 2\cdots n)} \\
 &= \frac{(2n)!}{(n+1)n!n!}
 \end{aligned}$$

# By the Binomial Theorem...

$$\begin{aligned}
 b_n &= \binom{1/2}{n+1} (-1)^n 2^{2n+1} \\
 &= \frac{\frac{1}{2} \left(\frac{1}{2}-1\right) \left(\frac{1}{2}-2\right) \cdots \left(\frac{1}{2}-n\right)}{(n+1)n(n-1)(n-2)\cdots 2\cdot 1} (-1)^n 2^{2n+1} \\
 &= \frac{\frac{1}{2} \left(1-\frac{1}{2}\right) \left(2-\frac{1}{2}\right) \cdots \left(n-\frac{1}{2}\right)}{(n+1)n(n-1)(n-2)\cdots 2\cdot 1} 2^{2n+1} \\
 &= \left(\frac{1}{2}\right)^{n+1} \cdot \frac{1(2-1)(4-1)\cdots(2n-1)}{(n+1)n(n-1)(n-2)\cdots 2\cdot 1} 2^{2n+1} \\
 &= \frac{1(2-1)(4-1)\cdots(2n-1)}{(n+1)n(n-1)(n-2)\cdots 2\cdot 1} \cdot 2^n \\
 &= \frac{1(1)(3)\cdots(2n-1)}{(n+1)n(n-1)(n-2)\cdots 2\cdot 1} \cdot 2^n \cdot \frac{2\cdot 4\cdot 6\cdots(2n)}{2^n(1\cdot 2\cdots n)} \\
 &= \frac{(2n)!}{(n+1)n!n!} = \frac{1}{n+1} \binom{2n}{n}.
 \end{aligned}$$

# By the Binomial Theorem...

$$\begin{aligned}
 b_n &= \binom{1/2}{n+1} (-1)^n 2^{2n+1} \\
 &= \frac{\frac{1}{2} \left(\frac{1}{2}-1\right) \left(\frac{1}{2}-2\right) \cdots \left(\frac{1}{2}-n\right)}{(n+1)n(n-1)(n-2)\cdots 2\cdot 1} (-1)^n 2^{2n+1} \\
 &= \frac{\frac{1}{2} \left(1-\frac{1}{2}\right) \left(2-\frac{1}{2}\right) \cdots \left(n-\frac{1}{2}\right)}{(n+1)n(n-1)(n-2)\cdots 2\cdot 1} 2^{2n+1} \\
 &= \left(\frac{1}{2}\right)^{n+1} \cdot \frac{1(2-1)(4-1)\cdots(2n-1)}{(n+1)n(n-1)(n-2)\cdots 2\cdot 1} 2^{2n+1} \\
 &= \frac{1(2-1)(4-1)\cdots(2n-1)}{(n+1)n(n-1)(n-2)\cdots 2\cdot 1} \cdot 2^n \\
 &= \frac{1(1)(3)\cdots(2n-1)}{(n+1)n(n-1)(n-2)\cdots 2\cdot 1} \cdot 2^n \cdot \frac{2 \cdot 4 \cdot 6 \cdots (2n)}{2^n(1 \cdot 2 \cdots n)} \\
 &= \frac{(2n)!}{(n+1)n!n!} = \frac{1}{n+1} \binom{2n}{n}.
 \end{aligned}$$

\* supplementary: Stirling's approximation

# Outline

1 Counting Binary Trees

2 Selection Trees

# Scenarios of Using the Selection Trees

- External sorting.
- Data stored in each queue (run) is sorted.

# Winner Selection Tree

- In the following figure, computing the first winner takes




# Winner Selection Tree

- In the following figure, computing the first winner takes  $k - 1$  comparisons.



# Winner Selection Tree


- In the following figure, computing the first winner takes  $k - 1$  comparisons.  
(not better?)
- But wait, how about the following iterations?



# Winner Selection Tree

(Winner) Selection Tree:


- Setting up the selection tree:  $O(k)$  time.
- Restructuring:  $O(\lg k)$  time.
- merging all  $n$  items:  $O(n \lg k)$  time.



# Winner Selection Tree

(Winner) Selection Tree:


- Setting up the selection tree:  $O(k)$  time.
- Restructuring:  $O(\lg k)$  time.
- merging all  $n$  items:  $O(n \lg k)$  time.



# Loser Selection Tree

Loser Tree (after the winner tree has been built)


- Sibling nodes represent the losers.
- The restructuring process can be simplified.



# Loser Selection Tree

Loser Tree (after the winner tree has been built)

- Sibling nodes represent the losers.
- The restructuring process can be simplified.



# Note for the Loser Selection Tree

- Comparison with the sibling is required for the first step construction.
- After the first construction, we only need to compare each node with its parent; “push” the smaller key value upward and left the “larger” key value as the **loser**.

# Discussions

