

# Mathematics for Machine Learning

## — Linear Algebra: Projections & Gram-Schmidt Orthogonalization

Joseph Chuang-Chieh Lin

Department of Computer Science & Information Engineering,  
Tamkang University

Fall 2023

## Credits for the resource

- The slides are based on the textbooks:
  - *Marc Peter Deisenroth, A. Aldo Faisal, and Cheng Soon Ong: Mathematics for Machine Learning. Cambridge University Press. 2020.*
  - *Howard Anton, Chris Rorres, Anton Kaul: Elementary Linear Algebra. Wiley. 2019.*
- We could partially refer to the monograph:  
*Francesco Orabona: A Modern Introduction to Online Learning.*  
<https://arxiv.org/abs/1912.13213>

# Outline

- 1 Orthogonal Projections
- 2 Gram-Schmidt Orthogonalization

# Outline

- 1 Orthogonal Projections
- 2 Gram-Schmidt Orthogonalization

# Motivations (1/2)

- In machine learning, we often need to deal with *high-dimensional* data.

## Motivations (1/2)

- In machine learning, we often need to deal with *high-dimensional* data.
- High-dimensional data is often hard to analyze or visualize.

## Motivations (1/2)

- In machine learning, we often need to deal with *high-dimensional* data.
- High-dimensional data is often hard to analyze or visualize.
- Sometimes, only a few dimensions contain most information.

## Motivations (1/2)

- In machine learning, we often need to deal with *high-dimensional* data.
- High-dimensional data is often hard to analyze or visualize.
- Sometimes, only a few dimensions contain most information.
- We might try to project the original high-dimensional data onto a lower-dimensional space and work on it.



## Motivations (1/2)

- In machine learning, we often need to deal with *high-dimensional* data.
- High-dimensional data is often hard to analyze or visualize.
- Sometimes, only a few dimensions contain most information.
- We might try to project the original high-dimensional data onto a lower-dimensional space and work on it.
- Note: When we compress or visualize high-dimensional data, we will lose information.

# Motivations (2/2)

Examples (dimensionality reduction)

- Principal component analysis (PCA)

## Motivations (2/2)

Examples (dimensionality reduction)

- Principal component analysis (PCA)
- Deep neural networks

## Motivations (2/2)

Examples (dimensionality reduction)

- Principal component analysis (PCA)
- Deep neural networks
- Classification

# Motivations (2/2)

Examples (dimensionality reduction)

- Principal component analysis (PCA)
- Deep neural networks
- Classification
- linear Regression

# Projection

## Projection

Let  $V$  be a vector space and  $U \subseteq V$  be a subspace of  $V$ . A linear mapping  $\pi : V \mapsto U$  is called a **projection** if  $\pi^2 = \pi \circ \pi = \pi$ .

# Projection

## Projection

Let  $V$  be a vector space and  $U \subseteq V$  be a subspace of  $V$ . A linear mapping  $\pi : V \mapsto U$  is called a **projection** if  $\pi^2 = \pi \circ \pi = \pi$ .

- Recall that linear mappings can be expressed by transformation matrices.

# Projection


## Projection

Let  $V$  be a vector space and  $U \subseteq V$  be a subspace of  $V$ . A linear mapping  $\pi : V \mapsto U$  is called a **projection** if  $\pi^2 = \pi \circ \pi = \pi$ .


- Recall that linear mappings can be expressed by transformation matrices.
- The projection matrices  $P_\pi$  exhibits the property that  $P_\pi^2 = P_\pi$ .



## Illustration of projections onto 1-D



(a) Projection of  $x \in \mathbb{R}^2$  onto a subspace  $U$  with basis vector  $b$ .



(b) Projection of a two-dimensional vector  $x$  with  $\|x\| = 1$  onto a one-dimensional subspace spanned by  $b$ .

# Illustration of projections onto 1-D

- $\pi_U(\mathbf{x})$ : closest to  $\mathbf{x}$ .
  - $\|\mathbf{x} - \pi_U(\mathbf{x})\|$  is minimal.

# Illustration of projections onto 1-D

- $\pi_U(\mathbf{x})$ : closest to  $\mathbf{x}$ .
  - $\|\mathbf{x} - \pi_U(\mathbf{x})\|$  is minimal.
  - $\langle \pi_U(\mathbf{x}) - \mathbf{x}, \mathbf{b} \rangle = 0$ .
- Projection  $\pi_U(\mathbf{x})$  of  $\mathbf{x}$  onto  $U$  must be an element in  $U$ .
  - $\pi_U(\mathbf{x}) = \lambda \mathbf{b}$  for some  $\lambda \in \mathbb{R}$ .

# Illustration of projections onto 1-D

- $\pi_U(\mathbf{x})$ : closest to  $\mathbf{x}$ .
  - $\|\mathbf{x} - \pi_U(\mathbf{x})\|$  is minimal.
  - $\langle \pi_U(\mathbf{x}) - \mathbf{x}, \mathbf{b} \rangle = 0$ .
- Projection  $\pi_U(\mathbf{x})$  of  $\mathbf{x}$  onto  $U$  must be an element in  $U$ .
  - $\pi_U(\mathbf{x}) = \lambda \mathbf{b}$  for some  $\lambda \in \mathbb{R}$ .
- Determining the coordinates:

## Illustration of projections onto 1-D

- $\pi_U(\mathbf{x})$ : closest to  $\mathbf{x}$ .
  - $\|\mathbf{x} - \pi_U(\mathbf{x})\|$  is minimal.
  - $\langle \pi_U(\mathbf{x}) - \mathbf{x}, \mathbf{b} \rangle = 0$ .
- Projection  $\pi_U(\mathbf{x})$  of  $\mathbf{x}$  onto  $U$  must be an element in  $U$ .
  - $\pi_U(\mathbf{x}) = \lambda \mathbf{b}$  for some  $\lambda \in \mathbb{R}$ .
- Determining the coordinates:  
Since  $\pi_U(\mathbf{b}) = \lambda \mathbf{b}$ :

$$\langle \mathbf{x} - \pi_U(\mathbf{x}), \mathbf{b} \rangle = 0 \Leftrightarrow \langle \mathbf{x} - \lambda \mathbf{b}, \mathbf{b} \rangle = 0.$$

# Illustration of projections onto 1-D

- $\pi_U(\mathbf{x})$ : closest to  $\mathbf{x}$ .
  - $\|\mathbf{x} - \pi_U(\mathbf{x})\|$  is minimal.
  - $\langle \pi_U(\mathbf{x}) - \mathbf{x}, \mathbf{b} \rangle = 0$ .
- Projection  $\pi_U(\mathbf{x})$  of  $\mathbf{x}$  onto  $U$  must be an element in  $U$ .
  - $\pi_U(\mathbf{x}) = \lambda \mathbf{b}$  for some  $\lambda \in \mathbb{R}$ .
- Determining the coordinates:  
Since  $\pi_U(\mathbf{b}) = \lambda \mathbf{b}$ :

$$\langle \mathbf{x} - \pi_U(\mathbf{x}), \mathbf{b} \rangle = 0 \Leftrightarrow \langle \mathbf{x} - \lambda \mathbf{b}, \mathbf{b} \rangle = 0.$$

$$\Leftrightarrow \langle \mathbf{x}, \mathbf{b} \rangle - \lambda \langle \mathbf{b}, \mathbf{b} \rangle = 0 \Leftrightarrow \lambda = \frac{\langle \mathbf{x}, \mathbf{b} \rangle}{\langle \mathbf{b}, \mathbf{b} \rangle} = \frac{\langle \mathbf{b}, \mathbf{x} \rangle}{\|\mathbf{b}\|^2}$$

# Illustration of projections onto 1-D

- $\pi_U(\mathbf{x})$ : closest to  $\mathbf{x}$ .
  - $\|\mathbf{x} - \pi_U(\mathbf{x})\|$  is minimal.
  - $\langle \pi_U(\mathbf{x}) - \mathbf{x}, \mathbf{b} \rangle = 0$ .
- Projection  $\pi_U(\mathbf{x})$  of  $\mathbf{x}$  onto  $U$  must be an element in  $U$ .
  - $\pi_U(\mathbf{x}) = \lambda \mathbf{b}$  for some  $\lambda \in \mathbb{R}$ .
- Determining the coordinates:  
Since  $\pi_U(\mathbf{b}) = \lambda \mathbf{b}$ :

$$\langle \mathbf{x} - \pi_U(\mathbf{x}), \mathbf{b} \rangle = 0 \Leftrightarrow \langle \mathbf{x} - \lambda \mathbf{b}, \mathbf{b} \rangle = 0.$$

$$\Leftrightarrow \langle \mathbf{x}, \mathbf{b} \rangle - \lambda \langle \mathbf{b}, \mathbf{b} \rangle = 0 \Leftrightarrow \lambda = \frac{\langle \mathbf{x}, \mathbf{b} \rangle}{\langle \mathbf{b}, \mathbf{b} \rangle} = \frac{\langle \mathbf{b}, \mathbf{x} \rangle}{\|\mathbf{b}\|^2} = \frac{\mathbf{b}^\top \mathbf{x}}{\mathbf{b}^\top \mathbf{b}}.$$

- Finding the projection  $\pi_U(\mathbf{x}) \in U$ :

$$\pi_U(\mathbf{x}) = \lambda \mathbf{b} = \frac{\langle \mathbf{x}, \mathbf{b} \rangle}{\|\mathbf{b}\|^2} \mathbf{b} = \frac{\mathbf{b}^\top \mathbf{x}}{\|\mathbf{b}\|^2} \mathbf{b},$$



- Finding the projection  $\pi_U(\mathbf{x}) \in U$ :

$$\pi_U(\mathbf{x}) = \lambda \mathbf{b} = \frac{\langle \mathbf{x}, \mathbf{b} \rangle}{\|\mathbf{b}\|^2} \mathbf{b} = \frac{\mathbf{b}^\top \mathbf{x}}{\|\mathbf{b}\|^2} \mathbf{b},$$

Note that  $\|\pi_U(\mathbf{x})\| = \|\lambda \mathbf{b}\| = |\lambda| \|\mathbf{b}\|$ .

- Finding the projection  $\pi_U(\mathbf{x}) \in U$ :

$$\pi_U(\mathbf{x}) = \lambda \mathbf{b} = \frac{\langle \mathbf{x}, \mathbf{b} \rangle}{\|\mathbf{b}\|^2} \mathbf{b} = \frac{\mathbf{b}^\top \mathbf{x}}{\|\mathbf{b}\|^2} \mathbf{b},$$

Note that  $\|\pi_U(\mathbf{x})\| = \|\lambda \mathbf{b}\| = |\lambda| \|\mathbf{b}\|$ .

- If we use the dot product as the inner product and let  $\theta$  be the angle between  $\mathbf{x}$  and  $\mathbf{b}$ :

$$\|\pi_U(\mathbf{x})\| = \frac{\mathbf{b}^\top \mathbf{x}}{\|\mathbf{b}\|^2} \|\mathbf{b}\| = |\cos \theta| \|\mathbf{x}\| \|\mathbf{b}\| \frac{\|\mathbf{b}\|}{\|\mathbf{b}\|^2} = |\cos \theta| \|\mathbf{x}\|.$$

- Finding the projection matrix  $P_\pi$ :
  - Recall: projection is a linear mapping.
  - With the dot product as the inner product,

$$\|\pi_U(\mathbf{x})\| = \lambda \mathbf{b} = \mathbf{b} \lambda = \mathbf{b} \frac{\mathbf{b}^\top \mathbf{x}}{\|\mathbf{b}\|^2} = \frac{\mathbf{b} \mathbf{b}^\top}{\|\mathbf{b}\|^2} \mathbf{x}.$$

- So,

$$P_\pi = \frac{\mathbf{b} \mathbf{b}^\top}{\|\mathbf{b}\|^2}.$$

Note  $\mathbf{b} \mathbf{b}^\top$  is a symmetric matrix.

- Finding the projection matrix  $P_\pi$ :
  - Recall: projection is a linear mapping.
  - With the dot product as the inner product,

$$\|\pi_U(\mathbf{x})\| = \lambda \mathbf{b} = \mathbf{b} \lambda = \mathbf{b} \frac{\mathbf{b}^\top \mathbf{x}}{\|\mathbf{b}\|^2} = \frac{\mathbf{b} \mathbf{b}^\top}{\|\mathbf{b}\|^2} \mathbf{x}.$$


- So,

$$P_\pi = \frac{\mathbf{b} \mathbf{b}^\top}{\|\mathbf{b}\|^2}.$$

Note  $\mathbf{b} \mathbf{b}^\top$  is a symmetric matrix.

# Projection onto General Subspaces (1/4)

Orthogonal projections of  $\mathbf{x} \in \mathbb{R}^n$  onto  $U \subseteq \mathbb{R}^n$  with  $\dim(U) = m \geq 1$ .



## Projection onto General Subspaces (2/4)

- Any projection can be represented as a linear combination of the basis vectors  $\mathbf{b}_1, \dots, \mathbf{b}_m$  of  $U$ .
  - $\pi_U(\mathbf{x}) = \sum_{i=1}^m \lambda_i \mathbf{b}_i$ .
- Find the coordinates  $\lambda_1, \dots, \lambda_m$ :

$$\pi_U(\mathbf{x}) = \sum_{i=1}^m \lambda_i \mathbf{b}_i = \mathbf{B}\boldsymbol{\lambda}$$

for  $\mathbf{B} = [\mathbf{b}_1, \dots, \mathbf{b}_m] \in \mathbb{R}^{n \times m}$ ,  $\boldsymbol{\lambda} = [\lambda_1, \dots, \lambda_m]^\top \in \mathbb{R}^m$ .

## Projection onto General Subspaces (2/4)

- Any projection can be represented as a linear combination of the basis vectors  $\mathbf{b}_1, \dots, \mathbf{b}_m$  of  $U$ .
  - $\pi_U(\mathbf{x}) = \sum_{i=1}^m \lambda_i \mathbf{b}_i$ .
- Find the coordinates  $\lambda_1, \dots, \lambda_m$ :

$$\pi_U(\mathbf{x}) = \sum_{i=1}^m \lambda_i \mathbf{b}_i = \mathbf{B}\boldsymbol{\lambda} \quad (\text{closest to } \mathbf{x} \text{ on } U)$$

for  $\mathbf{B} = [\mathbf{b}_1, \dots, \mathbf{b}_m] \in \mathbb{R}^{n \times m}$ ,  $\boldsymbol{\lambda} = [\lambda_1, \dots, \lambda_m]^\top \in \mathbb{R}^m$ .

Note:  $\mathbf{x} \perp \{\mathbf{b}_1, \dots, \mathbf{b}_m\}$  ( $\because$  minimum distance)

$$\begin{aligned} \langle \mathbf{b}_1, \mathbf{x} - \pi_U(\mathbf{x}) \rangle &= \mathbf{b}_1^\top (\mathbf{x} - \pi_U(\mathbf{x})) = 0 \\ &\vdots \\ \langle \mathbf{b}_m, \mathbf{x} - \pi_U(\mathbf{x}) \rangle &= \mathbf{b}_m^\top (\mathbf{x} - \pi_U(\mathbf{x})) = 0 \end{aligned}$$

## Projection onto General Subspaces (3/4)

- Any projection can be represented as a linear combination of the basis vectors  $\mathbf{b}_1, \dots, \mathbf{b}_m$  of  $U$ .
  - $\pi_U(\mathbf{x}) = \sum_{i=1}^m \lambda_i \mathbf{b}_i$ .
- Find the coordinates  $\lambda_1, \dots, \lambda_m$ :

$$\pi_U(\mathbf{x}) = \sum_{i=1}^m \lambda_i \mathbf{b}_i = \mathbf{B}\boldsymbol{\lambda} \quad (\text{closest to } \mathbf{x} \text{ on } U)$$

for  $\mathbf{B} = [\mathbf{b}_1, \dots, \mathbf{b}_m] \in \mathbb{R}^{n \times m}$ ,  $\boldsymbol{\lambda} = [\lambda_1, \dots, \lambda_m]^\top \in \mathbb{R}^m$ .

Note:  $\mathbf{x} \perp \{\mathbf{b}_1, \dots, \mathbf{b}_m\}$  ( $\because$  minimum distance)

$$\mathbf{b}_1^\top (\mathbf{x} - \mathbf{B}\boldsymbol{\lambda}) = 0$$

$$\vdots$$

$$\mathbf{b}_m^\top (\mathbf{x} - \mathbf{B}\boldsymbol{\lambda}) = 0$$



## Projection onto General Subspaces (4/4)

Since

$$\mathbf{b}_1^\top (\mathbf{x} - \mathbf{B}\boldsymbol{\lambda}) = 0$$

$$\vdots$$

$$\mathbf{b}_m^\top (\mathbf{x} - \mathbf{B}\boldsymbol{\lambda}) = 0$$

We have

$$\begin{bmatrix} \mathbf{b}_1^\top \\ \vdots \\ \mathbf{b}_m^\top \end{bmatrix} [\mathbf{x} - \mathbf{B}\boldsymbol{\lambda}] = \mathbf{0} \Leftrightarrow \mathbf{B}^\top (\mathbf{x} - \mathbf{B}\boldsymbol{\lambda}) = \mathbf{0}$$

$$\Leftrightarrow \mathbf{B}^\top \mathbf{B}\boldsymbol{\lambda} = \mathbf{B}^\top \mathbf{x}$$

**Note:**  $\mathbf{B}^\top \mathbf{B}$  is invertible

## Projection onto General Subspaces (4/4)

Since

$$\begin{aligned} \mathbf{b}_1^\top (\mathbf{x} - \mathbf{B}\boldsymbol{\lambda}) &= 0 \\ &\vdots \\ \mathbf{b}_m^\top (\mathbf{x} - \mathbf{B}\boldsymbol{\lambda}) &= 0 \end{aligned}$$

We have

$$\begin{aligned} \begin{bmatrix} \mathbf{b}_1^\top \\ \vdots \\ \mathbf{b}_m^\top \end{bmatrix} [\mathbf{x} - \mathbf{B}\boldsymbol{\lambda}] &= \mathbf{0} \Leftrightarrow \mathbf{B}^\top (\mathbf{x} - \mathbf{B}\boldsymbol{\lambda}) = \mathbf{0} \\ &\Leftrightarrow \mathbf{B}^\top \mathbf{B}\boldsymbol{\lambda} = \mathbf{B}^\top \mathbf{x} \end{aligned}$$

**Note:**  $\mathbf{B}^\top \mathbf{B}$  is invertible  $\Rightarrow \boldsymbol{\lambda} = (\mathbf{B}^\top \mathbf{B})^{-1} \mathbf{B}^\top \mathbf{x}$ .

## Projection onto General Subspaces (4/4)

Since

$$\begin{aligned}\mathbf{b}_1^\top (\mathbf{x} - \mathbf{B}\lambda) &= 0 \\ \vdots \\ \mathbf{b}_m^\top (\mathbf{x} - \mathbf{B}\lambda) &= 0\end{aligned}$$

We have

$$\begin{aligned}\begin{bmatrix} \mathbf{b}_1^\top \\ \vdots \\ \mathbf{b}_m^\top \end{bmatrix} [\mathbf{x} - \mathbf{B}\lambda] = \mathbf{0} &\Leftrightarrow \mathbf{B}^\top (\mathbf{x} - \mathbf{B}\lambda) = \mathbf{0} \\ &\Leftrightarrow \mathbf{B}^\top \mathbf{B}\lambda = \mathbf{B}^\top \mathbf{x}\end{aligned}$$

**Note:**  $\mathbf{B}^\top \mathbf{B}$  is invertible  $\Rightarrow \lambda = (\mathbf{B}^\top \mathbf{B})^{-1} \mathbf{B}^\top \mathbf{x}$ .

- $\pi_U(\mathbf{x}) = \mathbf{B}(\mathbf{B}^\top \mathbf{B})^{-1} \mathbf{B}^\top \mathbf{x}$

## Projection onto General Subspaces (4/4)

Since

$$\mathbf{b}_1^\top (\mathbf{x} - \mathbf{B}\lambda) = 0$$

$$\vdots$$

$$\mathbf{b}_m^\top (\mathbf{x} - \mathbf{B}\lambda) = 0$$

We have

$$\begin{bmatrix} \mathbf{b}_1^\top \\ \vdots \\ \mathbf{b}_m^\top \end{bmatrix} [\mathbf{x} - \mathbf{B}\lambda] = \mathbf{0} \Leftrightarrow \mathbf{B}^\top (\mathbf{x} - \mathbf{B}\lambda) = \mathbf{0}$$
$$\Leftrightarrow \mathbf{B}^\top \mathbf{B}\lambda = \mathbf{B}^\top \mathbf{x}$$

**Note:**  $\mathbf{B}^\top \mathbf{B}$  is invertible  $\Rightarrow \lambda = (\mathbf{B}^\top \mathbf{B})^{-1} \mathbf{B}^\top \mathbf{x}$ .

•  $\pi_U(\mathbf{x}) = \mathbf{B}(\mathbf{B}^\top \mathbf{B})^{-1} \mathbf{B}^\top \mathbf{x} \Rightarrow$  Projection matrix  $\mathbf{P}_\pi = \mathbf{B}(\mathbf{B}^\top \mathbf{B})^{-1} \mathbf{B}^\top$ .

## Example

### Example

For a subspace  $U = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix} \right\} \subseteq \mathbb{R}^3$  and  $\mathbf{x} = \begin{bmatrix} 6 \\ 0 \\ 0 \end{bmatrix} \in \mathbb{R}^3$ .

Find

- the coordinates  $\lambda$  of  $\mathbf{x}$  in terms of  $U$
- the projection point  $\pi_U(\mathbf{x})$
- the projection matrix  $\mathbf{P}_\pi$ .

p. 87; on the black board.

What if  $B = (\mathbf{b}_1, \dots, \mathbf{b}_m)$  is orthonormal?

- $\pi_U(\mathbf{x}) = B(B^\top B)^{-1}B^\top \mathbf{x}$

What if  $B = (\mathbf{b}_1, \dots, \mathbf{b}_m)$  is orthonormal?

- $\pi_U(\mathbf{x}) = B(\mathbf{B}^\top B)^{-1} \mathbf{B}^\top \mathbf{x} \Rightarrow \pi_U(\mathbf{x}) = BB^\top \mathbf{x}.$ 
  - $\because \mathbf{B}^\top B = I.$
- Coordinates:  $\lambda = (B^\top B)^{-1} B^\top \mathbf{x} = B^\top \mathbf{x}.$

# Outline

- 1 Orthogonal Projections
- 2 Gram-Schmidt Orthogonalization




# Illustration of Gram-Schmidt Orthogonalization


- Goal:** Transform any basis  $(\mathbf{b}_1, \dots, \mathbf{b}_n)$  of an  $n$ -dimensional vector space  $V$  into an orthogonal/orthonormal basis of  $V$ .

$$\mathbf{u}_1 := \mathbf{b}_1$$


$$\mathbf{u}_k := \mathbf{b}_k - \pi_{\text{span}(\{\mathbf{u}_1, \dots, \mathbf{u}_{k-1}\})}(\mathbf{b}_k), \quad k = 2, \dots, n.$$



(a) Original non-orthogonal basis vectors  $\mathbf{b}_1, \mathbf{b}_2$ .



(b) First new basis vector  $\mathbf{u}_1 = \mathbf{b}_1$  and projection of  $\mathbf{b}_2$  onto the subspace spanned by  $\mathbf{u}_1$ .



(c) Orthogonal basis vectors  $\mathbf{u}_1$  and  $\mathbf{u}_2 = \mathbf{b}_2 - \pi_{\text{span}[\mathbf{u}_1]}(\mathbf{b}_2)$ .

## Example

### Example


Consider a basis  $(\mathbf{b}_1, \mathbf{b}_2)$  of  $\mathbb{R}^2$ , where  $\mathbf{b}_1 = \begin{bmatrix} 2 \\ 0 \end{bmatrix}$ ,  $\mathbf{b}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ .

Apply the Gram-Schmidt method to construct an orthonormal basis  $(\mathbf{u}_1, \mathbf{u}_2)$  of  $\mathbb{R}^2$  (assuming the dot product as the inner product).


p. 89; on the black board.

# Projection onto Affine Spaces


- Given an affine space  $L = \mathbf{x}_0 + U$ .
  - $U$  is a low-dimensional subspace of  $V$ .
- $\pi_L(\mathbf{x}) = \mathbf{x}_0 + \pi_U(\mathbf{x} - \mathbf{x}_0)$



(a) Setting.



(b) Reduce problem to projection  $\pi_U$  onto vector subspace.



(c) Add support point back in to get affine projection  $\pi_L$ .

# Discussions