

Worksheet 3

Observables and Operators

Quantum States • Observables • Operators • Measurement

1. Consider a *quantum mouse* – this is like a normal mouse but its properties behave in a quantum way.¹ There are lots of things we can observe about the mouse, like its colour, or tail length, or eye size, or mood, although for this worksheet we'll stick to making measurements of the mouse's

- tail length, symbolized by T (with the corresponding operator $\hat{T}$); observations always measure one of two possible values, 5 cm (short-tailed) or 10 cm (long-tailed), and
- mood, symbolized by M (with the corresponding operator $\hat{M}$); observations always measure one of two possible moods, either happy (which we'll measure as $M = +1$) or sad (for which we'd measure $M = -1$).

If you measure the tail length of a mouse and get $T = 5$ cm, we'll say the mouse is in state $|-\rangle$, and if you instead get $T = 10$ cm, it'll correspond to state $|-\rangle$. On the other hand, if you measure the mood of the mouse and get $M = +1$, we call the state $|\odot\rangle$, and the state $|\odot\rangle$ will correspond to getting a measurement of $M = -1$.

- (a) Write out the eigenvalue equations for each operator, $\hat{T}$ and $\hat{M}$. Identify which parts of the equations are the operator, the eigenvector, and the eigenvalue.

- (b) What do you think the operation $\langle - | - \rangle$ should be? What about $\langle \odot | \odot \rangle$?

¹This idea was borrowed from the tutorials produced at the University of Colorado Boulder.

2. We're going to work in the "mood" basis for the rest of this worksheet – that is, we'll express every operator and every state in terms of the two basis states of $\hat{M}$ (the kets $|\odot\rangle$ and $|\ominus\rangle$).

- (a) Write out the two basis states $|\odot\rangle$ and $|\ominus\rangle$ in matrix notation, and write out the operator $\hat{M}$ in matrix notation.

- (b) What about the two eigenstates of the tail length operator $\hat{T}$? Well, I'll tell you that

$$|-\rangle = \frac{1}{\sqrt{5}}|\odot\rangle + \frac{2}{\sqrt{5}}|\ominus\rangle.$$

Would say that mice with short tails are more happy or more sad?

- (c) What about the ket $|--\rangle$? Use the properties of orthonormality and completeness to figure out what it is (in other words, write out $|--\rangle = a|\odot\rangle + b|\ominus\rangle$ and figure out a and b).

- (d) Is your values of a and b unique? Does it matter if they're not?

3. Now let's make some observations. Suppose you have a quantum mouse, you measure the tail length, and find $T = 5$ cm.

(a) What quantum state is the mouse in now? Is there any ambiguity about the *state* at this point?

(b) You then re-measure the tail length T . What results would you get, and with what probabilities? What state is the mouse in after this second measurement?

(c) If you now measure the mood M of the mouse, what are the possible results and probabilities?

5. Last question, time permitting. We've written the operator $\hat{M}$ and all the state in the mood basis. What about the operator $\hat{T}$? That's a little trickier.

(a) First, write the matrix representation of $\hat{T}$ in the M basis explicitly like this:

$$\hat{T} \rightarrow \begin{pmatrix} \langle \ominus | \hat{T} | \ominus \rangle & \langle \ominus | \hat{T} | \oplus \rangle \\ \langle \oplus | \hat{T} | \ominus \rangle & \langle \oplus | \hat{T} | \oplus \rangle \end{pmatrix}$$

- (b) Then work out the action of the operator $\hat{T}$ on the states $|\ominus\rangle$ and $|\oplus\rangle$. Careful with this – we only know how $\hat{T}$ operates on its eigenstates (see question 1 (a) above), so you'll have to write $|\ominus\rangle$ and $|\oplus\rangle$ in terms of the states $|-\rangle$ and $|+\rangle$.
- (c) Then complete the matrix elements by doing the inner product with the bras.