

Contents

1	Probability	4
1.0.0.1	Sample Space and Outcome	4
1.0.0.2	Event	4
1.0.0.3	Disjunction of Events	4
1.0.0.4	Conjunction of Events	4
1.0.0.5	Mutually Exclusive Events	4
1.0.0.6	Axioms of Probability	4
1.0.0.7	Properties	5
1.0.0.8	Conditional Probability	5
1.0.0.9	Joint Probability	6
1.0.0.10	Marginal Probability	6
2	Stochastic Processes	7
2.0.1	Classifications	7
2.0.1.1	State Space	7
2.0.1.2	Discrete State Space	7
2.0.1.3	Continuous State Space	7
2.0.1.4	Time Parameter	7
2.0.1.5	Discrete Time Parameter	7
2.0.1.6	Continuous Time Parameter	7
2.1	Discrete State Space and Time	7
2.1.0.7	Markov Chain	8
2.1.0.8	Homogeneous Markov Chains	8
2.1.0.9	State Transistion Diagram	8
2.1.0.10	Chapman-Kolmogorov Equation	8
2.1.0.11	Irreducible Markov Chain	9
2.1.0.12	Recurrent State	9
2.1.0.13	Recurrent Non-null	10
2.1.0.14	Recurrent Null	10
2.1.0.15	Periodicity	10
2.1.0.16	State Probability	10
2.1.0.17	Steady State Therom	11
2.1.0.18	Residence Time in a State	12
2.2	Continuous Time	12
2.2.0.19	Assumption: Continuos Time Markov Chain .	12
2.2.0.20	Transistion Probability	13

2.2.0.21	Apply Chapman-Kolmogorov Equation	13
2.2.0.22	Partial of $P_{ij}(v, t)$	13
2.2.0.23	Rate Definitions for Simplificaiton	14
2.2.0.24	State Probabilitiy	14
2.2.0.25	Steady State Results	14
2.2.0.26	Residence Time in State	14
3	Derivations	15
3.1	Probability	15
3.2	Stochastic Processes	15
3.2.0.27	Chapman-Kolmogorov Equation	15
3.2.0.28	Partial of $P_{ij}(v, t)$	15
3.2.0.29	Derivative of $\pi_j(t)$	15
4	Results	16
4.1	Analytical Results	16
4.1.1	Single Server	16
4.1.2	Open Network Model	17
4.1.3	Finite Population	17
4.1.4	Closed Network Model: No Users (ie: Cycle)	18
4.1.5	Closed Network Model: Web Application	19
4.2	Probability	20
4.2.0.1	Sample Space	20
4.2.0.2	Event	20
4.2.0.3	Disjunction of Events	20
4.2.0.4	Conjunction of Events	20
4.2.0.5	Mutually Exclusive	20
4.2.0.6	Axioms of Probability	20
4.2.0.7	Corollaries from Axioms	20
4.2.0.8	Conditional Probability	20
4.2.0.9	Joint Probability	20
4.2.0.10	Marginal Probability	20
4.2.0.11	Independent Events	20
4.2.0.12	Total Probability	20
4.2.0.13	Bayes Rule	20
4.2.0.14	Generalized Bayes Rule	20
4.2.1	Discrete	21
4.2.1.1	Random Variable	21

4.2.1.2	Probability Mass Function	21
4.2.1.3	Mean	21
4.2.1.4	Variance	21
4.2.1.5	Cumulative Distribution Function	21
4.2.1.6	Bernoulli	21
4.2.1.7	Binomial	21
4.2.1.8	Poisson	21
4.2.1.9	Geometric	21
4.2.1.10	Memoryless Property	21
4.2.2	Continuous	21
4.2.2.1	Probability Density Function	21
4.2.2.2	21
4.2.2.3	21
4.2.2.4	21
4.3	Stochastic Processes	22
4.4	Statistics	22
5	Useful Formulas	23
5.0.1	Finite Sums	23
5.0.2	Infinite Sums	23

This book is separated into three parts. The first, are explanation of the results, and informal proofs. The second part is the formula proof. The final part is all the results enumerated.

1 Probability

1.0.0.1 Sample Space and Outcome We perform random experiments and the sample space is the set of possible outcomes. For example, consider rolling a die. The set of possible outcomes are:

$$S = \{1, 2, 3, 4, 5, 6\}$$

1.0.0.2 Event An event is a subset of the sample space. An example event is rolling a die and getting an even odd outcome:

$$E = \{1, 3, 5\}$$

1.0.0.3 Disjunction of Events The event E occurs if E_1 or E_2 occur. Another way to imagine this is the union of events: $E = E_1 \cup E_2$.

1.0.0.4 Conjunction of Events The event E occurs if E_1 and E_2 occur. Another way to imagine this is the intersection of events: $E = E_1 \cap E_2$. Some alternative ways of writing this are:

$$P(E) = P(E_1 \cap E_2)$$

$$P(E) = P(E_1 \wedge E_2)$$

$$P(E) = P(E_1, E_2)$$

$$P(E) = P(E_1 E_2)$$

1.0.0.5 Mutually Exclusive Events Events E_1 and E_2 are mutually exclusive if only one of them can occur in a single experiment. For example, the event rolling an even number and the event rolling an odd number on a die are mutually exclusive events:

$$E_{\text{even}} \cap E_{\text{odd}} = \{2, 4, 6\} \cap \{1, 3, 5\} = \emptyset$$

1.0.0.6 Axioms of Probability These are the rules we accept as truth without proof. We build probability atop of these axioms.

1. $0 \leq P(E) \leq 1$, for any event E . In the smallest case, the event cannot occur which is indicated by a probability of 0. In the largest case, the event always occurs, which is indicated by the probability of 1.

2. $P(S) = 1$, where S is the sample space. The sample space contains all possible outcomes for each experiment. It's reasonable to accept that an event from the sample space always occurs.
3. For a potentially infinite set of mutually exclusive events E_1, E_2, \dots

$$P(\cup_{i=1}^{\infty} E_i) = \sum_{i=1}^{\infty} P(E_i)$$

It makes sense that events that do not share outcomes for a single event, can have their probabilities added to arrive at the probability of combining the outcomes from the events.

1.0.0.7 Properties From the above axioms, we get the following useful properties (TODO proof):

1. For any event E , let \bar{E} be the complement of E . More concretely, $\bar{E} = S - E$, where S is the sample space. Then E and \bar{E} are mutually exclusive.
2. $P(\emptyset) = 0$ You can never get none of the outcomes of the sample space.
3. If E_1 and E_2 are mutually exclusive events then

$$P(E_1 \cup E_2) = P(E_1) + P(E_2) - P(E_1, E_2)$$

1.0.0.8 Conditional Probability We use conditional probability to model the probability given knowing some circumstance has happened. Given two event E and F , the conditional probability, the probability of F given E has occurred, is defined as ($P(E) \neq 0$):

$$P(F|E) = \frac{P(E, F)}{P(E)}$$

An example is what is the probability of rolling a 3, given that we rolled an odd number. Let $F = \{3\}$ and $E = \{1, 3, 5\}$:

$$P(F|E) = \frac{P(E, F)}{P(E)} = \frac{\frac{1}{6}}{\frac{1}{2}} = \frac{1}{3}$$

1.0.0.9 Joint Probability In queuing theory, we often have to use multiple sample sapce. The theory in this book so far has covered only a single probability space.

Suppose we have two sample spaces S_1 and S_2 . The outcomes of the joint probability space are the tuples that result from the cross product of the two sample spaces:

$$S_{joint} = S_1 \times S_2$$

For example, consider rolling a die and a coin

$$\begin{aligned} S_{die} &= \{1, 2, 3, 4, 5, 6\} \\ S_{coin} &= \{H, T\} \\ S &= S_{die} \times S_{coin} \\ S &= \{(1, H), (2, H), (3, H), (4, H), (5, H), (6, H), \\ &\quad (1, T), (2, T), (3, T), (4, T), (5, T), (6, T)\} \end{aligned}$$

1.0.0.10 Marginal Probability Given the joint probabilties, we might want to compute the probabilties of only one of the sample spaces.

For example, suppose that we know the joint probability of the number of jobs at server 1 and server two and we want to compute the probability of the number of jobs at server 1 only. We can apply Marginal probability to determine that.

2 Stochastic Processes

A family of random variables, indexed by time

2.0.1 Classifications

2.0.1.1 State Space The set of possible values (states).

2.0.1.2 Discrete State Space Example: the number of jobs in the system. ($S = 0, 1, 2, 3, \dots$). We will only deal with the discrete case in this class. To make notation easier the state is usually identified by the number.

2.0.1.3 Continuous State Space Example: motion of a particle. We will not be studying this in this class.

2.0.1.4 Time Parameter There are two ways to observe times in Stochastic processes.

2.0.1.5 Discrete Time Parameter We consider the states at X_0, X_1, \dots, X_n , and so on. For example, looking at the state of the system at the i^{th} hour.

2.0.1.6 Continuous Time Parameter The states are function of time t , $X(t)$.

2.1 Discrete State Space and Time

There might be a dependency between the previous time interval X_i 's and the states those time interval can be in that need to be model in the current X_n .

$$P(X_{n+1} = j | X_n = i_n, X_{n-1} = i_{n-1}, \dots, X_1 = i_1, X_0 = i_0)$$

The number of dependency combinations is exponential because X_{n+1} depends on X_n to X_0
and X_n depends on X_{n-1} to X_0
and so on.

2.1.0.7 Markov Chain As a result of the exponential size, we make a simplifying assumption. We only use the latest information. X_{n+1} only depends on X_n . Now we are left with transition probabilities:

$$P(X_{n+1}|X_n) = P(X_{n+1} = j|X_n = i_n, X_{n-1} = i_{n-1}, \dots, X_1 = i_1, X_0 = i_0)$$


Given $P(X_0 = i)$ for all i 's, we can compute any state. However, notice that the formula depends on n , the discrete time that has passed so far, which make analysis difficult still. For example, in the 9am one hour interval the number of jobs in a login system tends to be higher than at 2am.

2.1.0.8 Homogeneous Markov Chains We now make the assumption that the transition probabilities do not depend on time. For example, the transition probabilities for the number of webcrawling robots requesting a webpage remain the same despite the time. So we can write

$$P(X_{n+1} = j|X_n = i) = P(X_n = j|X_{n-1} = i), \forall n, i, j \geq 0$$

Which is abbreviated to P_{ij} , the probability of going from state i to state j .

2.1.0.9 State Transition Diagram It's a diagram that enumerates all possible state transitions, and annotates the edges with the probabilities of going from one state to the next. Below is an example of a state transition diagram.



2.1.0.10 Chapman-Kolmogorov Equation Let $P_{ij}^{(m)}$ be the m step transition probability from state i to j defined as:

$$P_{ij}^{(m)} = P(X_{n+m} = j|X_n = i)$$

To take m steps to go from (possibly visiting m in an intermediate state multiple itmes), you can take $m-1$ to steps. At step $m-1$, you can arrive at any state k . To go from each state k at the $m-1$ step to the m step, you just apply the transistion probability.

We can sum the probabilities because the probabillites of going from state k in the $m-1$ step to state j are mutually exclusive (they are mutually exclusive because they are different states).

Lastly, we can mulitple the probability of going to state k in $m-1$ steps by the probability of going to state j because the probabilities are independent.

$$P_{ij}^{(m)} = \sum_k p_{ik}^{(m-1)} p_{kj}$$

Which is exactly the same as taking 1 step, and then $m-1$ steps.

$$P_{ij}^{(m)} = \sum_k p_{ik} p_{kj}^{(m-1)}$$

For a derivation of this, see (3.2.0.27).

2.1.0.11 Irreducible Markov Chain allows every state to be reached from every other state for all pairs of states i and j . More concretely:

$$\forall i \forall j \neq i : \exists m_{ij} : P_{ij}^{(m_{ij})} > 0$$

2.1.0.12 Recurrent State : State j is recurrent if after leaving state j then you are guarenteed to eventually comeback.

Let $f_j^{(n)}$ be the probability that you first return to state j in n steps.

Notice that $f_j^{(0)} = 0$ because it's impossible to comeback without taking any steps. Also notice that $f_j^{(1)} > 0$ is only possible is the transistion pointing to itself $P_{jj} > 0$.

j is recurrent if and only if

$$f_j = \sum_{n=1}^{\infty} f_j^{(n)} = 1$$

2.1.0.13 Recurrent Non-null A state j is recurrent non-null if and only if we get back to state j , but it does not take forever. More concretely:

j is recurrent non-null if and only if j is recurrent ($f_j = 1$) and

$$M_j = \sum_{n=1}^{\infty} n f_j^{(n)} < \infty$$

Where M_j is the expected number of steps to come back.

2.1.0.14 Recurrent Null j is recurrent null if and only if j is recurrent and j is not recurrent non-null.

TODO EXAMPLE WITH DIAGRAM

TODO EXAMPLE WITH DIAGRAM

2.1.0.15 Periodicity A state j is periodic if and only if the only way to come back to state j is to take $r, 2r, 3r, \dots, cr$, steps.

If a state j is not periodic, it's called aperiodic

If state j as a self loop ($p_{jj} > 0$), then state j is aperiodic

If the system is a irreducible Markov Chain, and contains a self loop, then all states j are aperiodic.

2.1.0.16 State Probability Let X_n be the random variable for the state at interval n , then in a homogeneous Markov Chain we have that:

$$\pi_j^{(n)} = P(X_n = j) \text{ - at step } j$$

Let p_{ij} be the transition probability for going from state i to state j (independent of time, so it's the same for all intervals (X_n)), then

$$\pi_j^{(n+1)} = \sum_{i=0}^{\infty} p_{ij} \pi_i^{(n)} \text{ (by applying total probability)}$$

Given initial conditions $\pi_j^{(0)} \forall j$, we can compute all $\pi_j^{(i)}$ apply the above formula recursively.

2.1.0.17 Steady State Theorem (Equilibrium Probability Theorem) (I made up this name. It seems better than 'Fundamental Theorem.)

If a homogeneous Markov Chain is irreducible and aperiodic, then there exists a limiting probability (equilibrium):

$$\pi_j = \lim_{n \rightarrow \infty} \pi_j^{(n)}$$


which is independent of the initial conditions $\pi_j^{(0)}$.

Moreover, if all states j are recurrent non-null, then π_j is non zero and can be uniquely determined from the equations:

$$\pi_j = \sum_i p_{ij} \pi_i, \forall j$$

$$\sum_j \pi_j = 1$$

2.1.0.17.1 Example



Let the initial state be (1). This is the 'initial condition'

n	0	1	2	3	4	...	∞
$\pi_1^{(n)}$	1	0	.25	.187	.203		.2
$\pi_2^{(n)}$	0	.25	.062	.359	.254		.28
$\pi_3^{(n)}$	0	.75	.688	.454	.543		.52

To get the steady state column, you have to solve the system of equations that results from the above theorem. From the table we can see that the empirical basis for the steady state theorem. As you compute more columns, it approaches the values computed from the steady state equations.

2.1.0.18 Residence Time in a State The residence time, is the amount of time that the system spends in state j for m steps, given that we are already in state j .

$$\begin{aligned} &P(\text{in state } j \text{ for } m \text{ steps} | \text{already in state } j) \\ &= P(\text{in state } j \text{ for } m \text{ steps} \wedge \text{step } m+1 \text{ is not } j | \text{already in state } j) \end{aligned}$$

We have that the transition probabilities are independent of time (homogeneous), so we can multiple the individual probabilities. p_{jj} is the homogeneous transition probability of going from state j to state j .

$$= p_{jj}^m (1 - p_{jj})$$

2.2 Continuous Time

In continuous time, state transitions happen independent of clock ticks. We can extend the discrete time theory we have developed by considering the times $t_0, t_1, t_2, \dots, t_n$. These are the times when state transitions occur.

Note that, if we can accept accuracy within a time interval, then we can just use discrete time. Arrival and departure models generally use continuous time. This distribution is heavily connected with the exponential function.

2.2.0.19 Assumption: Continuous Time Markov Chain Let $X(t)$ be a random variable, where the distribution of the random variable depends on the real parameter t . For instance, at $t = 1.0$ the distribution of X might be exponential, but at $t = \pi$ the distribution might be normal. Given the above notation for state transition, we have the following expression for the probability of the $(n + 1)^{th}$ state transition.

$$P(X(t_{n+1}) = j | X(t_n) = i_n, \dots, X(t_0) = i_0)$$

Applying the Markov assumption that the next state transition only depends on the previous state gives us:

$$P(X(t_{n+1}) = j | X(t_n) = i_n)$$

All the definitions we created in discrete time for recurrent, irreducible, etc apply to continuous time as well by considering only the times when state transitions occur.

2.2.0.20 Transition Probability Let the transition probability be the probability of going from one state i at time v , to another state at time t .

$$P_{ij}(v, t) = P(X(t) = j | X(v) = i)$$

There is a special for the above formula:

$$P_{ij}(t, t) = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$$

2.2.0.21 Apply Chapman-Kolmogorov Equation We can apply the Chapman-Kolmogorov Equation to the transition probability. The probability of going

1. from state i at time v
2. to state j at time t

is the same as going

1. from state i at time v
2. to state k at some intermediate time u (summed over all possible k 's)
3. to state j at time t

gives us the following equation:

$$P_{ij}(v, t) = \sum_{k=1}^{\infty} P_{ik}(v, u) P_{kj}(u, t)$$

2.2.0.22 Partial of $P_{ij}(v, t)$

$$\frac{\partial P_{ij}(v, t)}{\partial t} = \lim_{\Delta t \rightarrow 0} \sum_{k \neq j} P_{ik}(v, t) \frac{P_{kj}(t, t + \Delta t) - P_{kj}(t, t)}{\Delta t} + P_{ij}(v, t) \frac{P_{jj}(t, t + \Delta t) - 1}{\Delta t}$$

For a derivation of this, see (3.2.0.28).

2.2.0.23 Rate Definitions for Simplification We define q_{ij} in order to simplify the partial expression above. Additionally, we define q_{ij} in such a way that $\sum_j q_{ij}(t) = 0 \forall i, t$

$$q_{kj}(t) = \lim_{\Delta t \rightarrow 0} \frac{P_{kj}(t, t + \Delta t) - 1}{\Delta t} \quad k \neq j$$

$$q_{jj}(t) = \lim_{\Delta t \rightarrow 0} \frac{P_{jj}(t, t + \Delta t) - 1}{\Delta t}$$

This gives us the intended simplification of the partial:

$$\frac{\partial P_{ij}(v, t)}{\partial t} = \lim_{\Delta t \rightarrow 0} \sum_{k \neq j} P_{ik}(v, t) q_{kj}(t) + P_{ij}(v, t) q_{jj}(t)$$

These q_{ij} 's become useful later on as they become the arrival and departure rates in a birth-death process.

2.2.0.24 State Probability Let $\pi_j(t)$ be the probability of being in state j , at time t .

$$\pi_j(t) = P(X(t) = j)$$

By applying total probability, over some earlier time v we get:

$$\pi_j(t) = \sum_i P_{ij}(v, t) \pi_i(v) \quad (\text{which is identical to the discrete case})$$

Taking the derivative we arrive at (see derivation (3.2.0.29)):

$$\frac{d\pi_j(t)}{dt} = \sum_{k \neq j} \pi_k(t) q_{kj}(t) + \pi_j(t) q_{jj}(t)$$

Applying the assumption that the transition probabilities are homogeneous (independent over time), we have that $q_{ij}(t) = q_{ij}$

$$\frac{d\pi_j(t)}{dt} = \sum_{k \neq j} \pi_k(t) q_{kj} + \pi_j(t) q_{jj}$$

2.2.0.25 Steady State Results

2.2.0.26 Residence Time in State

3 Derivations

3.1 Probability

3.2 Stochastic Processes

3.2.0.27 Chapman-Kolmogorov Equation For context see (2.1.0.10). TODO

3.2.0.28 Partial of $P_{ij}(v, t)$ For context see (2.2.0.22). TODO

3.2.0.29 Derivative of $\pi_j(t)$ For context see (2.2.0.24).

4 Results

4.1 Analytical Results

4.1.1 Single Server

Let λ be the arrival rate. Then the average inter-arrival time (time between successive arrivals) is given by:

$$InterarrivalTime = \frac{1}{\lambda}$$

Let μ be the the service rate. Then the average service time is given by:

$$ServiceTime = \frac{1}{\mu}$$

The service time of a job is computed by:

$$ServiceTime = \frac{ServiceRequirement}{ServerCapacity}$$

Let n be the number of jobs that arrived in time period $(0,L)$.

Assume that n is also the number of jobs processed in time period $(0,L)$.

Let x_j be the service time of the j^{th} job.

Let Y be the throughput, the rate at which jobs are completed. Then

$$Y = \lambda = \frac{n}{L}$$

Let S be the mean service time. Then:

$$S = \frac{1}{n} \sum_{j=1}^n x_j$$

Let U be the utilization, the percentage of time that the server is busy.

Then:

$$U = \frac{1}{L} \sum_{j=1}^n x_j = \lambda S = YS$$

Let r_j be the response time of the j^{th} job.

Let R be the mean response time. Then

$$R = \frac{1}{n} \sum_{j=1}^n r_j$$

Let Q be the mean number of jobs in the system. Then

$$Q = \frac{1}{L} \sum_{j=1}^n r_j = \lambda R = YR \text{ (Little's Law)}$$

4.1.2 Open Network Model

Let p_{ij} be the probability goes from server i to server j after finishing at i .
The $(M + 1)^{th}$ $p_{i,(M+1)}$ is the probability for leaving the system from server i .

The probability of take any of the edges when leaving a server i is one, so:

$$\sum_{j=1}^{M+1} p_{ij} = 1 \quad \forall i$$

Let γ_i be the arrival rate of jobs coming from outside the system.

The arrival rates at each server can be solved using the following equations (M unknowns are the λ_i 's).

$$\lambda_i = \gamma_i + \sum_{j=1}^M \lambda_j p_{ji}$$

Let U_i be the utilization of server i .

Let Q_i be the mean number of jobs at server i .

Let R_i be the mean response time at server i .

Let γ be the total arrival rate of jobs entering the system.

Let Y be the system throughput.

Let R be the system response time. Then:

$$U_i = \lambda_i S_i$$

$$Q_i = \lambda_i R_i$$

$$Q = \sum_{i=1}^M Q_i$$

$$Y = \sum_{j=1}^M \lambda_j P_{j,(M+1)} = \sum_{i=1}^M \gamma_i$$

$$R = \frac{Q}{\gamma} = \frac{1}{\gamma} \sum_{i=1}^M \lambda_i R_i$$

4.1.3 Finite Population

Let N be the number of users.

Let Z be the mean think time.

Recall that S is the mean service time.

Recall that L is the length of the observation period.

Recall that we assume n is the number of jobs that enter equal to the number that leave the system.

$$R = \frac{N}{Y - Z}$$

We know that

$$U = YS \leq 1$$

So

$$\begin{aligned} N^* &= \frac{S + Z}{Z} \\ R &\geq \begin{cases} S & 1 \leq N \leq N^* \\ NS - Z & N > N^* \end{cases} \\ Y &\leq \begin{cases} \frac{N}{S + Z} & 1 \leq N \leq N^* \\ \frac{1}{S} & N > N^* \end{cases} \end{aligned}$$

4.1.4 Closed Network Model: No Users (ie: Cycle)

Let p_{ij} be the probability goes from server i to server j after finishing at i .

Notice there is no $p_{i,0}$ or $p_{i,(M+1)}$

The probability of take any of the edges when leaving a server i is one, so:

$$\sum_{j=1}^M p_{ij} = 1 \quad \forall i$$

Recall λ_i is the arrival rate of jobs at server i

We get the follow set if linearly **dependant** :

$$\lambda_i = \sum_{j=1}^M \lambda_j p_{ji}$$

We can only determine:

$$\begin{aligned} \frac{\lambda_i}{\lambda_j} &= \frac{Y_i}{Y_j} \\ \frac{U_i}{U_j} &= \frac{\lambda_i S_i}{\lambda_j S_j} \end{aligned}$$

Let U_b be the server with highest utilization. Then:

$$U_i \leq \frac{\lambda_i S_i}{\lambda_b S_b}$$

4.1.5 Closed Network Model: Web Application

Let p_{ij} be the probability goes from server i to server j after finishing at i .

The 0^{th} $p_{i,0}$ is the probability of the request returning to the user.

The probability to take any of the edges when leaving a server i is one, so:

$$\sum_{j=0}^M p_{ij} = 1 \quad \forall i$$

We get the follow set if linearly **dependant** :

$$\lambda_i = \sum_{j=0}^M \lambda_j p_{ji}$$

Let V_i the the visit ration:

$$V_i = \frac{\lambda_i}{\lambda_0}$$

Let $D_i = V_i S_i$ (???)

The throughput of the system is total arrival rate from the users. This gives us:

$$\begin{aligned} Y &= \lambda_0 \\ \lambda_i &= Y V_i \\ U_i &= \lambda_i S_i = Y V_i S_i = Y D_i \\ \frac{U_i}{U_j} &= \frac{Y D_i}{Y D_j} = \frac{D_i}{D_j} \end{aligned}$$

Let Z be the mean think time.

Let b the the server with the highest utilization. Then:

$$\begin{aligned} U_i &\leq \frac{D_i}{D_b} \\ N^* &= \frac{D + Z}{D_b} \\ R(N) &\geq \begin{cases} D & 1 \leq N \leq N^* \\ ND_b - Z & N > N^* \end{cases} \\ Y(N) &\leq \begin{cases} \frac{N}{D + Z} & 1 \leq N \leq N^* \\ \frac{1}{D_b} & N > N^* \end{cases} \end{aligned}$$

4.2 Probability

4.2.0.1 Sample Space : set of all possible outcomes

4.2.0.2 Event : a subset of the space space

4.2.0.3 Disjunction of Events : $E = E_1 \cup E_2$

4.2.0.4 Conjunction of Events : $E = E_1 \cap E_2$ also written as $E_1 E_2$

4.2.0.5 Mutually Exclusive : E_1 and E_2 are mutually exclusive
 $\iff E_1 \cap E_2 = \emptyset$

4.2.0.6 Axioms of Probability

$$0 \leq P(event) \leq 1$$

$$P(SampleSpace) = 1$$

$$P(\cup_{i=1}^{\infty} E_i) = \sum_{i=1}^{\infty} P(E_i) \text{ where all } E_i \text{ are mutually exclusive}$$

4.2.0.7 Corollaries from Axioms Let E^c be the compliment of E, then E and E^c are mutually exclusive

$$P(\emptyset) = 0$$

If E_1 and E_2 are not mutually exclusive, then

$$P(E_1 \cup E_2) = P(E_1) + P(E_2) - P(E_1 E_2).$$

4.2.0.8 Conditional Probability $P(F|E) = \frac{P(EF)}{P(E)}$

4.2.0.9 Joint Probability You have n samples space to draw from.

The Joint Probability is the cross product of those spaces. IE:

$S_1 \times S_2 \times \dots \times S_n$ and joint probability given by

$E_1 \in S_1, E_2 \in S_2, \dots, E_n \in S_n$ and $P(E_1, E_2, \dots, E_n)$

4.2.0.10 Marginal Probability : Sum over all other sample spaces:

$$P(S_1) = \sum_{j_2=1}^{N^2} \dots \sum_{j_n=1}^{N^n} P(E_1, E_2, \dots, E_n)$$

4.2.0.11 Independent Events Events E and F are independent

$$\iff P(EF) = P(E)P(F) \text{ equivalently } P(F|E) = P(F)$$

4.2.0.12 Total Probability $P(F) = \sum_{i=1}^n P(F|E_i)P(E_i)$

4.2.0.13 Bayes Rule $P(E|F) = \frac{P(F|E)P(E)}{P(F)}$

4.2.0.14 Generalized Bayes Rule $P(E_i|F) = \frac{P(F|E_i)P(E_i)}{\sum_j P(F|E_j)P(E_j)}$

4.2.1 Discrete

4.2.1.1 Random Variable A variable that can take on any value from the sample space, along with a probability distribution on which it takes those values.

4.2.1.2 Probability Mass Function $P(X = x_i)$ shortened sometimes to $P(x_i)$

4.2.1.3 Mean $E[X] = \sum_{i=1}^n x_i P(x_i)$

4.2.1.4 Variance $var(X) = \sum_{i=1}^n (x_i - E[X])^2 P(x_i)$

4.2.1.5 Cumulative Distribution Function

$$F(x) = \sum_{i \leq x} P(i)$$

$$F(-\infty) = 0$$

$$F(\infty) = 1$$

$$F(b) - F(a) = P(a < X \leq b) = \sum_{a < i \leq b} P(i)$$

4.2.1.6 Bernoulli $P(1) = a, P(0) = 1 - a, E[X] = a, var(X) = a(1 - a)$

4.2.1.7 Binomial n independant Bernoulli trials:
 $P(i) = \binom{n}{i} a^i (1 - a)^{n-i}, E[X] = na, var(X) = na(1 - a)$

4.2.1.8 Poisson $P(i) = \frac{e^{-\lambda} \lambda^i}{i!}, E[X] = \lambda, var(X) = \lambda$

4.2.1.9 Geometric Prob that the i th Bernoulli trial is a success:
 $P(i) = (1 - a)^{i-1} a, E[X] = \frac{1}{a}, var(X) = \frac{1-a}{a^2}$, has the memoryless property

4.2.1.10 Memoryless Property $P(X = i + k | X > k) = P(X = i)$

4.2.2 Continuous**4.2.2.1 Probability Density Function****4.2.2.2****4.2.2.3****4.2.2.4**

4.3 Stochastic Processes

4.4 Statistics

5 Useful Formulas

5.0.1 Finite Sums

$$\begin{aligned}\sum_{k=1}^n k &= \frac{n(n+1)}{2} \\ \sum_{k=1}^n k^2 &= \frac{n(n+1)(2n+1)}{6} \\ \sum_{k=0}^n z^k &= \frac{1-z^{n+1}}{1-z} \\ \sum_{k=0}^n z^k &= \frac{1-z^{n+1}}{1-z}\end{aligned}$$

5.0.2 Infinite Sums

$$\begin{aligned}\sum_{n=k}^{\infty} x^n &= \frac{x^k}{1-x} & 0 \leq x < 1 \\ \sum_{n=k}^{\infty} nx^n &= \frac{kx^k}{1-x} + \frac{x^{k+1}}{(1-x)^2} & 0 \leq x < 1 \\ \sum_{k=0}^{\infty} \frac{z^k}{k!} &= e^z \\ \sum_{k=0}^{\infty} k \frac{z^k}{k!} &= ze^z \\ \binom{a}{k} &= \frac{a!}{k!(a-k)!} \\ \sum_{k=0}^{\infty} \binom{a}{k} z^k &= (1+z)^a\end{aligned}$$