

4. Air Resonance

PURPOSE AND BACKGROUND

The concept of *resonance* in a pipe is similar to that of a string. The waves in pipes consist of compressions and rarefactions of the air, with back-and-forth motion of the air molecules in the direction of propagation or against it. The waves in air thus are *longitudinal* waves. In this laboratory we study standing waves in a pipe. They are the result of two waves traveling in opposite directions inside the pipe, with each wave being reflected at the ends of the pipe. In this way the superposition of two waves yields a standing wave, provided that in addition the *resonance conditions* are met.

For a pipe with both ends open, resonance at the *lowest* frequency (*fundamental frequency* or *first harmonic*) occurs when there are anti-nodes of the air motion at the ends – and only there, with a single velocity node at the center, see Figure 1. The motion of air molecules is highest at the anti-nodes and lowest at the nodes. For a pipe with one end closed and one end open, resonance




Figure 1: Air molecule displacement in an open and closed pipe.

at the lowest frequency occurs when we have a velocity node at the closed end and an anti-node at the open end. Plotted in Figure 1 is the displacement or velocity of air molecules as a function of position along the pipe. The two curves for each pipe in Figure 1 are one-half period of oscillation apart.

For the pipe with both ends open, we have $L = \lambda/2$ according to Figure 1. For the closed pipe we have $L = \lambda/4$. The fundamental frequency is given by

$$f_1 = \frac{v}{\lambda} = \frac{v}{2L} \quad (\text{both ends open}), \quad f_1 = \frac{v}{\lambda} = \frac{v}{4L} \quad (\text{one end closed}), \quad (1)$$

where v is the velocity of sound.

Questions

1. For a pipe with both ends open, what are the formulas for the fundamental frequency and the frequencies f_2 , f_3 , f_4 of the next three overtones or harmonics? (Hint: Higher harmonics have frequencies that are integer multiples of the fundamental, and all integers are allowed for a pipe with both ends open.)

- For a pipe with one end closed and one end open, we have $L = \lambda/4$ according to Figure 1. Write down the equations for the fundamental frequency and for the first three existing overtones. (Hint: *Only odd integers are allowed* as you can see by extending the drawings in Figure 1 to higher harmonics.)

EXPERIMENTAL PROCEDURE

In the lab, we can record the frequency spectrum of sound in a pipe using the setup shown in Figure 2. (This figure is specifically for a tube that is open at both ends.) The speaker produces *white noise* and the “Electroacoustic Toolbox” can be used to record spectrum. Figure 3 shows sample recorded frequency spectra for both types of tubes.




Figure 2: Set up of the resonance tube in the “open tube” configuration. White noise from the Mac mini is applied to the speaker. The sound enters the tube on the left and excites the resonances. The microphone on the right records them for display in the computer. In the “closed tube” configuration the speaker and microphone must be on the same (right) side.

I Pipe Length Correction

If you were to compare the calculated and observed fundamental frequencies from an actual experiment, you would find that they do not agree very well. This has to do with the fact that in pipes, waves reflect from the ends of the tube by sticking out a little bit. There is an end correction that increases the wavelength. This correction is proportional to the radius of the tube. Therefore, the larger the tube radius, the more the wave will “stick out” and cause an increase in wavelength. The correction results in an extra length ΔL , given from theory by $\Delta L = 0.61R$ for each open end, where R is the radius of the pipe. Thus for a closed pipe and open pipe of length L and radius R , the effective lengths are, respectively,

$$L_{\text{eff, closed}} = L + 0.61R, \quad L_{\text{eff, open}} = L + 1.22R. \quad (2)$$

Questions

- Calculate the effective lengths of a closed and open tube if $L = 131$ cm and the diameter of the tube is $D = 14.4$ cm. (Recall: The diameter D equals twice the radius R .)

II Determination of the Velocity of Sound




Figure 3: Resonances of a PASCO resonance tube excited with white noise. Upper figure: Tube open at both ends with an effective length $L_{\text{eff, open}} = 1.18$ m. Lower figure: Tube closed at one end with an effective length $L_{\text{eff, closed}} = L_{\text{eff, open}}/2 = 0.59$ m (with a plug in the tube to shorten its length). The fundamental frequency for both tubes is $f_1 = 146$ Hz, but only the odd harmonics are observed in the tube closed at one end.




Figure 4: Frequency spectrum from a cylindrical PASCO packing tube, tapped on the floor with the closed end. The large peak is the $N = 1$ harmonic (fundamental frequency), the small peak is the $N = 3$ harmonic. The $N = 2$ harmonic is missing, as is to be expected for a tube closed at one end.




Figure 5: Decaying waveform of the packing tube.




Figure 6: Experimental setup for spherical Helmholtz resonator.




Figure 7: Helmholtz resonance curve from a large aluminum sphere. The measured and calculated values of the resonance frequency at the peak are 92 Hz and 93 Hz, respectively.




Figure 8: Helmholtz resonance from a 0.75 liter wine bottle.