

Notes: Thurs 8/27

- 1) Elliptic Functions
- 2) Simple pendulum

go beyond small  
angle approx

Elliptic Functions:

$$\int \frac{dx}{\sqrt{1-x^2}} = ? = \sin^{-1}(x) + \text{const}$$

"   
 arcsin(x)

$$\rightarrow \sin^2 \theta + \cos^2 \theta = 1$$

subst:  $x = \sin \theta \rightarrow \sqrt{1-x^2} = \sqrt{1-\sin^2 \theta} = \sqrt{\cos^2 \theta} = \cos \theta$

$$dx = \cos \theta d\theta$$

$$\int \frac{\cos \theta d\theta}{\cos \theta} = \int d\theta = \theta = \sin^{-1}(x)$$

$$x = \sin \theta$$
$$\theta = \sin^{-1}(x)$$


$$\frac{d}{d\theta} \sin \theta = \cos \theta$$

$$\lim_{h \rightarrow 0} \left( \frac{\sin(\theta+h) - \sin \theta}{h} \right)$$

$$\sin x = x - \frac{1}{3!} x^3 + \frac{1}{5!} x^5 - \dots$$

$$\sin(A+B) = \sin A \cos B + \cos A \sin B$$

# circular functions..



Def:

$$\sin \theta = \frac{y}{a}, \quad \cos \theta = \frac{x}{a}$$

$s$ : arc length from  $(a, 0)$  to  $(x, y)$

$$s = a\theta \quad \left| \quad \theta = \frac{s}{a}$$

$$= \frac{1}{a} \int ds$$

$$= \frac{1}{a} \int \sqrt{dx^2 + dy^2}$$

$$= \int d\theta$$

$$\sqrt{dx^2 + dy^2} = ds$$

Given:  $x^2 + y^2 = a^2$

Follows: (i)  $a^2 \cos^2 \theta + a^2 \sin^2 \theta = a^2 \rightarrow \boxed{\cos^2 \theta + \sin^2 \theta = 1}$

(ii)  $\boxed{\frac{d}{d\theta} \sin \theta} = \frac{1}{a} \frac{dy}{d\theta} = \frac{1}{a} \frac{dy}{\frac{1}{a} \sqrt{dx^2 + dy^2}} = \frac{1}{\sqrt{\left(\frac{y}{x}\right)^2 + 1}}$

$x dx + y dy = 0 \rightarrow dx = \frac{-y}{x} dy \quad \left| \quad = \frac{1}{\sqrt{\frac{y^2 + x^2}{x^2}}} = \frac{x}{a} = \boxed{\cos \theta}$

$$\frac{d(\sin \theta)}{d\theta} = \cos \theta \rightarrow \int \frac{d(\sin \theta)}{\cos \theta} = \int d\theta$$


$$x = \sin \theta$$

$$\cos \theta = \sqrt{1 - \sin^2 \theta}$$

$$= \sqrt{1 - x^2}$$


$$\int \frac{dx}{\sqrt{1-x^2}} = \theta = \sin^{-1}(x)$$

Elliptic  
function




$$\left(\frac{x}{a}\right)^2 + \left(\frac{y}{b}\right)^2 = 1$$

eccentricity :  $e = 0$  (for circle)



$$d_1 + d_2 = 2a$$




$$(ae)^2 + b^2 = a^2$$

$$a^2 e^2 + b^2 = a^2$$

$$e^2 = \frac{a^2 - b^2}{a^2} = 1 - \left(\frac{b}{a}\right)^2$$

$$e = \sqrt{1 - \left(\frac{b}{a}\right)^2} \equiv k$$

notation used  
in elliptic  
function



$$\left(\frac{x}{a}\right)^2 + \left(\frac{y}{b}\right)^2 = 1$$

$$x^2 + y^2 = r^2$$

$$x = r \cos \theta, \quad y = r \sin \theta$$

Define:

$$\operatorname{cn}(u; k) \equiv \frac{x}{a}, \quad \operatorname{sn}(u; k) \equiv \frac{y}{b}$$

$$\operatorname{dn}(u; k) \equiv \frac{r}{a} \quad \text{--- } (=1 \text{ for a circle})$$

where  $u = \frac{1}{b} \int_0^\theta r d\theta$   $y = r \sin \theta$   
(=  $\theta$  for a circle)

$$ds = \sqrt{dx^2 + dy^2} = \sqrt{dr^2 + r^2 d\theta^2}$$

Given:  $\left(\frac{x}{a}\right)^2 + \left(\frac{y}{b}\right)^2 = 1$  ,  $x^2 + y^2 = r^2$

$$\operatorname{dn}(u; k) = \frac{r}{a}$$

Follows: (i)  $\operatorname{cn}^2(u; k) + \operatorname{sn}^2(u; k) = 1$

(ii)  $\operatorname{dn}^2(u; k) + k^2 \operatorname{sn}^2(u; k) = 1$

$$u = \frac{1}{b} \int_0^\theta r d\theta$$

(iii)  $\frac{d}{du} \operatorname{sn}(u; k) = \operatorname{cn}(u; k) \operatorname{dn}(u; k)$

$$\frac{d}{du} \operatorname{cn}(u; k) = -\operatorname{sn}(u; k) \operatorname{dn}(u; k)$$

$$\frac{d}{du} \operatorname{dn}(u; k) = -k^2 \operatorname{sn}(u; k) \operatorname{cn}(u; k)$$

Analogous to  
 $\frac{d \sin \theta}{d\theta} = \cos \theta$

$$\frac{d \sin \theta}{\cos \theta} = d\theta$$

$$x = \sin \theta$$

$$\int \frac{dx}{\sqrt{1-x^2}} = \int d\theta = \theta = \sin^{-1} x$$

Integrate:  $\frac{d \operatorname{sn}(u; k)}{du} = \operatorname{cn}(u; k) \operatorname{dn}(u; k)$

$$\int \frac{d \operatorname{sn}(u; k)}{\operatorname{cn}(u; k) \operatorname{dn}(u; k)} = \int du = u$$

$$\int \frac{dx}{\sqrt{1-x^2} \sqrt{1-k^2 x^2}} = u + \text{const} = \operatorname{sn}^{-1}(x; k) + \text{const}$$

$x \equiv \operatorname{sn}(u; k)$

$$\int \frac{dx}{\sqrt{1-x^2}} = \sin^{-1} x + \text{const}$$

$$\int_0^1 \frac{dx}{\sqrt{1-x^2} \sqrt{1-k^2 x^2}}$$

$$\equiv K(k)$$

(complete elliptic  
integral of 1st  
kind)

related to  
period of a pendulum


going beyond  
small-angle  
approximation

$$\int_0^1 \frac{\sqrt{1-k^2 x^2} dx}{\sqrt{1-x^2}}$$

$$\equiv E(k)$$

(complete elliptic  
integral of  
2nd kind)

circumference  
around an ellipse



circle:  $C = 2\pi a$



Notes: Tuesday 9/1

- 1) Review of elliptic functions,
- 2) Simple pendulum


$$u = \frac{1}{b} \int_{(a,0)}^{(x,y)} r d\theta$$

$$b u = \int_0^\theta r d\theta \leq \int_0^\theta ds = s$$

$$ds = \sqrt{dx^2 + dy^2} = \sqrt{dr^2 + r^2 d\theta^2} \geq r d\theta$$



Simple pendulum:



Torque =  $I \alpha = m l^2 \ddot{\phi}$

(i) 'Freshman physics'

forces, free-body diagram

→ EOM, tension

tangential:

$$-mg \sin \phi = m a_{\text{tangential}}$$

$$-mg \sin \phi = m l \ddot{\phi}$$

$\phi$ : angular displacement [rad]

$\dot{\phi}$ : angular velocity [rad/sec]

$\ddot{\phi}$ : angular accel [rad/sec<sup>2</sup>]

$$\boxed{\ddot{\phi} = -\frac{g}{l} \sin \phi} \quad (\text{EOM})$$

radial:  $T - mg \cos \phi = m a_{\text{centripetal}}$

$$T - mg \cos \phi = m \dot{\phi}^2 l$$

$$\boxed{T = mg \cos \phi + m \dot{\phi}^2 l}$$



(ii) Small angle approximation:

$$\sin \phi \approx \phi \leftarrow \phi \ll 1 = 57 \text{ degrees}$$

radian

$$\ddot{\phi} = -\frac{g}{l} \sin \phi \approx -\frac{g}{l} \phi \rightarrow$$

$$\ddot{\phi} = -\frac{g}{l} \phi$$

$$\phi(t) = \boxed{a} \cos(\omega t) + \boxed{b} \sin(\omega t)$$


where  $\omega \equiv \sqrt{\frac{g}{l}}$

determined by  
initial condition

small angle  
approx.

ICs: If  $\phi(0) = \phi_0$  (at rest)

then  $\boxed{\phi(t) = \phi_0 \cos(\omega t)}$



Period:  $P = \frac{2\pi}{\omega} = 2\pi \sqrt{\frac{l}{g}}$

independent of  $\phi_0$  !!

(iii) Lagrangian approach

$$L = T - U$$


$T \equiv$  Kinetic energy

$U \equiv$  Potential energy

$$T = \frac{1}{2} m l^2 \dot{\phi}^2 \quad \left( = \frac{1}{2} m (\dot{x}^2 + \dot{y}^2) \right)$$

$$y = l \cos \phi$$

$$x = l \sin \phi$$



$$U = -mgl \cos \phi + \text{const}$$

$$U = -mgy \quad \text{action}$$

$$U = mgl(1 - \cos \phi)$$

$$L = \frac{1}{2} m l^2 \dot{\phi}^2 + mgl \cos \phi$$

$$S = \int_{t_1}^{t_2} dt \mathcal{L}(\phi, \dot{\phi}, t)$$

$$\frac{d}{dt} \left( \frac{\partial L}{\partial \dot{\phi}} \right) = \frac{\partial L}{\partial \phi}$$

Lagrange's equation

$$\frac{d}{dt} (m l^2 \dot{\phi}) = -mgl \sin \phi$$

$$m l^2 \ddot{\phi} = -mgl \sin \phi$$

$\rightarrow$

$$\ddot{\phi} = -\frac{g}{l} \sin \phi \quad (\text{Eom})$$

(iv) solving  $\ddot{\phi} = -\frac{g}{l} \sin \phi$  (2<sup>nd</sup> order non-linear ODE)   
 $E = \text{const}$    
 $= T + U$    
 $= \frac{1}{2} m l^2 \dot{\phi}^2 - m g l \cos \phi$    
 hard!!

$E = 0 - m g l \cos \phi_0$  release from rest   
 $= -m g l \cos \phi_0$  from  $\phi = \phi_0$

$$-m g l \cos \phi_0 = \frac{1}{2} m l^2 \dot{\phi}^2 - m g l \cos \phi$$

$$-m g l (\cos \phi_0 - \cos \phi) = \frac{1}{2} m l^2 \dot{\phi}^2$$

$$\frac{d\phi}{dt} = \dot{\phi} = \sqrt{-2 \frac{g}{l} (\cos \phi_0 - \cos \phi)}$$

$$|\phi| \leq \phi_0$$

$$\int \frac{d\phi}{\sqrt{-2 \frac{g}{l} (\cos \phi_0 - \cos \phi)}} = \int dt = t + \text{const}$$

separable  
1<sup>st</sup> order  
ODE

$$t + t_0 = \int \frac{d\phi}{\sqrt{-\frac{2g}{r} (\cos \phi_0 - \cos \phi)}} \quad \left| \quad \frac{1}{\sqrt{a+bx^2}} \right.$$

substitution:

$$\cos \phi = 1 - 2 \sin^2\left(\frac{\phi}{2}\right)$$

$$\cos \phi_0 = 1 - 2 \sin^2\left(\frac{\phi_0}{2}\right)$$

$$\cos \phi = \cos\left(2\left(\frac{\phi}{2}\right)\right)$$

$$= \cos^2\left(\frac{\phi}{2}\right) - \sin^2\left(\frac{\phi}{2}\right)$$

$$= 1 - 2 \sin^2\left(\frac{\phi}{2}\right)$$

$$\rightarrow \cos \phi_0 - \cos \phi = -2 \left( \sin^2\left(\frac{\phi_0}{2}\right) - \sin^2\left(\frac{\phi}{2}\right) \right)$$

$$t + t_0 = \int \frac{d\phi}{2 \sqrt{\frac{g}{r}} \sqrt{\sin^2\left(\frac{\phi_0}{2}\right) - \sin^2\left(\frac{\phi}{2}\right)}}$$

$$= \frac{1}{2 \sqrt{\frac{g}{r}}} \int \frac{d\phi}{\sin\left(\frac{\phi_0}{2}\right) \sqrt{1 - \frac{\sin^2\left(\frac{\phi}{2}\right)}{\sin^2\left(\frac{\phi_0}{2}\right)}}}$$

$$|\phi| \leq \phi_0$$

$$\text{let } x = \frac{\sin\left(\frac{\phi}{2}\right)}{\sin\left(\frac{\phi_0}{2}\right)}$$


$$x = \frac{\sin(\frac{\phi}{2})}{\sin(\frac{\phi_0}{2})}$$

$$\rightarrow dx = \frac{1}{\sin(\frac{\phi_0}{2})} \frac{1}{2} \cos(\frac{\phi}{2}) d\phi$$

$$\sqrt{1-x^2}$$


↑  
denominator

Find this out

①  $\phi(t) =$  

② Period = ??

③ Redo the analysis using Lagrange multiplier for find tension in string

$t + t_0 = \int$   — integrated for  $\sin^{-1}(x; k)$

$$d\phi = \frac{2 \sin(\frac{\phi_0}{2}) dx}{\cos(\frac{\phi}{2})}$$

$$= \frac{2 \sin(\frac{\phi_0}{2}) dx}{\sqrt{1 - \sin^2(\frac{\phi}{2})}}$$

$$= \frac{2 \sin(\frac{\phi_0}{2}) dx}{\sqrt{1 - \underbrace{\sin^2(\frac{\phi_0}{2})}_{k^2} x^2}}$$

$k = \sin(\frac{\phi_0}{2})$

Lec #4 : Thurs 9/3

$$\phi(t) = 2 \sin^{-1} \left[ k \operatorname{sn} \left( \omega_0 \left( t + \frac{P}{4} \right); k \right) \right] \star$$

$$k = \sin \left( \frac{\phi_0}{2} \right), \quad \omega_0 = \sqrt{\frac{g}{L}}$$

$$P = 4 \sqrt{\frac{L}{g}} K(k) = 4 \sqrt{\frac{L}{g}} \int_0^1 \frac{dx}{\sqrt{1-x^2} \sqrt{1-k^2 x^2}}$$

small angle  
approx

$$P_0 = 2\pi \sqrt{\frac{L}{g}}$$

$$\frac{\sin \left( \frac{\phi}{2} \right)}{\sin \left( \frac{\phi_0}{2} \right)} = x = \operatorname{sn} \left[ \sqrt{\frac{g}{L}} (t + t_0); k \right]$$


$$\sqrt{\frac{L}{g}} \int \frac{dx}{\sqrt{1-x^2} \sqrt{1-k^2 x^2}} = \sqrt{\frac{L}{g}} \operatorname{sn}^{-1}(x; k) = t + t_0$$

$$\operatorname{sn}^{-1}(x; k) = \sqrt{\frac{g}{L}} (t + t_0)$$

$$P = 4\sqrt{\frac{L}{g}} K(k) = 2\pi\sqrt{\frac{L}{g}} \left( 1 + \frac{1}{16} \phi_0^2 + \dots \right)$$

Problem Landau 11.1

Lagrange multiplier:



$$\varphi(x, y) = x^2 + y^2 - l^2 = 0$$

constraint function

$$x^2 + y^2 = l^2 \sin^2 \phi + l^2 \cos^2 \phi = l^2$$

$$T = \frac{1}{2} m (\dot{x}^2 + \dot{y}^2) = \frac{1}{2} m l^2 \dot{\phi}^2$$

$$L = T - U + \lambda \phi$$

↑  $\lambda, *$

— Lagrange multiplier

$$L(x, \dot{x}, y, \dot{y}, t)$$

$$q = (x, y) \quad \dot{q} = (\dot{x}, \dot{y})$$

$$L(r, \dot{r}, \phi, \dot{\phi}, t)$$

$$q = (r, \phi) \quad \dot{q} = (\dot{r}, \dot{\phi})$$

$$\begin{pmatrix} \lambda(t) \\ r(t) \\ \phi(t) \end{pmatrix}$$

$$L(r, \dot{r}, \phi, \dot{\phi}, t)$$

$$\phi(x, y) = x^2 + y^2 - l^2 = 0$$

$$\phi(r, \phi) = r - l = 0$$

$$L = \frac{1}{2} m (\dot{r}^2 + r^2 \dot{\phi}^2) + \underbrace{m g r \cos \phi}_y + \lambda (r - l)$$

$$r: \quad \frac{d}{dt} \left( \frac{\partial L}{\partial \dot{r}} \right) = \frac{\partial L}{\partial r}$$

$$\rightarrow \cancel{m \ddot{r}} = m r \dot{\phi}^2 + m g \cos \phi + \lambda$$

$$\phi: \quad \frac{d}{dt} \left( \frac{\partial L}{\partial \dot{\phi}} \right) = \frac{\partial L}{\partial \phi}$$

$$\rightarrow \frac{d}{dt} (m r^2 \dot{\phi}) = -m g r \sin \phi$$

$$\lambda: \quad r - l = 0 \quad \checkmark$$

$$\cancel{2 m r \dot{r} \dot{\phi}} + m r^2 \ddot{\phi} = -m g r \sin \phi$$



$$r - l = 0 \rightarrow \begin{aligned} r &= l \\ \dot{r} &= 0 \\ \ddot{r} &= 0 \end{aligned}$$

---

$$\rightarrow \ddot{\phi} = -\frac{g}{l} \sin \phi$$

---

$$m l \dot{\phi}^2 + m g \cos \phi + \lambda = 0$$

$$\lambda = - \underbrace{(m g \cos \phi + m l \dot{\phi}^2)}_T$$

$$\lambda = -T$$

$$L = T - U + \lambda \varphi$$

$$\frac{d}{dt} \left( \frac{\partial L}{\partial \dot{x}} \right) = \frac{\partial L}{\partial x}$$

$$\frac{d}{dt} (m \dot{x}) = -\frac{\partial U}{\partial x} + \lambda \frac{\partial \varphi}{\partial x}$$

$$U = U(x, t)$$

$$T = \frac{1}{2} m (\dot{x}^2 + \dot{y}^2)$$

$$\varphi(x, y, t)$$

$$U(x, y, t)$$

$$\frac{d}{dt} (m \dot{y}) = -\frac{\partial U}{\partial y} + \lambda \frac{\partial \varphi}{\partial y}$$

$$\frac{d\vec{p}}{dt} = \vec{F}_{\text{net}}$$


applied force      force

$$\frac{dp_x}{dt} = F_x + \lambda \frac{\partial \varphi}{\partial x}$$

$$\frac{dp_y}{dt} = F_y + \lambda \frac{\partial \varphi}{\partial y}$$

$$\frac{d\vec{p}}{dt} = \vec{F} + \lambda \overset{\text{constraint}}{\nabla} \varphi$$

$$\left( \frac{\partial \varphi}{\partial x}, \frac{\partial \varphi}{\partial y} \right) = \vec{\nabla} \varphi$$




$$\vec{\nabla} \varphi \perp \varphi = \text{const}$$


$$\varphi = r - l$$

$$\vec{\nabla} \varphi = \vec{r}$$

$$\vec{\nabla} \varphi$$

$$\int_0 \vec{F}_c \cdot d\vec{r} = W_{0 \rightarrow r}$$






$$U = -m_1 g y_1 - m_2 g y_2$$

$$\phi_1, \phi_2$$

$$E \sim m$$



$$\lambda \phi$$

$$\left( \lambda_1 \phi_1 + \lambda_2 \phi_2 + \dots \right)$$

$$L \rightarrow E \circ M$$

$$L \rightarrow L + \frac{d}{dt} (f(q, t))$$



$L'$

~~$f(q, \dot{q}, t)$~~

