


Soc 5, prob 1



$$x_1 = l_1 \sin \phi_1$$

$$y_1 = l_1 \cos \phi_1$$

$$x_2 = x_1 + l_2 \sin \phi_2 = l_1 \sin \phi_1 + l_2 \sin \phi_2$$

$$y_2 = y_1 + l_2 \cos \phi_2 = l_1 \cos \phi_1 + l_2 \cos \phi_2$$

$$U = -m_1 g y_1 - m_2 g y_2$$

$$= -m_1 g l_1 \cos \phi_1 - m_2 g (l_1 \cos \phi_1 + l_2 \cos \phi_2)$$

$$= -(m_1 + m_2) g l_1 \cos \phi_1 - m_2 g l_2 \cos \phi_2$$

$$T = \frac{1}{2} m_1 (\dot{x}_1^2 + \dot{y}_1^2) + \frac{1}{2} m_2 (\dot{x}_2^2 + \dot{y}_2^2)$$

$$\dot{x}_1 = l_1 \cos \phi_1 \dot{\phi}_1 \quad \dot{y}_1 = -l_1 \sin \phi_1 \dot{\phi}_1$$

$$\dot{x}_1^2 = l_1^2 \cos^2 \phi_1 \dot{\phi}_1^2 \quad \dot{y}_1^2 = l_1^2 \sin^2 \phi_1 \dot{\phi}_1^2$$

$$\text{Thus, } \dot{x}_1^2 + \dot{y}_1^2 = l_1^2 (\sin^2 \phi_1 + \cos^2 \phi_1) \dot{\phi}_1^2$$

$$= l_1^2 \dot{\phi}_1^2$$

$$\dot{x}_2 = l_1 \cos \phi_1 \dot{\phi}_1 + l_2 \cos \phi_2 \dot{\phi}_2$$

$$\rightarrow \dot{x}_2^2 = l_1^2 \cos^2 \phi_1 \dot{\phi}_1^2 + l_2^2 \cos^2 \phi_2 \dot{\phi}_2^2 + 2l_1 l_2 \cos \phi_1 \cos \phi_2 \dot{\phi}_1 \dot{\phi}_2$$

$$\dot{y}_2 = -l_1 \sin \phi_1 \dot{\phi}_1 - l_2 \sin \phi_2 \dot{\phi}_2$$

$$\rightarrow \dot{y}_2^2 = l_1^2 \sin^2 \phi_1 \dot{\phi}_1^2 + l_2^2 \sin^2 \phi_2 \dot{\phi}_2^2 + 2l_1 l_2 \sin \phi_1 \sin \phi_2 \dot{\phi}_1 \dot{\phi}_2$$

From,

$$\dot{x}_2^2 + \dot{y}_2^2 = l_1^2 \dot{\phi}_1^2 + l_2^2 \dot{\phi}_2^2 + 2l_1 l_2 (\cos \phi_1 \cos \phi_2 + \sin \phi_1 \sin \phi_2) \dot{\phi}_1 \dot{\phi}_2$$

$$= l_1^2 \dot{\phi}_1^2 + l_2^2 \dot{\phi}_2^2 + 2l_1 l_2 \cos(\phi_1 - \phi_2) \dot{\phi}_1 \dot{\phi}_2$$

$$\text{so, } T = \frac{1}{2} m_1 (\dot{x}_1^2 + \dot{y}_1^2) + \frac{1}{2} m_2 (\dot{x}_2^2 + \dot{y}_2^2)$$

$$= \frac{1}{2} m_1 l_1^2 \dot{\phi}_1^2 + \frac{1}{2} m_2 l_1^2 \dot{\phi}_1^2 + \frac{1}{2} m_2 l_2^2 \dot{\phi}_2^2$$

$$+ m_2 l_1 l_2 \cos(\phi_1 - \phi_2) \dot{\phi}_1 \dot{\phi}_2$$

$$= \frac{1}{2} (m_1 + m_2) l_1^2 \dot{\phi}_1^2 + \frac{1}{2} m_2 l_2^2 \dot{\phi}_2^2 + m_2 l_1 l_2 \cos(\phi_1 - \phi_2) \dot{\phi}_1 \dot{\phi}_2$$


$$U = -(m_1 + m_2) g l_1 \cos \phi_1 - m_2 g l_2 \cos \phi_2$$

$$L = T - U$$

$$= \frac{1}{2} (m_1 + m_2) l_1^2 \dot{\phi}_1^2 + \frac{1}{2} m_2 l_2^2 \dot{\phi}_2^2 + m_2 l_1 l_2 \cos(\phi_1 - \phi_2) \dot{\phi}_1 \dot{\phi}_2$$

$$+ (m_1 + m_2) g l_1 \cos \phi_1 + m_2 g l_2 \cos \phi_2$$

Sec 5 Prob. 2



Generalised coords: x, ϕ

$$(x_1, y_1) = (x, 0)$$

$$(x_2, y_2) = (x + l \sin \phi, l \cos \phi)$$

$$U = -m_1 g y_1 - m_2 g y_2$$

$$= -m_2 g l \cos \phi$$

$$T = \frac{1}{2} m_1 (\dot{x}_1^2 + \dot{y}_1^2) + \frac{1}{2} m_2 (\dot{x}_2^2 + \dot{y}_2^2)$$

$$\text{Now: } \dot{x}_1^2 + \dot{y}_1^2 = \dot{x}^2$$

$$\begin{aligned}\dot{x}_2^2 + \dot{y}_2^2 &= (\dot{x} + l \cos \phi \dot{\phi})^2 + (-l \sin \phi \dot{\phi})^2 \\ &= \dot{x}^2 + l^2 \cos^2 \phi \dot{\phi}^2 + 2 l \cos \phi \dot{x} \dot{\phi}\end{aligned}$$

$$= \dot{x}^2 + l^2 \dot{\phi}^2 + 2 l \cos \phi \dot{x} \dot{\phi}$$


$$\rightarrow T = \frac{1}{2} m_1 \dot{x}^2 + \frac{1}{2} m_2 (\dot{x}^2 + l^2 \dot{\phi}^2 + 2 l \cos \phi \dot{x} \dot{\phi})$$

$$= \frac{1}{2} (m_1 + m_2) \dot{x}^2 + \frac{1}{2} m_2 l^2 \dot{\phi}^2 + m_2 l \cos \phi \dot{x} \dot{\phi}$$

$$\begin{aligned}L &= \frac{1}{2} (m_1 + m_2) \dot{x}^2 + \frac{1}{2} m_2 l^2 \dot{\phi}^2 + m_2 l \cos \phi \dot{x} \dot{\phi} \\ &\quad + m_2 g l \cos \phi\end{aligned}$$

Sec 5, Prob 3

(a)



point of support O moves along circle.

$$x_0 = a \sin \alpha \quad , \quad y_0 = a \cos \alpha$$

where $\alpha = \omega t$, $\omega = \text{const}$

Pendulum bob:

$$\begin{aligned}(x, y) : \quad x &= x_0 + l \sin \phi \\ y &= y_0 + l \cos \phi\end{aligned}$$

$$U = -m g y = -m g y_0 - m g l \cos \phi$$

Specified function of time.

[can ignore in L]

$$T = \frac{1}{2} m (\dot{x}^2 + \dot{y}^2)$$

$$\dot{x} = \dot{x}_0 + l \cos \phi \dot{\phi}$$

$$\dot{x}^2 = \dot{x}_0^2 + l^2 \cos^2 \phi \dot{\phi}^2 + 2 l \cos \phi \dot{x}_0 \dot{\phi}$$

$$\dot{y} = \dot{x}_0 - l \sin \phi$$

$$\dot{y}^2 = \dot{x}_0^2 + l^2 \sin^2 \phi \dot{\phi}^2 - 2l \sin \phi \dot{x}_0 \dot{\phi}$$

thus,

$$T = \frac{1}{2} m(\dot{x}_0^2 + \dot{y}^2)$$

$$= \frac{1}{2} m (\dot{x}_0^2 + l^2 \cos^2 \phi \dot{\phi}^2 + 2l \cos \phi \dot{x}_0 \dot{\phi})$$

$$+ \dot{y}^2 + l^2 \sin^2 \phi \dot{\phi}^2 - 2l \sin \phi \dot{x}_0 \dot{\phi})$$

$$= \frac{1}{2} m(\dot{x}_0^2 + \dot{y}^2) + \frac{1}{2} m l^2 \dot{\phi}^2 + m l \dot{\phi} (\dot{x}_0 \cos \phi - \dot{y}_0 \sin \phi)$$

$$\text{NOTE: } \dot{x}_0^2 + \dot{y}^2 = a^2 \dot{x}^2 = a^2 \gamma^2$$

since this is a specified function of time, we can ignore it in the Lagrangian;

$$\text{thus, } L = \frac{1}{2} m l^2 \dot{\phi}^2 + m l \dot{\phi} (\dot{x}_0 \cos \phi - \dot{y}_0 \sin \phi) + m g l \cos \phi$$

We can rewrite the second term:

$$x_0 = a \sin \alpha \rightarrow \dot{x}_0 = a \cos \alpha \dot{\alpha} \quad (\alpha = \gamma)$$

$$y_0 = a \cos \alpha \rightarrow \dot{y}_0 = -a \sin \alpha \dot{\alpha}$$

thus,

$$m l \dot{\phi} (\dot{x}_0 \cos \phi - \dot{y}_0 \sin \phi) = m l \dot{\phi} a \gamma (\cos \alpha \cos \phi + \sin \alpha \sin \phi) \\ = m l \dot{\phi} a \gamma \cos(\phi - \alpha) \\ = m l \dot{\phi} a \gamma \cos(\phi - \gamma t)$$

$$\text{Now: } \frac{d}{dt} [m l a \gamma \sin(\phi - \gamma t)]$$

$$= m l a \gamma \cos(\phi - \gamma t) (\dot{\phi} - \gamma)$$


$$= m l a \dot{\phi} \gamma \cos(\phi - \gamma t) - m l a \gamma^2 \cos(\phi - \gamma t)$$

thus,

$$m l a \dot{\phi} \gamma \cos(\phi - \gamma t) = \frac{d}{dt} [m l a \gamma \sin(\phi - \gamma t)] + m l a \gamma^2 \cos(\phi - \gamma t)$$

(and we can ignore the total time derivative in the Lagrangian)

$$\rightarrow L = \frac{1}{2} m l^2 \dot{\phi}^2 + m g l \cos \phi + m l a \gamma^2 \cos(\phi - \gamma t)$$



point O moving according to $x_0 = a \cos \gamma t$

$$x = x_0 + l \sin \phi$$

$$y = l \cos \phi$$

$$v = -m g y = -m g l \cos \phi$$

$$T = \frac{1}{2} m (\dot{x}^2 + \dot{y}^2)$$

$$\dot{x} = \dot{x}_0 + l \cos \phi \dot{\phi}, \quad x_0 = a \cos \gamma t$$

$$= -a \sin(\gamma t) \dot{\gamma} + l \cos \phi \dot{\phi}$$

$$\rightarrow \dot{x}^2 = a^2 \dot{\gamma}^2 \sin^2 \gamma t + l^2 \cos^2 \phi \dot{\phi}^2$$

$$- 2al\dot{\gamma} \dot{\phi} \sin(\gamma t) \cos \phi$$

$$\dot{y} = -l \sin \phi \dot{\phi}$$

$$\rightarrow \dot{y}^2 = l^2 \sin^2 \phi \dot{\phi}^2$$

Thus, $T = \frac{1}{2} m (a^2 \dot{\gamma}^2 \sin^2 \gamma t + l^2 \cos^2 \phi \dot{\phi}^2)$

$$- 2al\dot{\gamma} \dot{\phi} \sin(\gamma t) \cos \phi + l^2 \sin^2 \phi \dot{\phi}^2)$$

$$= \frac{1}{2} m l^2 \dot{\phi}^2 + \frac{1}{2} m a^2 \dot{\gamma}^2 \sin^2 \gamma t$$

specified
function
of
time
(ignore)

$$L = \frac{1}{2} m l^2 \dot{\phi}^2 - mal\dot{\gamma} \dot{\phi} \sin(\gamma t) \cos \phi + mgl \cos \phi$$

2nd term:

$$- \frac{d}{dt} [mal\dot{\gamma} \sin(\gamma t) \sin \phi]$$

$$= -mal\dot{\gamma}^2 \cos(\gamma t) \sin \phi - mal\dot{\gamma} \dot{\phi} \sin(\gamma t) \cos \phi$$


$$- mal\dot{\gamma} \dot{\phi} \sin(\gamma t) \cos \phi = - \frac{d}{dt} [] + mal\dot{\gamma}^2 \cos(\gamma t) \sin \phi$$

is no se

Thus, ignoring total time derivatives,

$$L = \frac{1}{2} m l^2 \dot{\phi}^2 + mgl \cos \phi + mal\dot{\gamma}^2 \cos(\gamma t) \sin \phi$$

(c)



$$y_0 = a \cos \gamma t$$

(point of support)

$$x = l \sin \phi$$

$$y = y_0 + l \cos \phi$$

$$= a \cos \gamma t + l \cos \phi$$

$$U = -mgy$$

$$= -mg(a \cos \gamma t + l \cos \phi)$$

specified function of time [can ignore]

$$T = \frac{1}{2} m (\dot{x}^2 + \dot{y}^2)$$

$$\dot{x} = l \cos \phi \dot{\phi}$$

$$\dot{x}^2 = l^2 \cos^2 \phi \dot{\phi}^2$$

$$\dot{y} = -a \dot{\gamma} \sin(\gamma t) - l \sin \phi \dot{\phi}$$

$$\dot{y}^2 = a^2 \dot{\gamma}^2 \sin^2(\gamma t) + l^2 \sin^2 \phi \dot{\phi}^2 + 2al\dot{\gamma} \sin(\gamma t) \sin \phi \dot{\phi}$$

specified function of time [can ignore]

thus, ignoring this function of time

$$T = \frac{1}{2} m l^2 \dot{\phi}^2 + m g l \dot{\phi} \sin(\gamma t) \sin \phi$$

$$\rightarrow L = T - U$$

$$= \frac{1}{2} m l^2 \dot{\phi}^2 + m a l \dot{\phi} \sin(\gamma t) \sin \phi + m g l \cos \phi$$

Rewrite 2nd term:

$$-\frac{d}{dt} [m a l \dot{\phi} \sin(\gamma t) \cos \phi] = -m a l \gamma^2 \cos(\gamma t) \cos \phi$$

$$+ m a l \dot{\gamma} \sin(\gamma t) \sin \phi \dot{\phi}$$

$$\text{so } m a l \dot{\phi} \sin(\gamma t) \sin \phi = -\frac{d}{dt} [] + m a l \gamma^2 \cos(\gamma t) \cos \phi$$


thus, ignoring total time derivative

$$L = \frac{1}{2} m l^2 \dot{\phi}^2 + m g l \cos \phi + m a l \gamma^2 \cos(\gamma t) \cos \phi$$

Sec 7, Prob 1

#1

$$U_1 = \text{const}$$



#2

$$U_2 = \text{const}$$

- Energy conserved, since no time dependence.
- Also momentum in x-direction (\parallel to interface) is conserved, since no x-dependence of the potential

$$U(x, y) = \begin{cases} U_1 & y \geq 0 \\ U_2 & y < 0 \end{cases}$$

V_1 : given

$$E = \frac{1}{2} m V_1^2 + U_1 = \frac{1}{2} m V_2^2 + U_2$$

$$\rightarrow \frac{1}{2} m V_2^2 = \frac{1}{2} m V_1^2 + (U_1 - U_2)$$

$$V_2^2 = V_1^2 + \frac{2(U_1 - U_2)}{m}$$

$$\text{so, } V_2 = V_1 \sqrt{1 + \frac{(U_1 - U_2)}{\frac{1}{2} m V_1^2}}$$

The angles θ_1, θ_2 are related by

$$p_{1x} = p_{2x}$$

$$\mu v_1 \sin \theta_1 = \mu v_2 \sin \theta_2$$

$$\text{thus, } \frac{\sin \theta_1}{\sin \theta_2} = \frac{v_2}{v_1} = \sqrt{1 + \frac{(v_1 - v_2)^2}{2m v_1^2}}$$

Sec 8; Prob 1

Transformations of action $S = \int L dt$

K, K' : two inertial frames

K' moves with velocity \vec{V} wrt K

Assume that K, K' coincide at $t=0$ so
 $\vec{r}_a = \vec{r}'_a$ wrt these two frames

$$\text{Now: } \vec{v}_a = \vec{V} + \vec{v}'_a$$

$$L = T - U$$

$$= \sum_a \frac{1}{2} m_a |\vec{v}_a|^2 - U(\vec{r}_1, \vec{r}_2, \dots, t)$$

$$|\vec{v}_a|^2 = |\vec{V} + \vec{v}'_a|^2$$

$$= |\vec{V}|^2 + |\vec{v}'_a|^2 + 2 \vec{V} \cdot \vec{v}'_a$$

so

$$L = \sum_a \frac{1}{2} m_a (|\vec{V}|^2 + |\vec{v}'_a|^2 + 2 \vec{V} \cdot \vec{v}'_a) - U$$

$$= \frac{1}{2} \mu V^2 + T' + \vec{V} \cdot \sum_a \vec{v}'_a - U$$

$$= T' - U + \frac{1}{2} \mu V^2 + \vec{P}' \cdot \vec{V}$$

$$= L' + \frac{1}{2} \mu V^2 + \vec{P}' \cdot \vec{V}$$

where \vec{P}' = total momentum wrt K'

$$\mu = \sum_a m_a \leftarrow \text{total mass}$$

$$\begin{aligned}
 S &= \int_{t_1}^{t_2} \int \rho dt \\
 &= \int_{t_1}^{t_2} (\bar{L}' + \frac{1}{2}\mu V^2 + \bar{\rho}' \cdot \vec{V}) dt \\
 &= S' + \underbrace{\frac{1}{2}\mu V^2(t_2 - t_1)}_{\text{doesnt change}} + \vec{V} \cdot \int_{t_1}^{t_2} \bar{\rho}' dt
 \end{aligned}$$

EOMs

$$\begin{aligned}
 \vec{V} \cdot \int_{t_1}^{t_2} \bar{\rho}' dt &= \vec{V} \cdot \int_{t_1}^{t_2} \sum_m \vec{v}_a' dt \\
 &= \vec{V} \cdot \sum_a \int_{t_1}^{t_2} \left(\frac{d\vec{r}_a}{dt} \right) dt \\
 &= \vec{V} \cdot \sum_a \vec{r}_a \Big|_{t_1}^{t_2} \\
 &= \vec{V} \cdot \left(\mu \vec{R}(t_2) - \mu \vec{R}(t_1) \right) \\
 &= \mu \vec{V} \cdot \left(\vec{R}(t_2) - \vec{R}(t_1) \right)
 \end{aligned}$$


Difference in positions

$$\text{So: } S = S' + \frac{1}{2}\mu V^2(t_2 - t_1) + \mu \vec{V} \cdot (\vec{R}(t_2) - \vec{R}(t_1))$$

Sec 9, Prob 1.

cylindrical coordinates, (s, ϕ, z)

$$s^2 = x^2 + y^2$$



$$x = s \cos \phi$$

$$y = s \sin \phi$$

$$z = z$$

$$\vec{M} = \vec{r} \times \vec{p} = m \vec{r} \times \vec{r}$$

$$\text{Now, } M_x = m(y \dot{z} - z \dot{y})$$

$$M_y = m(z \dot{x} - x \dot{z})$$

$$M_z = m(x \dot{y} - y \dot{x})$$

$$M = \sqrt{M_x^2 + M_y^2 + M_z^2}$$

$$\dot{z} = \dot{z}$$

$$\dot{y} = s \sin \phi + s \cos \phi \dot{\phi}$$

$$\dot{x} = s \cos \phi - s \sin \phi \dot{\phi}$$

$$\text{Thus, } M_x = m(s \sin \phi \dot{z} - z(s \sin \phi + s \cos \phi \dot{\phi}))$$

$$= m(s \sin \phi \dot{z} - z \sin \phi s - z \cos \phi \dot{\phi})$$

$$M_y = m(z(s \cos \phi - s \sin \phi \dot{\phi}) - s \cos \phi \dot{z})$$

$$= m(z \cos \phi s - z \sin \phi \dot{\phi} - s \cos \phi \dot{z})$$

$$M_z = m(s \cos \phi (s \sin \phi + s \cos \phi \dot{\phi})$$

$$- s \sin \phi (s \cos \phi - s \sin \phi \dot{\phi})]$$

$$= m s^2 \dot{\phi}$$

$$\bar{M}^2 = M_x^2 + M_y^2 + M_z^2$$

$$= m^2 \left\{ (\sin \phi (sz - z's) - z s \cos \phi \dot{\phi})^2 + (\cos \phi (sz - z's) - z s \sin \phi \dot{\phi})^2 + (s^2 \dot{\phi})^2 \right\}$$

$$= m^2 \left\{ \sin^2 \phi (sz - z's)^2 + z^2 s^2 \cos^2 \phi \dot{\phi}^2 - 2 z s \sin \phi \cos \phi (sz - z's) + \cos^2 \phi (sz - z's)^2 + z^2 s^2 \sin^2 \phi \dot{\phi}^2 + 2 z s \sin \phi \cos \phi (sz - z's) + s^4 \dot{\phi}^2 \right\}$$

$$= m^2 \left\{ (sz - z's)^2 + z^2 s^2 \dot{\phi}^2 + s^4 \dot{\phi}^2 \right\}$$

$$= m^2 [(sz - z's)^2 + s^2 \dot{\phi}^2 (z^2 + s^2)]$$

Sec 9, Prob 2

repeat for spherical polar coords.

$$M_x = m(yz - zy)$$

$$M^2 = M_x^2 + M_y^2 + M_z^2$$

$$\text{Now: } x = r \sin \theta \cos \phi$$

$$y = r \sin \theta \sin \phi$$

$$z = r \cos \theta$$

$$\rightarrow \dot{x} = r \sin \theta \cos \phi + r \cos \theta \cos \phi \dot{\theta} - r \sin \theta \sin \phi \dot{\phi}$$

$$\dot{y} = r \sin \theta \sin \phi + r \cos \theta \sin \phi \dot{\theta} + r \sin \theta \cos \phi \dot{\phi}$$

$$\dot{z} = r \cos \theta - r \sin \theta \dot{\phi}$$

Then,

$$M_x = m(yz - zy)$$

$$= m \left\{ r \sin \theta \sin \phi (r \cos \theta - r \sin \theta \dot{\phi}) - r \cos \theta (r \sin \theta \sin \phi + r \cos \theta \sin \phi \dot{\theta} + r \sin \theta \cos \phi \dot{\phi}) \right\}$$

$$= m \left\{ -r^2 \sin^2 \theta \sin \phi \dot{\theta} - r^2 \cos^2 \theta \sin \phi \dot{\theta} - r^2 \sin \theta \cos \theta \cos \phi \dot{\phi} \right\}$$

$$= m \left\{ -r^2 \sin \phi \dot{\theta} - r^2 \sin \theta \cos \theta \cos \phi \dot{\phi} \right\}$$

$$= -mr^2 [\sin \phi \dot{\theta} + \sin \theta \cos \theta \cos \phi \dot{\phi}]$$

$$\begin{aligned}
 M_y &= m(z\dot{x} - x\dot{z}) \\
 &= m \{ r\cos\theta (r\sin\theta \cos\phi \dot{\theta} + r\cos\theta \cos\phi \dot{\phi} - r\sin\theta \sin\phi \dot{\phi}) \\
 &\quad - r\sin\theta \cos\phi (r\cos\theta \dot{\theta} - r\sin\theta \dot{\phi}) \} \\
 &= m \{ r^2 \cos^2\theta \cos\phi \dot{\theta} - r^2 \sin\theta \cos\theta \sin\phi \dot{\phi} \\
 &\quad + r^2 \sin^2\theta \cos\phi \dot{\theta} \} \\
 &= m \{ r^2 \cos\phi \dot{\theta} - r^2 \sin\theta \cos\theta \sin\phi \dot{\phi} \} \\
 &= mr^2 [\cos\phi \dot{\theta} - \sin\theta \cos\theta \sin\phi \dot{\phi}]
 \end{aligned}$$

$$\begin{aligned}
 M_z &= m(xy' - yx') \\
 &= m \{ r\sin\theta \cos\phi (r\sin\theta \sin\phi + r\cos\theta \cos\phi \dot{\theta} \\
 &\quad + r\sin\theta \cos\phi \dot{\phi}) \\
 &\quad - r\sin\theta \sin\phi (r\sin\theta \cos\phi + r\cos\theta \cos\phi \dot{\theta} \\
 &\quad - r\sin\theta \sin\phi \dot{\phi}) \} \\
 &= m [r^2 \sin^2\theta \cos^2\phi \dot{\theta} + r^2 \sin^2\theta \sin^2\phi \dot{\phi}] \\
 &= mr^2 \sin^2\theta \dot{\phi}
 \end{aligned}$$


$$\begin{aligned}
 M^2 &= M_x^2 + M_y^2 + M_z^2 \\
 &= m^2 r^4 [\sin^2\phi \dot{\theta}^2 + \sin^2\theta \cos^2\theta \cos^2\phi \dot{\phi}^2] \\
 &\quad + m^2 r^4 [\cos^2\phi \dot{\theta}^2 - \sin^2\theta \cos^2\theta \sin^2\phi \dot{\phi}^2] \\
 &\quad + m^2 r^4 \sin^4\theta \dot{\phi}^2
 \end{aligned}$$

(cross terms will cancel)

$$\begin{aligned}
 M^2 &= m^2 r^4 \{ \sin^2\phi \dot{\theta}^2 + \sin^2\theta \cos^2\theta \cos^2\phi \dot{\phi}^2 \\
 &\quad + \cos^2\phi \dot{\theta}^2 + \sin^2\theta \cos^2\theta \sin^2\phi \dot{\phi}^2 \\
 &\quad + \sin^4\theta \dot{\phi}^2 \} \\
 &= m^2 r^4 [\dot{\theta}^2 + \sin^2\theta \cos^2\theta \dot{\phi}^2 + \sin^4\theta \dot{\phi}^2] \\
 &= m^2 r^4 [\dot{\theta}^2 + \sin^2\theta \phi^2 / (\cos^2\theta + \sin^2\theta)] \\
 &= m^2 r^4 [\dot{\theta}^2 + \sin^2\theta \dot{\phi}^2]
 \end{aligned}$$

Sec 9, Prob 3


a) Infinite homogeneous plane



P_x, P_y conserved

M_z conserved where origin is anywhere in (x, y) plane


b) Infinite homogeneous cylinder



P_z conserved



M_z conserved, origin anywhere on Z -axis

c) Infinite homog prism




P_z conserved

d) two points : M_z conserved, origin at midpoint of line connecting the two points




P_y conserved




homogeneous cone

M_z conserved, origin anywhere on Z -axis.



homogeneous circular torus

M_z conserved with \hat{z} directed as shown and origin at O.



Field constant along helix:

$\Delta Z = b$ when $\Delta\phi = 2\pi$
at $s^2 = x^2 + y^2 = a^2$

$$\frac{\Delta\phi}{2\pi} = \frac{\Delta Z}{b}$$

$$\rightarrow \Delta Z = \left(\frac{b}{2\pi}\right) \Delta\phi$$

$$\begin{aligned} \vec{E} &= a\Delta\phi \hat{x} + \Delta Z \hat{z} \\ &= \cancel{\Delta\phi} \left(a\hat{x} - \hat{y} \right) + \cancel{\Delta Z} \hat{z} \end{aligned}$$

$$\vec{E} = \Delta\phi (x\hat{y} - y\hat{x}) + \left(\frac{b}{2\pi}\right) \Delta\phi \hat{z}$$

$$= \Delta\phi \left[x\hat{y} - y\hat{x} + \left(\frac{b}{2\pi}\right) \hat{z} \right]$$

Field unchanged if you move along \vec{t}
thus, $\vec{P} \cdot \vec{t} = \text{const}$

$$\vec{P} \cdot \vec{t} = \Delta\phi [xP_y - yP_x + \left(\frac{b}{2\pi}\right) P_z]$$

$$= \Delta\phi \left[M_z + \frac{b}{2\pi} P_z \right]$$

$$\text{so } M_z + \frac{b}{2\pi} P_z = \text{const}$$

where $z = \text{axis of helix}$

$$b = 4z \text{ for } \Delta\phi = 2\pi \text{ at } s = a$$

Sec 10, prob 1

Different masses, same path, same potential energy

$$L_1 = \frac{1}{2} m_1 v_1^2 - U$$

$$L_2 = \frac{1}{2} m_2 v_2^2 - U$$

$$\text{Thus, } m_1 v_1^2 = m_2 v_2^2$$

$$\frac{m_1}{t_1^2} = \frac{m_2}{t_2^2}$$

$$\rightarrow \left(\frac{t_2}{t_1} \right)^2 = \frac{m_2}{m_1}$$

$$\boxed{\frac{t_2}{t_1} = \sqrt{\frac{m_2}{m_1}}}$$

Sec 10, Prob 2:

Same path, mass but potential energies differing by a constant

$$L_1 = \frac{1}{2}mV_1^2 - U_1$$

$$L_2 = \frac{1}{2}mV_2^2 - U_2$$

$$\frac{T_{b1}}{T_{b2}} = \frac{V_1^2}{V_2^2} = \frac{U_1}{U_2}$$

$$\rightarrow \frac{(1/t_1)^2}{(t/t_2)^2} = \frac{U_1}{U_2}$$

$$\text{so } \frac{t_2}{t_1} = \sqrt{\frac{U_1}{U_2}}$$

Sec 40 - Prob 1

single particle in a constant external field

$$L = \frac{1}{2}m\vec{v}^2 - U(\vec{r})$$

a) Cartesian coords (x, y, z)

$$L = \frac{1}{2}m(\dot{x}^2 + \dot{y}^2 + \dot{z}^2) - U(x, y, z)$$

$$\rightarrow p_x = \frac{\partial L}{\partial \dot{x}} = m\dot{x} \rightarrow \dot{x} = p_x/m$$

$$p_y = \frac{\partial L}{\partial \dot{y}} = m\dot{y} \rightarrow \dot{y} = p_y/m$$

$$p_z = \frac{\partial L}{\partial \dot{z}} = m\dot{z} \rightarrow \dot{z} = p_z/m$$

$$H = \left(\frac{1}{2m} \vec{p}^2 - L \right) / \left. \vec{e} = \vec{e}(e_ip) \right|$$

$$= \left(p_x \dot{x} + p_y \dot{y} + p_z \dot{z} - \frac{1}{2}m(\dot{x}^2 + \dot{y}^2 + \dot{z}^2) - U(x, y, z) \right) / \left. \vec{e} = \vec{e}(e_ip) \right|$$

$$= p_x \left(\frac{p_x}{m} \right) + p_y \left(\frac{p_y}{m} \right) + p_z \left(\frac{p_z}{m} \right) - \frac{1}{2}m \left(\left(\frac{p_x}{m} \right)^2 + \left(\frac{p_y}{m} \right)^2 + \left(\frac{p_z}{m} \right)^2 \right) + U(x, y, z)$$

$$= \frac{1}{2m} (\vec{p}^2) + U(x, y, z)$$

b) cylindrical coords (s, ϕ, z) , $s^2 = x^2 + y^2$

$$L = \frac{1}{2}m(s^2\dot{s}^2 + \dot{s}^2\phi^2 + \dot{z}^2) - U(s, \phi, z)$$

$$\rightarrow p_s = \frac{\partial L}{\partial \dot{s}} = m\dot{s} \rightarrow \dot{s} = p_s/m$$

$$p_\phi = \frac{\partial L}{\partial \dot{\phi}} = m s^2 \dot{\phi} \rightarrow \dot{\phi} = p_\phi / m s^2$$

$$p_z = \frac{\partial L}{\partial \dot{z}} = m\dot{z} \rightarrow \dot{z} = p_z/m$$

$$\begin{aligned}
 H &= \left(p_r \dot{r} + p_\theta \dot{\theta} + p_z \dot{z} - \frac{1}{2} m (\dot{r}^2 + r^2 \dot{\theta}^2 + \dot{z}^2) + U(r, \theta, z) \right) \\
 &= p_r \left(\frac{p_r}{m} \right) + p_\theta \left(\frac{p_\theta}{mr^2} \right) + p_z \left(\frac{p_z}{m} \right) \\
 &\quad - \frac{1}{2} m \left(\left(\frac{p_r}{m} \right)^2 + r^2 \left(\frac{p_\theta}{mr^2} \right)^2 + \left(\frac{p_z}{m} \right)^2 \right) + U(r, \theta, z) \\
 &= \frac{1}{2m} \left(p_r^2 + \frac{p_\theta^2}{r^2} + p_z^2 \right) + U(r, \theta, z)
 \end{aligned}$$

c) spherical polar coords. (r, θ, ϕ)

$$\begin{aligned}
 L &= \frac{1}{2} m (r^2 \dot{\theta}^2 + r^2 \sin^2 \theta \dot{\phi}^2 + \dot{r}^2) - U(r, \theta, \phi) \\
 \rightarrow p_r &= \frac{\partial L}{\partial \dot{r}} = m \dot{r} \rightarrow \dot{r} = p_r/m \\
 p_\theta &= \frac{\partial L}{\partial \dot{\theta}} = m r^2 \dot{\theta} \rightarrow \dot{\theta} = p_\theta / mr^2 \\
 p_\phi &= \frac{\partial L}{\partial \dot{\phi}} = m r^2 \sin^2 \theta \dot{\phi} \rightarrow \dot{\phi} = p_\phi / m r^2 \sin^2 \theta \\
 H &= \left(p_r \dot{r} + p_\theta \dot{\theta} + p_\phi \dot{\phi} - \frac{1}{2} m (r^2 \dot{\theta}^2 + r^2 \sin^2 \theta \dot{\phi}^2) + U(r, \theta, \phi) \right) \\
 &= p_r \left(\frac{p_r}{m} \right) + p_\theta \left(\frac{p_\theta}{mr^2} \right) + p_\phi \left(\frac{p_\phi}{mr^2 \sin^2 \theta} \right) \\
 &\quad - \frac{1}{2} m \left(\left(\frac{p_r}{m} \right)^2 + r^2 \left(\frac{p_\theta}{mr^2} \right)^2 + r^2 \sin^2 \theta \left(\frac{p_\phi}{mr^2 \sin^2 \theta} \right)^2 \right) + U(r, \theta, \phi) \\
 &= \frac{1}{2m} \left(p_r^2 + \frac{p_\theta^2}{r^2} + \frac{p_\phi^2}{r^2 \sin^2 \theta} \right) + U(r, \theta, \phi)
 \end{aligned}$$

sec 40 - prob 2

$$L = \frac{1}{2} m v^2 + m \vec{v} \cdot (\vec{\Omega} \times \vec{r}) + \frac{1}{2} m |\vec{\Omega} \times \vec{r}|^2 - m \vec{W} \cdot \vec{r} - U$$

restrict to uniformly rotating frame of reference $\vec{W} = 0$, $\vec{\Omega} = \vec{\omega}$

$$\begin{aligned}
 \rightarrow L &= \frac{1}{2} m v^2 + m \vec{v} \cdot (\vec{\Omega} \times \vec{r}) \\
 &\quad + \frac{1}{2} m |\vec{\Omega} \times \vec{r}|^2 - U(r)
 \end{aligned}$$

$$\text{Now: } H = \vec{p} \cdot \vec{v} - L$$

$$\begin{aligned}
 \vec{p} &= \frac{\partial L}{\partial \vec{v}} = m \vec{v} + m \vec{\Omega} \times \vec{r} = m (\vec{v} + \vec{\Omega} \times \vec{r}) \\
 \rightarrow \vec{v} &= \frac{\vec{p}}{m} - \vec{\Omega} \times \vec{r}
 \end{aligned}$$

thus,


$$\begin{aligned}
 H &= \vec{p} \cdot \left(\frac{\vec{p}}{m} - \vec{\Omega} \times \vec{r} \right) - \frac{1}{2} m \left| \frac{\vec{p}}{m} - \vec{\Omega} \times \vec{r} \right|^2 \\
 &\quad - m \left(\frac{\vec{p}}{m} - \vec{\Omega} \times \vec{r} \right) \cdot (\vec{\Omega} \times \vec{r}) \\
 &\quad - \frac{1}{2} m |\vec{\Omega} \times \vec{r}|^2 + U(r)
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{|\vec{p}|^2}{m} - \vec{p} \cdot (\vec{\Omega} \times \vec{r}) - \frac{1}{2} m \left(\frac{|\vec{p}|^2}{m^2} + |\vec{\Omega} \times \vec{r}|^2 \right) - \cancel{2 \vec{p} \cdot (\vec{\Omega} \times \vec{r})} \\
 &\quad - \vec{p} \cdot (\vec{\Omega} \times \vec{r}) + m |\vec{\Omega} \times \vec{r}|^2 - \frac{1}{2} m |\vec{\Omega} \times \vec{r}|^2 + U(r) \\
 &= \frac{|\vec{p}|^2}{2m} - \vec{p} \cdot (\vec{\Omega} \times \vec{r}) + U(r) \\
 &= \frac{|\vec{p}|^2}{2m} - \vec{\Omega} \cdot (\vec{r} \times \vec{p}) + U(r)
 \end{aligned}$$

\vec{p} is angular momentum

Sec. 11, Prob 1

$$\text{Simple pendulum: } U = -mg\gamma \\ = -mgl \cos\phi$$



$$\text{Also, } T = \frac{1}{2}ml^2\dot{\phi}^2$$

$$\text{Thus, } E = T+U \\ = \frac{1}{2}ml^2\dot{\phi}^2 - mgl \cos\phi$$

Release from rest at $\phi = \phi_0$

$$E = -mgl \cos\phi_0$$

$$\rightarrow \frac{1}{2}ml^2\dot{\phi}^2 - mgl \cos\phi = -mgl \cos\phi_0$$

$$T(E) = 4\sqrt{\frac{ml^2}{2}} \int_0^{\phi_0} \frac{d\phi}{\sqrt{E - U(\phi)}}$$

$$= 4\sqrt{\frac{ml^2}{2}} \int_0^{\phi_0} \frac{d\phi}{\sqrt{-mgl \cos\phi_0 + mgl \cos\phi}}$$

$$= 4\sqrt{\frac{l}{2g}} \int_0^{\phi_0} \frac{d\phi}{\sqrt{\cos\phi - \cos\phi_0}}$$

$\phi < \phi_0$

$$\begin{aligned} \text{Now: } \cos\phi &= \cos\left(2\frac{\phi}{2}\right) \\ &= \cos^2\left(\frac{\phi}{2}\right) - \sin^2\left(\frac{\phi}{2}\right) \\ &= 1 - 2\sin^2\left(\frac{\phi}{2}\right) \end{aligned}$$

$$\text{Also, } \cos\phi_0 = 1 - 2\sin^2\left(\frac{\phi_0}{2}\right)$$

$$T(E) = 4\sqrt{\frac{l}{2g}} \int_0^{\phi_0} \frac{d\phi}{\sqrt{2 \left(\sin^2\left(\frac{\phi_0}{2}\right) - \sin^2\left(\frac{\phi}{2}\right) \right)}}$$

$$= 2\sqrt{\frac{l}{g}} \int_0^{\phi_0} \frac{d\phi}{\sqrt{\sin^2\left(\frac{\phi_0}{2}\right) \left(1 - \frac{\sin^2\left(\frac{\phi}{2}\right)}{\sin^2\left(\frac{\phi_0}{2}\right)} \right)}}$$

$$\text{Let } x = \frac{\sin\left(\frac{\phi}{2}\right)}{\sin\left(\frac{\phi_0}{2}\right)} \rightarrow dx = \frac{1}{2} \frac{\cos\left(\frac{\phi}{2}\right)}{\sin^2\left(\frac{\phi_0}{2}\right)}$$

$$= \frac{d\phi}{2} \frac{\sqrt{1 - \sin^2\left(\frac{\phi}{2}\right)}}{\sin\left(\frac{\phi_0}{2}\right)}$$

$$= \frac{d\phi}{2} \frac{\sqrt{1 - \sin^2\left(\frac{\phi_0}{2}\right)x^2}}{\sin\left(\frac{\phi_0}{2}\right)}$$

$$\begin{aligned} \text{Thus, } T(E) &= 2\sqrt{\frac{l}{g}} \int_0^1 \frac{2dx \cdot \sin\left(\frac{\phi_0}{2}\right)}{\sqrt{1 - \sin^2\left(\frac{\phi_0}{2}\right)x^2} \cdot \sin\left(\frac{\phi_0}{2}\right) \sqrt{1 - x^2}} \\ &= 4\sqrt{\frac{l}{g}} \int_0^1 \frac{dx}{\sqrt{1 - x^2} \sqrt{1 - k^2x^2}}, \quad K = \sin\left(\frac{\phi_0}{2}\right) \\ &= 4\int_{\frac{1}{2}}^1 K(k) \end{aligned}$$

where $K(k) = \text{complete elliptic integral of the 1st kind.}$

A problem from:

$$T(E) = 4\sqrt{\frac{E}{J}} \int_0^1 \frac{dx}{\sqrt{1-x^2} \sqrt{1-H^2 x^2}}$$

For $H \ll 1$: $\frac{1}{\sqrt{1-H^2 x^2}} \approx 1 + \frac{1}{2} H^2 x^2$

$$\int_0^1 \frac{dx}{\sqrt{1-x^2} \sqrt{1-H^2 x^2}} \approx \int_0^1 \frac{dx}{\sqrt{1-x^2}} \left(1 + \frac{1}{2} H^2 x^2 \right)$$

Now: $\int_0^1 \frac{dx}{\sqrt{1-x^2}} = \arcsin(1) = \left[\frac{\pi}{2} \right]$

$$\frac{1}{2} H^2 \int_0^1 \frac{x^2 dx}{\sqrt{1-x^2}} = \frac{1}{2} H^2 \int_0^{\pi/2} \frac{\sin^2 \theta \cos \theta d\theta}{\sqrt{1-\sin^2 \theta}}$$

Let: $x = \sin \theta$ $\cos 2\theta = 1 - 2\sin^2 \theta$
 $dx = \cos \theta d\theta$ $\rightarrow \sin^2 \theta = \frac{1}{2}(1 - \cos 2\theta)$
 $x^2 = \sin^2 \theta$


$$\begin{aligned} \rightarrow \frac{1}{2} H^2 \int_0^1 \frac{x^2 dx}{\sqrt{1-x^2}} &= \frac{1}{2} H^2 \frac{1}{2} \int_0^{\pi/2} (1 - \cos 2\theta) d\theta \\ &= \frac{1}{4} H^2 \left[\frac{\pi}{2} - \frac{\sin 2\theta}{2} \Big|_0^{\pi/2} \right] \\ &= \boxed{\frac{1}{8} H^2 \pi} \quad |T = \sin \left(\frac{\phi_0}{2} \right) \approx \frac{\phi_0}{2}} \end{aligned}$$

$$Therefore, T(E) = \frac{4\sqrt{\frac{E}{J}}}{\sqrt{2}} \left(\frac{\pi}{2} + \frac{1}{8} H^2 \pi + \dots \right) = \boxed{2\sqrt{\frac{E}{J}} \left(1 + \frac{1}{16} \phi_0^2 + \dots \right)}$$

Sec II, Prob 2:

(a) $U = A|x|^n \quad x_0(E)$

$$T(E) = 2\sqrt{2m} \int_0^{\infty} \frac{dx}{\sqrt{E - Ax^n}}$$



where $U(x_0) = E$

$$Ax_0^n = E$$

$$\rightarrow x_0 = \left(\frac{E}{A} \right)^{\frac{1}{n}}$$

$$\sqrt{E - Ax^n} = \sqrt{E} \sqrt{1 - \left(\frac{A}{E} \right)x^n}$$

$$= \sqrt{E} \sqrt{1 - t^n} \quad \text{where } t = \left(\frac{A}{E} \right)^{\frac{1}{n}} x$$

$$dt = \left(\frac{A}{E} \right)^{\frac{1}{n}} dx$$

$$t_0 = \left(\frac{A}{E} \right)^{\frac{1}{n}} x_0 = \left(\frac{A}{E} \right)^{\frac{1}{n}} \left(\frac{E}{A} \right)^{\frac{1}{n}} = 1$$

Thus,

$$T(E) = 2\sqrt{2m} \int_0^1 \left(\frac{E}{A} \right)^{\frac{1}{n}} \frac{dt}{\sqrt{1-t^n}}$$

Now let $u = t^n$

$$\begin{aligned} du &= n t^{n-1} dt = n u^{\frac{n-1}{n}} dt \\ &= n u^{1-\frac{1}{n}} dt \end{aligned}$$

$$dt = \frac{1}{n} u^{\frac{1}{n}-1} du, \quad t=0, 1 \rightarrow u=0, 1$$

Thus

$$T(E) = 2 \sqrt{\frac{2m}{E}} \left(\frac{E}{A}\right)^{\frac{1}{n}} \int_0^1 \frac{1}{n} \frac{u^{\frac{1}{n}-1}}{\sqrt{1-u}} du$$
$$= \frac{2}{n} \sqrt{\frac{2m}{E}} \left(\frac{E}{A}\right)^{\frac{1}{n}} \int_0^1 u^{\frac{1}{n}-1} (1-u)^{-\frac{1}{2}} du$$

Recall: Beta Function

$$B(x, y) = \int_0^1 dt t^{x-1} (1-t)^{y-1} = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}$$




(where $\operatorname{Re}(x) > 0, \operatorname{Re}(y) > 0$)

Thus,

$$\boxed{T(E)} = \frac{2}{n} \sqrt{\frac{2m}{E}} \left(\frac{E}{A}\right)^{\frac{1}{n}} \frac{\Gamma\left(\frac{1}{n}\right) \Gamma\left(\frac{1}{2}\right)}{\Gamma\left(\frac{1}{n} + \frac{1}{2}\right)} \quad \Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$$
$$= \frac{2}{n} \sqrt{\frac{2\pi m}{E}} \left(\frac{E}{A}\right)^{\frac{1}{n}} \frac{\Gamma\left(\frac{1}{n}\right)}{\Gamma\left(\frac{1}{n} + \frac{1}{2}\right)}$$

Gamma Function

$$(b) U = -U_0 / \cosh^2 \alpha x$$



$$-U_0 \leq E < 0$$

Thus, $E = -|E|$

$$|E| < U_0$$

$$T(E) = 2 \sqrt{2m} \int_0^{\infty} \frac{dx}{\sqrt{E - U(x)}}$$

where x_0 given by

$$E = U(x_0)$$
$$= \frac{-U_0}{\cosh^2(\alpha x_0)}$$
$$\rightarrow \cosh^2(\alpha x_0) = -\frac{U_0}{E}$$

$$\cosh(\alpha x_0) = \sqrt{\frac{U_0}{|E|}}$$

$$\alpha x_0 = \cosh^{-1}\left(\sqrt{\frac{U_0}{|E|}}\right)$$

$$\boxed{x_0 = \frac{1}{\alpha} \cosh^{-1}\left(\sqrt{\frac{U_0}{|E|}}\right)}$$

$$\begin{aligned}
 \sqrt{E - U(x)} &= \sqrt{E + \frac{U_0}{\cosh^2(\alpha x)}} \quad \cosh^2 - \sinh^2 = 1 \\
 &= \frac{1}{\cosh(\alpha x)} \sqrt{E \cosh^2(\alpha x) + U_0} \\
 &= \frac{1}{\cosh(\alpha x)} \sqrt{E(1 + \sinh^2(\alpha x)) + U_0} \\
 &= \frac{1}{\cosh(\alpha x)} \sqrt{(E+U_0) + E \sinh^2(\alpha x)} \\
 &= \frac{1}{\cosh(\alpha x)} \sqrt{(U_0 - |E|) - |E| \sinh^2(\alpha x)}
 \end{aligned}$$

thus


$$\begin{aligned}
 T(E) &= 2\sqrt{2m} \int_0^{x_0} \frac{dx}{\sqrt{E - U(x)}} \\
 &= 2\sqrt{2m} \int_0^{x_0} \frac{\cosh(\alpha x) dx}{\sqrt{(U_0 - |E|) - |E| \sinh^2(\alpha x)}}
 \end{aligned}$$

$$\text{Let: } t = \sinh(\alpha x) \rightarrow t_0 = \sinh(\alpha x_0)$$

$$dt = \alpha \cosh(\alpha x) dx$$

$$so \quad t_0 = \sinh^{-1} \left(\frac{U_0}{|E|} \right)$$

$$= \frac{U_0 - |E|}{|E|}$$



$$U_0 - y^2 = |E| \\ y = \sqrt{U_0 - |E|}$$

$$T(E) \approx 2\sqrt{2m} \int_0^{\sqrt{\frac{U_0 - |E|}{|E|}}} dt$$

$$= 2\sqrt{2m} \int_0^{\sqrt{\frac{U_0 - |E|}{|E|}}} \frac{dt}{\sqrt{U_0 - |E|} \sqrt{1 - \frac{|E|}{U_0 - |E|} t^2}}$$

$$\text{Def: } \sin u = \sqrt{\frac{|E|}{U_0 - |E|}} t$$

$$du \cos u = \sqrt{\frac{|E|}{U_0 - |E|}} dt$$

$$T(E) = \frac{2\sqrt{2m}}{\sqrt{U_0 - |E|}} \int_0^{\pi/2} \frac{\cos u du}{\sqrt{\frac{U_0 - |E|}{|E|}} \sqrt{1 - \frac{|E|}{U_0 - |E|} \sin^2 u}}$$

$$= \frac{\pi}{\alpha} \sqrt{\frac{2m}{|E|}} \frac{\pi}{2}$$

$$= \boxed{\frac{\pi}{\alpha} \sqrt{\frac{2m}{|E|}}}$$

$$(c) U = U_0 + \frac{1}{2} m \alpha x^2$$

$$T = 2\sqrt{2m} \int_0^{x_0} \frac{dx}{\sqrt{E - U(x)}}$$

where x_0 determined by

$$E = U(x_0)$$

$$= U_0 + \frac{1}{2} m \alpha x_0^2$$

$$\Rightarrow x_0 = \tan^{-1} \left(\sqrt{\frac{E}{U_0}} \right)$$

$$\boxed{x_0 = \pm \tan^{-1} \left(\sqrt{\frac{E}{U_0}} \right)}$$

$$\sqrt{E - U(x)} = \sqrt{E - U_0 - \frac{1}{2} m \alpha x^2}$$



$$= \sqrt{E - U_0 - \frac{1}{2} m \sin^2(\alpha x)}$$

$$= \pm \sqrt{E \cos^2(\alpha x) - U_0 \sin^2(\alpha x)}$$

$$= \pm \sqrt{E (1 - \sin^2(\alpha x)) - U_0 \sin^2(\alpha x)}$$

$$= \pm \sqrt{E - (U_0 + E) \sin^2(\alpha x)}$$

$$= \frac{\sqrt{E}}{\cos \alpha x} \sqrt{1 - \frac{(U_0 + E)}{E} \sin^2(\alpha x)}$$



Thus

$$T(E) = 2\sqrt{2m} \int_0^{x_0} \frac{\cos \alpha x \, dx}{\sqrt{E - \frac{U_0 + E}{E} \sin^2 \alpha x}}$$

$$\text{Let } u = \sin \alpha x$$

$$du = \alpha \cos \alpha x \, dx$$

$$x=0, x_0 \rightarrow u=0$$

$$u_0 = \sin(\alpha x_0)$$

$$= \sin \left(\tan^{-1} \sqrt{\frac{E}{U_0}} \right)$$

$$= \sqrt{\frac{E}{E+U_0}}$$

$$u_0$$

$$\rightarrow T(E) = 2\sqrt{2m} \frac{1}{\sqrt{E}} \int_0^{u_0} \frac{du}{\sqrt{1 - \frac{U_0 + E}{E} u^2}}$$

$$\text{Let } \sqrt{\frac{U_0 + E}{E}} u = \sin \theta \rightarrow \sqrt{\frac{U_0 + E}{E}} du = \cos \theta \, d\theta$$

$$u=0 \rightarrow \theta=0$$

$$u=u_0 \rightarrow \sin \theta_0 = \sqrt{\frac{U_0 + E}{E}} \sqrt{\frac{E}{U_0 + E}} = 1 \Rightarrow \theta_0 = \frac{\pi}{2}$$

$$\rightarrow T(E) = 2\sqrt{2m} \frac{1}{\sqrt{E}} \int_0^{\pi/2} \frac{\cos \theta \, d\theta}{\sqrt{\frac{U_0 + E}{E}} \sqrt{1 - \sin^2 \theta}}$$

$$= \frac{2\sqrt{2m}}{\alpha \sqrt{E}} \sqrt{\frac{E}{U_0 + E}} \frac{\pi}{2} = \boxed{\frac{\pi \sqrt{2m}}{\alpha \sqrt{E}} \sqrt{\frac{E}{U_0 + E}}}$$

Sec 13, Prob 1

\vec{X} : position vector of M

\vec{x}_a : $a=1, 2, \dots, n$ position vector of n masses
all with mass m .

Closed system \rightarrow linear momentum conserved
 \rightarrow COM frame

$$M\vec{X} + m(\vec{x}_1 + \vec{x}_2 + \dots + \vec{x}_n) = 0$$

$$\text{or } M\vec{X} + m \sum_{a=1}^n \vec{x}_a = 0 \quad (1)$$

$$\text{Taking time derivative } \rightarrow M\dot{\vec{X}} + \sum_a \dot{\vec{x}}_a = 0. \quad (2)$$

Define relative position vectors:

$$\vec{r}_1 \equiv \vec{x}_1 - \vec{X}$$

$$\vec{r}_2 \equiv \vec{x}_2 - \vec{X}$$

etc.

$$\text{or } \vec{r}_a \equiv \vec{x}_a - \vec{X}, \quad a=1, 2, \dots, n. \quad (3)$$

Summing up (3):

$$\begin{aligned} \sum \vec{r}_a &= \sum (\vec{x}_a - \vec{X}) \\ &= \sum \vec{x}_a - n \vec{X} \quad \text{total mass} \\ &= -\frac{M}{m} \vec{X} - n \vec{X} \\ &= -\frac{(M+n)m}{m} \vec{X} = -\mu \vec{X} \end{aligned}$$

$$\text{Thus, } \vec{X} = -\frac{m}{\mu} \sum \vec{r}_a$$

$$\text{and } \vec{x}_a = \vec{r}_a + \vec{X}$$

give you \vec{x}_a, \vec{X} in terms of \vec{r}_a :

$$\begin{aligned} \text{Hence: } T &= \frac{1}{2} \sum_a m |\vec{x}_a|^2 + \frac{1}{2} M |\vec{X}|^2 \\ &= \frac{1}{2} m \sum_a |\vec{r}_a + \vec{X}|^2 + \frac{1}{2} M |\vec{X}|^2 \\ &= \frac{1}{2} m \sum_a |\vec{r}_a|^2 + \frac{1}{2} \sum_a m 2\vec{r}_a \cdot \vec{X} \\ &\quad + \frac{1}{2} \sum_a m |\vec{X}|^2 + \frac{1}{2} M |\vec{X}|^2 \\ &= \frac{1}{2} m \sum_a |\vec{r}_a|^2 + m \left(\sum_a \vec{r}_a \right) \cdot \vec{X} \\ &\quad + \frac{1}{2} nm |\vec{X}|^2 + \frac{1}{2} M |\vec{X}|^2 \end{aligned}$$

$$\begin{aligned} &= \frac{1}{2} m \sum_a |\vec{r}_a|^2 - \mu |\vec{X}|^2 + \frac{1}{2} (nm + M) |\vec{X}|^2 \\ &= \frac{1}{2} m \sum_a |\vec{r}_a|^2 - \frac{1}{2} \mu |\vec{X}|^2 \end{aligned}$$

Now rewrite in t form:

$$-\frac{1}{2} \mu |\vec{X}|^2 = -\frac{1}{2} \mu \frac{m^2}{\mu^2} \sum_a |\vec{r}_a|^2 = -\frac{1}{2} \frac{m^2}{\mu} \sum_a |\vec{r}_a|^2$$

$$\text{Thus, } T = \frac{1}{2} m \sum_a |\vec{r}_a|^2 - \frac{1}{2} \frac{m^2}{\mu} \sum_a |\vec{r}_a|^2$$

Potential energy

$$U = U(|\vec{x}_1 - \vec{x}_2|, |\vec{x}_1 - \vec{x}_3|, \dots, |\vec{x}_1 - \vec{x}_n|)$$

$$|\vec{x}_1 - \vec{x}|, |\vec{x}_2 - \vec{x}|, \dots, |\vec{x}_n - \vec{x}|)$$

$$= \cup \{ |\vec{r}_1 - \vec{r}_2|, |\vec{r}_1 - \vec{r}_3|, \dots, |\vec{r}_1 - \vec{r}_{n_1}| \}$$

$$(\vec{r}_1), (\vec{r}_2), \dots, (\vec{r}_n)$$

which depends only on the relative position vectors $\vec{r}_1, \vec{r}_2, \dots, \vec{r}_n$.

Thus,

$$L = \frac{1}{2} m \sum_a |\vec{r}_a|^2 - \frac{1}{2} \frac{m^2}{M} \left| \sum_a \vec{r}_a \right|^2 - U(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_n)$$


Sec 14, Prob. 1

Spherical pendulum

$$T = \frac{1}{2} m l^2 (\dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2)$$

$$U = -mgz$$

$$r = -mg \cdot l \cdot \cos \theta$$



Thus

$$f = T \cup$$

$$= \frac{1}{2} m l^2 (\dot{\theta}^2 + m^2 \dot{\theta} \dot{\phi}^2) + m g l \cos \theta$$

$$\text{No } t\text{-dependence} \rightarrow E = T + U = \text{const}$$

$$\text{No } \phi\text{-dependence} \rightarrow \frac{\partial L}{\partial \dot{\phi}} = m \ell^2 \sin^2 \theta \phi \equiv M_2 = \text{const}$$

$$E = \frac{1}{2} m l^2 (\dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2) - mg l \cos \theta$$

$$= \frac{1}{2} m l^2 (\dot{\theta}^2 + \sin \theta \frac{M_z^2}{m^2 l^4 \sin^2 \theta}) - m g l \cos \theta$$

$$= \frac{1}{2} m l^2 \dot{\theta}^2 + \frac{M_z^2}{2m/l^2 \sin^2 \theta} - mg l \cos \theta$$

$$U_{eff}(\theta)$$

$$E = \frac{1}{2} m \lambda^2 \theta^2 + \frac{M_2^2}{2 m \lambda^2 \sin^2 \theta} + U(\theta)$$

$$\rightarrow \frac{1}{2} m l^2 \dot{\theta}^2 = E - U(\theta) = \frac{M z^2}{2 m l^2 r_m^2 \theta}$$

$$\textcircled{2} = \sqrt{\frac{2}{m_e^2} (E - U(\theta)) - \frac{M_e^2}{m_e^2 \ell^4 s_i h^2 \theta}}$$

$$\frac{d\theta}{dt} = \dot{\theta} = \sqrt{\frac{2}{ml^2} (E + mgl\cos\theta) - \frac{M_z^2}{m^2 l^4 \sin^2\theta}}$$

$$\rightarrow dt = \frac{d\theta}{\sqrt{\dots}}$$

$$t = \int \frac{d\theta}{\sqrt{\frac{2}{ml^2} (E + mgl\cos\theta) - \frac{M_z^2}{m^2 l^4 \sin^2\theta}}} + C_1 t$$

Path: use $M_z = ml^2 \sin^2\theta \dot{\phi}$

$$\text{Thus, } \frac{d\theta}{dt} = \frac{d\theta}{d\phi} \frac{d\phi}{dt} \\ = \frac{d\theta}{d\phi} \frac{M_z}{ml^2 \sin^2\theta}$$

$$\text{Thus, } \frac{d\theta}{d\phi} = \frac{d\theta}{dt} \frac{ml^2 \sin^2\theta}{M_z} = \sqrt{\frac{ml^2 \sin^2\theta}{M_z}}$$

$$d\phi = \frac{d\theta M_z}{\sqrt{ml^2 \sin^2\theta}}$$

$$\phi = \int \frac{M_z d\theta / ml^2 \sin^2\theta}{\sqrt{\frac{2}{ml^2} (E + mgl\cos\theta) - \frac{M_z^2}{m^2 l^4 \sin^2\theta}}} + C_2 \phi$$

Turning points: (where $\dot{\theta} = 0$)

$$E = V_{\text{eff}}(r)$$

$$= \frac{M_z^2}{2ml^2 \sin^2\theta} - mgl\cos\theta$$

$$\rightarrow 2Eml^2 \sin^2\theta = M_z^2 - 2mgl^3 \sin^2\theta \cos\theta$$

$$2Eml^2 (1 - \cos^2\theta) = M_z^2 - 2mgl^3 (1 - \cos^2\theta) \cos\theta$$

$$2Eml^2 - 2Eml^2 \cos^2\theta$$

$$= M_z^2 - 2mgl^3 \cos\theta + 2mgl^3 \cos^3\theta$$

$$\text{Thus, } 2mgl^3 \cos^3\theta + 2Eml^2 \cos^2\theta - 2mgl^3 \cos\theta$$



$$+ (M_z^2 - 2Eml^2) = 0$$

Divide by $2mgl^3$:


$$\rightarrow \cos^3\theta + \frac{E}{mgl} \cos^2\theta - \cos\theta + \left(\frac{M_z^2}{2mgl^3} - \frac{E}{mgl} \right) = 0$$

Cubic equation for $\cos\theta$:

Effective potential looks like:



Sec 14, Prob. 2



spherical polar coords (r, θ, ϕ)

constraint $\theta = \alpha$

Generalized coords: (r, ϕ)

$$T = \frac{1}{2}m(r^2 + r^2\sin^2\alpha\dot{\phi}^2)$$

$$= \frac{1}{2}m(r^2 + r^2\sin^2\alpha\dot{\phi}^2)$$

$$U = mgz$$

$$= mg r \cos \alpha$$

$$L = T - U$$

$$= \frac{1}{2}m(r^2 + r^2\sin^2\alpha\dot{\phi}^2) - mg r \cos \alpha$$

$E = \text{const}$ (since no explicit t dependence)

$$\frac{dL}{d\phi} = mr^2\sin^2\alpha\dot{\phi} = M_Z = \text{const}$$

(since no explicit ϕ dependence)

$$E = T + U$$


$$= \frac{1}{2}m(r^2 + r^2\sin^2\alpha\dot{\phi}^2) + mg r \cos \alpha$$

$$= \frac{1}{2}mr^2 + \frac{1}{2}mr^2\sin^2\alpha\left(\frac{M_Z^2}{m^2r^4\sin^4\alpha}\right) + mg r \cos \alpha$$

$$= \frac{1}{2}mr^2 + \frac{M_Z^2}{2mr^2\sin^2\alpha} + mg r \cos \alpha$$


$$= \frac{1}{2}mr^2 + U_{\text{eff}}(r)$$

$$U_{\text{eff}}(r) = \frac{M_Z^2}{2mr^2\sin^2\alpha} + mg r \cos \alpha$$



Bound orbits for $E > U_{\text{eff}, \min} = U_{\text{eff}}(r_0)$

r_0 : stable circular orbit



t-equation:

$$t = \int \frac{dr}{\sqrt{\frac{2}{m}(E - mg r \cos \alpha) - \frac{M_Z^2}{m^2 r^2 \sin^2 \alpha}}} + \text{const}$$

Using $M_Z = mr^2 \sin^2 \alpha \dot{\phi}$

$$\rightarrow \frac{dr}{dt} = \frac{dr}{d\phi} \frac{d\phi}{dt} = \frac{dr}{d\phi} \frac{M_Z}{mr^2 \sin^2 \alpha}$$

ϕ -equation:

$$\phi = \int \frac{\left(\frac{M_Z}{\sin^2 \alpha}\right) dr}{\sqrt{\frac{2m(E - mg r \cos \alpha)}{r^2 \sin^2 \alpha} - \frac{M_Z^2}{r^2 \sin^2 \alpha}}} + \text{const}$$

Turning points: $r = r_{\min}, r_{\max}$
Determined by effective potential

$$E = U_{\text{eff}}(r)$$


$$= \frac{M_2^2}{2m r^2 \sin^2 \alpha} + m g r \cos \alpha$$

$$\rightarrow 2m E r^2 \sin^2 \alpha = M_2^2 + 2m^2 g r^3 \sin^2 \alpha \cos \alpha$$


$$0 = 2m^2 g r^3 \sin^2 \alpha \cos \alpha - 2m E r^2 \sin^2 \alpha + M_2^2$$

$$= r^3 - \left(\frac{E}{m g \cos \alpha} \right) r^2 + \frac{M_2^2}{2m^2 g \sin^2 \alpha \cos \alpha}$$

cubic equation again



Sec 14, Prob 3



From Sec 5, Prob 2 we have

$$L = \frac{1}{2}(m_1 + m_2)\dot{x}^2 + \frac{1}{2}m_2 l^2 \dot{\phi}^2 + 2l\dot{x}\dot{\phi} \cos \phi + m_2 g l \cos \phi$$

No dependence on x :

$$\rightarrow p_x = \frac{\partial L}{\partial \dot{x}} = (m_1 + m_2)\dot{x} + m_2 l \dot{\phi} \cos \phi = \text{const}$$

(x -component of total momentum)

No explicit t -dependence

$$\begin{aligned} \rightarrow E &= T + U = \text{const} \\ &= \frac{1}{2}(m_1 + m_2)\dot{x}^2 + \frac{1}{2}m_2(l^2 \dot{\phi}^2 + 2l\dot{x}\dot{\phi} \cos \phi) - m_2 g l \cos \phi \end{aligned}$$

Want in frame where $\text{com}_x = 0$:

$$\begin{aligned} \text{com}_x &= m_1 x + m_2(x + l \sin \phi) \\ &= (m_1 + m_2)x + m_2 l \sin \phi \end{aligned}$$

$$\text{com}_x = 0 \rightarrow x = -\left(\frac{m_2}{m_1 + m_2}\right)l \sin \phi$$

$$\dot{x} = -\left(\frac{m_2}{m_1+m_2}\right) l \cos\phi \dot{\phi}$$

t^{loss}

$$E = \frac{1}{2}(m_1+m_2)\dot{x}^2 + \frac{1}{2}m_2(l^2\dot{\phi}^2 + 2l\dot{x}\dot{\phi}\cos\phi) - m_2gl\cos\phi$$

$$= \frac{1}{2}(m_1+m_2) \frac{m_2^2}{(m_1+m_2)^2} l^2 \cos^2\phi \dot{\phi}^2$$

$$+ \frac{1}{2}m_2l^2\dot{\phi}^2 - m_2\left(\frac{m_2}{m_1+m_2}\right)l^2\cos^2\phi \dot{\phi}^2$$

$$- m_2gl\cos\phi$$

$$= \frac{1}{2}m_2l^2\dot{\phi}^2 \left[1 - \left(\frac{m_2}{m_1+m_2}\right)\cos^2\phi \right]$$

$$- m_2gl\cos\phi$$

1-d problem:

$$\underline{E + m_2gl\cos\phi} = \frac{1}{2}m_2l^2\dot{\phi}^2$$

$$\left[1 - \left(\frac{m_2}{m_1+m_2}\right)\cos^2\phi \right]$$

$$\frac{d\phi}{dt} = \dot{\phi} = \sqrt{\frac{2}{m_2l^2} \left(E + m_2gl\cos\phi \right)} \over \sqrt{1 - \left(\frac{m_2}{m_1+m_2}\right)\cos^2\phi}$$

$$\rightarrow dt = \frac{d\phi}{\sqrt{}}$$

$$= d\phi \sqrt{1 - \left(\frac{m_2}{m_1+m_2}\right)\cos^2\phi} \\ \frac{2}{m_2l^2} (E + m_2gl\cos\phi)$$

$$= d\phi \sqrt{\frac{m_2l^2}{2} \frac{1}{m_1+m_2}} \sqrt{\frac{(m_1+m_2) - m_2\cos^2\phi}{E + m_2gl\cos\phi}}$$

$$= d\phi \sqrt{\frac{m_2}{m_1+m_2}} \sqrt{\frac{l^2}{2}} \sqrt{\frac{m_1+m_2\sin^2\phi}{E + m_2gl\cos\phi}}$$

so

$$t = \sqrt{\left(\frac{m_2}{m_1+m_2}\right)\frac{l^2}{2}} \int d\phi \sqrt{\frac{m_1+m_2\sin^2\phi}{E + m_2gl\cos\phi}} + \text{const}$$

$$\text{Now: } x_2 = x + l\sin\phi$$

$$y_2 = l\cos\phi$$

$$\text{using } x = -\left(\frac{m_2}{m_1+m_2}\right)l\sin\phi$$

$$\rightarrow x_2 = \left[-\left(\frac{m_2}{m_1+m_2}\right)l\sin\phi + l\cos\phi \right] = \left(\frac{m_1}{m_1+m_2}\right)l\sin\phi$$


$$x_2 = \left(\frac{m_1}{m_1 + m_2} \right) l \sin \phi = b \sin \phi$$

$$y_2 = l \cos \phi = a \cos \phi$$

$$\left(\frac{y_2}{a}\right)^2 + \left(\frac{x_2}{b}\right)^2 = \cos^2 \phi + \sin^2 \phi = 1$$

which is an ellipse with semi-major and semi-minor axes:

$$a = l, \quad b = l \left(\frac{m_1}{m_1 + m_2} \right)$$



If $m_1 \gg m_2$
then $a = b \approx l$
so that m_2 moves
along a circular
arc of radius l :

Sec 15, Prob 1

$$U = -\frac{\infty}{r}, \quad E = 0 \rightarrow e = 1$$

$$\frac{p}{r} = 1 + \cos \phi$$

when $\phi = 0, \quad p = 2 \cdot r$

$$\text{so } r_{min} = \frac{p}{2}$$


NOTE: $p = r + r \cos \phi$
 $= \sqrt{r^2 + y^2} + x$

$$\rightarrow (p-x)^2 = x^2 + y^2$$

$$p^2 + x^2 - 2px = x^2 + y^2$$

$$p^2 - y^2 = 2px$$

$$\rightarrow \boxed{x = \frac{p^2 - y^2}{2p}} \quad \text{parabola}$$



when $y = 0, \quad x = \frac{1}{2}P$

$$x = 0, \quad y = \pm P$$

Time equation:

$$t = \int \frac{dr}{\sqrt{\frac{2}{m} [E - U(r)] - \frac{p^2}{m^2 r^2}}} \quad t \propto \sqrt{d}$$

$$U(r) = -\frac{\alpha}{r}, \quad p = \frac{mv^2}{m\alpha}, \quad e=1, \quad E=0$$

$$\rightarrow t = \int \frac{dr}{\sqrt{\frac{2\alpha}{mr} - \frac{mv^2}{m^2 r^2}}} + \text{const}$$

$$= \int \frac{dr}{\sqrt{\frac{2\alpha}{mr} - \frac{mv^2}{m^2 r^2}}} + \text{const}$$

$$= \int \frac{dr}{\sqrt{\frac{2}{r} - \frac{p^2}{r^2}}} + \text{const}$$

$$= \int \frac{r dr}{\sqrt{2r-p}} + \text{const}$$

$$= \int \frac{r dr}{\sqrt{\frac{2r-1}{p}}} + \text{const}$$

Defn: $\frac{2r-1}{p} = \xi^2 > 0 \quad (-\infty < \xi < \infty)$

$$2r = p(1+\xi^2)$$

$$\boxed{r = \frac{p}{2}(1+\xi^2)} \rightarrow dr = p\xi d\xi$$

$$r dr = \frac{p^2}{2}(\xi + \xi^3) d\xi$$

Thus,

$$t = \sqrt{\frac{m}{\alpha p}} \int \frac{\frac{p^2}{2}(\xi + \xi^3) d\xi}{\sqrt{\xi^2}} + \text{const}$$

$$= \sqrt{\frac{mp^3}{\alpha}} \frac{1}{2} \int (1+\xi^2) d\xi + \text{const}$$

$$= \frac{1}{2} \sqrt{\frac{mp^3}{\alpha}} \left(\xi + \frac{1}{3}\xi^3 \right) + \text{const}$$

choose const so that $t=0 \Leftrightarrow \xi = 0$ (const=0)

$$\text{so, } \boxed{t = \frac{1}{2} \sqrt{\frac{mp^3}{\alpha}} \left(\xi + \frac{1}{3}\xi^3 \right)}$$

Now:

$$\frac{p}{r} = 1 + \cos\phi$$

$$p = r(1 + \cos\phi)$$

$$\phi = \frac{p}{2}(1+\xi^2)(1+\cos\phi)$$

$$Z = 1 + \xi^2 + \cos\phi + \xi^2 \cos\phi$$

$$1 - \xi^2 = (1+\xi^2) \cos\phi$$

$$\rightarrow \boxed{\cos\phi = \frac{1-\xi^2}{1+\xi^2}}$$

$$\begin{aligned} X &= r \cos \phi \\ &= \frac{p}{2} (1 + \xi^2) \left(\frac{1 - \xi^2}{1 + \xi^2} \right) \\ &= \frac{p}{2} (1 - \xi^2) \end{aligned}$$

Also,

$$\begin{aligned} x^2 + y^2 &= r^2 \\ \rightarrow y^2 &= r^2 - x^2 \\ &= \frac{p^2}{4} (1 + \xi^2)^2 - \frac{p^2}{4} (1 - \xi^2)^2 \\ &= \frac{p^2}{4} (x + \xi)^2 + 2\xi^2 - (x - \xi)^2 + 2\xi^2 \\ &= p^2 \xi^2 \end{aligned}$$

so $y = p \xi$

Sec 15, Prob 3:

$$\Delta\phi = 2 \int_{r_{\min}}^{r_{\max}} \frac{M dr / r^2}{\sqrt{2m(E-U) - M^2/r^2}}$$

This is the change in ϕ as r goes from r_{\min} to r_{\max} and then back to r_{\min} .

A closed bound orbit would have $\Delta\phi = 2\pi m/n$ for m, n integers.

Consider: $U = -\frac{\alpha}{r} + \delta U$ where $|\delta U| \ll |\frac{\alpha}{r}|$

$$\text{For } \delta U = 0, \Delta\phi = 2\pi$$

$$\Delta\phi = 2 \int_{r_{\min}}^{r_{\max}} \frac{M dr / r^2}{\sqrt{2m(E - (U + \delta U)) - M^2/r^2}}$$

$$= 2 \int_{r_{\min}}^{r_{\max}} \frac{M dr / r^2}{\sqrt{2m(E-U) - 2m\delta U - M^2/r^2}}$$

$$= 2 \int_{r_{\min}}^{r_{\max}} \frac{M dr / r^2}{\sqrt{(2m(E-U) - M^2/r^2)} \left(1 - \frac{2m\delta U}{2m(E-U) - M^2/r^2} \right)}$$

$$\approx 2 \int_{r_{\min}}^{r_{\max}} \frac{M dr / r^2}{\sqrt{2m(E-U) - M^2/r^2}} \left[1 + \frac{m\delta U}{2m(E-U) - M^2/r^2} \right]$$

For $V = -\alpha/r$:

$$\Delta\phi \approx 2\pi + 2 \int_{r_{min}}^{r_{max}} \frac{Md\tau/r^2}{\sqrt{2m(E-U) - M^2/r^2}} d\delta U$$

$$\approx 2\pi + \delta\phi$$

where

$$r_{max}$$

$$\delta\phi \equiv \int_{r_{min}}^{r_{max}} \frac{2m\delta U M dr/r^2}{\sqrt{2m(E-U) - M^2/r^2}}^{3/2}$$

$$= \frac{2}{2M} \left[\int_{r_{min}}^{r_{max}} \frac{2m\delta U dr}{\sqrt{2m(E-U) - M^2/r^2}} \right]$$

Evaluate terms in integrand along unperturbed path since δU is already small

$$\frac{p}{r} = 1 + e\cos\phi \quad , \quad p = q(1-e^2)$$

$$\rightarrow -\frac{p}{r^2} dr = -e\sin\phi d\phi \rightarrow \boxed{dr = \frac{r^2 e \sin\phi}{p} d\phi}$$

$$\sqrt{ } = \sqrt{2mE + 2m\alpha - \frac{M^2}{r^2}}$$

$$= \sqrt{-2m|E| + \frac{4m|E|\alpha}{r} - \frac{2m|E|p^2}{(1-e^2)^2}}$$

$$\begin{aligned} \sqrt{ } &= \sqrt{2m|E|} \sqrt{-1 + \frac{2}{(1-e^2)}(1+e\cos\phi) - \frac{1}{(1-e^2)}(1+e\cos\phi)^2} \\ &= \frac{\sqrt{2m|E|}}{\sqrt{1-e^2}} \sqrt{-1/(1-e^2) + 2/(1+e\cos\phi) - (1+2e\cos\phi + e^2\cos^2\phi)} \\ &= \frac{\sqrt{2m|E|}}{\sqrt{1-e^2}} \sqrt{e^2(1-\cos^2\phi)} \\ &= \frac{\sqrt{2m|E|}}{\sqrt{1-e^2}} e \sin\phi \end{aligned}$$

Thus,

$$\begin{aligned} |\delta\phi| &= \frac{2}{2M} \left[2m \int_0^\pi \frac{\delta U r^2 e \sin\phi d\phi}{\sqrt{2m|E|} e \sin\phi} \right] \\ &= \frac{2}{2M} \left[\frac{2m}{p \sqrt{\frac{2m|E|}{1-e^2}}} \int_0^\pi d\phi r^2 \delta U \right] \\ &= \frac{2}{2M} \left[\frac{2m}{M} \int_0^\pi d\phi r^2 \delta U \right] \end{aligned}$$

$$\text{using (15.6): } \frac{M}{\sqrt{2m|E|}} = \frac{p}{\sqrt{1-e^2}}$$

Evaluate:

$$\delta\phi = \frac{\partial}{\partial M} \left[\frac{2m}{M} \int_0^\pi d\phi r^2 \delta U \right]$$

For (a) $\delta U = \beta/r^2$, (b) $\delta U = \gamma/r^3$

(a) $\delta\phi = \frac{\partial}{\partial M} \left[\frac{2m}{M} \int_0^\pi d\phi \beta \right]$

$$= 2\pi\beta m \frac{\partial}{\partial M} \left(\frac{1}{M} \right)$$

$$= -\frac{2\pi\beta m}{M^2}$$

$$= \boxed{-\frac{2\pi\beta}{\alpha p}}$$

Recall: $j^0 = \frac{M^2}{m\alpha}$

$$\leftarrow \quad \alpha p = \frac{M^2}{m}$$

(b) $\delta\phi = \frac{\partial}{\partial M} \left[\frac{2m}{M} \int_0^\pi d\phi \frac{\gamma}{r} \right]$

$$= \frac{\partial}{\partial M} \left[\frac{2m\gamma}{M} \int_0^\pi d\phi \left(\frac{1 + e^{i\phi}}{r} \right) \right]$$

$$= 2m\gamma \frac{\partial}{\partial M} \left[\frac{1}{mp} \left(\pi + e^{i\phi} \right) \right]$$

Now: $\frac{1}{mp} = \frac{m\alpha}{M^3} \rightarrow \frac{\partial}{\partial M} \left(\frac{1}{mp} \right) = -\frac{3m\alpha}{M^4}$

so $\delta\phi = -6\pi\gamma \frac{m^2\alpha}{M^4}$

$$= -6\pi\gamma\alpha \left(\frac{1}{p\alpha} \right)^2$$

$$= \boxed{-\frac{6\pi\gamma}{p^2\alpha}}$$