Calculus

Joseph Le

Contents

Ι	Preliminaries	7				
1	Functions 1.1 Sets 1.2 What is a function? 1.3 Common Functions 1.4 Types of Functions 1.4.1 Even and Odd Functions 1.4.2 Increasing and Decreasing Functions 1.4.3 Periodic Functions 1.5 Inverse Function 1.6 Function Compositions 1.7 Function Transformations and Operations					
2	2.1 Linear Functions	18 18 18 18 18 18				
3	3.1 Conic Sections	15 15 15 15				
II	Differential Calculus	17				
4	4.1 Definition of a Limit	19 19 19 19				
5		21 21 21 21 21 21 21				

4 CONTENTS

	5.4	5.4.1 Implicit Differentiation	21 21 21 21
6	Apr	plications of Derivatives	23
Ū	6.1		23
	6.2		23
	6.3	•	23
		•	
			_
II	1 1	ntegral Calculus 2	25
7	Inte	gration and Integrals	27
	7.1		27
	7.2	Riemann Sums	27
	7.3	The Fundamental Theorem of Calculus	27
_	.		
8		0 1	29
	8.1		29
	8.2		29
	8.3 8.4		29 29
	8.5		29 29
	0.0	Try per bone Substitutions	29
9	App	olication of Integrals	31
	9.1	Areas Between Curves	31
	9.2	Valumes of 3D Shapes	31
		9.2.1 The Disk Method	31
		9.2.2 The Washer Method	31
		9.2.3 The Shell Method	31
	9.3	Arc Length	31
10	T	nonen Intermela and Numerical Intermetion	
10	-	. 0	33 33
			ээ 33
	10.2		33
		<u>.</u>	33
		10.2.2 Shinpson's Itule).)
11	Sequ	, , ,	35
		1	35
	11.2		35
			35
			35
		1	35
	44.0	S .	35
	11.3	3	35
		9	35
		<u>.</u>	35
		1	35
		Θ	35 25
			35 35
	11 /		35 35
			ээ 35
	TT.0	INTO DOLLO	v

CONTENTS 5

IV	Multivariable Calculus	37
12 '	Vectors and Vector Spaces	39
	2.1 Vectors in 2D/3D	39
	2.2 Vector Products	39
	12.2.1 Dot Products	39
	12.2.2 Cross Products	39
	2.3 Lines and Planes in Space	39
13	Functions of Several Variables	41
	3.1 Parametric Curves	41
	3.2 Derivatives and Integrals of Vector Functions	41
	3.3 Curvature	41
14]	Partial Derivatives	43
	4.1 Limits and Continuity in Higher Dimensions	43
	4.2 Partial Derivatives	43
	4.3 The Chain Rule	43
	4.4 The Gradient	43
-	4.5 Extremal Values	43
15 I	Multiple Integrals	45
	5.1 Double Integrals	45
	5.2 Coordinate Systems	45
	15.2.1 Polar coordinates	45
	15.2.2 Cylindrical Coordinates	45
	15.2.3 Spherical Coordinates	45
	5.3 Surface Area and Volume	45
	5.4 Triple Integrals	45
	5.5 Change of Variables	45
16 '	Vector Calculus	47
	6.1 Line Integrals	47
	6.2 Green's Theorem	47
	6.3 Divergence and Curl	47
	6.4 Surface Integrals	47
	6.5 Stokes' Theorem	47
	6.6 The Divergence Theorem	47
-		41
	Applications of Multivariable Calculus	49
	7.1 Lagrange Multipliers and Optimization	49
	7.2 The Jacobian	49
	7.3 Center of Mass	49
	7.4 Fluid Flows	49
${f V}$	Annondix	K 1
V	Appendix	51

6 CONTENTS

Part I Preliminaries

Functions

1.1 Sets

Before defining what a function is or what it does, it is important to briefly discuss what goes into function and what comes out. Simply, *sets* are a collection of items and each one of those items are usually referred to as *elements*. Without getting into the weeds of set theory, sets can contain pretty much anything from numbers, functions, and other sets [3].

Some common sets that you may be familiar with are the *natural numbers* $\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$, the *integers* $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$, and the *real numbers* \mathbb{R} , which is usually represented via a number line. Sets can also be intervals on the real line (i.e. [a,b) is an interval on \mathbb{R} containing a but not b) or even the possible results of flipping a coin $C = \{H, T\}$.

We will now define the basic notation when dealing with sets and the operations that can be performed on sets. We say that x is an element of a set A with the notation $x \in A$ and when x is not in A, we say $x \notin A$. For example, given the set $A = \{1, 2, 3, 4\}$, we can say that $1 \in A$ is true as well as $5 \notin A$.

The notion of combining sets comes with *unions* and *intersections*. Given A and B are sets, the union of A and B is denoted as $A \cup B$ and is equal to the set the contains elements in either A or B. Similarly, the intersection between A and B is denoted as $A \cap B$ and is the set that contains elements that are in both A and B. For example, given the sets $A = \{1, 2, 3, 4\}$ and $B = \{3, 4, 5, 6\}$, the union and intersection between A and B is

$$A \cup B = \{1, 2, 3, 4, 5, 6\}$$

$$A \cap B = \{3, 4\}$$
 (1.1)

Subsets are often used to relate different sets and their elements. A set A is said to be a subset of another set B if all of the elements of A are also within B and is denoted as $A \subseteq B$ and a set A is equal to a set B if and only if $A \subseteq B$ and $B \subseteq A$. An important subset is the empty set represented by the symbols \emptyset , \emptyset , or simply $\{\}$. It is important to note that the empty set is also a subset of all sets.

Using sets by listing them out can become cumbersome and sometimes confusing, instead set builder notation is used to build a set based on a rule. For example, the set of all positive even integers can be written as

$$A = \{2, 4, 6, 8, \dots\} = \{z : z \text{ is an positive even integer}\} = \{z : z = 2n, n \in \mathbb{Z} \text{ and } n > 0\}$$
 (1.2)

Here, the : stands for "such that" which indicates the rule (the words "such that" or the symbol | is also often used). The rational numbers $\mathbb Q$ can also be constructed via the integers with

$$\mathbb{Q} = \{ \frac{p}{q} : p, q \in \mathbb{Z} \text{ and } q \neq 0 \}$$
 (1.3)

As mentioned previously, intervals on the real number line can be represented as sets. Given two values a and b and assuming that $a \le b$, intervals on the real line are represented as

• Closed interval:
$$[a,b] = \{x \in \mathbb{R} : a \le x \le b\}$$

a b

• Open interval: $(a,b) = \{x \in \mathbb{R} : a < x < b\}$

• Half-open interval:

$$-(a,b] = \{x \in \mathbb{R} : a < x \le b\}$$

$$-[a,b] = \{x \in \mathbb{R} : a \le x < b\}$$

$$a \qquad b$$

• Infinite interval:

$$-(a, \infty) = \{x \in \mathbb{R} : a < x\}$$

$$-[a, \infty) = \{x \in \mathbb{R} : a \le x\}$$

$$-(-\infty, b) = \{x \in \mathbb{R} : x < b\}$$

$$-(-\infty, b] = \{x \in \mathbb{R} : x \le b\}$$

$$-(-\infty, \infty) = \mathbb{R}$$

$$\leftarrow (-\infty, \infty) = \mathbb{R}$$

Here, the open brackets (and) indicate that the respective endpoint is not included, while the closed brackets [and] indicate that the respective endpoint is included in the interval.

The Cartesian product (also known as the direct product) is used often to describe ordered pairs or even higher-dimension coordinates. Some examples include the xy-plane also known as \mathbb{R}^2 , 3D space with \mathbb{R}^3 . The Cartesian product for two sets A and B is defined as

$$A \times B = \{(a, b) : a \in A, b \in B\}$$
 (1.4)

Note that, in general, $A \times B \neq B \times A$ as the operation is order dependent. Higher-order products are defined with sets A_1, A_2, \ldots, A_n as

$$A_1 \times A_2 \times \dots \times A_n = \{(a_1, a_2, \dots, a_n) : a_i \in A_i, i \in \{1, 2, \dots, n\}\}$$
(1.5)

As previously mentioned, the 2D plane as well as the 3D plane can be constructed using Cartesian products as well and is done as such with the set of real numbers \mathbb{R}

$$\mathbb{R}^{2} = \mathbb{R} \times \mathbb{R} = \{(x, y) : x, y \in \mathbb{R}\}$$

$$\mathbb{R}^{3} = \mathbb{R} \times \mathbb{R} \times \mathbb{R} = \{(x, y, z) : x, y, z \in \mathbb{R}\}$$

$$\vdots$$

$$\mathbb{R}^{n} = \mathbb{R} \times \mathbb{R} \times \dots \times \mathbb{R} = \{(x_{1}, x_{2}, \dots, x_{n}) : x_{i} \in \mathbb{R}, i \in \{1, 2, \dots, n\}\}$$

$$(1.6)$$

For further and more formal discussion on sets and naive set theory as well as formal logic and proof writing, refer to [2] or [3].

1.2 What is a function?

Functions are objects in math that describe a relationship or mapping between two sets. Given two sets X and Y, a function f maps the unique elements of X, called the *domain* of the function, to elements in the set Y called the *codomain* of the function and the relationship is denoted as $f: X \to Y$ ("f maps from X to Y") or y = f(x) ("f equals f of f and f is known as the *dependent variable* with f as the range of the function and f is known as the *independent variable* or the *argument* of the function. The range of a function is the set of all possible values that f(f) is able to output with f as its domain, note that the range is a subset of the codomain f but is not always equal. Formally, the definition is as follows

Definition 1.1. A function is a mapping or rule that assigns each element from a set called the domain X of the function to a unique element in the range R = f(X) of the function which is a subset of the codomain $R \subseteq Y$ and the element $x \in X$ is mapped to an element in $y \in R \subseteq Y$ with $x \mapsto y$.

Further emphasis must be made on the elements y = f(x). The function f(x) maps the element $x \in X$ to exactly one element $y \in Y$ [4]. For single variable functions, this can be tested using the vertical line test on a function's graph (covered shortly). If a formula results in two answers with one input (i.e. the equation for a circle: $x^2 + y^2 = r$ gives two points of y for each x) it is no longer considered a function. Figure 1.1 shows how a function maps elements of the domain to the range.




Figure 1.1: Function mapping diagram showing the domain X, codomain Y, and range f(X)

Example 1.2. Some real-world examples of functions are

• The area of a circle: $A(r) = \pi r^2$ • The height of a falling ball: $h(t) = h_0 + v_0 t - (1/2) g t^2$ • Compound interest: $A(t) = P(1 + \frac{r}{n})^{nt}$ • Temperature conversion: $C(F) = \frac{5}{9}(F - 32)$

It can be useful to think of functions as machines that take in an input and produces an output. The examples above show such cases using mathematical formulas that give a set of instructions on how the input is transformed from the element in the domain to the element in the range.

Another way to represent functions is with its graph which is a set of points consisting of the input and its corresponding output and is represented with the set of ordered pairs $\{(x,y):y=f(x),x\in\mathbb{R}\}$ for functions that accept real numbers.

Example 1.3. A familiar example is the linear function f(x) = mx + b where m is the slope of the line and b is the vertical offset. Suppose m = 1 and b = 0, we get the following graph for the function f(x) = x in figure 1.3

Example 1.4. Another example is the quadratic function of the form $f(x) = ax^2 + bx + c$, where $a, b, c \in \mathbb{R}$ are the coefficients of the function and determine its shape. Suppose that a = 1, b = 0, c = -1 resulting in the function $f(x) = x^2 - 1$. Using knowledge of the roots a quadratic function, it can be deduced that this will result in a parabola with x-intersections at x = -1 and x = 1 which can be confirmed in the graph in figure 1.4




Figure 1.2: Graph of the function f(x) = x




Figure 1.3: Graph of the function $f(x) = x^2 - 1$

1.3 Common Functions

1.4 Types of Functions

- 1.4.1 Even and Odd Functions
- 1.4.2 Increasing and Decreasing Functions
- 1.4.3 Periodic Functions

1.5 Inverse Function

1.6 Function Compositions

1.7 Function Transformations and Operations

Special Functions

- 2.1 Linear Functions
- 2.2 Quadratic Functions
- 2.3 Polynomial and Rational Functions
- 2.4 Exponential and Logarithmic Functions
- 2.5 Absolute Value and Piecewise Functions
- 2.6 Systems of Equations

Analytic Geometry

- 3.1 Conic Sections
- 3.2 Parametric Equations
- 3.3 Polar Coordinates

Part II Differential Calculus

Limits and Continuity

- 4.1 Defnition of a Limit
- 4.2 One-Sided Limits and Limits to Infinity
- 4.3 Continuity and the Intermediate Value Theorem
- 4.4 The Squeeze Theorem

Differentiation and Derivatives

- 5.1 The Limit Definition of the Derivative
- 5.2 Differential Rules
- 5.2.1 The Power Rule
- 5.2.2 The Product Rule
- 5.2.3 The Quotient Rule
- 5.2.4 The Chain Rule
- 5.3 Common and Special Derivatives
- 5.4 Advanced Differential Techniques
- 5.4.1 Implicit Differentiation
- ${\bf 5.4.2}\quad {\bf Logarithmic\ Differentiation}$
- 5.4.3 Higher-Order Derivatives

Applications of Derivatives

- 6.1 Related Rates
- 6.2 Optimization
- 6.3 L'Hôpital's Rule Rule

Part III Integral Calculus

Integration and Integrals

- 7.1 Antiderivatives
- 7.2 Riemann Sums
- 7.3 The Fundamental Theorem of Calculus

Integration Techniques

- 8.1 Substitution
- 8.2 Integration by Parts
- 8.3 Partial Fractions
- 8.4 Trigonometric Substitutions
- 8.5 Hyperbolic Substitutions

Application of Integrals

- 9.1 Areas Between Curves
- 9.2 Valumes of 3D Shapes
- 9.2.1 The Disk Method
- 9.2.2 The Washer Method
- 9.2.3 The Shell Method
- 9.3 Arc Length

Improper Integrals and Numerical Integration

- 10.1 Improper Integrals and Their Convergence
- 10.2 Numerical Integration
- 10.2.1 Trapezoidal Rule
- 10.2.2 Simpson's Rule

Sequences, Series, and Convergence

- 11.1 Sequences
- 11.2 Infinite Series
- 11.2.1 Algebraic Series
- 11.2.2 Geometric Series
- 11.2.3 p-Series
- 11.2.4 Alternating Series
- 11.3 Convergence Tests
- 11.3.1 The Divergence Test
- 11.3.2 The Comparison Test
- 11.3.3 The Limit Comparison Test
- 11.3.4 The Integral Test
- 11.3.5 The Ratio Test
- 11.3.6 The Root Test
- 11.4 Power Series
- 11.5 Taylor Series

Part IV Multivariable Calculus

Vectors and Vector Spaces

- 12.1 Vectors in 2D/3D
- 12.2 Vector Products
- 12.2.1 Dot Products
- 12.2.2 Cross Products
- 12.3 Lines and Planes in Space

Functions of Several Variables

- 13.1 Parametric Curves
- 13.2 Derivatives and Integrals of Vector Functions
- 13.3 Curvature

Partial Derivatives

- 14.1 Limits and Continuity in Higher Dimensions
- 14.2 Partial Derivatives
- 14.3 The Chain Rule
- 14.4 The Gradient
- 14.5 Extremal Values

Multiple Integrals

- 15.1 Double Integrals
- 15.2 Coordinate Systems
- 15.2.1 Polar coordinates
- 15.2.2 Cylindrical Coordinates
- 15.2.3 Spherical Coordinates
- 15.3 Surface Area and Volume
- 15.4 Triple Integrals
- 15.5 Change of Variables

Vector Calculus

- 16.1 Line Integrals
- 16.2 Green's Theorem
- 16.3 Divergence and Curl
- 16.4 Surface Integrals
- 16.5 Stokes' Theorem
- 16.6 The Divergence Theorem

Applications of Multivariable Calculus

- 17.1 Lagrange Multipliers and Optimization
- 17.2 The Jacobian
- 17.3 Center of Mass
- 17.4 Fluid Flows

$\begin{array}{c} {\rm Part~V} \\ {\rm Appendix} \end{array}$

Bibliography

- [1] Stephen Abbott. Understanding Analysis. 2nd Edition. Springer, 2015.
- [2] Paul R. Halmos. Naive Set Theory. Springer, 1998.
- [3] Richard Hammack. Book of Proof. 3rd Edition. Richard Hammack, 2018.
- [4] Georgi E. Shilov. Elementary Real and Complex Analysis. Trans. by Richard A. Silverman. Dover, 1973.

Index

```
function, 11
argument, 11
codomain, 11
dependent variable, 11
domain, 11
graph, 11
independent variable, 11
range, 11

set, 9
Cartesian product, 10
direct product, 10
element, 9
intersection, 9
union, 9
```