

Calculus

Joseph Le

Contents

I	Preliminaries	7
1	Functions	9
1.1	Sets	9
1.2	What is a function?	11
1.3	Common Functions	12
1.4	Types of Functions	12
1.4.1	Even and Odd Functions	12
1.4.2	Increasing and Decreasing Functions	12
1.4.3	Periodic Functions	12
1.5	Inverse Function	12
1.6	Function Compositions	12
1.7	Function Transformations and Operations	12
2	Special Functions	13
2.1	Linear Functions	13
2.2	Quadratic Functions	13
2.3	Polynomial and Rational Functions	13
2.4	Exponential and Logarithmic Functions	13
2.5	Absolute Value and Piecewise Functions	13
2.6	Systems of Equations	13
3	Analytic Geometry	15
3.1	Conic Sections	15
3.2	Parametric Equations	15
3.3	Polar Coordinates	15
II	Differential Calculus	17
4	Limits and Continuity	19
4.1	Definition of a Limit	19
4.2	One-Sided Limits and Limits to Infinity	19
4.3	Continuity and the Intermediate Value Theorem	19
4.4	The Squeeze Theorem	19
5	Differentiation and Derivatives	21
5.1	The Limit Definition of the Derivative	21
5.2	Differential Rules	21
5.2.1	The Power Rule	21
5.2.2	The Product Rule	21
5.2.3	The Quotient Rule	21
5.2.4	The Chain Rule	21
5.3	Common and Special Derivatives	21

5.4	Advanced Differential Techniques	21
5.4.1	Implicit Differentiation	21
5.4.2	Logarithmic Differentiation	21
5.4.3	Higher-Order Derivatives	21
6	Applications of Derivatives	23
6.1	Related Rates	23
6.2	Optimization	23
6.3	L'Hôpital's Rule Rule	23
III	Integral Calculus	25
7	Integration and Integrals	27
7.1	Antiderivatives	27
7.2	Riemann Sums	27
7.3	The Fundamental Theorem of Calculus	27
8	Integration Techniques	29
8.1	Substitution	29
8.2	Integration by Parts	29
8.3	Partial Fractions	29
8.4	Trigonometric Substitutions	29
8.5	Hyperbolic Substitutions	29
9	Application of Integrals	31
9.1	Areas Between Curves	31
9.2	Volumes of 3D Shapes	31
9.2.1	The Disk Method	31
9.2.2	The Washer Method	31
9.2.3	The Shell Method	31
9.3	Arc Length	31
10	Improper Integrals and Numerical Integration	33
10.1	Improper Integrals and Their Convergence	33
10.2	Numerical Integration	33
10.2.1	Trapezoidal Rule	33
10.2.2	Simpson's Rule	33
11	Sequences, Series, and Convergence	35
11.1	Sequences	35
11.2	Infinite Series	35
11.2.1	Algebraic Series	35
11.2.2	Geometric Series	35
11.2.3	p-Series	35
11.2.4	Alternating Series	35
11.3	Convergence Tests	35
11.3.1	The Divergence Test	35
11.3.2	The Comparison Test	35
11.3.3	The Limit Comparison Test	35
11.3.4	The Integral Test	35
11.3.5	The Ratio Test	35
11.3.6	The Root Test	35
11.4	Power Series	35
11.5	Taylor Series	35

IV	Multivariable Calculus	37
12	Vectors and Vector Spaces	39
12.1	Vectors in 2D/3D	39
12.2	Vector Products	39
12.2.1	Dot Products	39
12.2.2	Cross Products	39
12.3	Lines and Planes in Space	39
13	Functions of Several Variables	41
13.1	Parametric Curves	41
13.2	Derivatives and Integrals of Vector Functions	41
13.3	Curvature	41
14	Partial Derivatives	43
14.1	Limits and Continuity in Higher Dimensions	43
14.2	Partial Derivatives	43
14.3	The Chain Rule	43
14.4	The Gradient	43
14.5	Extremal Values	43
15	Multiple Integrals	45
15.1	Double Integrals	45
15.2	Coordinate Systems	45
15.2.1	Polar coordinates	45
15.2.2	Cylindrical Coordinates	45
15.2.3	Spherical Coordinates	45
15.3	Surface Area and Volume	45
15.4	Triple Integrals	45
15.5	Change of Variables	45
16	Vector Calculus	47
16.1	Line Integrals	47
16.2	Green's Theorem	47
16.3	Divergence and Curl	47
16.4	Surface Integrals	47
16.5	Stokes' Theorem	47
16.6	The Divergence Theorem	47
17	Applications of Multivariable Calculus	49
17.1	Lagrange Multipliers and Optimization	49
17.2	The Jacobian	49
17.3	Center of Mass	49
17.4	Fluid Flows	49
V	Appendix	51

Part I

Preliminaries

Chapter 1

Functions

1.1 Sets

Before defining what a function is or what it does, it is important to briefly discuss what goes into function and what comes out. Simply, *sets* are a collection of items and each one of those items are usually referred to as *elements*. Without getting into the weeds of set theory, sets can contain pretty much anything from numbers, functions, and other sets [3].

Some common sets that you may be familiar with are the *natural numbers* $\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$, the *integers* $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$, and the *real numbers* \mathbb{R} , which is usually represented via a number line. Sets can also be intervals on the real line (i.e. $[a, b)$ is an interval on \mathbb{R} containing a but not b) or even the possible results of flipping a coin $C = \{H, T\}$.

We will now define the basic notation when dealing with sets and the operations that can be performed on sets. We say that x is an element of a set A with the notation $x \in A$ and when x is not in A , we say $x \notin A$. For example, given the set $A = \{1, 2, 3, 4\}$, we can say that $1 \in A$ is true as well as $5 \notin A$.

The notion of combining sets comes with *unions* and *intersections*. Given A and B are sets, the union of A and B is denoted as $A \cup B$ and is equal to the set that contains elements in either A or B . Similarly, the intersection between A and B is denoted as $A \cap B$ and is the set that contains elements that are in both A and B . For example, given the sets $A = \{1, 2, 3, 4\}$ and $B = \{3, 4, 5, 6\}$, the union and intersection between A and B is

$$\begin{aligned} A \cup B &= \{1, 2, 3, 4, 5, 6\} \\ A \cap B &= \{3, 4\} \end{aligned} \tag{1.1}$$

Subsets are often used to relate different sets and their elements. A set A is said to be a subset of another set B if all of the elements of A are also within B and is denoted as $A \subseteq B$ and a set A is equal to a set B if and only if $A \subseteq B$ and $B \subseteq A$. An important subset is the empty set represented by the symbols \emptyset , \emptyset , or simply $\{\}$. It is important to note that the empty set is also a subset of all sets.

Using sets by listing them out can become cumbersome and sometimes confusing, instead set builder notation is used to build a set based on a rule. For example, the set of all positive even integers can be written as

$$A = \{2, 4, 6, 8, \dots\} = \{z : z \text{ is an positive even integer}\} = \{z : z = 2n, n \in \mathbb{Z} \text{ and } n > 0\} \tag{1.2}$$

Here, the $:$ stands for "such that" which indicates the rule (the words "such that" or the symbol $|$ is also often used). The *rational numbers* \mathbb{Q} can also be constructed via the integers with


$$\mathbb{Q} = \left\{ \frac{p}{q} : p, q \in \mathbb{Z} \text{ and } q \neq 0 \right\} \tag{1.3}$$

As mentioned previously, intervals on the real number line can be represented as sets. Given two values a and b and assuming that $a \leq b$, intervals on the real line are represented as

- Closed interval: $[a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$



- Open interval: $(a, b) = \{x \in \mathbb{R} : a < x < b\}$



- Half-open interval:

$$- (a, b] = \{x \in \mathbb{R} : a < x \leq b\}$$




$$- [a, b) = \{x \in \mathbb{R} : a \leq x < b\}$$




- Infinite interval:


$$- (a, \infty) = \{x \in \mathbb{R} : a < x\}$$




$$- [a, \infty) = \{x \in \mathbb{R} : a \leq x\}$$




$$- (-\infty, b) = \{x \in \mathbb{R} : x < b\}$$



$$- (-\infty, b] = \{x \in \mathbb{R} : x \leq b\}$$



$$- (-\infty, \infty) = \mathbb{R}$$



Here, the open brackets (and) indicate that the respective endpoint is not included, while the closed brackets [and] indicate that the respective endpoint is included in the interval.

The *Cartesian product* (also known as the *direct product*) is used often to describe ordered pairs or even higher-dimension coordinates. Some examples include the xy -plane also known as \mathbb{R}^2 , 3D space with \mathbb{R}^3 . The Cartesian product for two sets A and B is defined as

$$A \times B = \{(a, b) : a \in A, b \in B\} \quad (1.4)$$

Note that, in general, $A \times B \neq B \times A$ as the operation is order dependent. Higher-order products are defined with sets A_1, A_2, \dots, A_n as

$$A_1 \times A_2 \times \dots \times A_n = \{(a_1, a_2, \dots, a_n) : a_i \in A_i, i \in \{1, 2, \dots, n\}\} \quad (1.5)$$

As previously mentioned, the 2D plane as well as the 3D plane can be constructed using Cartesian products as well and is done as such with the set of real numbers \mathbb{R}

$$\begin{aligned} \mathbb{R}^2 &= \mathbb{R} \times \mathbb{R} = \{(x, y) : x, y \in \mathbb{R}\} \\ \mathbb{R}^3 &= \mathbb{R} \times \mathbb{R} \times \mathbb{R} = \{(x, y, z) : x, y, z \in \mathbb{R}\} \\ &\vdots \\ \mathbb{R}^n &= \mathbb{R} \times \mathbb{R} \times \dots \times \mathbb{R} = \{(x_1, x_2, \dots, x_n) : x_i \in \mathbb{R}, i \in \{1, 2, \dots, n\}\} \end{aligned} \quad (1.6)$$

For more discussion on sets and set theory, refer to [2] or [3].

1.2 What is a function?

Functions are objects in math that describe a relationship or mapping between two sets. Given two sets X and Y , a function f maps the unique elements of X , called the *domain* of the function, to elements in the set Y called the *codomain* of the function and the relationship is denoted as $f : X \rightarrow Y$ (" f maps from X to Y ") or $y = f(x)$ (" y equals f of x " [1]), where $y \in R \subseteq Y$ is known as the *dependent variable* with R as the range of the function and $x \in X$ is known as the *independent variable* or the *argument* of the function. The *range* of a function is the set of all possible values that $f(x)$ is able to output with X as its domain, note that the range is a subset of the codomain Y but is not always equal. Formally, the definition is as follows

Definition 1.1. A *function* is a mapping or rule that assigns each element from a set called the domain X of the function to a unique element in the range $R = f(X)$ of the function which is a subset of the codomain $R \subseteq Y$ and the element $x \in X$ is mapped to an element in $y \in R \subseteq Y$ with $x \mapsto y$.

Further emphasis must be made on the elements $y = f(x)$. The function $f(x)$ maps the element $x \in X$ to exactly one element $y \in Y$ [4]. For single variable functions, this can be tested using the vertical line test on a function's graph (covered shortly). If a formula results in two answers with one input (i.e. the equation for a circle: $x^2 + y^2 = r$ gives two points of y for each x) it is no longer considered a function. Figure 1.1 shows how a function maps elements of the domain to the range.




Figure 1.1: Function mapping diagram showing the domain X , codomain Y , and range $f(X)$

Example 1.2. Some real-world examples of functions are

- The area of a circle: $A(r) = \pi r^2$
- The height of a falling ball: $h(t) = h_0 + v_0 t - (1/2)gt^2$
- Compound interest: $A(t) = P(1 + \frac{r}{n})^{nt}$
- Temperature conversion: $C(F) = \frac{5}{9}(F - 32)$
-

It can be useful to think of functions as machines that take in an input and produces an output. The examples above show such cases using mathematical formulas that give a set of instructions on how the input is transformed from the element in the domain to the element in the range.

Another way to represent functions is with its *graph* which is a set of points consisting of the input and its corresponding output and is represented with the set of ordered pairs $\{(x, y) : y = f(x), x \in \mathbb{R}\}$ for functions that accept real numbers.

Example 1.3. A familiar example is the linear function $f(x) = mx + b$ where m is the slope of the line and b is the vertical offset. Suppose $m = 1$ and $b = 0$, we get the following graph for the function $f(x) = x$ in figure 1.3

Example 1.4. Another example is the quadratic function of the form $f(x) = ax^2 + bx + c$, where $a, b, c \in \mathbb{R}$ are the coefficients of the function and determine its shape. Suppose that $a = 1, b = 0, c = -1$ resulting in the function $f(x) = x^2 - 1$. Using knowledge of the roots a quadratic function, it can be deduced that this will result in a parabola with x -intersections at $x = -1$ and $x = 1$ which can be confirmed in the graph in figure 1.4




Figure 1.2: Graph of the function $f(x) = x$




Figure 1.3: Graph of the function $f(x) = x^2 - 1$

1.3 Common Functions

1.4 Types of Functions

1.4.1 Even and Odd Functions

1.4.2 Increasing and Decreasing Functions

1.4.3 Periodic Functions

1.5 Inverse Function

1.6 Function Compositions

1.7 Function Transformations and Operations

Chapter 2

Special Functions

2.1 Linear Functions

2.2 Quadratic Functions

2.3 Polynomial and Rational Functions

2.4 Exponential and Logarithmic Functions

2.5 Absolute Value and Piecewise Functions

2.6 Systems of Equations

Chapter 3

Analytic Geometry

3.1 Conic Sections

3.2 Parametric Equations

3.3 Polar Coordinates

Part II

Differential Calculus

Chapter 4

Limits and Continuity

4.1 Definition of a Limit

4.2 One-Sided Limits and Limits to Infinity

4.3 Continuity and the Intermediate Value Theorem

4.4 The Squeeze Theorem

Chapter 5

Differentiation and Derivatives

5.1 The Limit Definition of the Derivative

5.2 Differential Rules

5.2.1 The Power Rule

5.2.2 The Product Rule

5.2.3 The Quotient Rule

5.2.4 The Chain Rule

5.3 Common and Special Derivatives

5.4 Advanced Differential Techniques

5.4.1 Implicit Differentiation

5.4.2 Logarithmic Differentiation

5.4.3 Higher-Order Derivatives

Chapter 6

Applications of Derivatives

6.1 Related Rates

6.2 Optimization

6.3 L'Hôpital's Rule Rule

Part III

Integral Calculus

Chapter 7

Integration and Integrals

7.1 Antiderivatives

7.2 Riemann Sums

7.3 The Fundamental Theorem of Calculus

Chapter 8

Integration Techniques

8.1 Substitution

8.2 Integration by Parts

8.3 Partial Fractions

8.4 Trigonometric Substitutions

8.5 Hyperbolic Substitutions

Chapter 9

Application of Integrals

9.1 Areas Between Curves

9.2 Volumes of 3D Shapes

9.2.1 The Disk Method

9.2.2 The Washer Method

9.2.3 The Shell Method

9.3 Arc Length

Chapter 10

Improper Integrals and Numerical Integration

10.1 Improper Integrals and Their Convergence

10.2 Numerical Integration

10.2.1 Trapezoidal Rule

10.2.2 Simpson's Rule

Chapter 11

Sequences, Series, and Convergence

11.1 Sequences

11.2 Infinite Series

11.2.1 Algebraic Series

11.2.2 Geometric Series

11.2.3 p-Series

11.2.4 Alternating Series

11.3 Convergence Tests

11.3.1 The Divergence Test

11.3.2 The Comparison Test

11.3.3 The Limit Comparison Test

11.3.4 The Integral Test

11.3.5 The Ratio Test

11.3.6 The Root Test

11.4 Power Series

11.5 Taylor Series

Part IV

Multivariable Calculus

Chapter 12

Vectors and Vector Spaces

12.1 Vectors in 2D/3D

12.2 Vector Products

12.2.1 Dot Products

12.2.2 Cross Products

12.3 Lines and Planes in Space

Chapter 13

Functions of Several Variables

13.1 Parametric Curves

13.2 Derivatives and Integrals of Vector Functions

13.3 Curvature

Chapter 14

Partial Derivatives

14.1 Limits and Continuity in Higher Dimensions

14.2 Partial Derivatives

14.3 The Chain Rule

14.4 The Gradient

14.5 Extremal Values

Chapter 15

Multiple Integrals

15.1 Double Integrals

15.2 Coordinate Systems

15.2.1 Polar coordinates

15.2.2 Cylindrical Coordinates

15.2.3 Spherical Coordinates

15.3 Surface Area and Volume

15.4 Triple Integrals

15.5 Change of Variables

Chapter 16

Vector Calculus

16.1 Line Integrals

16.2 Green's Theorem

16.3 Divergence and Curl

16.4 Surface Integrals

16.5 Stokes' Theorem

16.6 The Divergence Theorem

Chapter 17

Applications of Multivariable Calculus

17.1 Lagrange Multipliers and Optimization

17.2 The Jacobian

17.3 Center of Mass

17.4 Fluid Flows

Part V

Appendix

Bibliography

- [1] Stephen Abbott. *Understanding Analysis*. 2nd Edition. Springer, 2015.
- [2] Paul R. Halmos. *Naïve Set Theory*. Springer, 1998.
- [3] Richard Hammack. *Book of Proof*. 3rd Edition. Richard Hammack, 2018.
- [4] Georgi E. Shilov. *Elementary Real and Complex Analysis*. Trans. by Richard A. Silverman. Dover, 1973.

Index

- function, 11
 - argument, 11
 - codomain, 11
 - dependent variable, 11
 - domain, 11
 - graph, 11
 - independent variable, 11
 - range, 11
- set, 9
 - Cartesian product, 10
 - direct product, 10
 - element, 9
 - intersection, 9
 - union, 9