

Calculus

Joseph Le

Contents

I	Preliminaries	7
1	Functions	9
1.1	Sets	9
1.1.1	Exercises	11
1.2	What is a function?	11
1.2.1	Graphs of Functions	11
1.2.2	Table Functions	12
1.2.3	Exercises	13
1.3	Common Functions	13
1.3.1	Linear Functions	13
1.3.2	Quadratic Functions	13
1.3.3	Polynomial and Rational Functions	14
1.3.4	Trigonometric Functions	16
1.3.5	Power, Exponential, and Logarithmic Functions	16
1.3.6	Systems of Equations	16
1.4	Types of Functions	16
1.4.1	Piecewise Functions and Absolute Values	16
1.4.2	Even and Odd Functions	16
1.4.3	Increasing and Decreasing Functions	17
1.4.4	Periodic Functions	17
1.5	Function Transformations and Operations	17
1.5.1	Function Arithmetic	17
1.5.2	Function Translation	17
1.5.3	Function Composition	17
1.5.4	Function Inverses	17
2	Analytic Geometry	19
2.1	Conic Sections	19
2.2	Parametric Equations	19
2.3	Polar Coordinates	19
3	Limits and Continuity	21
3.1	Definition of a Limit	21
3.2	One-Sided Limits and Limits to Infinity	21
3.3	Continuity and the Intermediate Value Theorem	21
3.4	The Squeeze Theorem	21
II	Differential Calculus	23
4	Differentiation and Derivatives	25
4.1	The Limit Definition of the Derivative	25
4.2	Differential Rules	25

4.2.1	The Power Rule	25
4.2.2	The Product Rule	25
4.2.3	The Quotient Rule	25
4.2.4	The Chain Rule	25
4.3	Common and Special Derivatives	25
4.4	Advanced Differential Techniques	25
4.4.1	Implicit Differentiation	25
4.4.2	Logarithmic Differentiation	25
4.4.3	Higher-Order Derivatives	25
5	Applications of Derivatives	27
5.1	Related Rates	27
5.2	Optimization	27
5.3	L'Hôpital's Rule Rule	27
6	Integration and Integrals	29
6.1	Antiderivatives	29
6.2	Riemann Sums	29
6.3	The Fundamental Theorem of Calculus	29
III	Integral Calculus	31
7	Integration Techniques	33
7.1	Substitution	33
7.2	Integration by Parts	33
7.3	Partial Fractions	33
7.4	Trigonometric Substitutions	33
7.5	Hyperbolic Substitutions	33
8	Application of Integrals	35
8.1	Areas Between Curves	35
8.2	Volumes of 3D Shapes	35
8.2.1	The Disk Method	35
8.2.2	The Washer Method	35
8.2.3	The Shell Method	35
8.3	Arc Length	35
9	Improper Integrals and Numerical Integration	37
9.1	Improper Integrals and Their Convergence	37
9.2	Numerical Integration	37
9.2.1	Trapezoidal Rule	37
9.2.2	Simpson's Rule	37
10	Sequences, Series, and Convergence	39
10.1	Sequences	39
10.2	Infinite Series	39
10.2.1	Algebraic Series	39
10.2.2	Geometric Series	39
10.2.3	p-Series	39
10.2.4	Alternating Series	39
10.3	Convergence Tests	39
10.3.1	The Divergence Test	39
10.3.2	The Comparison Test	39
10.3.3	The Limit Comparison Test	39

10.3.4	The Integral Test	39
10.3.5	The Ratio Test	39
10.3.6	The Root Test	39
10.4	Power Series	39
10.5	Taylor Series	39
11	Vectors and Vector Spaces	41
11.1	Vectors in 2D/3D	41
11.2	Vector Products	41
11.2.1	Dot Products	41
11.2.2	Cross Products	41
11.3	Lines and Planes in Space	41
IV	Multivariable Calculus	43
12	Functions of Several Variables	45
12.1	Parametric Curves	45
12.2	Derivatives and Integrals of Vector Functions	45
12.3	Curvature	45
13	Partial Derivatives	47
13.1	Limits and Continuity in Higher Dimensions	47
13.2	Partial Derivatives	47
13.3	The Chain Rule	47
13.4	The Gradient	47
13.5	Extremal Values	47
14	Multiple Integrals	49
14.1	Double Integrals	49
14.2	Coordinate Systems	49
14.2.1	Polar coordinates	49
14.2.2	Cylindrical Coordinates	49
14.2.3	Spherical Coordinates	49
14.3	Surface Area and Volume	49
14.4	Triple Integrals	49
14.5	Change of Variables	49
15	Vector Calculus	51
15.1	Line Integrals	51
15.2	Green's Theorem	51
15.3	Divergence and Curl	51
15.4	Surface Integrals	51
15.5	Stokes' Theorem	51
15.6	The Divergence Theorem	51
16	Applications of Multivariable Calculus	53
16.1	Lagrange Multipliers and Optimization	53
16.2	The Jacobian	53
16.3	Center of Mass	53
16.4	Fluid Flows	53
V	Appendix	55

Part I

Preliminaries

Chapter 1

Functions

1.1 Sets

Before defining what a function is or what it does, it is important to briefly discuss what goes into function and what comes out. Simply, *sets* are a collection of items and each one of those items are usually referred to as *elements*. Without getting into the weeds of set theory, sets can contain pretty much anything from numbers, functions, and other sets [3].

Some common sets that you may be familiar with are the *natural numbers* $\mathbb{N} = \{1, 2, 3, 4, 5, \dots\}$, the *integers* $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$, and the *real numbers* \mathbb{R} , which is usually represented via a number line. Sets can also be intervals on the real line (i.e. $[a, b)$ is an interval on \mathbb{R} containing a but not b) or even the possible results of flipping a coin $C = \{H, T\}$.

We will now define the basic notation when dealing with sets and the operations that can be performed on sets. We say that x is an element of a set A with the notation $x \in A$ and when x is not in A , we say $x \notin A$. For example, given the set $A = \{1, 2, 3, 4\}$, we can say that $1 \in A$ is true as well as $5 \notin A$.

The notion of combining sets comes with *unions* and *intersections*. Given A and B are sets, the union of A and B is denoted as $A \cup B$ and is equal to the set that contains elements in either A or B . Similarly, the intersection between A and B is denoted as $A \cap B$ and is the set that contains elements that are in both A and B . For example, given the sets $A = \{1, 2, 3, 4\}$ and $B = \{3, 4, 5, 6\}$, the union and intersection between A and B is

$$\begin{aligned} A \cup B &= \{1, 2, 3, 4, 5, 6\} \\ A \cap B &= \{3, 4\} \end{aligned} \tag{1.1}$$

Subsets are often used to relate different sets and their elements. A set A is said to be a subset of another set B if all of the elements of A are also within B and is denoted as $A \subseteq B$ and a set A is equal to a set B if and only if $A \subseteq B$ and $B \subseteq A$. An important subset is the empty set represented by the symbols \emptyset , \emptyset , or simply $\{\}$. It is important to note that the empty set is also a subset of all sets.

Using sets by listing them out can become cumbersome and sometimes confusing, instead set builder notation is used to build a set based on a rule. For example, the set of all positive even integers can be written as

$$A = \{2, 4, 6, 8, \dots\} = \{z : z \text{ is an positive even integer}\} = \{z : z = 2n, n \in \mathbb{Z} \text{ and } n > 0\} \tag{1.2}$$

Here, the $:$ stands for "such that" which indicates the rule (the words "such that" or the symbol $|$ is also often used). The *rational numbers* \mathbb{Q} can also be constructed via the integers with


$$\mathbb{Q} = \left\{ \frac{p}{q} : p, q \in \mathbb{Z} \text{ and } q \neq 0 \right\} \tag{1.3}$$

As mentioned previously, intervals on the real number line can be represented as sets. Given two values a and b and assuming that $a \leq b$, intervals on the real line are represented as

- Closed interval: $[a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$



- Open interval: $(a, b) = \{x \in \mathbb{R} : a < x < b\}$



- Half-open interval:

$$- (a, b] = \{x \in \mathbb{R} : a < x \leq b\}$$



$$- [a, b) = \{x \in \mathbb{R} : a \leq x < b\}$$




- Infinite interval:


$$- (a, \infty) = \{x \in \mathbb{R} : a < x\}$$




$$- [a, \infty) = \{x \in \mathbb{R} : a \leq x\}$$




$$- (-\infty, b) = \{x \in \mathbb{R} : x < b\}$$



$$- (-\infty, b] = \{x \in \mathbb{R} : x \leq b\}$$



$$- (-\infty, \infty) = \mathbb{R}$$



Here, the open brackets (and) indicate that the respective endpoint is not included, while the closed brackets [and] indicate that the respective endpoint is included in the interval.

The *Cartesian product* (also known as the *direct product*) is used often to describe ordered pairs or even higher-dimension coordinates. Some examples include the xy -plane also known as \mathbb{R}^2 , 3D space with \mathbb{R}^3 . The Cartesian product for two sets A and B is defined as

$$A \times B = \{(a, b) : a \in A, b \in B\} \quad (1.4)$$

Note that, in general, $A \times B \neq B \times A$ as the operation is order dependent. Higher-order products are defined with sets A_1, A_2, \dots, A_n as

$$A_1 \times A_2 \times \dots \times A_n = \{(a_1, a_2, \dots, a_n) : a_i \in A_i, i \in \{1, 2, \dots, n\}\} \quad (1.5)$$

As previously mentioned, the 2D plane as well as the 3D plane can be constructed using Cartesian products as well and is done as such with the set of real numbers \mathbb{R}

$$\begin{aligned} \mathbb{R}^2 &= \mathbb{R} \times \mathbb{R} = \{(x, y) : x, y \in \mathbb{R}\} \\ \mathbb{R}^3 &= \mathbb{R} \times \mathbb{R} \times \mathbb{R} = \{(x, y, z) : x, y, z \in \mathbb{R}\} \\ &\vdots \\ \mathbb{R}^n &= \mathbb{R} \times \mathbb{R} \times \dots \times \mathbb{R} = \{(x_1, x_2, \dots, x_n) : x_i \in \mathbb{R}, i \in \{1, 2, \dots, n\}\} \end{aligned} \quad (1.6)$$

For further and more formal discussion on sets and naive set theory as well as formal logic and proof writing, refer to [2] or [3].

1.1.1 Exercises

1.2 What is a function?

Functions are objects in math that describe a relationship or mapping between two sets. Given two sets X and Y , a function f maps the unique elements of X , called the *domain* of the function, to elements in the set Y called the *codomain* of the function and the relationship is denoted as $f : X \rightarrow Y$ (" f maps from X to Y ") or $y = f(x)$ (" y equals f of x " [1]), where $y \in R \subseteq Y$ is known as the *dependent variable* with R as the range of the function and $x \in X$ is known as the *independent variable* or the *argument* of the function. The *range* of a function is the set of all possible values that $f(x)$ is able to output with X as its domain, note that the range is a subset of the codomain Y but is not always equal. Formally, the definition is as follows

Definition 1.1. A *function* is a mapping or rule that assigns each element from a set called the domain X of the function to a unique element in the range $R = f(X)$ of the function which is a subset of the codomain $R \subseteq Y$ and the element $x \in X$ is mapped to an element in $y \in R \subseteq Y$ with $x \mapsto y$.

Further emphasis must be made on the elements $y = f(x)$. The function $f(x)$ maps the element $x \in X$ to exactly one element $y \in Y$ [4]. For single variable functions, this can be tested using the vertical line test on a function's graph (covered shortly). If a formula results in two answers with one input (i.e. the equation for a circle: $x^2 + y^2 = r$ gives two points of y for each x) it is no longer considered a function. Figure 1.1 shows how a function maps elements of the domain to the range.




Figure 1.1: Function mapping diagram showing the domain X , codomain Y , and range $f(X)$

Example 1.2. Some real-world examples of functions are


- The area of a circle: $A(r) = \pi r^2$
- The height of a falling ball: $h(t) = h_0 + v_0 t - (1/2)gt^2$
- Compound interest: $A(t) = P(1 + \frac{r}{n})^{nt}$
- Temperature conversion: $C(F) = \frac{5}{9}(F - 32)$
-

It can be useful to think of functions as machines that take in an input and produces an output. The examples above show such cases using mathematical formulas that give a set of instructions on how the input is transformed from the element in the domain to the element in the range.


1.2.1 Graphs of Functions

Another way to represent functions is with its *graph* which is a set of points consisting of the input and its corresponding output and is represented with the set of ordered pairs $\{(x, y) : y = f(x), x \in \mathbb{R}\}$ for functions that accept real numbers.

Example 1.3. A familiar example is the linear function $f(x) = mx + b$ where m is the slope of the line and b is the vertical offset. Suppose $m = 1$ and $b = 0$, we get the following graph for the function $f(x) = x$ in Figure 1.3

Figure 1.2: Graph of the function $f(x) = x$

Example 1.4. Another example is the quadratic function of the form $f(x) = ax^2 + bx + c$, where $a, b, c \in \mathbb{R}$ are the coefficients of the function and determine its shape. Suppose that $a = 1, b = 0, c = -1$ resulting in the function $f(x) = x^2 - 1$. Using knowledge of the roots a quadratic function, it can be deduced that this will result in a parabola with x -intersections at $x = -1$ and $x = 1$ which can be confirmed in the graph in Figure 1.4

Figure 1.3: Graph of the function $f(x) = x^2 - 1$

1.2.2 Table Functions

The last representation of a function that will be discussed are functions that are in the form of tables. Consider two finite sets $X = \{1, 2, 3, 4, 5\}$ and $Y = \{2, 4, 6, 8, 10\}$ and a function that maps the elements of X to the elements of Y . A table can be created that shows the assignment of the elements (Table 1.1) via the function $f(x)$.

The function shown in Table 1.1 is constructed from a mathematical formula ($y = 2x$), this is not necessary, as long as there is only one output for any given input, then, the table is considered a function.

Example 1.5. For example, consider the following table that shows the function that maps $X = \{1, 3, 5, 8, 4\}$ to $Y = \{3, 5, 2, 1, 1\}$ in Table 1.2. Though a value in Y repeats once, it still passes the vertical line test as the function maps only one output y from any input x .

$x \in X$	1	2	3	4	5
$y \in Y$	2	4	6	8	10

Table 1.1: Function Mapping from X to Y

$x \in X$	1	3	5	8	4
$y \in Y$	3	5	2	1	1

Table 1.2: Function Mapping from X to Y

This conceptual foundation of what a function is prepares us for the introduction and review of commonly used functions and function families (functions with similar forms and properties).

1.2.3 Exercises

1.3 Common Functions

Functions, namely those defined via mathematical formulas, can be classified into function families whose members share common features, forms, and properties. Some such families often differ only in coefficients and others may change the base in which they operate. The following types of functions are used frequently in the study of calculus.

1.3.1 Linear Functions

Beginning with the most elementary form of functions created using mathematical formulas is the *linear function* or the equation for a line.

Definition 1.6. Functions of the form $f(x) = mx + b$, where m is the slope and b is the y -intercept, are *linear functions*.

The graphs of linear functions are straight lines, if the domain of the function is \mathbb{R} , then the graph of f extends from $-\infty$ to $+\infty$. Its slope determines the angle of the line and can be reconstructed with any two points on the line with the following formula

$$m = \frac{y_2 - y_1}{x_2 - x_1} \quad (1.7)$$

where y_i corresponds to x_i and $x_2 > x_1$. The y -intercept the value of the y -coordinate of the function when $x = 0$ and denotes where the line crosses the y -axis on the graph. An example for the graph of a linear function is shown in Figure 1.3 where $f(x) = x$.

Example 1.7. Linear functions can model the conversion between temperatures in degrees Celsius and Fahrenheit with the function $C(F) = \frac{5}{9}(F - 32) = \frac{5}{9}F - \frac{160}{9}$. Here the independent variable is F , the slope is $\frac{5}{9}$, and the y -intercept is $\frac{160}{9}$ and the graph is shown in Figure 1.7.

1.3.2 Quadratic Functions

Quadratic functions are the next natural progression after linear functions. The simplest form of a quadratic function $f(x) = x^2$ consists of x multiplied by itself. The graph of a quadratic equation is a *parabola* and can be seen in Figure 1.4.

Definition 1.8. Functions of the form $f(x) = ax^2 + bx + c$ are *quadratic functions*, this is the standard form for a quadratic function. Alternative forms to the standard quadratic function are the root form




Figure 1.4: Graph of the Fahrenheit to Celsius Conversion, $C(F) = \frac{5}{9}(F - 32)$

$f(x) = (x-a)(x-b)$ where a and b are the x -intercepts of the function and the vertex form $f(x) = a(x-h)^2 + k$ where the point h, k is the vertex of the parabola.

Quadratic functions have some useful properties such as the root(s) of the parabola (the x -intercept(s)) as well as having an explicit formula for computing the root(s) of the function known as the *quadratic formula*. The quadratic formula solves for the x -coordinate(s) of the roots for a given quadratic function (where $y = 0$)

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad (1.8)$$

where a , b , and c are the coefficients in the standard form of a quadratic function. This formula gives real roots when the *discriminant* $b^2 - 4ac > 0$, a single real root (with a multiplicity of two) when $b^2 - 4ac = 0$, complex roots when $b^2 - 4ac < 0$, and purely imaginary roots when $b = 0$ and $4ac > 0$.

Example 1.9. An example of a quadratic function is *vertical projectile motion* under gravity: Given an object with initial height h_0 and initial vertical velocity v_0 , its height at time t is modeled by

$$h(t) = -\frac{1}{2}gt^2 + v_0t + h_0 \quad (1.9)$$

where $h(t)$ is the height at time t , g is the acceleration due to gravity ($g \approx 9.81 \text{ m/s}^2$) at Earth's surface, and air resistance is neglected. The graph showing the height of a particle thrown from a height of 20 meters and at an initial vertical velocity of 2 m/s² is shown in Figure 1.9.

1.3.3 Polynomial and Rational Functions

Generalizing upon the linear and quadratic functions further, polynomial functions are the next step.

Definition 1.10. Functions of the form $f(x) = a_nx^n + a_{n-1}x^{n-1} + \cdots + a_2x^2 + a_1x + a_0$ are *polynomial functions*. Where n is the *degree* of the polynomial (where $a_n \neq 0$) and a_i for $i \in 0, 1, \dots, n-1$, n are the coefficients.

Similar to quadratic functions, polynomials have up to n roots but do not have an explicit formula to compute those roots for polynomials with degree $n > 3$. Linear functions and quadratic functions can be considered special cases of polynomial functions with degree $n = 1$ and $n = 2$ respectively. Common higher-order polynomials are *cubic functions* $n = 3$, *quartic functions* $n = 4$, and *quintic functions* $n = 5$. A useful property of polynomials is that there are at most n roots for a polynomial of degree n , note that when a polynomial has less (real and complex) roots than its degree, one of its roots may have a multiplicity greater than one meaning that it is repeated. For example, the polynomial (in root form)




Figure 1.5: Graph of a Particle Falling Due to Gravity over time, $h(t) = -\frac{1}{2}gt^2 + v_0t + h_0$

$f(x) = (x+1)(x+1)(x+1) = (x+1)^3 = x^3 + 3x^2 + 3x + 1$ has all three roots at $x = -1$ which means that the root $(x = -1, y = 0)$ has multiplicity of 3.

Example 1.11. To show the differences between polynomials with varying order, Figure 1.6 graphs polynomials of the form x^n up to the 6th degree.




Figure 1.6: Polynomials of varying degrees $n \in \{1, 2, 3, 4, 5, 6\}$

Rational functions branch from polynomials in a similar way to how rational numbers branch from the integers as they are constructed by dividing one polynomial by another.

Definition 1.12. The function of the form $f(x) = \frac{p(x)}{q(x)}$ where $p(x)$ and $q(x) \neq 0$ are polynomial functions are known as *rational functions*.

Like polynomials, rational functions have *zeros* or roots when the numerator is equal to zero, but also zeros when the denominator of the function is equal to zero called *poles*. These describe discontinuous points as the the denominators reach zero resulting in holes (when a zero and a pole cancel) and an *asymptote* when the pole is not cancelled.

Example 1.13. An example of a rational function that has both holes and asymptotes is

$$f(x) = \frac{(x-1)(x+1)(x+2)}{(x-1)(x+1)(x-2)} \quad (1.10)$$

This function has holes at $x = \pm 1$ where the poles and zeros cancel out asymptotes at $x = 2$ where the denominator becomes zero and $y = 1$ for the horizontal asymptote. The graph for this function is shown in Figure 1.7.




Figure 1.7: Rational function $f(x) = \frac{(x-1)(x+1)(x+2)}{(x-1)(x+1)(x-2)}$ with holes at $(1, -3)$ and $(-1, -\frac{1}{3})$, vertical asymptote at $x = 2$, and horizontal asymptote at $y = 1$

1.3.4 Trigonometric Functions

Breaking away from the progression found from starting with linear functions and ending with polynomials

1.3.5 Power, Exponential, and Logarithmic Functions

1.3.6 Systems of Equations

1.4 Types of Functions

It is important to be able to distinguish functions and classify their behaviors into distinct characteristics in order to properly analyze them. Characteristics such as symmetry (evenness and oddness of a function) and periodicity (repeating functions) are defined in such ways that may give insight into solving certain problems or giving us the opportunity to exploit their definitions.

1.4.1 Piecewise Functions and Absolute Values

1.4.2 Even and Odd Functions

Definition 1.14. A function $f(x)$ is said to be an *even function* of the independent variable x if and only if

$$f(-x) = f(x)$$

for all x in the function's domain.

Definition 1.15. A function $f(x)$ is said to be an *odd function* of the independent variable x if and only if

$$f(-x) = -f(x)$$

for all x in the function's domain.

1.4.3 Increasing and Decreasing Functions

Definition 1.16. A function f is *increasing* on an interval I if for all $x_1, x_2 \in I$ with $x_1 < x_2$,

$$f(x_1) \leq f(x_2).$$

If the inequality is strict ($f(x_1) < f(x_2)$), we say f is *strictly increasing*.

Definition 1.17. A function f is *decreasing* on an interval I if for all $x_1, x_2 \in I$ with $x_1 < x_2$,

$$f(x_1) \geq f(x_2).$$

If the inequality is strict ($f(x_1) > f(x_2)$), we say f is *strictly decreasing*.

1.4.4 Periodic Functions

Definition 1.18. A function is said to be a *period function* if and only if for some value a ,

$$f(x) = f(x + a)$$

where a is the period of the function. The period defines the magnitude of the independent variable required for the function to repeat.

1.5 Function Transformations and Operations

1.5.1 Function Arithmetic

1.5.2 Function Translation

1.5.3 Function Composition

1.5.4 Function Inverses

Chapter 2

Analytic Geometry

2.1 Conic Sections

2.2 Parametric Equations

2.3 Polar Coordinates

Chapter 3

Limits and Continuity

3.1 Definition of a Limit

3.2 One-Sided Limits and Limits to Infinity

3.3 Continuity and the Intermediate Value Theorem

3.4 The Squeeze Theorem

Part II

Differential Calculus

Chapter 4

Differentiation and Derivatives

4.1 The Limit Definition of the Derivative

4.2 Differential Rules

4.2.1 The Power Rule

4.2.2 The Product Rule

4.2.3 The Quotient Rule

4.2.4 The Chain Rule

4.3 Common and Special Derivatives

4.4 Advanced Differential Techniques

4.4.1 Implicit Differentiation

4.4.2 Logarithmic Differentiation

4.4.3 Higher-Order Derivatives

Chapter 5

Applications of Derivatives

5.1 Related Rates

5.2 Optimization

5.3 L'Hôpital's Rule Rule

Chapter 6

Integration and Integrals

6.1 Antiderivatives

6.2 Riemann Sums

6.3 The Fundamental Theorem of Calculus

Part III

Integral Calculus

Chapter 7

Integration Techniques

7.1 Substitution

7.2 Integration by Parts

7.3 Partial Fractions

7.4 Trigonometric Substitutions

7.5 Hyperbolic Substitutions

Chapter 8

Application of Integrals

8.1 Areas Between Curves

8.2 Volumes of 3D Shapes

8.2.1 The Disk Method

8.2.2 The Washer Method

8.2.3 The Shell Method

8.3 Arc Length

Chapter 9

Improper Integrals and Numerical Integration

9.1 Improper Integrals and Their Convergence

9.2 Numerical Integration

9.2.1 Trapezoidal Rule

9.2.2 Simpson's Rule

Chapter 10

Sequences, Series, and Convergence

10.1 Sequences

10.2 Infinite Series

10.2.1 Algebraic Series

10.2.2 Geometric Series

10.2.3 p-Series

10.2.4 Alternating Series

10.3 Convergence Tests

10.3.1 The Divergence Test

10.3.2 The Comparison Test

10.3.3 The Limit Comparison Test

10.3.4 The Integral Test

10.3.5 The Ratio Test

10.3.6 The Root Test

10.4 Power Series

10.5 Taylor Series

Chapter 11

Vectors and Vector Spaces

11.1 Vectors in 2D/3D

11.2 Vector Products

11.2.1 Dot Products

11.2.2 Cross Products

11.3 Lines and Planes in Space

Part IV

Multivariable Calculus

Chapter 12

Functions of Several Variables

12.1 Parametric Curves

12.2 Derivatives and Integrals of Vector Functions

12.3 Curvature

Chapter 13

Partial Derivatives

13.1 Limits and Continuity in Higher Dimensions

13.2 Partial Derivatives

13.3 The Chain Rule

13.4 The Gradient

13.5 Extremal Values

Chapter 14

Multiple Integrals

14.1 Double Integrals

14.2 Coordinate Systems

14.2.1 Polar coordinates

14.2.2 Cylindrical Coordinates

14.2.3 Spherical Coordinates

14.3 Surface Area and Volume

14.4 Triple Integrals

14.5 Change of Variables

Chapter 15

Vector Calculus

15.1 Line Integrals

15.2 Green's Theorem

15.3 Divergence and Curl

15.4 Surface Integrals

15.5 Stokes' Theorem

15.6 The Divergence Theorem

Chapter 16

Applications of Multivariable Calculus

16.1 Lagrange Multipliers and Optimization

16.2 The Jacobian

16.3 Center of Mass

16.4 Fluid Flows

Part V

Appendix

Bibliography

- [1] Stephen Abbott. *Understanding Analysis*. 2nd Edition. Springer, 2015.
- [2] Paul R. Halmos. *Naïve Set Theory*. Springer, 1998.
- [3] Richard Hammack. *Book of Proof*. 3rd Edition. Richard Hammack, 2018.
- [4] Georgi E. Shilov. *Elementary Real and Complex Analysis*. Trans. by Richard A. Silverman. Dover, 1973.

Index

- function, 11
 - argument, 11
 - codomain, 11
 - decreasing, 17
 - dependent variable, 11
 - domain, 11
 - even, 16
 - graph, 11
 - increasing, 17
 - independent variable, 11
 - linear, 13
 - odd, 16
 - periodic, 17
 - polynomial, 14
 - cubic, 14
 - degree, 14
 - quartic, 14
 - quintic, 14
 - quadratic, 13
 - discriminant, 14
 - formula, 14
 - parabola, 13
 - range, 11
 - rational, 15
 - asymptote, 15
 - poles, 15
 - zeros, 15
 - strictly decreasing, 17
 - strictly increasing, 17
- set, 9
 - Cartesian product, 10
 - direct product, 10
 - element, 9
 - intersection, 9
 - union, 9