

# Math Homework

José Porcar

Semester 2 2025

## Contents

1	Riemann and Trapezoidal sums	2
2	Volume of Solids of Revolutions (Disk Method)	4
3	Volume of Solids of Revolutions (Washer Method)	6
4	Separation of Variables	7

# 1 Riemann and Trapezoidal sums

## Left Riemann Sums

$f(x) = x + 6$ ;  $[1, 5]$ ; 4 rectangles

$x$	1	2	3	4	5
$y$	7	8	9	10	11

$$\Delta x = \frac{|a - b|}{r}$$

$$\Delta x = \frac{|1 - 5|}{4}$$

$$\Delta x = 1$$

$$A = \sum_{n=a}^{b-1} f(n) \Delta x$$

$$A = \sum_{n=1}^4 f(n)$$

$$A = 7 + 8 + 9 + 10$$

$$A = 34$$

$f(x) = x + 4$ ;  $[-2, 2]$ ; 4 rectangles

$x$	-2	-1	0	1	2
$y$	2	3	4	5	6

$$\Delta x = \frac{|a - b|}{r}$$

$$\Delta x = \frac{|-2 - 2|}{4}$$

$$\Delta x = 1$$

$$A = \sum_{n=a}^{b-1} f(n) \Delta x$$

$$A = \sum_{n=-2}^1 f(n)$$

$$A = 2 + 3 + 4 + 5$$

$$A = 14$$

## Right Riemann Sums

$f(x) = -x^2 - 2x + 9$ ;  $[-3, 2]$ ; 5 rectangles

$x$	-3	-2	-1	0	1	2
$y$	6	9	10	9	6	1

$$\Delta x = \frac{|a - b|}{r}$$

$$\Delta x = \frac{|-3 - 2|}{5}$$

$$\Delta x = 1$$

$$A = \sum_{n=a+1}^b f(n) \Delta x$$

$$A = \sum_{n=-2}^2 f(n)$$

$$A = 9 + 10 + 9 + 6 + 1$$

$$A = 35$$

$f(x) = \frac{2}{x}$ ;  $[2, 7]$ ; 5 rectangles

$x$	2	3	4	5	6	7
$y$	1	$\frac{2}{3}$	$\frac{1}{2}$	$\frac{2}{5}$	$\frac{1}{3}$	$\frac{2}{7}$

$$\Delta x = \frac{|a - b|}{r}$$

$$\Delta x = \frac{|2 - 7|}{5}$$

$$\Delta x = 1$$

$$A = \sum_{n=a+1}^b f(n) \Delta x$$

$$A = \sum_{n=3}^7 f(n)$$

$$A = \frac{2}{3} + \frac{1}{2} + \frac{2}{5} + \frac{1}{3} + \frac{2}{7}$$

$$A = 2.19$$

## Trapezoidal Sums

$$y = -\frac{x^2}{2} + x + 5; [0, 3]; 5 \text{ Trapezoids}$$

$x$	0	0.6	1.2	1.8	2.4	3
$y$	5	5.42	5.48	5.18	4.52	3.5

$$\Delta x = \frac{|a - b|}{r}$$

$$\Delta x = \frac{|0 - 3|}{5}$$

$$\Delta x = 0.6$$

$$A = \sum_{n=1}^{rectangles} \frac{f(n\Delta x) + f((n-1)\Delta x)}{2} \Delta x$$

$$A = \sum_{n=1}^5 \frac{f(0.6n) + f(0.6(n-1))}{2} 0.6$$

$$A = 3.13 + 3.27 + 3.23 + 2.91 + 2.41$$

$$A = 14.95$$

$$f(x) = \frac{x^2}{2} + x + 1; [-2, 1]; 5 \text{ Trapezoids}$$

$x$	-2	-1.4	-0.8	-0.2	0.4	1
$y$	1	0.58	0.52	0.82	1.48	2.5

$$\Delta x = \frac{|a - b|}{r}$$

$$\Delta x = \frac{|-2 - 1|}{5}$$

$$\Delta x = 0.6$$

$$A = \sum_{n=1}^{rectangles} \frac{f(n\Delta x) + f((n-1)\Delta x)}{2} \Delta x$$

$$A = \sum_{n=1}^5 \frac{f(0.6n) + f(0.6(n-1))}{2} 0.6$$

$$A = 0.474 + 0.33 + 0.402 + 0.69 + 1.194$$

$$A = 3.09$$

## 2 Volume of Solids of Revolutions (Disk Method)

3. Find the volume of the solid of revolution generated by revolving the region bounded by  $y = \sqrt{1 - x^2}$  and  $y = 0$  about the  $x$ -axis.

$$\begin{aligned} V &= \pi \int_b^a [r^2] dx \\ &= \pi \int_{-1}^1 \left[ \left( \sqrt{1 - x^2} \right)^2 \right] dx \\ &= \pi \int_{-1}^1 [1 - x^2] dx \\ &= \pi \left[ x - \frac{x^3}{3} \right]_{-1}^1 \\ &= \pi \left( 1 - \frac{1}{3} - \left( -1 - \frac{-1}{3} \right) \right) \\ V &= \frac{4\pi}{3} \end{aligned}$$

8. Find the volume of the solid formed by revolving the region bounded by the graphs of  $y = \sqrt{\sin x}$  and the  $x$ -axis ( $0 \leq x \leq \pi$ ), about the  $x$ -axis.

$$\begin{aligned} V &= \pi \int_0^\pi (\sqrt{\sin x})^2 dx \\ &= \pi \int_0^\pi \sin x dx \\ &= \pi [-\cos x]_0^\pi \\ &= \pi (-(-1) + 1) \\ V &= 2\pi \end{aligned}$$

11. The region bounded by the parabola  $y = 4x - x^2$  and the  $x$ -axis is revolved about the  $x$ -axis. Find the volume of the solid.

$$\begin{aligned} \{a, b\} &= x \quad \text{where } 4x - x^2 = 0 \\ a &= 0; \quad b = 4 \\ V &= \pi \int_0^4 (4x - x^2)^2 dx \\ &= \pi \int_0^4 (16x^2 - 8x^3 + x^4) dx \\ &= \pi \left[ \frac{16}{3}x^3 - 2x^4 + \frac{1}{5}x^5 \right]_0^4 \\ &= \pi \left( \left( \frac{1024}{3} - 512 + \frac{1024}{5} \right) - 0 \right) \\ V &= \frac{512\pi}{15} \end{aligned}$$

- 13a. Find the volume of the solid generated when the area bounded by the curves  $y = x^3 - x + 1$ ,  $x = -1$ , and  $x = 1$  is revolved around the  $x$ -axis.

$$\begin{aligned}
 V &= \pi \int_{-1}^1 (x^3 - x + 1)^2 dx \\
 &= \pi \int_{-1}^1 (x^6 - 2x^4 + 2x^3 + x^2 - 2x + 1) dx \\
 &= \pi \left[ \frac{1}{7}x^7 - \frac{2}{5}x^5 + \frac{1}{2}x^4 + \frac{1}{3}x^3 - x^2 + x \right]_{-1}^1 \\
 &= \pi \left( \left( \frac{1}{7} - \frac{2}{5} + \frac{1}{2} + \frac{1}{3} \right) - \left( -\frac{1}{7} + \frac{2}{5} + \frac{1}{2} - \frac{1}{3} - 2 \right) \right) \\
 V &= \frac{226\pi}{105} \text{ source: deepseek I aint doing them fractions :b}
 \end{aligned}$$

- 13b. Find the volume of the solid generated when the area bounded by the curve  $y = x - x^2$  and the  $x$ -axis is revolved around the  $x$ -axis.

$$\begin{aligned}
 V &= \pi \int_0^1 (x - x^2)^2 dx \\
 &= \pi \int_0^1 (x^2 - 2x^3 + x^4) dx \\
 &= \pi \left[ \frac{1}{3}x^3 - \frac{1}{2}x^4 + \frac{1}{5}x^5 \right]_0^1 \\
 &= \pi \left( \frac{1}{3} - \frac{1}{2} + \frac{1}{5} \right) \\
 V &= \frac{\pi}{30}
 \end{aligned}$$

### 3 Volume of Solids of Revolutions (Washer Method)

Find the volume of the solid that results when the region enclosed by the curves is revolved about the given axis.

$$f(x) = x^2 - 3, \quad g(x) = \sqrt{x} - 3; \quad \text{axis: } y = 2$$

$$\begin{aligned} f(x) &= g(x) \\ x^2 - 3 &= \sqrt{x} - 3 \end{aligned}$$

$$x^2 = \sqrt{x}$$

$$x^4 = x$$

$$x(x^3 - 1) = 0$$

$$x = 0 \quad x^3 = 1$$

$$a = 0 \quad b = 1$$

$$\begin{aligned} V &= \pi \int_a^b \left| (f(x) - y)^2 - (g(x) - y)^2 \right| dx \\ &= \pi \int_0^1 \left| (x^2 - 5)^2 - (\sqrt{x} - 5)^2 \right| dx \\ &= \pi \int_0^1 \left| x^4 - 10x^2 + 25 - (x - 10\sqrt{x} + 25) \right| dx \\ &= \pi \left| \frac{x^5}{5} - \frac{10x^3}{3} - \frac{x^2}{2} + \frac{20x^{\frac{3}{2}}}{3} \right|_0^1 \\ &= \pi \left| \frac{1}{5} - \frac{10}{3} - \frac{1}{2} + \frac{20}{3} \right| \\ V &= \frac{91\pi}{30} \end{aligned}$$

$$f(x) = x^2 - 3, \quad g(x) = \sqrt{x} - 3; \quad \text{axis: } y = 1$$

$$\begin{aligned} f(x) &= g(x) \\ x^2 - 3 &= \sqrt{x} - 3 \end{aligned}$$

$$x^2 = \sqrt{x}$$

$$x^4 = x$$

$$x(x^3 - 1) = 0$$

$$x = 0 \quad x^3 = 1$$

$$a = 0 \quad b = 1$$

$$\begin{aligned} V &= \pi \int_a^b \left| (f(x) - y)^2 - (g(x) - y)^2 \right| dx \\ &= \pi \int_0^1 \left| (x^2 - 4)^2 - (\sqrt{x} - 4)^2 \right| dx \\ &= \pi \int_0^1 \left| x^4 - 8x^2 + 16 - (x - 8\sqrt{x} + 16) \right| dx \\ &= \pi \left| \frac{x^5}{5} - \frac{8x^3}{3} - \frac{x^2}{2} + \frac{16x^{\frac{3}{2}}}{3} \right|_0^1 \\ &= \pi \left| \frac{1}{5} - \frac{8}{3} - \frac{1}{2} + \frac{16}{3} \right| \\ V &= \frac{71\pi}{30} \end{aligned}$$

$$f(x) = x^2 + 4, \quad g(x) = 2; \quad a = 0, \quad b = 1; \\ \text{axis: } y = -1$$

$$\begin{aligned} V &= \pi \int_a^b \left| (f(x) - y)^2 - (g(x) - y)^2 \right| dx \\ &= \pi \int_0^1 \left| (x^2 + 5)^2 - 3^2 \right| dx \\ &= \pi \int_0^1 \left| x^4 + 10x^2 + 16 \right| dx \\ &= \pi \left| \frac{x^5}{5} + \frac{10x^3}{3} + 16x \right|_0^1 \\ &= \pi \left| \frac{1}{5} + \frac{10}{3} + 16 \right| \\ V &= \frac{293\pi}{15} \end{aligned}$$

$$f(x) = x^2 + 3, \quad g(x) = 3; \quad a = 0, \quad b = 2; \\ \text{axis: } y = 1$$

$$\begin{aligned} V &= \pi \int_a^b \left| (f(x) - y)^2 - (g(x) - y)^2 \right| dx \\ &= \pi \int_0^2 \left| (x^2 + 2)^2 - 2^2 \right| dx \\ &= \pi \int_0^2 \left| x^4 + 4x^2 \right| dx \\ &= \pi \left| \frac{x^5}{5} + \frac{4x^3}{3} \right|_0^2 \\ &= \pi \left| \frac{32}{5} + \frac{32}{3} \right| \\ V &= \frac{256\pi}{15} \end{aligned}$$

## 4 Separation of Variables

### General Solution

1.	2.	3.	4.
$\frac{dy}{dx} = \frac{x^3}{y^2}$	$\frac{dy}{dx} = \frac{1}{\sec^2 y}$	$\frac{dy}{dx} = 3e^{x-y}$	$\frac{dy}{dx} = \frac{2x}{e^{2y}}$
$y^2 dy = x^3 dx$	$\sec^2 y dy = dx$	$e^y dy = 3e^x dx$	$e^{2y} dy = 2x dx$
$\int y^2 dy = \int x^3 dx$	$\int \sec^2 y dy = \int dx$	$\int e^y dy = \int 3e^x dx$	$\int e^{2y} dy = \int 2x dx$
$\frac{y^3}{3} = \frac{x^4}{4} + C$	$\tan y = x + C$	$e^y = 3e^x + C$	$\frac{e^{2y}}{2} = x^2 + C$
$y^3 = \frac{3x^4}{4} + 3C$	$y = \tan^{-1}(x + C)$	$y = \ln(3e^x + C)$	$e^{2y} = 2(x^2 + C)$
$y = \sqrt[3]{\frac{3x^4}{4} + 3C}$			$2y = \ln 2 + \ln(x^2 + C)$
			$y = \ln(\sqrt{x^2 + C}) + \frac{\ln 2}{2}$