

Tratamiento de Señales


Version 2022-2


Convolución 1D

[Capítulo 4]


Dr. José Ramón Iglesias

DSP-ASIC BUILDER GROUP Director Semillero TRIAC Ingenieria Electronica Universidad Popular del Cesar






$$y(t) = \int_{-\infty}^{+\infty} g(t - \tau) x(\tau) d\tau$$




$$y(t) = \int_{-\infty}^{+\infty} g(t - \tau) x(\tau) d\tau$$


$$y(t) = x(t) * g(t)$$


$$y(t) = \int_{-\infty}^{+\infty} g(t - \tau) x(\tau) d\tau$$






$$y(t) = \int_{-\infty}^{+\infty} g(t - \tau) x(\tau) d\tau$$







$$y(t) = \int_{-\infty}^{+\infty} g(t - \tau) x(\tau) d\tau$$




$$y(t) = \int_{-\infty}^{+\infty} g(t - \tau) x(\tau) d\tau$$




$$y(t) = \int_{-\infty}^{+\infty} g(t - \tau) x(\tau) d\tau$$




$$y(t) = \int_{-\infty}^{+\infty} g(t - \tau) x(\tau) d\tau$$




$$y(t) = \int_{-\infty}^{+\infty} g(t - \tau) x(\tau) d\tau$$




$$y(t) = \int_{-\infty}^{+\infty} g(t - \tau) x(\tau) d\tau$$




$$y(t) = \int_{-\infty}^{+\infty} g(t - \tau) x(\tau) d\tau$$




$$y(t) = \int_{-\infty}^{+\infty} g(t - \tau) x(\tau) d\tau$$




$$y(t) = \int_{-\infty}^{+\infty} g(t - \tau) x(\tau) d\tau$$




$$y(t) = \int_{-\infty}^{+\infty} g(t - \tau) x(\tau) d\tau$$




$$y(t) = \int_{-\infty}^{+\infty} g(t - \tau) x(\tau) d\tau$$



$$y(t) = \int_{-\infty}^{+\infty} g(t - \tau) x(\tau) d\tau$$





Continua:
$$y(t) = \int_{-\infty}^{+\infty} g(t-\tau)x(\tau)d\tau$$

Discreta:
$$y(k) = \sum_{i=-\infty}^{+\infty} g(k-i)x(i)$$