

Silhouette plot of clara(x = tenencia3[, -c(1:2)], k = n, metric = "euclidean", stand = FALSE, samples = 100, sampsize = 30

n = 300

2 clusters C_j

$j : n_j \mid \text{ave}_{i \in C_j} s_i$

1 : 32 | 0.60

2 : 268 | 1.00



0.0 0.2 0.4 0.6 0.8 1.0

Silhouette width s_i

Average silhouette width : 0.95

Silhouette plot of clara(x = tenencia3[, -c(1:2)], k = n, metric = "euclidean", stand = FALSE, samples = 100, sampsize = 300)

n = 300



Average silhouette width : 1

Silhouette plot of clara(x = tenencia3[, -c(1:2)], k = n, metric = "euclidean", stand = FALSE, samples = 100, sampsize = 300

n = 300



Average silhouette width : 0.99

Silhouette plot of clara(x = tenencia3[, -c(1:2)], k = n, metric = "euclidean", stand = FALSE, samples = 100, sampsize = 300

n = 300



Average silhouette width : 0.98

Silhouette plot of clara(x = tenencia3[, -c(1:2)], k = n, metric = "euclidean", stand = FALSE, samples = 100, sampsize = 300

n = 300



Average silhouette width : 0.99

Silhouette plot of clara(x = tenencia3[, -c(1:2)], k = n, metric = "euclidean", stand = FALSE, samples = 100, sampsize = 30

n = 300



Average silhouette width : 0.98

Silhouette plot of clara(x = tenencia3[, -c(1:2)], k = n, metric = "euclidean", stand = FALSE, samples = 100, sampsize = 30

n = 300



Average silhouette width : 0.99

Silhouette plot of clara(x = tenencia3[, -c(1:2)], k = n, metric = "euclidean", stand = FALSE, samples = 100, sampsize = 30

n = 300



Average silhouette width : 0.98

Silhouette plot of clara(x = tenencia3[, -c(1:2)], k = n, metric = "euclidean", stand = FALSE, samples = 100, sampsize = 30

n = 300



Average silhouette width : 0.98