

Efficacy and Safety of Oral Ziltravir in High-Risk Outpatients with COVID-19: Results from a Phase 3 Randomized Controlled Trial

Julian Smith, Amelia Doe, Robert Johnson

ABSTRACT

Background: We evaluated the efficacy and safety of oral ziltravir in non-hospitalized adults with early-stage COVID-19 to determine its impact on disease progression.

Methods: In this phase 3, double-blind trial, 1,042 unvaccinated adults with mild-to-moderate COVID-19 and at least one risk factor for progression were randomized 1:1 to receive a 5-day course of ziltravir (800 mg twice daily) or placebo. The primary endpoint was COVID-19–related hospitalization or all-cause mortality through day 28.

Results: Hospitalization or death occurred in 15 of 521 (2.9%) ziltravir recipients compared with 38 of 521 (7.3%) placebo recipients (relative risk reduction, 60.3%; 95% CI, 25.1 to 80.2; $P = 0.002$). No deaths occurred in the ziltravir group, while two occurred in the placebo group. Adverse events were comparable between groups (18.2% ziltravir vs. 20.5% placebo), consisting mostly of mild gastrointestinal symptoms.

Conclusion: A 5-day course of oral ziltravir safely and significantly reduced the risk of hospitalization or death in high-risk outpatients with mild-to-moderate COVID-19.

Trial Context

Underlying phase 3, double-blind randomized controlled trial evaluating an oral antiviral (ziltravir) versus placebo in high-risk outpatients with mild-to-moderate COVID-19.