

**Using a qualitative model to explore impacts of ecosystem  
and anthropogenic drivers upon declining marine survival in  
Pacific salmon**

Journal:	<i>Environmental Conservation</i>
Manuscript ID	EC-17-05-111.R1
Manuscript Type:	Research Paper
Date Submitted by the Author:	n/a
Complete List of Authors:	Sobocinski, Kathryn; NOAA Northwest Fisheries Science Center, Fish Ecology; Long Live the Kings Greene, Correigh; NOAA Northwest Fisheries Science Center, Fish Ecology Schmidt, Michael; Long Live the Kings
Keywords:	Network model, Indicators, Ecosystem, Salmon, Qualitative

SCHOLARONE™  
Manuscripts

Review

# Using a qualitative model to explore impacts of ecosystem and anthropogenic drivers upon declining marine survival in Pacific salmon

Kathryn L. Sobocinski<sup>\*,1,2</sup>, Correigh M. Greene<sup>1</sup>, Michael W. Schmidt<sup>2</sup>

<sup>1</sup>Northwest Fisheries Science Center  
National Marine Fisheries Service  
National Oceanic and Atmospheric Administration  
2725 Montlake Blvd. E  
Seattle, WA 98112

<sup>2</sup>Long Live the Kings  
1326 5th Ave. #450  
Seattle, WA 98101

\*Corresponding Author:  
kathryn.sobocinski@noaa.gov  
(206) 302-2466

Running Head: Salish Sea salmon qualitative network model

Word Count: 5,901, revised: 6,458

Keywords: Network model, indicators, ecosystem, salmon, qualitative

Submitted: May 12, 2017

Revised: August 19, 2017

## Summary

Coho salmon (*Oncorhynchus kisutch*), Chinook salmon (*O. tshawytscha*), and steelhead (*O. mykiss*) in Puget Sound and the Strait of Georgia have exhibited declines in marine survival over the last 40 years. While the cause of these declines is unknown, multiple factors, acting cumulatively or synergistically, have likely contributed. To evaluate the potential contribution of a broad suite of drivers on salmon survival, we used qualitative network modeling (QNM). QNM is a conceptually based tool that uses networks with specified relationships among the variables. In a simulation framework, linkages are weighted and then the models are subjected to user-specified perturbations. Our network had 33 variables, including: environmental and oceanographic drivers (e.g., temperature, precipitation), primary production variables, food web components from zooplankton to predators, and anthropogenic impacts (e.g., habitat loss, hatcheries). We included salmon traits (survival, abundance, residence time, fitness, and size) as response variables. We invoked perturbations to each node and to suites of drivers and evaluated the responses of these variables. The model showed that anthropogenic impacts resulted in the strongest negative responses in salmon survival and abundance. Additionally, feedbacks through the food web were strong, beginning with primary production, suggesting that several food web variables may be important in mediating effects on salmon survival within the system. With this model, we were able to compare the relative influence of multiple drivers on salmon survival.

22    **Introduction**

23            Problems of complex interactions are common in many fields, including medicine,  
24    economics, and ecology (Levins 1974). In ecology, much attention has been given to describing  
25    food webs and interactions among species (Paine 1966, May 1974, Pimm et al. 1991, Dunne et  
26    al. 2002a). But often, these food webs are nested within larger ecological or social-ecological  
27    contexts where exogenous forces influence components of the food web system. External  
28    forcings may include physical drivers, anthropogenic impacts, or ecosystem components that  
29    are not characterized within the focal network. In social-environmental systems, tools that  
30    incorporate ecological properties, abiotic variables, and management actions within the same  
31    analytical framework are needed to accurately understand the dynamics of complex systems  
32    and evaluate potential management actions (Liu et al. 2007). However, rarely are compatible  
33    datasets available for this type of analysis. Here we use a qualitative network model, a  
34    conceptually based modeling approach, and a suite of simulations to address questions about  
35    the relative impacts of human and natural influences on early marine survival of juvenile  
36    salmon.

37            In recent years, attention has turned to marine life-history stages of Pacific salmon  
38    (*Oncorhynchus* spp.) in an effort to understand population declines and the subsequent failure  
39    to rebound, given myriad conservation and restoration efforts in freshwater streams. In  
40    Chinook salmon, coho salmon, and steelhead (*O. tshawytscha*, *O. kisutch*, and *O. mykiss*,  
41    respectively), declines in marine survival have been evidenced within the Salish Sea (Puget  
42    Sound, WA, USA and the Strait of Georgia, BC, Canada) that have not been seen in coastal  
43    populations (Beamish et al. 2010, Johannessen and McCarter 2010, Zimmerman et al. 2015,

Ruff et al. 2017, Kendall et al. 2017). These inland water bodies serve as habitats for juvenile salmon as they pass from natal streams to ocean waters during their outmigration. Yet, because of complex anthropogenic changes brought about by population increases and the associated human activity in these waters, it is likely that a number of factors and their cumulative—synergistic or additive—effects are contributing to early-marine mortality. Other salmon species, such as chum, pink and sockeye salmon (*O. keta*, *O. gorbuscha*, and *O. nerka*, respectively), have not experienced similar declines (Debertin et al. 2017), suggesting that life-history characteristics may also contribute to increased mortality for some species in this region. Teasing apart which factors have negatively impacted survival of juvenile salmon in marine waters is of concern to local, regional, and federal governments and other stakeholders (e.g., Salish Sea Marine Survival Project, <http://marinesurvivalproject.com/>) and management actions are being sought to lessen impacts and increase survival during this period.

Correlative studies of salmon abundance and environmental factors have been ongoing in the greater region for many years (Pearcy 1988, Beamish et al. 2000, Teo et al. 2009, Burke et al. 2013). Food web models for the Strait of Georgia (Priekshot 2008) and Puget Sound (Harvey et al. 2012) show primary production as an important driver in the biological system and also the effects of top predators in creating trophic cascades and influencing food web dynamics in the mid-trophic levels where time-series data are sparse (Harvey et al. 2012). Even with an understanding of the main variables in a given system, measuring abundances of each variable and the flux of energy among them often poses a logistical challenge (Christensen and Walters 2004). These models do not easily incorporate non-fisheries anthropogenic impacts, such as habitat loss or contaminant exposure, yet we understand that in many systems diverse but

66 cumulative impacts can play a role in species population change. For this reason, conceptually  
67 based models, incorporating a broader array of variables, are an important tool in providing an  
68 integrated picture of ecological and human drivers of ecosystem change.

69 One tool for evaluating the relative influence of ecosystem components is Qualitative  
70 Network Modeling (QNM, also called Qualitative Network Analysis, QNA, or Loop Analysis,  
71 Levins 1974, Puccia and Levins 1985, Raymond et al. 2011, Melbourne-Thomas et al. 2012,  
72 Harvey et al. 2016), which is advantageous for understanding a system of complex interactions  
73 which are not fully specified and when precise measurement is impossible, but when some  
74 mechanistic understanding of interactions exists. It allows for the testing of competing  
75 hypotheses, given different model structures or the invocation of perturbations to one or more  
76 of the model variables. QNM does not explicitly include non-linear direct effects, which occur in  
77 and influence social and ecological systems, however, it may help to determine the relative  
78 impacts of competing hypothesized factors or indicate where empirical work could be focused  
79 to improve system understanding (Levins 1974).

80 Researchers have used QNM for evaluating ecosystem response to ocean acidification in  
81 shellfish management (Reum et al. 2015) and the impacts of eutrophication and species  
82 management within a food web (Carey et al 2013), and for discerning the impact of  
83 management actions for species recovery (Harvey et al. 2016) in the Pacific Northwest. QNM is  
84 an important conceptual tool for discerning relative impacts of ecosystem components from  
85 which more complex, data-driven modeling efforts can stem. Here we apply this technique to  
86 evaluate a suite of potential drivers thought to be contributing to increased early marine  
87 mortality in a group of Pacific salmon in the Salish Sea. Using a simulation framework, we

invoke perturbations to each model variable and suites of variables based on salmon early marine survival hypotheses, and assess model response related to the salmon species of concern. This work is a foundational step in understanding the impacts of multiple drivers of marine survival declines in Salish Sea salmon.

## Methods

We used QNM to address our primary question regarding the relative impacts of various factors on salmon early marine survival. Our analysis had three main steps: (1) construct an enhanced conceptual model showing positive, negative, and neutral relationships; (2) generate a pool of stable simulated models with random weights applied to each model linkage; and (3) invoke one or more perturbations based upon mechanistic understanding of the system and determine the model response.

## Conceptual Model

To construct our conceptual model of the Salish Sea system, we gathered existing literature and experts on ecosystem components from within and outside of the project technical team. The technical team includes scientists from resource agencies, universities, and tribal entities, with expertise ranging from salmon genetics to disease ecology to numerical ocean modeling. We began by developing a list of over 40 possible variables drawn from hypotheses about the decline of Pacific salmon survival within the system (Salish Sea Marine Survival Project hypotheses, <http://marinesurvivalproject.com/the-project/key-hypotheses/>). The variables included: physical forcings, biological components from primary production to top predators and competitors, and anthropogenic variables. We drew a draft model based upon

our knowledge of the system and existing literature and then we conducted small meetings with experts on particular components, such as disease ecology or oceanography, and iteratively developed a working conceptual model. We sought out additional feedback from those working within the Salish Sea, but not on the technical team, on both the model components and structure, and received further feedback during public presentations to refine the conceptual model.

For the final model, we grouped model variables into several driver groups: environmental factors, primary production, food web interactions, and anthropogenic impacts (Table 1). While the conceptual model is not exhaustive, it does include many of the drivers identified in our working hypotheses and reflects known interactions within the ecosystem. The inclusion of model variables that are not biomass pools (e.g., temperature, habitat loss) highlights the flexibility of the qualitative modeling approach. Focusing on physical, bottom-up, top-down, and anthropogenic factors fit with the working hypotheses of the Marine Survival Project and enabled exploration of combinations of diverse variables.

The emphasis of the modeling effort was on understanding sources of decreased survival of the focal salmon species (Chinook, coho, and steelhead), which have shown an overall declining population trend (Figure 1, left panel, Appendix 1 for details), in addition to a decline in marine survival (Zimmerman et al. 2015, Ruff et al. 2017, Kendall et al. 2017). Central to our approach was specifying multiple salmon characteristics as modeled network nodes, namely: size, fitness, residence time, abundance, and survival. We used these traits as primary response variables throughout our analysis. While marine survival (herein ‘survival’) was our principal variable of interest, we included additional traits to evaluate the relative impact to



metrics of salmon performance. We included ‘other salmon’ as a model variable, representing pink, chum, and sockeye salmon, because the migration timing of all Pacific salmon species means competitive interactions occur. However, the species represented by the ‘other salmon’ variable have not experienced the same negative population trends (Irvine and Ruggerone 2016, Figure 1 and Appendix 1 for details) and are seen as important to the analysis but different from the focal species. While the emphasis was on representing the most direct impacts on the focal salmon traits, we recognize that many of the model variables (e.g., temperature) could potentially have direct connections to other model nodes; we have included these where interactions were important for understanding implications for the focal salmon variables or where existing literature has shown strong connections.

We defined relationships among variables as positive, negative, or null, based upon mechanistic understanding of the Salish Sea system and input from regional experts. To implement the simulation modeling, we developed a conceptual digraph using the directed graphing software, Dia (v.0.97.2) to represent the model system and the interactions among variables. This diagram served as the foundation for our qualitative modeling.

### Simulated Networks

We used the *QPress* package for Qualitative Network Analysis (Melbourne-Thomas et al. 2012) with custom modifications in R (R Core Team 2016) to interpret the conceptual digraph, construct simulated networks, and perform our analyses. The digraph is interpreted as an interaction matrix, **A**, where each directed pairwise interaction is represented as coefficients  $a_{ij}$ . **A** is treated similarly to a community interaction matrix, wherein the rate of change of any

given node is a continuous function of all other interacting nodes (Levins 1974; Puccia and Levins 1985). The interacting components (i.e., model nodes) are set up as a series of differential equations:

$$\frac{dx_i}{dt} = f_i(x_1, x_2, \dots, x_n; c_1, c_2, \dots, c_m)$$

where  $x_i$  is the density of model component (population)  $i$ , the  $c$ s are growth parameters, and  $f_i$  is a function describing per capita growth-rate of that population (Raymond et al. 2011). Therefore, the interaction coefficients  $a_{ij}$  describe the effect of a change in the level of component  $j$  on the level of component  $i$ , as defined by the partial derivative of  $f_i$  with respect to  $N_j$ :  $a_{ij} = \partial f_i / \partial x_j$  evaluated at the equilibrium (Levins 1974, Raymond et al. 2011, Melbourne-Thomas et al. 2012).

Given a network model and corresponding interaction matrix,  $\mathbf{A}$ , the negative of the inverse community matrix ( $-\mathbf{A}^{-1}$ ) yields estimated changes in the equilibrium abundances of each component  $x$  as a function of a sustained (press) perturbation of one or more system components (Puccia and Levins 1985). The *QPress* analysis package provides routines for evaluating the impact of a press perturbation to the system through simulation. For each simulation, a weight (drawn from a random uniform distribution, 0-1) was assigned to each linkage (edge). These weights were positive or negative depending upon the relationship between the two endpoints. If the resulting model with all assigned weights was stable (i.e., converged), the model was accepted. We simulated the network c. 122,000 times to result in 10,000 stable simulated networks. We then assessed the proportion of model runs with positive, negative, and neutral responses given changes to particular nodes (see below).

We assessed the sensitivity and robustness of the model. We experimented with changing both the distribution and the variance of the weighting scheme, but did not find large differences in results, so maintained the default weighting for our analyses. We explored the weights of linkages in the balanced models to look for anomalies (methods and results in Appendix 2). Additionally, we calculated distance to the Survival node via pathways from each model variable to check for effects of model structure, described network properties such as connectance and linkage density, and evaluated model behavior with the sequential addition of perturbed nodes and a set of ‘cumulative effects’ of both influential and neutral nodes.

### Invoking Perturbations

To test hypotheses regarding marine survival, we developed *a priori* perturbations to each model node (Table 2). The direction of the perturbation (increase or decrease) was based upon our understanding of the system, changes that have occurred concomitant with declines in salmon marine survival (since the 1970s), and expected impacts as a result of anthropogenic change (Appendix 3). We employed several cumulative effects scenarios and modified existing software functions to meet our analytical objectives.

First, we perturbed each node individually and observed outcomes to all other model variables. This allowed for a simple comparison of impacts on the focal salmon metrics from each variable and the ability to compare the extent of the impact to that from any other variable. Second, we evaluated the relative effects of different groups of drivers (Table 3). For example, we were interested in food web effects, so we simultaneously decreased forage fish, increased marine mammals, decreased piscivorous fish, and increased gelatinous

zooplankton—trends that have been observed in Puget Sound—and observed the impacts to the other model components. For each driver group, we selected four nodes to perturb, thereby standardizing the level of change invoked. By comparing impacts to salmon traits from primary production, food web, environmental, and anthropogenic drivers, we were able to query the relative impacts of each of these groups.

Finally, we developed scenarios based upon observed changes within three regions of Puget Sound to see how well the model reproduced cumulative impacts in terms of response to the focal salmon metrics, especially survival. The three regions were: (a) South Sound, with a known decline in salmon abundance and cumulative impacts including increased gelatinous zooplankton, nutrients, contaminants and hatchery production, and decreased forage fish abundance; (b) Hood Canal, which has had relatively stable salmon abundances, but impacts in oceanography, including increased stratification and temperature and low dissolved oxygen; and (c) Central Basin which has shown a decline in salmon abundance, relatively less than South Sound, but with a different suite of cumulative impacts including habitat loss, contaminant input, and decreased primary production. In reality, causes of declining survival are likely multifaceted, complex, and non-linear, and this modeling exercise allowed us to examine the relative influence of many factors within one modeling framework.

## Results

### Model

Our final conceptual model had 33 nodes including salmon traits and atmospheric, oceanographic, primary production, food web, and anthropogenic drivers (Table 1, Fig. 2,

Appendix 3). There were a total of 150 linkages out of 1089 potential linkages within the model. This gives a network density or connectance (realized linkages/potential linkages) of 0.138 and a linkage density (average # of linkages/node) of 4.55. Connectance has been linked with network stability in ecological networks (Dunne et al. 2002b) and in social network theory applied to behavioral ecology (Sih et al. 2009), and also to resilience in social-ecological systems (Janssen et al. 2006). The most highly connected nodes were temperature, diatoms, and (salmon) abundance with a total of 11 connections each; the other salmon and zooplankton nodes were both highly connected (10 linkages) and highly influenced by other variables (10 and 9 influencing nodes, respectively, Table 1). All nodes were a minimum distance of four nodes or fewer from survival, but the range of feedback linkages varied greatly, from 1 to 10. Each model node included a self-regulating feedback to better represent ecological limits and to aid in model convergence; the exception was survival, which was considered the primary variable of interest and was not constrained.

To assess whether the proximity of each model node to survival influenced the outcomes of our analysis (i.e., are nodes more directly connected to survival more likely to result in stronger outcomes?), we evaluated the proportion of negative results for survival with the minimum nodal distance to survival and found no relationship. Both closely connected nodes (minimum nodal distance of 1) and those more distant ( $\geq 2$  nodes away), resulted in a range of negative responses (<20% to >95%) with respect to survival. Thus, we do not believe the model structure strongly confounded our results.

234    **Perturbations**

235            The results of the press perturbations to each node showed that anthropogenic impacts  
236    resulted in the most consistent negative responses in salmon traits, specifically survival and  
237    abundance (Table 2, Appendix 4). Here we use consistency of response to refer to the relative  
238    proportion of outcomes that were positive or negative given a perturbation—a strongly  
239    consistent response was when proportionally more simulated models (here >80%) had positive  
240    or negative responses for the node of interest, while a neutral response resulted when the  
241    simulated outcomes were equally positive/negative in outcome.

242            Individual perturbations showed an increase in CO<sub>2</sub> resulted in a consistently positive  
243    response in survival and abundance; CO<sub>2</sub> positively influences diatoms in the model, with  
244    positive effects cascading through the food web. Conversely, a decrease in diatoms (primary  
245    production) resulted in a consistently negative response in survival and abundance, as well as  
246    negatives outcomes to zooplankton and turbidity. A direct perturbation to zooplankton  
247    (decrease) resulted in strongly consistent negative responses in fitness and size, but less  
248    consistently negative results in survival and abundance. On the other hand, a direct decrease to  
249    turbidity resulted in a consistent positive response in the individual traits, fitness and size, but  
250    slightly negative responses in survival and abundance, population-level traits. This is despite the  
251    fact that turbidity directly and positively affects survival in the model; this relationship is a  
252    result of the association of turbidity with primary production and the resulting predation  
253    dynamics in the model. A decrease in the predators (piscivorous fish and birds) resulted in  
254    positive responses in survival and abundance and more ambiguous impacts to size and fitness.  
255    Marine mammals, also known predators of salmon, but with increasing populations (and thus a

positive perturbation), had a neutral response on all salmon response variables. Unexpectedly, increased harvest had a positive effect on survival; harvest has a direct negative effect on abundance within the model, but the feedback to survival is mediated by the food web, specifically forage fish and zooplankton, which may moderate the harvest impacts to survival through reduced competition. This same result indicates that strong feedback mechanisms, like density dependence, were simulated by the model and highlights the importance of complex food web interactions for salmon survival.

The results of the driver group analysis, which evaluated cumulative impacts from one section of the network, showed anthropogenic impacts to have predominantly negative effects on survival, abundance, and fitness, with over 85% of the simulations having negative responses within these model groups (Fig. 3). For the environmental driver group, most simulation results were positive for survival and abundance, but neutral for the other response variables. The primary production group showed consistently negative results in survival and abundance, indicating that changes to primary production can have strong impacts on salmon via the food web. Interestingly, the food web manipulation yielded strong negative responses on the salmon individual traits (size and fitness) but more neutral response on the population-level traits (abundance and survival). The other salmon category had a consistently positive response in both the food web and anthropogenic driver simulations, indicating that conditions that are less favorable within the food web for the focal salmon species may not be detrimental for other species, with diets that tend to be more planktivorous and generally shorter rearing times within the Salish Sea.

For the regional differences, where we modeled three different regions of Puget Sound with respect to salmon survival, our model replicated the observed trends within these regions, with strong negative responses in focal salmon survival, abundance, and fitness in both South Sound and Central Basin. The results for Hood Canal were more weakly negative for salmon survival and abundance than in the other regions, and were neutral for fitness and size, suggesting that some of the oceanographic changes evidenced in Hood Canal may be less detrimental for salmon. Responses of other salmon were consistently positive in all three regions, reflecting observed population trends (Fig. 1). Therefore, although our model is a generalization of the processes occurring in the southern portion of the Salish Sea, it does replicate observed trends in the region.

We recognize that many of the perturbations invoked within the model are happening concurrently. To evaluate whether the model would maintain robustness when multiple interacting factors (e.g. cumulative effects) were invoked, we sequentially added disturbances to the model and evaluated the outcomes. When influential individual drivers were included (from Table 2), the results were strongly negative for survival and abundance. We compared 10 of the strongest responding nodes with 10 that showed neutral influence on survival and compared the results (Fig. 4). We observed that the influential individual nodes resulted in consistently negative impacts to survival, while the response was neutral for the weaker suite of disturbances.



## Discussion

Our model showed that a wide variety of drivers had negative effects on early marine survival of coho salmon, Chinook salmon and steelhead. The impacts on the other salmon variable (representing pink, chum, and sockeye) were neutral or positive. With only five drivers (precipitation, river flow, microbial detritivores, zooplankton, and ichthyoplankton) negatively influencing the other salmon node in our simulations, the model structure seemed to capture the reduced impacts to these species within the Salish Sea (Fig. 1, Debertin et al. 2017). The combination of drivers having negative effects on attributes of salmon fitness and survival suggests that a single sector of the network is insufficient for explaining increased marine mortality and that feedbacks and complex interactions may both exacerbate and mediate the effects of individual drivers.

Anthropogenic factors induced negative responses in salmon traits, especially survival, abundance, and fitness. The factors are both direct (e.g., contaminants and disease) and mediated by the food web (e.g., hatcheries, with increased production leading to competitive interactions; and habitat loss, which has a negative effect on salmon residency and fitness, but also on forage fish, because nearshore habitat is critical to forage fish spawning). There are likely indirect connections that were unaccounted for in our model that may make these impacts even stronger in the real world. The food web components individually did not yield strong responses in salmon survival and abundance, but did impact size and fitness. Considering that many of these linkages are both directly and indirectly tied to salmon, the negative outcomes, even where marginal, should be noted. Additionally, many of the feedbacks present in the model were through the food web. In the driver group analysis, the negative

318 responses from food web changes to the individual traits of size and fitness is notable. Our  
319 model does not have a temporal component, but negative impacts to individual traits would  
320 likely manifest in survival and population declines over time.

321 The conceptual model exhibited particularly strong sensitivity to changes in key linkages:  
322 the effects of fitness upon size and vice-versa, and of survival upon abundance. These results  
323 suggest that processes influencing these factors will strongly influence marine survival and  
324 point to the importance of monitoring these pathways. Size and fitness are relatively easy-to-  
325 monitor characteristics measured in standard salmon sampling programs. The measurement of  
326 size, combined with techniques that can measure growth and fitness, including the use of  
327 chemical-based indicators of fitness, such as fatty acid biomarkers (Hook et al. 2014) or  
328 hormone markers (Beckman 2011), would provide some indication of how these attributes are  
329 changing over time. However, understanding mechanisms for changes in size and fitness is  
330 more complex. As our model and existing literature suggest, several factors contribute to  
331 changes in size over time, with food web alteration being one of the likely mechanisms.  
332 However, these changes are not yet fully understood in the Salish Sea. Our model allowed for  
333 comparison of multiple factors and showed that a decrease in primary production had the  
334 strongest negative impact on salmon survival. Additional research on the impacts of changing  
335 primary production in this system and the links to salmon condition would aid in further teasing  
336 apart this relationship.

337 Our response metrics focused on model runs that converged on an equilibrium (i.e., only  
338 balanced models were used in the perturbation scenarios). The number of model runs needed  
339 (c. 122,000) to get a subset of converged models (10,000) suggests that most models did not

converge to a stable solution. The ‘real’ Salish Sea is likely represented by one of the many possible combinations and may in fact be unstable, not in a ‘converged’ state, as our model assumed. Such unstable states are predicted outcomes when ecosystems surpass tipping points (Carpenter and Brock 2006, Samhouri et al. 2017). Alternately, the Salish Sea ecosystem may have reached a new equilibrium: analysis of marine survival trends indicate a steep decline in the 1970s and 80s, leveling off at a low level that has persisted to the present (Zimmerman et al. 2015). Hence, the simulation framework with multiple sets of initial conditions may have allowed us to detect endpoints that include a new equilibrium for marine survival of salmon.

The conceptual model underlying the analysis represents a complex set of feedbacks. Additional interconnections that we did not represent would tend to stabilize the system even more (Dunne et al. 2002a, Ives and Carpenter 2007). And while nonlinear properties are fundamental to ecological systems, they are nearly always influenced by feedbacks, an essential component built into our model structure (DeAngelis and Waterhouse 1987, Scheffer et al. 2001). The lack of explicit spatial or temporal components within the model limits our ability to make predictions beyond a static snapshot. However, through our comparison of three sub-regions within our system, we were able to compare different starting conditions and gauge model response. While additional complexity in the temporal component would allow for detecting evidence of change over time, the lack of comprehensive empirical data across all ecosystem components currently limits the tractability of such a modeling approach.

The model results in and of themselves are informative for comparing among a suite of potential causes of declining marine survival in salmon and in evaluating cumulative effects of these drivers. Additionally, this conceptual model forms the foundation for additional research

efforts. Currently, development of multivariable time-series analysis and an end-to-end ecosystem model (Atlantis) is ongoing, via the Salish Sea Marine Survival Project. The development of the Atlantis ecosystem model has been aided by the conceptual underpinnings presented herein. Efforts have been made to develop data streams where the conceptual model has shown important linkages (e.g., a sub-project was initiated using satellite-derived data to better understand variability in primary production in the system). In this way, the qualitative network model is foundational for continuing quantitative modeling work, but is also an important tool for conveying the complexity of the system and the problem to diverse audiences.

Our results suggest that teasing out the causes of declines in marine survival will be challenging, multi-faceted, and will involve both understood and unknown feedbacks. Multiple singular factors led to declines in most of the simulations, and suites of ecosystem components had strong effects on marine survival and other salmon attributes. Nevertheless, our ability to distinguish causal factors will likely be improved by tracking multiple ecosystem indicators, especially those influencing salmon size and fitness. As with any model, ours is a simplified version of the ecosystem; however, our approach resulted in a complex representation of declining populations in an ecosystem context that served as a useful tool for identifying relative influences of numerous hypothesized drivers of marine mortality. Through this examination, we have identified some sectors of the ecosystem, such as the food web and anthropogenic impacts, which warrant further examination.

382 [Supplementary material](#)

383 For supplementary material accompanying this paper, visit:

384 <http://www.journals.cambridge.org/ENC>

385

386

Proof for Review

387    **Acknowledgements**

388    We thank members of the Salish Sea Marine Survival Project technical team for input to the  
389    conceptual model. B. Raymond provided assistance with modifying R code. I. Kaplan, C. Harvey,  
390    I. Perry, I. Pearsall, T. Curran, and I. Kemp provided insight on early versions of the modeling  
391    effort and results. A. Dufault and N. Kendall (Washington Department of Fish and Wildlife)  
392    assisted with data compilation for Fig. 1. The manuscript benefited from comments provided by  
393    I. Kaplan, C. Harvey, and R. Zabel (NOAA-Northwest Fisheries Science Center) and from the  
394    anonymous reviewers.

395    **Financial Support**

396    This work was paid for by the Pacific Salmon Commission's Southern Endowment Fund via Long  
397    Live the Kings. This is publication number 12 from the Salish Sea Marine Survival Project, an  
398    international research collaboration designed to determine the primary factors affecting the  
399    survival of juvenile salmon and steelhead in the Salish Sea.

400    **Conflict of Interest**

401    None

402    **Human and/or Animal Experimentation**

403    None

## Literature Cited

- Beamish, R.J., Noakes, D.J., McFarlane, G.A., Pinnix, W., Sweeting, R. & King, J. (2000) Trends in coho marine survival in relation to the regime concept. *Fisheries Oceanography* **9**:114–119.
- Beamish, R.J., Sweeting, R.M., Lange, K.L., Noakes, D.J., Preikshot, D. & Neville, C.M. (2010) Early marine survival of coho salmon in the Strait of Georgia declines to very low levels. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science* **2**: 424–439.
- Beckman, B.R. (2011) Perspectives on concordant and discordant relations between insulin-like growth factor 1 (IGF1) and growth in fishes. *General and Comparative Endocrinology*, **170**:233–252.
- Burke, B.J., Peterson, W.T., Beckman, B.R., Morgan, C., Daly, E.A. & Litz, M. (2013) Multivariate models of adult Pacific salmon returns. *PLoS ONE* **8**: e54134.doi:10.1371/journal.pone.0054134
- Carey, M.P., Levin, P.S., Townsend, H., Minello, T.J., Sutton, G.R., Francis, T.B., Harvey, C.J., Toft, J.E., Arkema, K.K., Burke, J.L., Kim, C., Guerry, A.D., Plummer, M., Spiridonov, G. & Ruckelshaus, M. (2014) Characterizing coastal foodwebs with qualitative links to bridge the gap between the theory and the practice of ecosystem-based management. *ICES Journal of Marine Science* **71**: 713–724.
- Carpenter, S.R., & Brock, W.A. (2006) Rising variance: a leading indicator of ecological transition. *Ecology Letters* **9**: 311–318.
- Christensen, V. & Walters, C.J. (2004) Ecopath with Ecosim: methods, capabilities and limitations. *Ecological Modelling* **172**:109–139.
- DeAngelis, D.L. & Waterhouse, J.C. (1987) Equilibrium and nonequilibrium concepts in ecological models. *Ecological Monographs* **57**: 1:21.
- Debertin, A.J., Irvine, J.R., Holt, C.A., Oka, G. & Trudel, M. (2017) Marine growth patterns of southern British Columbia chum salmon explained by interactions between density-dependent competition and changing climate. *Can. J. of Fish. and Aqua. Sci.* **74**: 1077–1087.
- Dunne, J.A., Williams, R.J. & Martinez, N.D. (2002a). Network structure and biodiversity loss in food webs: robustness increases with connectance. *Ecol. Lett.* **5**: 558–567.
- Dunne, J.A., Williams, R.J. & Martinez, N.D. (2002b). Food-web structure and network theory: the role of connectance and size. *Proceedings of the National Academy of Sciences of the United States of America.* **99**: 12917–12922.

- 445 Greene, C.M., Kuehne, L., Rice, C.A., Fresh, K.L. & Penttala, D. (2015) Forty years of change in  
 446 forage fish and jellyfish abundance across greater Puget Sound, Washington (USA):  
 447 anthropogenic and climate associations. *Marine Ecology Progress Series* **525**: 153-170.  
 448
- 449 Harvey, C.J., Williams, G.D. & Levin, P.S. (2012) Food web structure and trophic control in  
 450 central Puget Sound. *Estuaries and Coasts* **35**: 821-838. doi:10.1007/s12237-012-9483-1  
 451
- 452 Harvey, C.J., Reum, J.C.P., Poe, M.R., Williams, G.D. & Kim, S.J. (2016) Using conceptual models  
 453 and qualitative network models to advance integrative assessments of marine ecosystems.  
 454 *Coastal Management*. **44**: 486-503, DOI: 10.1080/08920753.2016.1208881  
 455
- 456 Hoekstra, J.M., Bartz, K.K., Ruckelshaus, M.H., Moslemi, J.M. & Harms, T.K. (2007) Quantitative  
 457 threat analysis for management of an imperiled species: Chinook salmon (*Oncorhynchus*  
 458 *tshawytscha*). *Ecological Applications* **17**:2061–2073.  
 459
- 460 Hook, S., Gallagher, E. & Batley, G. (2014) The role of biomarkers in the assessment of aquatic  
 461 ecosystem health. *Integrated Environmental Assessment and Monitoring*. **10**:327-41.  
 462
- 463 Holmes, E.E. (2001) Estimating risks in declining populations with poor data. *Proceedings of the*  
 464 *National Academy of Sciences of the United States of America*. **98**: 5072-5077.  
 465
- 466 Irvine, J.R. & Ruggerone, G.T. (2016) Provisional estimates of numbers and biomass for natural-  
 467 origin and hatchery-origin pink, chum, and sockeye salmon in the North Pacific, 1952-2015.  
 468 *NPAFC Doc. 1660*. 45 pp. Fisheries and Oceans Canada, Pacific Biological Station and Natural  
 469 Resources Consultants, Inc. (Available at <http://www.npafc.org>).  
 470
- 471 Ives, A.R. & Carpenter, S.R. (2007) Stability and Diversity of Ecosystems. *Nature* **317**: 58-62.  
 472
- 473 Janssen, M.A., Bodin, Ö., Anderies, J.M., Elmqvist, T., Ernstson, H., McAllister, R.R.J., Olsson, P.,  
 474 & Ryan, P. (2006) A network perspective on the resilience of social-ecological systems. *Ecology*  
 475 *and Society* **11**: 15.  
 476
- 477 Johannessen, S.C. & McCarter, B. (2010) Ecosystem Status and Trends Report for the Strait of  
 478 Georgia Ecozone. *DFO Can. Sci. Advis. Sec. Res. Doc.* 2010/010. vi + 45 p.  
 479
- 480 Kendall, N.W., Marston, G.W., & Klungle, M.M. (2017) Declining patterns of Pacific Northwest  
 481 steelhead trout (*Oncorhynchus mykiss*) adult abundance and smolt survival in the ocean. *Can. J.*  
 482 *of Fish. and Aqua. Sci.* **74**: 1275-1290.  
 483
- 484 Levins, R. (1974) The qualitative analysis of partially specified systems. *Annals of the New York*  
 485 *Academy of Sciences* **231**: 123–138.  
 486



- 487 Liu, J., Dietz, T., Carpenter, S.R., Folke, C., Alberti, M., Redman, C.L., Schneider, S.H., Ostrom, E.,  
 488 Pell, A.N., Lubchenco, J., Taylor, W.W., Ouyang, Z., Deadman, P., Kratz, T., & Provencher, W.  
 489 (2007) Coupled human and natural systems. *AMBIO*. **236**: 639–649.  
 490
- 491 Mauger, G.S., Casola, J.H., Morgan, H.A., Strauch, R.L., Jones, B., Curry, B., Busch Isaksen, T.M.,  
 492 Whitely Binder, L., Krosby, M.B., & Snover, A.K. (2015) *State of Knowledge: Climate Change in*  
 493 *Puget Sound*. Report prepared for the Puget Sound Partnership and the National Oceanic and  
 494 Atmospheric Administration. Climate Impacts Group, University of Washington, Seattle.  
 495 doi:10.7915/CIG93777D  
 496
- 497 May, R. M. (1974) Stability and Complexity in Model Ecosystems. *Monographs in Population*  
 498 *Biology* **6**, Princeton University Press, Princeton, New Jersey  
 499
- 500 Melbourne-Thomas, J., Wotherspoon, S., Raymond, B., & Constable, A. (2012) Comprehensive  
 501 evaluation of model uncertainty in qualitative network analyses. *Ecological Monographs*, **82**:  
 502 505–519.  
 503
- 504 Ogden, A.D., Irvine, J.R., English, K.K., Grant, S., Hyatt, K.D., Godbout, L. & Holt, C.A. (2015)  
 505 Productivity (recruits-per-spawner) data for sockeye, pink, and chum salmon from British  
 506 Columbia. *Can. Tech. Rep. Fish. Aquat. Sci.* 3130: vi + 57 p.  
 507
- 508 O'Neill, S.M. & West, J.E. (2009) Marine distribution, life history traits, and the accumulation of  
 509 polychlorinated biphenyls in Chinook salmon from Puget Sound, Washington. *Transactions of*  
 510 *the American Fisheries Society* **138**: 616-632.  
 511
- 512 Paine, R.T. (1966) Food web complexity and species diversity. *American Naturalist* **100**: 65-76.  
 513
- 514 Pearcy, W.G. (1988) Factors affecting survival of Coho Salmon off Oregon and Washington.  
 515 Pages 67–73 in W. J. McNeil, editor. *Salmon production, management, and allocation*. Oregon  
 516 State University Press, Corvallis.  
 517
- 518 Pimm, S.L., Lawton, J.H. & J.E. Cohen (1991) Food web patterns and their consequences.  
 519 *Nature* **350**, 669–674.  
 520
- 521 Preikshot, D.B. (2008) *Public Summary—Computer Modelling of Marine Ecosystems:*  
 522 *Applications to Pacific Salmon Management and Research*. Vancouver, BC: Pacific Fisheries  
 523 Resource Conservation Council.  
 524
- 525 Puccia, C.J. & Levins, R. (1985) *Qualitative modeling of complex systems: An introduction to loop*  
 526 *analysis and time averaging*. Cambridge, MA: Harvard University Press.  
 527
- 528 Puget Sound Ecosystem Monitoring Program (PSEMP) Marine Waters Workgroup (2016) *Puget*  
 529 *Sound Marine Waters: 2015 Overview*. S.K. Moore, R. Wold, K. Stark, J. Bos, P. Williams, K.

- 530 Dzinbal, C. Krembs and J. Newton (Eds). URL:  
 531 [www.psp.wa.gov/PSEMP/PSmarinewatersoverview.php](http://www.psp.wa.gov/PSEMP/PSmarinewatersoverview.php).  
 532
- 533 R Core Team (2016) R: A language and environment for statistical computing. R Foundation for  
 534 Statistical Computing, Vienna. [www.r-project.org](http://www.r-project.org)  
 535
- 536 Raymond, B., McInnes, J., Dambacher, J.M., Way, S. & Bergstrom, D.M. (2011) Qualitative  
 537 modelling of invasive species eradication on subantarctic Macquarie Island. *Journal of Applied*  
 538 *Ecology* **48**: 181–191.  
 539
- 540 Reum, J.C.P., Ferriss, B.E., McDonald, P.S., Farrell, D.M., Harvey, C.J., Klinger, T. & Levin, P.S.  
 541 (2015) Evaluating community impacts of ocean acidification using qualitative network models.  
 542 *Marine Ecology Progress Series* **536**: 11–24.  
 543
- 544 Roberts, M., Mohamedali, T., Sackmann, B., Khangaonkar, T. & Long, W. (2014) *Puget Sound*  
 545 *and the Straits Dissolved Oxygen Assessment: Impacts of Current and Future Nitrogen Sources*  
 546 *and Climate Change Through 2070*. Washington Department of Ecology, Publication No. 14-03-  
 547 007. Olympia, WA 151 pp.  
 548
- 549 Ruff, C.P., Anderson, J.H., Kemp, I.M., Kendall, N.W., Mchugh, P. Velez-Espino, A., Greene, C.M.,  
 550 Trudel, M., Holt, C.A., Ryding, K.E. & Rawson, K. (2017) Salish Sea Chinook salmon exhibit  
 551 weaker coherence in early marine survival trends than coastal populations. *Fisheries*  
 552 *Oceanography*. DOI: 10.1111/fog.12222  
 553
- 554 Samhuri, J.F., Andrews, K.S., Fay, G., Harvey, C.J., Hazen, E.L., Hennessey, S., Holsman, K.K.,  
 555 Hunsicker, M.E., Large, S.I., Marshall, K., Stier, A.C., Tam, J. & Zador, S. (2017) Defining  
 556 ecosystem thresholds for human activities and environmental pressures in the California  
 557 Current. *Ecosphere* **8**:e01860. 10.1002/ecs2.1860  
 558
- 559 Scheffer, M., Carpenter, S., Foley, J.A., Folke, C. & Walker, B. (2001) Catastrophic shifts in  
 560 ecosystems. *Nature* **413**: 591–596.  
 561
- 562 Sih, A., Hanser, S.F. & McHugh, K.A. (2009) Social network theory: new insights and issues for  
 563 behavioral ecologists. *Behavioral Ecology and Sociobiology* **63**: 975.  
 564
- 565 Teo, S.L.H., Botsford, L.W. & Hastings, A. (2009) Spatio-temporal covariability in Coho Salmon  
 566 (*Oncorhynchus kisutch*) survival, from California to southwest Alaska. *Deep Sea Research Part II:*  
 567 *Topical Studies in Oceanography* **56**: 2570–2578.  
 568
- 569 Zimmerman, M., Irvine, J.R., O'Neill, M., Anderson, J.H., Greene, C.M., Weinheimer, J., Trudel,  
 570 M. & Rawson, K. 2015. Spatial and temporal patterns in smolt survival of wild and hatchery  
 571 coho salmon in the Salish Sea. *Marine and Coastal Fisheries* **7**: 116–134.

## 572 Table and Figure Captions

### 573 Tables

574 Table 1. Descriptions of model nodes (variables) in the Salish Sea qualitative network model by  
575 driver group. Also shown are: connectivity among nodes, the number of influencing nodes, and  
576 the distance to the survival node.

577 Table 2. Model variables in the Salish Sea qualitative network model with the direction of the  
578 press perturbation invoked in the simulations. The response of perturbations to each individual  
579 model node on the focal salmon traits (survival, abundance, fitness, size, residence) and the  
580 other salmon model group are indicated by the colored boxes. The key to direction and  
581 strength of responses of model simulations is in the lower left portion of the table.

582 Table 3. Perturbations and responses by driver group. Total nodal distance is the sum of the  
583 nodal distances of each node to the survival node.

584 Table 4. Salish Sea subbasin analysis with perturbations invoked and outcomes.

### 585 Figures

586 Figure 1. Salmon population trends within Puget Sound (left most subbasins), the Strait of  
587 Georgia (center subbasins) and the Pacific Coast (right) for species with generally decreasing  
588 trends (Chinook, coho, and steelhead, left) and increasing or stable populations trends (chum,  
589 pink, and cockeye, right). Two abnormally high Pink salmon runs were omitted from the right  
590 panel for ease of presentation—these runs had trends of 1.05 and 0.43 and were both from  
591 Puget Sound subbasins. Data are from Washington Department of Fish and Wildlife, Pacific  
592 States Marine Fisheries Commission, Ogden et al. 2015, and Zimmerman et al. 2015.

593 Figure 2. Conceptual model of the Salish Sea related to salmon survival. Model variables  
594 (shapes) represent food web components (ovals), physical drivers (rectangles), anthropogenic  
595 impacts (diamonds) and salmon traits of interest (triangles). Survival is shown as a hexagon and  
596 was the primary variable of interest.

597 Figure 3. Results for the primary response variables for each of the four groups of drivers.  
598 Perturbations were according to Table 3 and in all four figures, bars represent the number of  
599 runs resulting in negative (dark bars) or positive (light bars) response in the response variable.

600 Figure 4. Comparison of multiple ( $n=10$ ) strongly influential (top) and neutral (bottom) variables  
601 on salmon survival. The influential variables (resulting from the analysis presented in Table 2)  
602 were: CO<sub>2</sub>, contaminants, diatoms, disease, gelatinous zooplankton, habitat loss, hatcheries,  
603 other salmon, temperature, and zooplankton. The neutral variables were: global warming,  
604 forage fish, ichthyoplankton, marine mammals, microbial detritivores, nutrients, precipitation,  
605 stratification, sunlight, and upwelling. The variables in each set were perturbed simultaneously  
606 to simulate cumulative impacts and assess model response. Dark bars show negative impacts to  
607 the response variable and light bars show positive responses.

Supplementary Material—Salish Sea Salmon Qualitative Network

Model

Appendix 1. Methods and Data Supporting Figure 1.

Methods

To arrive at population trends, we compiled available adult run sizes for salmon in rivers throughout the Salish Sea. Data were from Pacific Fishery Management Council Stock Assessment and Fishery Evaluation (SAFE) Documents (<http://www.pcouncil.org/salmon/stock-assessment-and-fishery-evaluation-safe-documents/review-of-2015-ocean-salmon-fisheries/>), Pacific Salmon Commission Technical Committee Reports (<http://www.psc.org/publications/technical-reports/technical-committee-reports/chinook/>), Washington Department of Fish and Wildlife, Ogden et al. 2015, and Zimmerman et al. 2015 (Table A1.1). We used the longest time series available (typically from 1970s to >2010) and used the methods of Holmes (2001) to estimate a population trend and confidence interval for each run. If total run size (TRS, a total of catch and escapement or number of spawners) was not available, we used escapement/spawner (E) data.

Table A1.1 Salmon runs used in the calculation of population trends. TRS=Total Run Size, E=Escapement.

Run	Species	Data	Years	Data Source	Subbasin	Population Trend	95% Confidence Interval
Lake Washington	Chinook	TRS	1975-2014	PSC Joint Tech. Comm. 2016	Central Puget Sound	0.004	0.009
Green	Chinook	TRS	1975-2014	PSC Joint Tech. Comm. 2016	Central Puget Sound	-0.027	0.006
Snohomish	Chinook	TRS	1975-2014	PSC Joint Tech. Comm. 2016	Whidbey	-0.035	0.004
Hood Canal	Chinook	TRS	1981-2014	PFMC 2016	Hood Canal	0.035	0.010
Juan de Fuca	Chinook	TRS	1981-2014	PFMC 2016	Juan de Fuca	-0.006	0.006
South Puget Sound	Chinook	TRS	1981-2014	PFMC 2016	South Sound	0.013	0.005
Nooksack	Chinook	TRS	1981-2014	PFMC 2016	San Juan and Gulf Islands	-0.046	0.007
Skagit Spring	Chinook	TRS	1975-2014	PSC Joint Tech. Comm. 2016	Whidbey	0.014	0.006
Skagit Summer/fall	Chinook	TRS	1975-2014	PSC Joint Tech. Comm. 2016	Whidbey	-0.026	0.005
Stillaguamish	Chinook	TRS	1975-2014	PSC Joint Tech. Comm. 2016	Whidbey	-0.030	0.011

Willapa	Chinook	TRS	1976-2014	PFMC 2016	Pacific Coast	0.021	0.007
Grays Harbor Spring/Summer	Chinook	TRS	1976-2014	PFMC 2016	Pacific Coast	0.018	0.005
Grays Harbor Fall	Chinook	TRS	1975-2014	PSC Joint Tech. Comm. 2016	Pacific Coast	0.013	0.005
Queets Spr/Summer	Chinook	TRS	1976-2015	PSC Joint Tech. Comm. 2016	Pacific Coast	-0.025	0.007
Queets fall	Chinook	TRS	1976-2014	PSC Joint Tech. Comm. 2016	Pacific Coast	0.002	0.004
Hoh Spring/Summer	Chinook	TRS	1976-2015	PSC Joint Tech. Comm. 2016	Pacific Coast	-0.019	0.006
Hoh fall	Chinook	TRS	1976-2015	PSC Joint Tech. Comm. 2016	Pacific Coast	-0.002	0.004
Quillayute spring/summer	Chinook	TRS	1976-2015	PSC Joint Tech. Comm. 2016	Pacific Coast	-0.029	0.005
Quillayute fall	Chinook	TRS	1976-2015	PFMC 2016	Pacific Coast	-0.005	0.005
Hoko fall term	Chinook	TRS	1986-2015	PSC Joint Tech. Comm. 2016	Pacific Coast	0.021	0.009
Fraser Spring/summer	Chinook	TRS	1975-2015	PSC Joint Tech. Comm. 2016	Central Strait of Georgia	0.017	0.002
Harrison	Chinook	TRS	1984-2015	PSC Joint Tech. Comm. 2016	Central Strait of Georgia	0.050	0.004
Lake Shuswap	Chinook	E	1975-2015	PSC Joint Tech. Comm. 2016	Central Strait of Georgia	0.026	0.006
Nanaimo	Chinook	E	1975-2015	PSC Joint Tech. Comm. 2016	South Strait of Georgia, San Juan and Gulf Islands	-0.018	0.007
Cowichan	Chinook	E	1975-2015	PSC Joint Tech. Comm. 2016	South Strait of Georgia, San Juan and Gulf Islands	-0.006	0.007
Nimpkish	Chinook	E	1975-2015	PSC Joint Tech. Comm. 2016	Johnstone Strait	-0.052	0.015
Klinaklini	Chinook	E	1975-2015	PSC Joint Tech. Comm. 2016	Johnstone Strait	0.030	0.009
Kakweiken	Chinook	E	1975-2015	PSC Joint Tech. Comm. 2016	Johnstone Strait	-0.043	0.011
Kingcome	Chinook	E	1975-2015	PSC Joint Tech. Comm. 2016	Johnstone Strait	-0.064	0.018
Wakeman	Chinook	E	1975-2015	PSC Joint Tech. Comm. 2016	Johnstone Strait	-0.151	0.028
Marble	Chinook	E	1975-2015	PSC Joint Tech. Comm. 2016	Pacific Coast	0.013	0.005
Burman	Chinook	E	1975-2015	PSC Joint Tech. Comm. 2016	Pacific Coast	0.061	0.012
Tahsis	Chinook	E	1975-2015	PSC Joint Tech. Comm. 2016	Pacific Coast	0.024	0.014
Artlish	Chinook	E	1975-2015	PSC Joint Tech. Comm. 2016	Pacific Coast	0.029	0.016
Kaouk	Chinook	E	1975-2015	PSC Joint Tech. Comm. 2016	Pacific Coast	0.041	0.026
Tahsish	Chinook	E	1975-2015	PSC Joint Tech. Comm. 2016	Pacific Coast	0.004	0.011
Cedar River Winter	Steelhead	TRS	1987-2013	N. Kendall, WDFW pers. comm	Central Puget Sound	-0.220	0.023
Green River Winter	Steelhead	TRS	1978-2013	N. Kendall, WDFW pers. comm	Central Puget Sound	-0.053	0.004
Puyallup/Carbon Winter	Steelhead	TRS	1983-2013	N. Kendall, WDFW pers. comm comm	Central Puget Sound	-0.077	0.006
White River (Puyallup) Winter	Steelhead	TRS	1986-2013	N. Kendall, WDFW pers. comm	Central Puget Sound	-0.042	0.008

Englishman River winter	Steelhead	E	1982-2015	K. Pellett, pers. comm	Central Strait of Georgia	-0.063	0.011
East Hood Canal Winter	Steelhead	TRS	1989-2013	N. Kendall, WDFW pers. comm	Hood Canal	0.088	0.016
Skokomish Winter	Steelhead	TRS	1985-2006	N. Kendall, WDFW pers. comm	Hood Canal	-0.020	0.013
South Hood Canal Winter	Steelhead	TRS	1988-2013	N. Kendall, WDFW pers. comm	Hood Canal	-0.004	0.009
West Hood Canal Winter	Steelhead	TRS	2003-2013	N. Kendall, WDFW pers. comm	Hood Canal	0.015	0.021
Tsitika River summer	Steelhead	E	1976-2015	K. Pellett, pers. comm	Johnstone Strait	0.008	0.007
Salmon River winter	Steelhead	TRS	1981-2015	N. Kendall, WDFW pers. comm	Johnstone Strait	0.021	0.036
Sequim and Discovery Bays Tributaries Winter	Steelhead	TRS	1987-2013	N. Kendall, WDFW pers. comm	Juan de Fuca	-0.101	0.100
Strait of Juan de Fuca Independent Tributaries Winter	Steelhead	TRS	1991-2009	N. Kendall, WDFW pers. comm	Juan de Fuca	-0.028	0.022
Glendale winter	Steelhead	E	1992-2014	K. Pellett, pers. comm	Pacific Coast	-0.178	0.085
Ahnuhati winter	Steelhead	E	1992-2014	K. Pellett, pers. comm	Pacific Coast	-0.186	0.065
Kakweiken winter	Steelhead	E	1992-2014	K. Pellett, pers. comm	Pacific Coast	-0.206	0.063
Atwaykellesse winter	Steelhead	E	1992-2014	K. Pellett, pers. comm	Pacific Coast	-0.116	0.062
Wahpeeto winter	Steelhead	E	1992-2014	K. Pellett, pers. comm	Pacific Coast	-0.201	0.066
Heber River summer	Steelhead	E	1975-2015	K. Pellett, pers. comm	Pacific Coast	-0.012	0.004
Gordon River summer	Steelhead	E	1998-2015	K. Pellett, pers. comm	Pacific Coast	0.030	0.042
Nisqually Winter	Steelhead	TRS	1980-2013	N. Kendall, WDFW pers. comm	South Sound	-0.107	0.016
Samish Winter	Steelhead	TRS	1979-2013	N. Kendall, WDFW pers. comm	South Strait of Georgia	0.039	0.011
Pilchuck Winter	Steelhead	E	1981-2014	N. Kendall, WDFW pers. comm	Whidbey	-0.020	0.009
Skagit River Summer and Winter	Steelhead	TRS	1978-2013	N. Kendall, WDFW pers. comm	Whidbey	-0.034	0.007
Snohomish System Winter	Steelhead	TRS	1987-2013	N. Kendall, WDFW pers. comm	Whidbey	-0.048	0.012
Snohomish/Skykomish Winter	Steelhead	TRS	1987-2013	N. Kendall, WDFW pers. comm	Whidbey	-0.061	0.010
Snoqualmie Winter	Steelhead	TRS	1987-2013	N. Kendall, WDFW pers. comm	Whidbey	-0.056	0.008
Stillaguamish Winter	Steelhead	TRS	1987-2013	N. Kendall, WDFW pers. comm	Whidbey	-0.075	0.022
Green	Coho	TRS	1973-2010	Zimmerman et al. 2015	Central Puget Sound	0.039	0.014
Puyallup	Coho	TRS	1974-2010	Zimmerman et al. 2015	Central Puget Sound	-0.038	0.008
Big Qualicum	Coho	TRS	1973-2010	Zimmerman et al. 2015	Central Strait of Georgia	-0.143	0.011
Inch	Coho	TRS	1984-2010	Zimmerman et al. 2015	Central Strait of Georgia	0.103	0.023
Big Beef Creek	Coho	TRS	1977-2010	Zimmerman et al. 2015	Hood Canal	-0.025	0.009
Quilcene	Coho	TRS	1979-2010	Zimmerman et al. 2015	Hood Canal	0.039	0.014
Skokomish	Coho	TRS	1973-2010	Zimmerman et al. 2015	Hood Canal	0.013	0.008
Straits	Coho	TRS	1985-2010	Zimmerman et al. 2015	Juan de Fuca	-0.034	0.012
Black	Coho	TRS	1978-2010	Zimmerman et al.	North Strait of	-0.052	0.021

				2015	Georgia		
Puntledge	Coho	TRS	1974-2010	Zimmerman et al. 2015	North Strait of Georgia	-0.151	0.020
Quinsam	Coho	TRS	1978-2004	Zimmerman et al. 2015	North Strait of Georgia	-0.104	0.010
Bingham Creek	Coho	TRS	1976-2010	Zimmerman et al. 2015	Pacific Coast	0.017	0.014
Cowlitz	Coho	TRS	1982-2010	Zimmerman et al. 2015	Pacific Coast	-0.126	0.030
Grays	Coho	TRS	1982-2010	Zimmerman et al. 2015	Pacific Coast	-0.036	0.009
Quinault	Coho	TRS	1977-2010	Zimmerman et al. 2015	Pacific Coast	0.079	0.015
Robertson	Coho	TRS	1975-2010	Zimmerman et al. 2015	Pacific Coast	-0.014	0.013
Satsop	Coho	TRS	1973-2010	Zimmerman et al. 2015	Pacific Coast	0.036	0.010
Washougal	Coho	TRS	1976-2010	Zimmerman et al. 2015	Pacific Coast	-0.102	0.023
Deschutes	Coho	TRS	1977-2008	Zimmerman et al. 2015	South Sound	-0.048	0.041
Chilliwack	Coho	TRS	1982-2004	Zimmerman et al. 2015	South Strait of Georgia	-0.066	0.060
Nooksack	Coho	TRS	1976-2009	Zimmerman et al. 2015	South Strait of Georgia	-0.071	0.028
Skagit	Coho	TRS	1991-2010	Zimmerman et al. 2015	Whidbey	-0.049	0.014
Stilliguamish	Coho	TRS	1981-2014	Zimmerman et al. 2015	Whidbey	-0.013	0.013
Snohomish	Coho	TRS	1981-2014	Zimmerman et al. 2015	Whidbey	-0.026	0.010
Baker	Coho	TRS	1983-2010	Zimmerman et al. 2015	Whidbey	-0.048	0.018
Skykomish	Coho	TRS	1978-2010	Zimmerman et al. 2015	Whidbey	0.012	0.010
Tulalip Bay	Coho	TRS	1974-2010	Zimmerman et al. 2015	Whidbey	0.011	0.017
Green River (Area 10A)	Pink	TRS	2001-2013	A. Dufault, WDFW, pers. comm.	Central Puget Sound	1.053	0.364
Puyallup River (Area 11A)	Pink	TRS	1959-2013	A. Dufault, WDFW, pers. comm.	Central Puget Sound	0.133	0.038
Chambers	Pink	TRS	1959-2013	A. Dufault, WDFW, pers. comm.	Central Puget Sound	0.096	0.206
Nisqually R. Drainage	Pink	TRS	1959-2013	A. Dufault, WDFW, pers. comm.	Central Puget Sound	0.167	0.088
Area 13A streams	Pink	TRS	1959-2013	A. Dufault, WDFW, pers. comm.	Central Puget Sound	-0.029	0.054
Area 13B streams	Pink	TRS	1959-2013	A. Dufault, WDFW, pers. comm.	Central Puget Sound	0.036	0.035
Port Gamble	Pink	TRS	1959-2013	A. Dufault, WDFW, pers. comm.	Hood Canal	0.425	0.593
Dosewallips	Pink	TRS	1959-2013	A. Dufault, WDFW, pers. comm.	Hood Canal	-0.095	0.028
Duckabush	Pink	TRS	1959-2013	A. Dufault, WDFW, pers. comm.	Hood Canal	-0.025	0.054
Hamma	Pink	TRS	1959-2013	A. Dufault, WDFW, pers. comm.	Hood Canal	0.091	0.052
Hoodspout Hatchery	Pink	TRS	1959-2013	A. Dufault, WDFW, pers. comm.	Hood Canal	0.063	0.024
Misc 12C	Pink	TRS	1959-2013	A. Dufault, WDFW, pers. comm.	Hood Canal	-0.002	0.041
Skokomish R. Drainage	Pink	TRS	1959-2013	A. Dufault, WDFW, pers. comm.	Hood Canal	0.093	0.055



Dungeness River	Pink	TRS	1959-2013	A. Dufault, WDFW, pers. comm.	Juan de Fuca	-0.009	0.053
Elwha River	Pink	TRS	1959-2013	A. Dufault, WDFW, pers. comm.	Juan de Fuca	-0.182	0.093
Nooksack River	Pink	TRS	1959-2013	A. Dufault, WDFW, pers. comm.	South Strait of Georgia	-0.012	0.035
Samish River	Pink	TRS	1959-2013	A. Dufault, WDFW, pers. comm.	South Strait of Georgia	0.221	0.068
Fraser	Pink	TRS	1959-2011	Ogden et al. 2015	South Strait of Georgia	0.070	0.030
Skagit River	Pink	TRS	1959-2013	A. Dufault, WDFW, pers. comm.	Whidbey	0.004	0.030
Snohomish River	Pink	TRS	1959-2013	A. Dufault, WDFW, pers. comm.	Whidbey	0.080	0.025
Stillaguamish River	Pink	TRS	1959-2013	A. Dufault, WDFW, pers. comm.	Whidbey	0.000	0.027
Lake Washington	Sockeye	E	1972-2015	A. Dufault, WDFW, pers. comm.	Central Puget Sound	-0.005	0.006
Fraser	Sockeye	TRS	1960-2012	Ogden et al. 2015	South Strait of Georgia	0.003	0.004
Baker Lake	Sockeye	E	1970-2015	A. Dufault, WDFW, pers. comm.	Whidbey	0.044	0.012
Misc 10 -Seattle	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm.	Central Puget Sound	0.084	0.009
Green-Duwamish River	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm.	Central Puget Sound	0.135	0.014
Misc 10e -Port Orchard	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm.	Central Puget Sound	0.022	0.007
Misc 11 -Tacoma	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm.	Central Puget Sound	0.020	0.006
Puyallup River	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm.	Central Puget Sound	0.062	0.012
Port Gamble Pens (9a)	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm.	Hood Canal	-0.016	0.009
N. Hood Canal (12)	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm.	Hood Canal	-0.015	0.008
Misc 12b -Hood Canal	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm.	Hood Canal	0.013	0.007
Quilcene River (12a)	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm.	Hood Canal	0.034	0.011
Misc 12c -Hood Canal	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm.	Hood Canal	0.063	0.006
Hoodspout Hatchery	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm.	Hood Canal	0.049	0.010
Skokomish River	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm.	Hood Canal	0.073	0.007
Misc 12d -Hood Canal	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm.	Hood Canal	0.025	0.007
Dungeness River	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm.	Juan de Fuca	0.052	0.012
Elwha River	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm.	Juan de Fuca	-0.061	0.014
Misc Strait Streams	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm.	Juan de Fuca	0.023	0.009
Grays Harbor Wild Runsize	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm.	Pacific Coast	-0.006	0.006
Willapa Bay Wild And Hatchery Escapement And Total Runsize	Chum	TRS	1980-1999	A. Dufault, WDFW, pers. comm.	Pacific Coast	-0.010	0.006
Inside South Coast (Non-Fraser)	Chum	TRS	1953-2012	Ogden et al. 2015	Pacific Coast	0.000	0.005
Hecate Lowlands	Chum	E	1954-2012	Ogden et al. 2015	Pacific Coast	-0.014	0.004



North Haida Gwaii	Chum	E	1954-2012	Ogden et al. 2015	Pacific Coast	-0.028	0.009
Portland Inlet	Chum	E	1954-2012	Ogden et al. 2015	Pacific Coast	-0.006	0.005
Rivers Inlet	Chum	E	1954-2012	Ogden et al. 2015	Pacific Coast	-0.009	0.014
Skidegate	Chum	E	1954-2012	Ogden et al. 2015	Pacific Coast	-0.013	0.006
Misc 13 -South Puget Sound	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm	South Sound	-0.018	0.050
Chambers Creek	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm	South Sound	0.004	0.032
Nisqually River	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm	South Sound	0.026	0.054
Misc 13a -Minter Creek	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm	South Sound	0.062	0.008
Misc 13b -Olympia	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm	South Sound	0.029	0.003
Nooksack River	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm	South Strait of Georgia	0.019	0.005
Samish River	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm	South Strait of Georgia	0.007	0.010
Misc 7b Streams	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm	South Strait of Georgia	0.082	0.011
Skagit River	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm	Whidbey	-0.026	0.006
Tulalip Tribe	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm	Whidbey	0.063	0.013
Snohomish River	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm	Whidbey	-0.001	0.011
Stillaguamish River	Chum	TRS	1968-2009	A. Dufault, WDFW, pers. comm	Whidbey	-0.011	0.008

19    Appendix 2. Model Sensitivity Analysis

20    Methods


21    To assess the sensitivity of the model linkages in the simulations, we calculated means and  
22    standard deviations of the weights for all linkages from the pool of accepted models. Our  
23    hypothesis was that some linkages would be more influential in model stability and that those  
24    with mean weights different than the expected mean ( $\mu=0.5$ , given assignments that were  
25    random ( $U(0,1)$ )) would provide some indication of linkage sensitivity.

26    Results

27            Our sensitivity analysis showed that most model edges (linkages between groups) were  
28    stable with regard to the weights applied in the simulation routine, with means from the pool  
29    of balanced models very close to the expected mean of 0.5 (>75% of 148 edges). However,  
30    some model linkages in the pool of accepted models had mean weights above or below the  
31    expected mean and we considered these model groups more sensitive (Fig. A2.1). Linkages  
32    among salmon traits were most sensitive, with means in the balanced models considerably  
33    higher or lower than the expected mean (e.g., Fitness to Size and Survival to Abundance, self-  
34    limiting loops on salmon traits). However, some food web components such as Ichthyoplankton  
35    and Forage Fish were also in the group of lower-than-expected model means, suggesting that  
36    certain food web pathway weights were sensitive in the model simulations. The links with  
37    bidirectional connectivity (positive feedbacks, e.g. Fitness to Size, arrows shown in royal blue in  
38    Fig. 2) tended to be the most sensitive to the weighting scheme, with lower-than-average  
39    weights resulting in balanced models. Model linkages with mean weights above the expected  
40    mean included many high trophic level linkages (e.g., Piscivorous Fish to Marine Mammals), as  
41    well as self-limiting loops (salmon traits especially), highly connected nodes (e.g. Zooplankton),  
42    and top-level predators (Marine Mammals and Piscivorous Birds). The linkages with both  
43    higher- and lower-than-expected means (that were not self-limiting loops) tended be highly  
44    connected food web components rather than anthropogenic impacts or physical drivers.

45

46



47

48

49

50

51

52

Figure A2.1. Calculated means and standard deviations from all model runs for edges (model linkages) with mean values different from the expected mean (0.5). The expected mean is denoted by the black dashed line. Positive linkages are indicated by an arrow and negative linkages are indicated by a dash and asterisk.

53    Appendix 3. Model Perturbations and References

Drivers	Variables	Reference	Invoked Perturbation
<i>Environmental</i>	Sunlight	Littell et al. 2009	↑
	Winter Storms	Littell et al. 2009	↑
	Precipitation	Mauger et al. 2015, Littell et al. 2009	↑
	Upwelling <sup>1</sup>	Mauger et al. 2015	↓
	Stratification <sup>2</sup>	Mauger et al. 2015	↑
	Temperature	Littell et al. 2009, Field et al. 2006, Hollowed et al. 2001	↑
	River Flow	Littell et al. 2009 <sup>3</sup>	↑
	Turbidity	PSEMP 2016, WADOE 2017	↓
	Dissolved Oxygen	Roberts et al. 2014	↓
<i>Production</i>	Nutrients	Roberts et al. 2014	↑
	Microplankton	Moore et al. 2015	↑
	Microbial Detritivores	PSEMP 2016	↑
	Diatoms	Brandenberger 2008, WADOE 2017	↓
<i>Food web</i>	Zooplankton <sup>4</sup>	Li et al. 2013	↓
	Gelatinous Zooplankton	Greene et al. 2015	↑
	Forage Fish	Greene et al. 2015	↓
	Ichthyoplankton	Palsson et al. 1998	↓
	Other Salmon	PFMC 2016, Figure 1	↑
	Piscivorous Fish	Palsson et al. 1998	↓
	Piscivorous Birds	Gaydos and Pearson 2011, Anderson et al. 2009	↓
	Marine Mammals	Chasco et al. 2017	↑
<i>Anthropogenic</i>	Hatcheries	Christie et al. 2012, Waples 1999	↑
	Harvest <sup>5</sup>		↑
	Habitat Loss	Puget Sound Water Quality Action Team 2002	↑
	CO2	Feely et al. 2010	↑
	Global Warming	IPCC 2014	↑
	Contaminants	O’Neill et al. 2009, Meador et al. 2006, Crecelius et al. 1995	↑
	Disease	Roon et al. 2015	↑

54    Literature Cited

<sup>1</sup> While impacts to upwelling are unknown, we invoked a decrease in upwelling because it would reduce the delivery of nutrients to Puget Sound and potentially disrupt primary production.

<sup>2</sup> Decreased mixing as a result of changes in freshwater flow could lead to increased stratification; while the exact response is unknown (see Mauger et al. 2015), we have invoked an increase in stratification because decreased mixing would likely lead to declines in primary productivity.

<sup>3</sup> River flow is projected to be higher during the winter/spring period of salmon outmigration, but lower in the summers due to decreased precipitation and higher temps. during this season (Littell et al. 2009).

<sup>4</sup> While it is unknown if total abundance of zooplankton has decreased, there is some indication that the community has shifted. We invoked a decrease in zooplankton to reflect this shift.

<sup>5</sup> Harvest has decreased over the time period we used to frame this analysis; however, we were interested in the impacts of increased harvest on salmon survival so invoked an increase in harvest as the perturbation

- Anderson, E.M., Bower, J.L., Nysewander, D.R., Evenson, J.R. and Lovvorn, J.R. 2009. Changes in avifaunal abundance in a heavily-used wintering and migration site in Puget Sound, Washington during 1966–2007. *Marine Ornithology* 37: 19-27.
- Burke, B. J., B. R. Beckman, W. T. Peterson, C. Morgan, E. A. Daly, M. Litz. 2013. Multivariate methods to forecast Pacific salmon returns. *PLoS ONE* 8(1): e54134.
- Chasco, B. I. Kaplan, A. Thomas, A. Alejandro-Gutiérrez, D.P. Noren, M.J. Ford, M.B. Hanson, J. Scordino, S.J. Jeffries, S.F. Pearson, K. Marshall, E.J. Ward. 2017. Estimates of Chinook salmon consumption in Washington State inland water by four marine mammal predators from 1970-2015. *Can. J. of Fish. and Aqua. Sci.* 10.1139/cjfas-2016-0203
- Christie, M.R., M.L. Marine, R.A. French, R.S. Waples and M.S. Blouin. 2012. Effective size of a wild salmonid population is greatly reduced by hatchery supplementation. *Heredity* 109: 254-260.
- Crecelius, E.A., V.I. Cullinan, L.F. Lefkovitz and C. Peden. 1995. Historical trends in the accumulation of chemicals in Puget Sound. 825-831, in Volume 2, Puget Sound Research '95 Proceedings, Puget Sound Water Quality Authority, Olympia.
- Feely, R. A., S. R. Alin, J. Newton, C. L. Sabine, M. Warner, A. Devol, C. Krembs, and C. Maloy. 2010. The combined effects of ocean acidification, mixing, and respiration on pH and carbonate saturation in an urbanized estuary. *Estuarine, Coastal and Shelf Science*, 88, 442-449, doi:10.1016/j.ecss.2010.05.004
- Field, D., D. Cayan, and F. Chavez. 2006. Secular warming in the California current and North Pacific. *California Cooperative Oceanic Fisheries Investigations Reports*, 47, 92-108.
- Hoekstra, J. M., K.K. Bartz, M.H. Ruckelshaus, J.M. Moslemi, and T.K. Harms. 2007. Quantitative threat analysis for management of an imperiled species: Chinook salmon (*Oncorhynchus tshawytscha*). *Ecological Applications*, 17: 2061-2073.
- Hollowed, A. B., S. R. Hare, and W. S. Wooster. 2001. Pacific Basin climate variability and patterns of Northeast Pacific marine fish production. *Progress in Oceanography*, 49, 257-282, doi:10.1016/S0079-6611(01)00026-X.
- Gaydos, J.K. and S.F. Pearson. 2011. Birds and mammals that depend on the Salish Sea: A compilation. *Northwest Naturalist*: 92:79–94.
- Intergovernmental Panel on Climate Change. 2014. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. IPCC, Geneva, Switzerland, 151 pp.

- Li, L., D. Mackas, B. Hunt, J. Schweigert, E. Pakhomov, I. Perry, M. Galbraith, T.J. Pitcher. 2013. Zooplankton communities in the Strait of Georgia, British Columbia, track large-scale climate forcing over the Pacific Ocean. *Progress in Oceanography*. doi:<http://dx.doi.org/10.1016/j.pocean.2013.05.025>
- Littell, J.S., M. McGuire Elsner, L.C. Whitely Binder, and A.K. Snover (eds). 2009. *The Washington Climate Change Impacts Assessment: Evaluating Washington's Future in a Changing Climate*. Climate Impacts Group, University of Washington, Seattle, Washington. Available at: [www.cses.washington.edu/db/pdf/wacciaexecsummary638.pdf](http://www.cses.washington.edu/db/pdf/wacciaexecsummary638.pdf)
- Mauger, G.S., J.H. Casola, H.A. Morgan, R.L. Strauch, B. Jones, B. Curry, T.M. Busch Isaksen, L. Whitely Binder, M.B. Krosby, and A.K. Snover. 2015. State of Knowledge: Climate Change in Puget Sound. Report prepared for the Puget Sound Partnership and the National Oceanic and Atmospheric Administration. Climate Impacts Group, University of Washington, Seattle. doi:10.7915/CIG93777D
- Meador, J.P., F.C. Sommers, G.M. Ylitalo, C.A. Sloan. 2006. Altered growth and related physiological response in juvenile Chinook salmon (*Oncorhynchus tshawytscha*) from dietary exposure to polycyclic aromatic hydrocarbons (PAHs). *Can. J. of Fish. and Aqua. Sci.* 63: 2364-2376.
- Moore, S.K. et al., 2011. Past trends and future scenarios for environmental conditions favoring the accumulation of paralytic shellfish toxins in Puget Sound shellfish. *Harmful Algae*, 10, 521-529.
- Moore, S.K. et al., 2015. Present-day and future climate pathways affecting the harmful algal blooms species *Alexandrium catenella* in Puget Sound, WA, USA. *Harmful Algae*, 48, 1-11. 25
- O'Neill, S.M., and West, J.E. 2009. Marine Distribution, Life History Traits, and the Accumulation of Polychlorinated Biphenyls in Chinook Salmon from Puget Sound, Washington. *Transactions of the American Fisheries Society* 138(3): 616-632.
- Pacific Fishery Management Council. 2016. Review of 2015 Ocean Salmon Fisheries: Stock Assessment and Fishery Evaluation Document for the Pacific Coast Salmon Fishery Management Plan. Pacific Fishery Management Council, Portland, Oregon. [http://www.pcouncil.org/wp-content/uploads/2016/02/Review\\_of\\_2015\\_Salmon\\_Fisheries\\_FullDocument.pdf](http://www.pcouncil.org/wp-content/uploads/2016/02/Review_of_2015_Salmon_Fisheries_FullDocument.pdf)
- Palsson, W.A., T.J. Northrup, and M.W. Baker. 1998. Puget Sound Groundfish Management Plan. Washington Department of Fish and Wildlife. Olympia, WA.
- PSEMP Marine Waters Workgroup. 2016. Puget Sound marine waters: 2015 overview. S. K. Moore, R. Wold, K. Stark, J. Bos, P. Williams, K. Dzinbal, C. Krembs and J. Newton (Eds). URL: [www.psp.wa.gov/PSEMP/PSmarinewatersoverview.php](http://www.psp.wa.gov/PSEMP/PSmarinewatersoverview.php)

Puget Sound Water Quality Action Team. 2002. *2002 Puget Sound Update: Eighth Annual Report of the Puget Sound Ambient Monitoring Program*. Puget Sound Water Quality Action Team. Olympia, Washington.

Roberts, M., T. Mohamedali, B. Sackmann, T. Khangaonkar, W. Long. 2014. Puget Sound and the Straits Dissolved Oxygen Assessment Impacts of Current and Future Human Nitrogen Sources and Climate Change through 2070. Washington State Department of Ecology, Publication No. 14-03-007. Olympia, WA.

Roon, S.R., J.D. Alexander, K.C. Jacobson, and J.L. Bartholomew. 2015. Effect of *Nanophyetus salmincola* and bacterial co-infection on mortality of juvenile Chinook salmon. *Journal of Aquatic Animal Health*, 27:4, 209-216.


Waples, R.S. 1999. Dispelling some myths about hatcheries. *Fisheries* 24: 12–21.


WADOE. 2017. Marine Waters Program, Washington Department of Ecology.  
[http://www.ecy.wa.gov/programs/eap/mar\\_wat/](http://www.ecy.wa.gov/programs/eap/mar_wat/)

Model output showing 6 model nodes of interest: salmon survival, abundance, size, residency, fitness and other salmon. Other salmon refers to the populations (chum, pink, and sockeye) which have not seen a noticeable decline in survival in recent decades. In each plot box, the model node that was perturbed is shown in the title, with the direction of the press (increase or decrease) shown below. The bar graphs indicate the proportion of model simulations with negative (red bars) and positive (blue bars) outcomes for that model node, given the invoked press perturbation. Where the bars are dark gray, there was no impact to those nodes.










Review

177  
178  
179  
180


























181  
182  
183  
184  
185

Review

<i>Drivers</i>	<i>Variables</i>	<i>Total # of Connected Nodes</i>	<i>Number of Influencing Nodes</i>	<i>Minimum Nodal Distance to Survival</i>	<i>Description</i>
<i>Environmental</i>	Sunlight	4	2	3	Solar radiation
	Winter Storms	5	1	4	Commonly occurring winter storm events
	Precipitation	5	2	3	Annual total precipitation
	Upwelling	5	2	3	Oceanographic upwelling driven by wind and currents in the coastal waters
	Stratification	8	6	3	Formation of layers in the water column resulting from ocean conditions
	Temperature	11	5	2	Water temperature within the Salish Sea (generalized, but upper portion of the water column where salmon occur)
	River Flow	4	2	2	Annual streamflow
	Turbidity	4	3	1	Relative clarity of the water within the Salish Sea
	Dissolved Oxygen	9	7	2	Amount of oxygen available in Salish Sea waters
<i>Production</i>	Nutrients	5	5	4	Total nutrients (generalized to be anthropogenic sources of N)
	Microplankton	9	6	4	Dinoflagellates, e.g. <i>Noctiluca</i> spp.
	Microbial Detritivores	7	6	3	Generalized microbes, including bacteria
	Diatoms	11	9	3	Autotrophic phytoplankton
<i>Food web</i>	Zooplankton	10	9	2	Energy-rich zooplankton (e.g., copepods, krill, amphipods)
	Gelatinous Zooplankton	6	5	3	Zooplankton including ctenophores, medusae, and salps
	Forage Fish	9	9	2	Herring, smelt, and other small-bodied fishes
	Ichthyoplankton	7	6	2	Immature stages of fish, residing in the water column
	Other Salmon	10	10	2	Chum, Pink, and Sockeye Salmon
	Piscivorous Fish	7	6	1	Any fish-eating fish; characterized by gadids and scorpaenids in the Salish Sea
	Piscivorous Birds	5	4	1	Any fish-eating bird, such as cormorants and auklets
	Marine Mammals	7	6	1	Generally harbor seals, sea lions, orcas, and dolphins
	Hatcheries	4	1	2	Production, through human intervention, of large numbers of juvenile fish through breeding programs, specifically salmon
<i>Anthropogenic</i>	Harvest	2	1	3	Catch of fish, specifically Steelhead, Coho, and Chinook salmon; generalized to include both recreational and commercial take
	Habitat Loss	5	1	2	Loss on intertidal and subtidal habitats for spawning or rearing
	CO2	5	5	4	Input of carbon dioxide via anthropogenic activities
	Global Warming	3	1	2	The general warming trend of the earth's atmosphere
	Contaminants	6	1	2	Exposure to common toxins like PCBs, PBDE, etc., as well as contaminants of emerging concern (e.g. pharmaceuticals)
	Disease	3	2	2	Exposure to diseases such as <i>Nanophyetus</i> and bacterial kidney disease
<i>Salmon Traits</i>	Residence Time	6	5	1	The amount of time an outmigrating salmon spends in the Salish Sea
	Size	6	5	1	Overall size of salmon
	Fitness	7	6	1	Overall health of salmon
	Abundance	11	4	2	Number or biomass of salmon
	Survival	8	7	-	Successful completion of the marine life stage by individuals of a population

Drivers	Variables	Invoked Perturbation	Response Variables					
			Survival	Abundance	Fitness	Size	Residence	Other Salmon
Environmental	Sunlight	↑						
	Winter Storms	↑						
	Precipitation	↑						
	Upwelling	↓						
	Stratification	↑						
	Temperature	↑						
	River Flow	↑						
	Turbidity	↓						
	Dissolved Oxygen	↓						
Production	Nutrients	↑						
	Microplankton	↑						
	Microbial Detritivores	↑						
	Diatoms	↓						
Food web	Zooplankton	↓						
	Gelatinous Zooplankton	↑						
	Forage Fish	↓						
	Ichthyoplankton	↓						
	Other Salmon	↑						
	Piscivorous Fish	↓						
	Piscivorous Birds	↓						
	Marine Mammals	↑						
	Disease	↑						
Anthropogenic	Hatcheries	↑						
	Harvest	↑						
	Habitat Loss	↑						
	CO2	↑						
	Global Warming	↑						
	Contaminants	↑						
	Disease	↑						
	Disease	↑						
Strong Neg. Effect (>80% of runs negative)								
Weak Neg. Effect (60-80% of runs negative)								
Neutral (40-60% of runs positive/negative)								
Weak Pos. Effect (60-80% of runs positive)								
Strong Pos. Effect (>80% of runs positive)								

Variable	Driver Group			
	Environmental	Primary Production	Food web	Anthropogenic
Winter Storms	↑			
Precipitation	↑			
Temperature	↑			
Dissolved Oxygen	↓			
Nutrients		↑		↑
Microplankton		↑		
Microbial Detritivores		↑		
Diatoms		↓		
Gelatinous Zooplankton			↑	
Forage Fish			↓	
Piscivorous Fish			↓	
Marine Mammals			↑	
Hatcheries				↑
Habitat Loss				↑
Contaminants				↑
Total Nodal Distance	11	14	8	11

<i>Drivers</i>	<i>Perturbations</i>	<i>References</i>	<i>South Sound</i>	<i>Hood Canal</i>	<i>Central Basin</i>
<i>Oceanographic</i>	Nutrients	Roberts et al. 2014	↑		
	Stratification	Mauger et al. 2015		↑	
	Dissolved Oxygen	Roberts et al. 2014		↓	
	Turbidity	PSEMP 2016		↓	
	Temperature	PSEMP 2016		↑	
<i>Food web</i>	Diatoms	PSEMP 2016			↓
	Gelatinous Zooplankton	Greene et al. 2015	↑		↑
	Forage Fish	Greene et al. 2015	↓		↓
	Other Salmon	Fig. 1, this paper		↑	
<i>Anthropogenic Impacts</i>	Contaminants	O'Neill et al. 2009	↑		↑
	Habitat Loss	Hoekstra et al. 2007			↑
	Hatcheries	Hoekstra et al. 2007	↑		
<i>Response Key</i>		<i>Responses</i>	<i>South Sound</i>	<i>Hood Canal</i>	<i>Central Basin</i>
Strong Neg. Effect (>80% of runs were negative)		Survival			
Weak Neg. Effect (60-80% of runs were negative)		Abundance			
Neutral (40-60% of runs positive/negative)		Fitness			
Weak Pos. Effect (60-80% of runs were positive)		Size			
Strong Pos. Effect (>80% of runs were positive)		Residency			
		Other Salmon			




Figure 1. Salmon population trends within Puget Sound (left most subbasins), the Strait of Georgia (center subbasins) and the Pacific Coast (right) for species with generally decreasing trends (Chinook, coho, and steelhead, left) and increasing or stable populations trends (chum, pink, and cockeye, right). Two abnormally high Pink salmon runs were omitted from the right panel for ease of presentation—these runs had trends of 1.05 and 0.43 and were both from Puget Sound subbasins. Data are from Washington Department of Fish and Wildlife, Pacific States Marine Fisheries Commission, Ogden et al. 2015, and Zimmerman et al. 2015.

139x66mm (300 x 300 DPI)



## Unable to Convert Image

The dimensions of this image (in pixels) are too large to be converted. For this image to convert, the total number of pixels (height x width) must be less than 40,000,000 (40 megapixels).

Figure 2. Conceptual model of the Salish Sea related to salmon survival. Model variables (shapes) represent food web components (ovals), physical drivers (rectangles), anthropogenic impacts (diamonds) and salmon traits of interest (triangles). Survival is shown as a hexagon and was the primary variable of interest.




Figure 3. Results for the primary response variables for each of the four groups of drivers. Perturbations were according to Table 3 and in all four figures, bars represent the number of runs resulting in negative (dark bars) or positive (light bars) response in the response variable.

101x58mm (300 x 300 DPI)




Figure 4. Comparison of multiple (n=10) strongly influential (top) and neutral (bottom) variables on salmon survival. The influential variables (resulting from the analysis presented in Table 2) were: CO<sub>2</sub>, contaminants, diatoms, disease, gelatinous zooplankton, habitat loss, hatcheries, other salmon, temperature, and zooplankton. The neutral variables were: global warming, forage fish, ichthyoplankton, marine mammals, microbial detritivores, nutrients, precipitation, stratification, sunlight, and upwelling. The variables in each set were perturbed simultaneously to simulate cumulative impacts and assess model response. Dark bars show negative impacts to the response variable and light bars show positive responses.

177x266mm (300 x 300 DPI)