

1. We prove that the two definitions given of lexicographic order are the same. Recall the definitions. The recursive definition:

BASIS.

1. $\epsilon < w$ for any string w other than ϵ itself.
2. If $c < d$, where c and d are characters, then for any strings w and x , we have $cw < dx$.

INDUCTION. If $w < x$ for strings w and x , then for any character c we have $cw < cx$.

The iterative definition. Let $C = c_1c_2 \cdots c_k$ and $D = d_1d_2 \cdots d_m$ be two strings. We say $C < D$ if either of the following holds:

1. That $k < m$ and for $i = 1, 2, \dots, k$ we have $c_i = d_i$.
2. For some value of $i > 0$, the first $i - 1$ characters of the two strings agree, but the i th character of the first string is less than the i th character of the second string.

We prove first that the recursive definition is the same as the iterative definition by complete induction on the number of times the recursive rule is applied to the strings.

STATEMENT $S(n)$: If it is necessary to apply the recursive rule n times to show that $w < x$, then w precedes x according to the iterative definition of 'lexicographic order'.

We say that there is a necessary number of times to apply the recursive rule to the strings to show that $w < x$. There is a minimum number, which is the lowest number of applications needed until either basis case is satisfied. There is a maximum number, which corresponds to the length of w . The minimum number here is exactly what we mean by the necessary number in the statement.


We say this as opposed to "under the recursive definition, $w < x$ after n applications of the recursive rule ...". The number n cannot be arbitrary since there is a minimum and maximum. Thus we specify that we must meet this necessary number to show that $w < x$.

BASIS. The basis is $n = 0$, that is when either basis case holds trivially. Then $w < x$ by the recursive definition. Thus rule (1) of the iterative definition holds where $w = \epsilon$, and rule (2) holds where the basis (2) applies. Therefore the basis is true.

INDUCTION. Assume that $S(i)$ is true for $0 \leq i \leq n$. We shall prove $S(n + 1)$. That is, we apply the recursive rule $n + 1$ times to show that $w < x$. Consider the $n + 1$ th application of the recursive rule, in which we took two strings cw_1 and cx_1 , where $w_1 < x_1$ is already known, and determined that $cw_1 = w < x = cx_1$. Since w_1 precedes x_1 without requiring more than n applications of the recursive rule, then the inductive hypothesis applies to both w_1 and x_1 . Therefore w_1 precedes x_1 according to the iterative definition of lexicographic order.


We now must prove that $cw_1 = w < x = cx_1$ under the iterative definition. We have that cw_1 and cx_1 are only one character longer than w_1 and x_1 . Hence in the iterative definition, we substitute k and m for $k + 1$ and $m + 1$, and rule (1) holds. For rule (2), we substitute i for $i + 1$, and thus the rule holds. Since the iterative

b) $()()()()$ is not profile-balanced, it does not satisfy rule (2).




c) $((()())()())$ is profile-balanced. By the recursive definition, we have

$$\begin{aligned} x = y = \epsilon &\rightarrow () \\ x = \epsilon, y = () &\rightarrow ()() \\ x = y = ()() &\rightarrow (()())()() \\ x = (()())()(), y = \epsilon &\rightarrow ((()())()()) \end{aligned}$$



d) $((()((()()))))$ is profile-balanced. By the recursive definition, we have

$$\begin{aligned} x = y = \epsilon &\rightarrow () \\ x = (), y = \epsilon &\rightarrow (() \\ x = \epsilon, y = (() &\rightarrow ()((() \\ x = ()(((), y = \epsilon &\rightarrow (()((() \\ x = \epsilon, y = (()((() &\rightarrow ()((()((() \\ x = ()((()(((), y = \epsilon &\rightarrow (()((()((()()) \end{aligned}$$



3. We shall prove by induction the following statement on the number of times the recursive rule of the definition of balanced parentheses is used.

STATEMENT $S(n)$: After n applications of the recursive rule of the definition of balanced parenthesis to construct a string S , then S is the string of parentheses of some arithmetic expression.

BASIS. The basis is $n = 0$, and ϵ is the string of balanced parentheses for any of the atomic operands that are arithmetic expressions. This proves the basis.

INDUCTION. Assume $S(i)$ for $0 \leq i \leq n$ and $n \geq 0$. We shall prove $S(n + 1)$. Consider the string of balanced parentheses S constructed from the $n + 1$ th application of the recursive rule on x and y . That is, $S = (x)y$, where x and y are strings of balanced parentheses, and we can apply the inductive hypothesis to both x and y . We can take arithmetic expressions E_1 and E_2 with some binary operator θ in either of the recursive rules to form $(E_1\theta E_2)$. We can also form $(-E)$ where E is also an arithmetic expression. Clearly S constitutes the string of parentheses of both expressions. This proves the inductive step. Therefore $S(n)$ for $n \geq 0$. ♦

4.

- a) $<$ is an infix binary operator.
- b) $\&$ is a prefix unary operator.
- c) $\%$ is an infix binary operator.

5. We give a “constructive” definition of the file system.

BASIS. A file is either

- 1) A regular file
- 2) A directory

INDUCTION. If F_1, \dots, F_n are files, then D is a directory containing these files.

6.

- a) We shall sieve this set. We remove all in the set for $n \geq 0$ the numbers $5n, 7n$, and $5n + 7, 5n + 7 \times 2, \dots, 5n + 7 \times 9$, and $5 + 7n, 5 \times 2 + 7n, \dots, 5 \times 9 + 7n$. The largest element not in S is 23.

b) We must prove that all integers > 23 are in S . That is, all integers $i \geq 24$ can be written in the form $5a + 7b = i$ where a and b are nonnegative. We know that $5 \times 2 + 7 \times 2 = 24$. Assume that $i \geq 24$ is in S , now we must show that $i + 1$ is in S . Write $5a + 7b = i + 1$. Then $5a + 7b - 1 \geq 24$