

Obesity, Adipose Tissue and Vascular Dysfunction

Mascha Koenen¹*, Michael A. Hill²*, Paul Cohen³, James R. Sowers

ABSTRACT: Cardiovascular diseases are the leading cause of death worldwide. Overweight and obesity are strongly associated with comorbidities such as hypertension and insulin resistance, which collectively contribute to the development of cardiovascular diseases and resultant morbidity and mortality. Forty-two percent of adults in the United States are obese, and a total of 1.9 billion adults worldwide are overweight or obese. These alarming numbers, which continue to climb, represent a major health and economic burden. Adipose tissue is a highly dynamic organ that can be classified based on the cellular composition of different depots and their distinct anatomical localization. Massive expansion and remodeling of adipose tissue during obesity differentially affects specific adipose tissue depots and significantly contributes to vascular dysfunction and cardiovascular diseases. Visceral adipose tissue accumulation results in increased immune cell infiltration and secretion of vasoconstrictor mediators, whereas expansion of subcutaneous adipose tissue is less harmful. Therefore, fat distribution more than overall body weight is a key determinant of the risk for cardiovascular diseases. Thermogenic brown and beige adipose tissue, in contrast to white adipose tissue, is associated with beneficial effects on the vasculature. The relationship between the type of adipose tissue and its influence on vascular function becomes particularly evident in the context of the heterogeneous phenotype of perivascular adipose tissue that is strongly location dependent. In this review, we address the abnormal remodeling of specific adipose tissue depots during obesity and how this critically contributes to the development of hypertension, endothelial dysfunction, and vascular stiffness. We also discuss the local and systemic roles of adipose tissue derived secreted factors and increased systemic inflammation during obesity and highlight their detrimental impact on cardiovascular health.

Key Words: adipose tissue ■ cardiovascular diseases ■ hypertension ■ obesity

The worldwide prevalence of obesity has tripled since 1975, with a parallel trend in type 2 diabetes.^{1,2} Globally, over 1.9 billion adults were overweight or obese in 2016 and >60% of people with obesity live in developing countries.³ Today, about 2 out of 3 adults (69%) are overweight or obese in the United States, and current projections suggest that nearly 50% of adults in the United States will be obese by 2030.⁴ Predictions made in 2008, estimated up to 3.3 billion individuals to become overweight and obese by 2030, if adjusted for secular trends.⁵ Nonadjusted predictions for 2030 generated by the same study predicted only 1.35 billion overweight and 573 million obese individuals for 2030,⁵ a number that was outdated already by 2016.³ While it is well documented that genetic and epigenetic factors contribute to obesity, environmental factors such as diet, physical activity, and environmental toxins also play a major role in the increased prevalence of this disorder (Figure 1). For example, the increase in obesity in the United States and

other industrialized nations is closely related to increased consumption of high fructose corn syrup and saturated fat and to reduced physical activity^{3,4} (Figure 1). Further, there is emerging evidence that consumption of high fructose corn syrup diets by pregnant women programs the offspring for the subsequent development of obesity and associated cardiometabolic and cardiovascular diseases (CVD) in later life (Figure 2).⁶ These maternal influences seem to be mediated through adverse effects of metabolic factors such as impaired insulin signaling, dyslipidemia, and altered blood supply on placental function and resultant fetal nutrition as well as epigenetic influences that originate from maternal obesity.⁶

There is considerable evidence that overweight and obesity and their comorbidities, hypertension and insulin resistance, increase CVD and overall morbidity and mortality rates.^{7–12} Indeed, a positive association even exists between a progressive increase in body mass index (BMI) within the normal and overweight range and the

Correspondence to: Paul Cohen, MD, PhD, Email pcohen@rockefeller.edu; or James R. Sowers, MD, Email sowersj@health.missouri.edu

*M. Koenen and M.A. Hill contributed equally.

For Sources of Funding and Disclosures, see page 962.

© 2021 American Heart Association, Inc.

Circulation Research is available at www.ahajournals.org/journal/res

Nonstandard Abbreviations and Acronyms

AMPKα1	5'AMP-activated protein kinase catalytic subunit alpha 1
Ang 1-7	angiotensin 1-7
Ang II	angiotensin II
ApoE	apolipoprotein E
AT	adipose tissue
BAT	brown adipose tissue
BMI	body mass index
EBPα	enhancer binding protein alpha
CVD	cardiovascular disease
EC	endothelial cells
EnNaC	endothelial Na ²⁺ channel
eNOS	endothelial nitric oxide synthase
FABP4	fatty acid-binding protein
HIF1α	hypoxia-inducible factor 1 alpha
IL	interleukin
KDM5C	lysine-specific demethylase 5C
NO	nitric oxide
PDGFRα	platelet-derived growth factor receptor alpha
PRDM16	PR domain containing 16
PVAT	perivascular adipose tissue
RAAS	renin-angiotensin-aldosterone-system
SGK-1	serum/glucocorticoid regulated kinase 1
TNFα	tumor necrosis factor alpha
UCP1	uncoupling protein 1
VSMC	vascular smooth muscle cell
WAT	white adipose tissue

risk of CVD.^{6,7} In this regard, an analysis of the Framingham Heart Study showed a positive association between overweight (BMI, 25–29.9 kg/m²) and the relative risks of hypertension and CVD.⁸ In addition, the presence of childhood obesity has been shown to increase the risk for development of type 2 diabetes, hypertension, dyslipidemia, and atherosclerosis and related CVD in adulthood.^{9–11} This review discusses the various factors that promote vascular dysfunction and CVD in obesity, with a focus on the role of dysfunctional adipose tissue.

TYPES OF ADIPOSE TISSUE

Functionally Distinct Adipose Tissue Depots in Mice and Humans

Adipose tissue (AT) is a dynamic organ distributed throughout the body with an almost unlimited capacity to expand during obesity. Several distinct depots can be defined by their location, size, cellular composition, and function. While many functions of AT are conserved

between mouse models and humans, their location and abundance can vary broadly. Mammals possess 2 major types of AT: white and brown (Figure 3). White adipose tissue (WAT) represents the largest proportion of whole-body AT and can be found around major organs and blood vessels in the abdominal cavity and subcutaneously (Figure 4). WAT stores excess energy in the form of triglycerides, and increased accumulation of WAT, particularly in visceral depots, is a key determinant of the relative risk for cardiometabolic disorders, hypertension, and CVD.^{12–17} To this point, fat distribution dictates CVD risk such that individuals with higher visceral AT and ectopic fat deposition have an increased prevalence of cardiometabolic disorders including hypertension,^{18,19} dyslipidemia, and insulin resistance^{15–17} compared with equally obese individuals with less visceral AT and relatively more subcutaneous fat. Thus, measurements limited to the determination of BMI do not reflect the actual risk for CVD conferred by obesity.

In contrast to WAT, brown AT (BAT) represents only \approx 4.3% of all AT in adult humans and can be found in cervical, supraclavicular, axillary, paraspinal, mediastinal, and abdominal depots^{20–22} (Figure 4). In addition, newborns possess interscapular BAT that decreases in size over time and is no longer detectable in adults.²³ BAT protects animals from hypothermia by dissipating energy as heat, via a process called nonshivering thermogenesis, and has more recently been found to also have anti-obesity and anti-diabetes properties and to confer broad cardiometabolic health benefits in humans.²⁴

The main functional cell type of AT is the adipocyte or fat cell. White adipocytes contain a single large lipid droplet (unilocular) and only possess a small number of mitochondria. Brown adipocytes, on the other hand, have multilocular lipid droplets and contain a large number of cristae-dense mitochondria, which uniquely express UCP1 (uncoupling protein 1) in the inner mitochondrial membrane (Figure 3). UCP1 uncouples oxidative phosphorylation from adenosine triphosphate production, ultimately resulting in the generation of heat.²⁵ More recently, several UCP1-independent thermogenic mechanisms have also been described.²⁶

In addition to developmentally preformed brown adipocytes, mice and humans also have inducible brown adipocytes, referred to as beige or brite adipocytes. These multilocular fat cells come from a distinct developmental lineage and tend to be interspersed within WAT but also express UCP1²⁷ (Figure 3). At baseline or during thermoneutrality, beige adipocytes display a more white-like phenotype with large lipid droplets and low expression of thermogenic genes,²⁸ but activation by cold exposure, beta-adrenergic stimulation, or exercise results in the robust upregulation of a thermogenic program in a process commonly called “browning.” While these cold-inducible brown-like adipocytes were first described almost 40 years ago,^{28–32} their developmental




Figure 1. Obesity, vascular stiffness, and cardiovascular disease (CVD): genetic/epigenetic and environment interactions. A food desert refers to an area with limited access to nutritious, affordable food.

origin, molecular properties, and physiological roles have only more recently been investigated. In mice, beige adipocytes are enriched within subcutaneous fat depots and are rarely detected in visceral depots. Intriguingly, due to their temperature-dependent epigenomic plasticity, beige adipocytes also have the capacity to whiten in a warm environment.³³

In light of their morphological and functional differences, it is not surprising that white and thermogenic brown/beige adipocytes are derived from distinct precursors.^{31,34–37} White adipocytes arise from mural precursors that are CD24, CD34,³⁸ and PDGFRα (platelet-derived

growth factor receptor alpha) positive,^{12,39} and subcutaneous and visceral white adipocytes seem to originate from distinct progenitor populations.⁴⁰ Developmentally preformed or classical brown fat is derived from a myogenic precursor expressing *Pax7*, *Engrailed-1*, and *Myf5* around embryonic days 9.5 to 11.5 in mice, even before white adipocytes develop.^{35,37,41} Beige adipocytes, in contrast, originate from a vascular smooth muscle lineage.⁴² Despite their distinct origins, the development of both brown and beige adipocytes is dependent on the transcriptional coregulatory protein PRDM16 (PR domain containing 16). Adult humans also have inducible




Figure 2. Prenatal programming and epigenetics in the genesis of obesity and cardiovascular disease (CVD) in offspring (Illustration credit: Ben Smith).

EC indicates endothelial cell; EnNaC, endothelial Na²⁺ channel; HFpEF, heart failure with preserved systolic function; IR, insulin receptor; MR, mineralocorticoid receptor; ROS, reactive oxygen species; and TPA, tissue plasminogen activator.

HYPERTENSION COMPENDIUM




			
	White Adipocyte	Beige Adipocyte	Brown Adipocyte
Depots	Visceral and subcutaneous and most PVAT depots	Cervical*, supraclavicular*, axillary*, paraspinal*, renal*, thoracic PVAT*, subcutaneous (only rodents)	Interscapular (human only infants), thoracic PVAT*
Function	Storage of triglycerides, endocrine (secretion of adipokines and vasoactive factors)	Thermogenesis, anti-inflammatory properties, cardioprotective	Thermogenesis, anti-inflammatory properties, cardioprotective
Progenitor	CD24 ⁺ , CD34 ⁺ , PDGFR α ⁺	Vascular smooth muscle origin	Myogenic origin Pax7 ⁺ , En-1 ⁺ , Myf5 ⁺
Changes during obesity	Hyperplasia, hypertrophy, Immune cell infiltration, secretion of vasoconstrictors	"Whitening", loss of UCP1 expression	Potentially resistant to obesity-induced inflammation
* The definition of these thermogenic AT depots in humans as brown or beige is still under debate			

Figure 3. Function and localization of different adipose tissue depots. Comparison of white, beige, and brown adipocytes in regard to their localization in specific depots in human and mice. Their major functions and progenitor cells are depicted. Major changes occurring during adipose tissue remodeling in obesity are highlighted. AT indicates adipose tissue; CD, cluster of differentiation; En-1, Engrailed-1; Myf5, myogenic factor 5; Pax7, paired box 7; PDGFR α , platelet-derived growth factor receptor alpha; PVAT, perivascular adipose tissue; and Ucp-1, uncoupling protein-1.

thermogenic adipocytes, and evidence suggests that these cells share properties with both murine brown and beige adipocytes.^{23,27,29,36,43} The relative proportion of brown versus beige adipocytes in different human depots in various contexts remains to be fully clarified.⁴⁴

Stromal Cell Composition of AT and Impact on Physiology

Although adipocytes account for most of the volume of AT, they only make up about 50% of the cellular content.^{45,46} Other cell types include immune cells such as macrophages,^{47–49} lymphocytes,^{50–53} eosinophils^{54,55} and mast cells,⁴⁹ as well as fibroblasts, adipocyte precursors, vascular cells,⁴⁵ multipotent mesenchymal stem-like cells,⁵⁶ and nerve processes.^{57,58} Visceral AT, in contrast to subcutaneous AT, tends to have a higher content of macrophages,⁴⁹ regulatory T cells,⁵² natural killer T cells,⁵¹ and eosinophils.⁵⁴ Further, visceral and subcutaneous AT display differences in angiogenesis^{59–63} and sympathetic innervation,^{58,64,65} which can modulate the propensity for energy storage versus dissipation. Finally, changes in macrophages,⁶⁶ eosinophils,^{66–68} and group 2 innate lymphoid cells⁶⁹ can regulate browning of AT.

Perivascular and Epicardial Adipose Tissue

In addition to the well-described white and brown adipose depots, AT is also located around most large blood vessels including the aorta and mesenteric vessels but not the pulmonary and brain vasculature or the

microcirculation⁷⁰ (Figure 4). Perivascular adipose tissue (PVAT) is a specialized local deposit of adipose tissue surrounding blood vessels that also provides mechanical protection and regulation of blood vessel tone.^{71–73} Ex vivo aortic ring experiments revealed a role for PVAT in the relaxation of mesenteric arteries and the thoracic aorta of rats in response to stretch-mediated.⁷⁴ The contractile response of isolated murine mesenteric arteries toward norepinephrine, on the other hand, is significantly reduced in the presence of PVAT.⁷⁵ Further, electrical field stimulation assays of mesenteric arteries demonstrated a role for sympathetic nerve activation⁷⁶ and sensory neurons⁷⁷ in the vasodilatory effects of PVAT. The anti-contractile effects of sympathetic stimulation are mediated by the stimulation of β_3 -adrenoreceptors in PVAT, and treatment with an antagonist of β_3 -adrenoreceptors reduces these effects.⁷⁶

Interestingly, PVAT is itself heterogeneous, with its phenotype strongly location-dependent.^{78–80} Because of its close proximity to the vasculature and direct contact with the adventitia,⁸¹ PVAT is thought to play a role in vascular function and pathology. PVAT surrounding the abdominal aorta and the mesenteric arteries displays a mostly white phenotype in humans⁸² and mice, with almost no UCP1 expressing thermogenic adipocytes.²⁸ On the other hand, rodent PVAT surrounding the thoracic aorta has a brown-like phenotype with multilocular adipocytes and UCP1 expression similar to classical brown adipocytes.^{83–86} This is supported by patterns of BAT detected by positron emission tomography–computed tomography in the para-aortic area and around




Figure 4. Changes in different adipose tissue depots in homeostasis and during obesity.

In states of normal body weight (left), thermogenic brown and beige adipocytes are found surrounding the thoracic aorta (perivascular adipose tissue [PVAT]) and can be detected in the cervical, supraclavicular, axillary, paraspinous, renal and epicardial area and in infants in the interscapular depot. These cells have a multilocular appearance and due to the high density of mitochondria appear brown. The abdominal aorta and mesenteric vasculature are surrounded by white adipocytes. These unilocular adipocytes are also found in visceral and subcutaneous adipose depots.

Adiponectin and angiotensin 1-7 are secreted by adipocytes and have a vasodilating effect on the vasculature. In the lean state, adipose tissue is populated with different immune cells important for homeostasis, which change dramatically during obesity. During obesity (right), T regulatory cells (Treg) are lost in visceral adipose tissue and inflammatory CD8 T cells and macrophages infiltrate the visceral, mesenteric, and to a lesser extent subcutaneous adipose depot. Thermogenic adipose tissue in proximity to the heart and the aorta downregulates thermogenic gene expression and becomes infiltrated with immune cells. Classical brown adipose tissue is potentially protected against obesity-induced immune cell infiltration. Secretion of vasodilatory factors from adipocytes are downregulated whereas leptin and Ang II (angiotensin II) are predominantly secreted, resulting in elevations in blood pressure (Illustration credit: Ben Smith). Ang 1-7 indicates angiotensin 1-7; CD, cluster of differentiation; and WAT, white adipose tissue.

the heart of humans.⁸⁷ In addition, autopsy studies of Siberian adults revealed clear UCP1 expression and multilocular and paucilocular appearance of about 40% of mediastinal periaortic vascular AT, with some individuals displaying up to 73%.⁸⁸ In mice, long-term moderate cold exposure (16°C) results in further browning of thoracic PVAT with a markedly increased expression of *Ucp1* and *Pgc1α* and β .⁸⁴

Thermogenesis of PVAT through cold exposure or genetic manipulation in mice supports a protective role of thoracic PVAT in inflammation and atherosclerosis. Overexpression of the mitochondrial membrane protein MitoNEET induces browning of WAT and thermogenic gene expression.^{89,90} *Ucp1*-driven overexpression of MitoNEET in BAT and PVAT prevented mice from an intravascular temperature drop during cold exposure and increased energy expenditure even after removal of interscapular BAT.⁹⁰ Further, cold exposure of atherosclerosis-prone ApoE (Apolipoprotein E)-deficient or ApoE-MitoNEET double-deficient mice with removed interscapular BAT

resulted in reduced atherosclerotic lesion sizes.^{84,90} In contrast, lack of PVAT in ApoE-deficient mice with an additional smooth muscle-specific deletion of PPAR γ (peroxisome proliferator-activated receptor γ) increased atherosclerotic lesions and abrogated the protective effects observed after cold exposure.⁸⁴ Although the potential contribution of cold-induced browning of WAT was not excluded, these studies imply a contribution of PVAT to whole-body thermogenesis and protection from atherosclerosis.

Several studies in humans have examined the phenotype of perivascular fat surrounding the internal thoracic arteries. While human internal thoracic artery PVAT has been reported to have a white phenotype in one study, it is important to note that 84% of the individuals examined in this study were overweight or obese, which might affect the appearance of their AT.⁹¹ Nevertheless, PVAT of human internal thoracic arteries attenuated the contractile response to the thromboxane A2/prostaglandin H2 receptor agonist U46619 and phenylephrine.⁹¹ Similar effects were observed in PVAT stripped arteries

through the transfer of PVAT-incubated supernatant.⁹¹ However, detailed analysis of human thoracic PVAT is limited due to difficulties with sample acquisition and is often isolated from patients with underlying cardiovascular complications, complicating phenotypic assessment.

Despite the close morphological relationship between tp BAT and tPVAT (thoracic PVAT) in mice, proteomics data revealed a depot-specific clustering and an only 43% overlap of their proteome on a standard diet.⁹² This is comparable to the overlap of 44% of detected proteins between tPVAT with visceral WAT or the overlap of 53% between visceral WAT and BAT, 2 very distinct depots with different functions⁹² suggesting a potentially unique PVAT composition. Interestingly, PVAT has been shown to regulate vascular tone^{83,93} through contact dependent and paracrine functions that are impaired during obesity in mice and humans.^{91,94,95} The contractile response of mesenteric arteries to norepinephrine, for example, is reduced in the presence of PVAT but compromised in diet-induced obesity.⁹⁵ Further, the expression of vasodilatory factors, such as angiotensin (1-7),⁹⁶⁻⁹⁸ adiponectin,^{75,76} and nitric oxide⁹⁹ is inhibited during obesity,^{94,95,99,100} and the expression of the vasoconstrictor Ang II (angiotensin II) is induced in PVAT.⁷⁰ Finally, a recent single-cell RNA sequencing study demonstrated the existence of 2 main clusters of mesenchymal stem/stromal cells in PVAT of the thoracic aorta of mice.¹⁰¹ One of the clusters was associated with angiogenic and adipogenic potential, whereas the other cluster was

enriched for genes associated with vascular smooth muscle cell differentiation.¹⁰¹ Transplantation of those PVAT-derived mesenchymal stem/stromal cells to a vein graft model significantly promoted neointima formation demonstrating a possible role of PVAT in vascular remodeling.¹⁰¹

PVAT is an important contributing factor to hypertension,^{18,19} endothelial dysfunction,¹⁰² and other vascular abnormalities in obesity^{71-73,94,103,104} (Figure 5). PVAT normally releases vasodilatory mediators, including adiponectin,^{75,76} and yet to be fully characterized molecules often acting on K⁺ channels, which exert an anticontractile activity and promote vascular relaxation.⁷⁰ However, in the setting of obesity and insulin resistance, oxidative stress and inflammation are increased in PVAT, thereby resulting in an increase in proinflammatory adipokines including TNF α (tumor necrosis factor alpha), and (ILs) interleukins (IL-6 and IL-8), leading to vascular insulin resistance, impaired relaxation, and vascular stiffness.⁷¹ IL-6 and TNF α also attenuate the vasodilation of mesenteric arteries ex vivo.⁹⁴ Other cytokines such as IL-18 are thought to have protective effects on PVAT and vascular function, and loss of IL-18 results in elevated blood pressure in mice associated with the whitening of thoracic PVAT.¹⁰⁵ However, the specific impact of IL-18 in PVAT needs to be addressed in AT-specific conditional knock out animals. The Framingham Offspring and Third Generation cohort studies showed that increased PVAT volume is associated with higher thoracic and abdominal




Figure 5. Effects of obesity on the vasculature which promote dysfunctional remodeling and stiffness of the vasculature (Illustration credit: Ben Smith).

Ang II indicates angiotensin II; CRP, C-reactive protein; EC, endothelial cell; ENaC, epithelial Na⁺ channel; IL, interleukin; IR, insulin receptor; MCP-1, monocyte chemoattractant protein-1; MMP, matrix metalloproteinase; MR, mineralocorticoid receptor; NO, nitric oxide; TG2, tissue transglutaminase; TGF- β , transforming growth factor- β ; TNF, tumor necrosis factor; and VSMC, vascular smooth muscle cell.

aortic dimensions and increased arterial stiffness, even after adjusting for age and CVD risk factors including BMI and visceral AT volume.¹⁰⁴

The heart is also directly associated with specific AT depots. Epicardial AT is located on the surface of the myocardium in direct contact with the coronary arteries, and pericardial AT is in contact with the pericardial sac.¹⁰⁶ Under physiological conditions, epicardial AT may supply energetic substrates to the heart and has a greater capacity for free fatty acid turnover than other visceral AT depots.¹⁰⁷ Although its cold-induced UCP1 expression does not reach levels of classical BAT,²⁸ human epicardial AT has a thermogenic phenotype and has been suggested to regulate the temperature of the myocardium.¹⁰⁸ Other studies described the portion of epicardial AT surrounding the coronary arteries in humans as a white-like depot despite the expression of some classical brown fat marker genes such as UCP1, PRDM16, and CPT1 β .¹⁰⁹ The same study found a lower expression of adipogenic marker genes PPAR γ , FABP4 (fatty acid-binding protein 4) and C/EBP α (enhancer binding protein alpha) but an increased expression of proinflammatory cytokines compared with subcutaneous AT.¹⁰⁹ This discrepancy might be explained by the reported whitening of epicardial AT after birth in humans, with only a distinct subset of multilocular UCP1 positive cells.¹¹⁰ Epicardial AT secretes polypeptides, such as adiponectin¹¹¹ and adrenomedullin,¹¹² which have cardioprotective effects, with low expression of adiponectin in epicardial AT being associated with hypertension.¹¹³ Healthy epicardial AT accounts for \approx 5% to 20% of the heart weight,¹¹⁴ and the thickness of epicardial AT is increased in hypertensive individuals.^{115–117} Under pathological conditions, epicardial AT becomes infiltrated with immune cells expressing proinflammatory genes (IL-1 β , IL-6, and TNF α)¹¹⁸ and can contribute to structural changes in the heart.^{119–121} Studies from epicardial AT derived from coronary artery bypass grafts showed significantly lower adiponectin expression compared with other visceral adipose depots and a marked increase in CD45 expression, suggesting increased immune cell infiltration compared with omental AT.¹²² Studies of mild cold exposure in humans and the analysis of epicardial AT could be beneficial to understanding the role of epicardial AT thermogenesis for CVD.¹¹⁰ Since mice do not have a comparable epicardial AT depot, a mechanistic understanding of how epicardial AT contributes to blood pressure modulation is lacking.

CARDIOVASCULAR CONSEQUENCES OF OBESITY AND ADIPOSE TISSUE DYSFUNCTION

Impact of AT on Blood Pressure Regulation

One of the central modes of blood pressure regulation is via the renin-angiotensin-aldosterone system (RAAS).

Its major bioactive component Ang II is produced from its precursor angiotensinogen by the activation of angiotensin-converting enzymes 1 and 2. Angiotensin-converting enzyme 2 can further process Ang II to generate angiotensin 1-7, which has vasodilatory properties.^{96–98} Ang II⁸³ and aldosterone are also secreted by adipocytes and can directly activate vascular smooth muscle cells (VSMCs) via the angiotensin type 1 receptor.¹²³ Ang II is a prominent regulator of vascular tone,¹²⁴ and its expression is spatially regulated in PVAT, with higher expression in mesenteric PVAT compared to thoracic PVAT.⁸³ Interestingly, studies in rats have demonstrated that fasting reduces angiotensinogen expression in visceral AT, whereas refeeding significantly induces its expression and results in elevated blood pressure.¹²⁵ A similar effect can be observed by overexpression of angiotensinogen in mice, which also results in hypertension.¹²⁶

All of the components of the RAAS are also secreted by human WAT.¹²⁷ However, there are conflicting data as to whether the basal expression of RAAS components differ in visceral and subcutaneous AT in lean individuals. One study reported a higher general expression of angiotensinogen, the precursor of Ang II, in visceral AT compared with subcutaneous AT.¹²⁸ A more recent, larger study, however, reported no changes in angiotensinogen expression between the 2 depots in lean individuals.¹²⁹ Nevertheless, visceral AT expressed higher amounts of renin, angiotensin-converting enzyme 2 and both angiotensin receptor types 1 and 2 in the same study, whereas ACE1 was not changed.¹²⁹ In rats, mesenteric PVAT expresses higher levels of Ang II and both angiotensin receptor subtypes than thermogenic thoracic PVAT.⁸³ This is in line with the reported downregulation of angiotensinogen after β -adrenergic stimulation of murine adipocytes in vitro.¹³⁰

Thermogenic brown and beige AT is considered to have protective effects on the vasculature, as individuals with detectable thermogenic AT have lower odds for hypertension and coronary artery disease relative to individuals without thermogenic AT.²⁴ Moreover, coding variants in PRDM16, the master regulator of thermogenic AT, are associated with hypertension in humans.¹³¹ Interestingly, components of the RAAS cascade can directly affect AT, and angiotensin 1-7, besides its vasodilatory actions^{96–98} also induces BAT and reduces diet-induced obesity in mice.^{132,133} Surprisingly, pharmacological activation of angiotensin receptor 2 and Ang II treatment can induce browning of subcutaneous white adipocytes in vivo and stimulation of brown precursor differentiation in vitro.^{134,135} This protective impact on BAT is assumed to be either mediated by increased sympathetic nerve activation¹³⁵ or through increased conversion of Ang II to angiotensin 1-7. Moreover, deletion of the type 1 angiotensin receptor results in increased appearance of multilocular beige adipocytes.¹³⁶ Taken together, it seems that angiotensin 1-7 and activation of the angiotensin

receptor 2 or inhibition of the type 1 angiotensin receptor can stimulate BAT, which in turn has beneficial effects on blood pressure and attenuates development of CVD. Further studies will be needed to investigate the direct impact and molecular basis of the protective impact of thermogenic AT on hypertension.

Adipose Tissue Remodeling During Obesity

Obesity results in a chronic low-grade inflammatory state in adipose tissue.^{137,138} Visceral obesity, in particular, is strongly associated with the development of CVD.^{13,14} Defining and understanding remodeling of different AT depots during obesity is thus of utmost importance to ultimately preventing deleterious sequelae. During obesity, AT can expand by either enlargement of existing adipocytes (hypertrophy) or by increasing the number of adipocytes (hyperplasia) (Figure 3), with the relative importance of either mechanism varying based on depot, sex, and age.³¹ At baseline, fed a standard diet, neither visceral nor subcutaneous AT exhibit significant new adipogenesis in adult humans or mice.^{31,139} Long-term high fat feeding of mice, on the other hand, resulted in increased adipogenesis and hypertrophy in the visceral AT, including mesenteric PVAT, whereas subcutaneous AT adapts to the higher energy intake by hypertrophy.³¹ The individual impact of hypertrophy versus hyperplasia in the development of the metabolic syndrome is still under debate.¹⁴⁰ Maximum hypertrophy in adipocytes in established obese conditions can result in the exhaustion of the lipid storing capacity in adipocytes, which in turn can induce ectopic storage of fat in other organs such as the liver, supporting the development of the metabolic syndrome.¹⁴¹ On the other hand, visceral AT, despite its ability to expand by hyperplasia is more susceptible to AT inflammation, which in turn contributes to metabolic and CVD outcomes.¹⁴² Sex-dependent differences in AT distribution have been reviewed elsewhere,^{143–145} but in short, females most often accumulate AT in the subcutaneous depot, whereas men and postmenopausal women tend to accumulate AT in central visceral depots.¹⁴³ Hormone replacement therapy in postmenopausal women prevents this central AT distribution,¹⁴⁶ highlighting the role of sex hormones in fat distribution. However, recent studies, using an elegant separation of gonadal sex and sex chromosomes demonstrated that the XX chromosomal sex results in increased weight gain independent of the gonadal sex.^{147,148} This was mediated through the X-chromosome-escaped dose-dependent expression differences of the histone demethylase KDM5C (lysine-specific demethylase 5C) in females compared with males, and lowering KDM5C levels in females to the same extend seen in males resulted in weight loss and reduced body fat content.¹⁴⁸

In obesity, the immune cell composition of different AT depots demonstrates dynamic changes^{70,142,149} (Figure 4). For example, adipose tissue macrophages

increase in obesity and their ablation improves insulin sensitivity and reduces inflammation.^{47,150–152} The recruitment⁴⁷ and proliferation¹⁵³ of proinflammatory macrophages during obesity is greater in visceral than in subcutaneous AT.^{154,155} Obesity further results in the loss of protective CD4 helper¹⁵⁶ and regulatory T cells^{52,157} and in the enrichment of CD8 T cells in visceral AT.⁵³ These variations in immune cell infiltration between visceral and subcutaneous AT result in a low-grade inflammatory environment that can contribute to CVD.^{158,159} Recently, eosinophils have gained attention for their role in promoting beige adipocyte activation,^{67,68} and their loss during obesity, especially in visceral and mesenteric AT, renders mice susceptible to diet-induced obesity⁵⁴ and abolishes the anticontractile effect of PVAT to norepinephrine.⁹⁵ However, some of these findings require further clarification and together with detailed information on PVAT immune cell content and changes during obesity are discussed elsewhere.⁷⁰

Thermogenic brown and beige fat, on the other hand, have antiobesity effects in humans,^{160,161} and depletion of UCP1 itself or ablation of UCP1 expressing thermogenic AT results in weight gain.^{162,163} In contrast to WAT, classical BAT of obese mice expresses lower levels of genes associated with immune cells, suggesting that thermogenic AT is resistant to diet-induced inflammation.⁸⁶ However, other studies have shown that macrophages¹⁶⁴ and B lymphocytes¹⁶⁵ infiltrate thermogenic AT during obesity, and together with increased inflammatory cytokines¹⁰⁹ are thought to suppress UCP1 expression in brown adipocytes.¹⁶⁴ Further, mice fed a high fat diet for 12 weeks show reduced expression of some thermogenic marker genes, and adipocytes shifted from a multilocular to an unilocular appearance with increased lipid accumulation in BAT and thoracic PVAT.⁹² The increased body and PVAT weight also impairs anticontractile effects of PVAT.⁹¹ High fat feeding further results in the upregulation of Notch1 specifically in thoracic PVAT compared with WAT or BAT.⁹² Genetic adipocyte-specific induction of Notch1 resulted in morphological changes of tPVAT comparable to high fat diet-induced effects.⁹² This is supported by another study showing that adipocyte-specific overexpression of Notch1 impairs thermogenesis and insulin sensitivity and results in whitening of classical BAT, whereas pharmacological inhibition of Notch1 results in browning of WAT and ameliorates high fat diet-induced obesity.¹⁶⁶

Remodeling of AT During Obesity and Its Impact on Blood Pressure Homeostasis

Obesity is strongly associated with the development of hypertension,¹³ a major risk factor for CVD morbidity and mortality.^{167,168} Compared with normal weight individuals, obese individuals also carry a greater risk for coronary artery calcification, carotid artery intimal media thickening,

and left ventricular hypertrophy, even after adjustment for traditional CVD risk factors.¹⁶⁹ Weight reduction significantly improves blood pressure,^{19,170,171} and therefore, suggests a direct link between AT phenotype and odds of developing CVD and hypertension. Visceral obesity in rodents and humans is particularly associated with the metabolic syndrome,¹⁷² which consists of several risk factors for CVD, including hypertension.¹⁷³ On the other hand, humans with thermogenic AT have lower odds for hypertension, coronary artery disease, and congestive heart failure, even when obese.²⁴

Angiotensinogen expression is significantly elevated in obese individuals and is also higher in visceral AT compared with subcutaneous AT^{128,174,175} (Figure 4). Interestingly, expression of Ang II is increased in subcutaneous AT in obese individuals with hypertension compared with normotensive obese individuals.¹²⁸ Diet-induced obesity did not affect angiotensinogen levels in BAT, liver, kidney, or heart in wild-type mice or in mice expressing the human angiotensinogen gene under its own promoter.¹⁷⁵ Importantly, adipocyte-specific deletion of angiotensinogen prevents increased Ang II in the circulation and blocks elevation of BP in obese mice,¹⁷⁶ suggesting a direct impact of AT-derived angiotensinogen on blood pressure. Moreover, angiotensin receptor type 1 inhibition reverses obesity-induced blood pressure elevation in rats.¹⁷⁷ Finally, angiotensinogen levels are negatively regulated by PRDM16, and deletion of PRDM16 and ablation of beige adipocytes results in increased angiotensinogen expression.^{178,179} Ablation of BAT in mice results in obesity as well as elevated blood pressure¹⁸⁰; however, whether this is a consequence of obesity induced changes in RAAS or can be directly linked to factors secreted by brown AT needs to be further determined. Aldosterone, another component of the RAAS secreted by adipocytes,¹²³ also positively correlates with BMI, and weight loss reduces serum aldosterone levels and reduces hypertension.¹⁸¹ Components of the RAAS can therefore affect VSMC and endothelial dependent regulation of vascular tone, both of which are adversely affected during obesity.

Leptin, an adipocyte-derived hormone that regulates food intake and energy expenditure, is significantly increased in obesity in mice and humans^{182,183} (Figure 4). In contrast to angiotensinogen, it may be expressed at higher levels in subcutaneous than in visceral AT,^{184–186} and its expression is correlated with adipocyte size.¹⁸⁵ Nevertheless, diet-induced obesity results in elevated leptin levels and attendant increases in heart rate and blood pressure in rodents.^{92,187,188} This induction is mediated by a leptin-stimulated increase in sympathetic nerve activity,^{189,190} and antibody blockade of leptin or inhibition of leptin receptors on hypothalamic neurons normalized blood pressure in obese rodents.¹⁸⁷ Finally, leptin-deficient mice¹⁹¹ and humans with loss of function mutations in leptin or the leptin receptor have lower blood pressure

despite severe obesity.¹⁸⁷ It is not well understood how the chronic increase of leptin in obese subjects results in leptin-resistance¹⁹² and whether this affects blood pressure. Based on the abovementioned data, reduced leptin signaling ameliorates blood pressure in mice, and therefore, leptin-resistant obesity should be beneficial in regard to blood pressure. Indeed, leptin also has some vasodilatory effects in healthy rodents, via induction of nitric oxide expression in endothelial cells (ECs)^{77,193} and in healthy humans by a mechanism independent of nitric oxide.¹⁹⁴ Further, leptin resistance was demonstrated to selectively affect neurons in the hypothalamus that regulate food intake, while affecting other neuronal circuits to a lesser extent,^{195,196} which could explain how obese individuals do not have beneficial effects on blood pressure when leptin resistant. In detail, agouti obese mice were resistant to food intake and body weight effects of systemic leptin administration but had a preserved induction of leptin-induced renal sympathetic activation.^{196,197} Similar results in diet-induced obese mice showed the preservation of leptin-induced renal sympathetic activation and blood pressure regulation despite the resistance to weight-reducing actions of leptin.¹⁸⁸

Resistin is enriched in visceral AT,¹⁹⁸ including epicardial AT¹⁹⁹ and PVAT,²⁰⁰ and is markedly increased during obesity.^{200,201} Resistin has an important role in type 2 diabetes and insulin resistance in mice.²⁰¹ In humans with type 2 diabetes, resistin expression was only elevated in combination with hypertension and not in patients without hypertension.²⁰² In hypertensive patients without type 2 diabetes, resistin levels did not correlate with blood pressure indicating a more complex connection of obesity, insulin resistance, and blood pressure regulation by resistin. In mice, resistin treatment induced hypertension through the induction of angiotensinogen.²⁰³ Finally, resistin treatment of isolated human VSMC similar to angiotensin, resulted in increased proliferation.²⁰⁴

Visfatin is also expressed in visceral AT, including PVAT,²⁰⁰ and increased through hypoxia-induced expression of HIF1 α ²⁰⁵ in obesity.²⁰⁰ Hypertensive patients have elevated serum visfatin levels²⁰⁶; however, newly diagnosed, nonobese hypertensive men did not show any association of plasma visfatin levels and hypertension.²⁰⁷ Importantly, visfatin is mostly enriched in adipose tissue macrophages in mice²⁰⁰ and humans,²⁰⁸ and therefore, its role in adipocyte specific regulation of blood pressure might be a secondary cause of increased immune cell infiltration in obesity. Nevertheless, it was shown that hypoxic conditions can induce visfatin in murine adipocyte cell lines and its adipocyte specific role in blood pressure regulation should be determined by adipocyte-specific deletion of visfatin.

Adiponectin is another endocrine factor secreted by AT that tends to be reduced during obesity^{209,210} (Figure 4). In humans, visceral adiposity inversely correlates with adiponectin secretion, whereas secretion of adiponectin by

subcutaneous AT is not affected by adiposity.²⁰⁹ Serum adiponectin levels are reduced in obese individuals with hypertension,²¹¹ and lifestyle intervention²¹² or antihypertensive therapy²¹¹ resulted in increased adiponectin levels and improved blood pressure.²¹² In addition, lower adiponectin levels correlate with the risk for development of hypertension in humans,^{213,214} independent of body fat distribution.²¹⁵ Mice on a standard diet that lack adiponectin display elevated blood pressure despite similar body weight,⁷⁶ whereas adiponectin overexpression in obese mice ameliorates elevated blood pressure.²¹⁰ To understand the direct impact of adiponectin without secondary metabolic effects such as insulin resistance, mice lacking adiponectin were fed a high salt diet. These mice developed hypertension, which could be rescued by adiponectin administration.²¹⁰ The observed elevation in blood pressure was associated with reduced eNOS (endothelial nitric oxide synthase) and prostaglandin I₂ synthase,²¹⁰ indicating a role for adiponectin in EC-mediated vasodilation.²¹⁶ Further, ex vivo stimulation of murine mesenteric arteries with norepinephrine was significantly reduced in the presence of PVAT or PVAT-derived supernatant and could be blocked by adiponectin blocking peptide or in vessels derived from adiponectin-deficient mice.⁹⁵ Adiponectin blocking peptide also blocked electrical field stimulation of mesenteric arteries depending on the presence of PVAT.⁷⁶ Adiponectin treatment of isolated mesenteric arteries stripped of PVAT restores the anticontractile effects,^{75,76} depending on the vascular large-conductance Ca²⁺-activated K⁺ channel on VSMC.⁷⁵ Finally, AMPK α 1 (5'AMP-activated protein kinase catalytic subunit alpha 1)-deficient mice secrete less adiponectin, and ex vivo stimulation of thoracic aortic rings from these mice displayed an impaired vasodilatory effect of PVAT after U46619 treatment.²¹⁷

Another factor enriched in human omental AT and detected in human serum is omentin.²¹⁸ Like adiponectin, it is reduced in obese conditions²¹⁹ and induced through weight reduction.²²⁰ In rats, omentin treatment ameliorates Ang II or noradrenalin-induced hypertension and reduces blood pressure in normotensive rats.^{221,222} Interestingly, omentin suppressed inflammatory mediators in various vascular cell types^{222–224} and induced adiponectin levels, which might result in the indirect regulation of blood pressure. This is also the case for adipolin,²²⁵ which is reduced in obese mice²²⁶ and has a protective role in vascular remodeling through the inhibition of VSMC proliferation and macrophage activation,²²⁷ and although associated with protective effects on CVD, its role in regulation of blood pressure needs to be further determined.

Several other factors secreted by different adipose tissue depots have been associated with a role in blood pressure regulation; however, functional and mechanistic proof is still sparse and will be required to understand the independent impact of those AT-derived mediators in the regulation of hypertension. IL-33

(Interleukin-33), for example, plays a pivotal role in the activation of eosinophils, and genetic loss or obesity-induced reduction of eosinophils in PVAT results in a reduced anticontractile response.⁹⁵ Further, activation of eosinophils by IL-33 treatment rescues obesity-induced high blood pressure to the level of control mice, dependent on an EC and nitric oxide synthase-mediated effect.²²⁸ Of note, patients with pulmonary hypertension showed elevated IL-33 levels,²²⁹ and deficiency of the IL-33 receptor attenuates the progression of pulmonary arterial hypertension in mice.²³⁰ Therefore, IL-33 could play a differential role in blood pressure regulation of the vasculature with and without PVAT.

Vascular Stiffening and CVD Risk

While vascular stiffening is a normal phenomenon with increasing age, obesity, and associated insulin resistance accelerates this process. To this point, a population study showed that skin-fold thickness is a predictor of arterial stiffness in hypertensive patients.²³¹ Another study found an association between abdominal obesity and increased vascular stiffness.^{232,233} Epidemiological studies have demonstrated that hyperinsulinemia or insulin resistance, as present in overweight and obese individuals, is an independent risk factor for vascular stiffening. This vascular stiffening in association with obesity and insulin resistance has been observed in all age groups, including children.^{234,235}

There is considerable evidence that the vascular stiffening that is increased in obesity is a powerful risk factor for CVD. Data from the Framingham Heart Study have established an increased incidence of CVD events with increasing weight in both men and women,⁸ and CVD has been strongly associated with vascular stiffness.^{235,236} Importantly, arterial stiffening is especially striking in obese and diabetic premenopausal females who tend to lose the normal protection afforded by female sex hormones against vascular disease and show an increase in CVD events relative to lean, nondiabetic, age-matched women.²³⁷ Indeed, vascular stiffness independently predicts heart disease, cerebrovascular disease, and renal disease, as increased vascular stiffness is significantly associated with damage to target organs such as the heart, kidney, and brain.²³⁸ For example, stiffening of central arteries increases systolic pressure and decreases diastolic pressure, resulting in increased pulse pressure and afterload leading to an increase in left ventricular mass and myocardial oxygen demand. Further, the decrease in diastolic pressure is associated with reduced coronary blood flow during diastole. These changes have been consistently associated with left ventricular remodeling and fibrosis together with left ventricular diastolic dysfunction and associated heart failure with preserved systolic function^{239,240} (Figure 5). While early detection of arterial stiffening in obese individuals certainly helps to

identify a powerful risk factor for CVD, definitive studies on the impact of weight loss on reversal of vascular stiffness have yet to be conducted.

Mechanisms in CV Stiffness With Obesity

Development of arterial stiffness is a complex process that is driven by the interaction of endocrine factors and AT-derived cytokines, as well as interactions between different vascular cellular components, the ECM (extracellular matrix), PVAT, and immune cells in the vasculature.^{6,94} The paragraphs that follow focus on mechanisms involved in CV stiffness in conditions of overnutrition and obesity. This includes a discussion of the role of vascular endothelial abnormalities, which lead to impaired eNOS activation and associated increases in vascular stiffness. We also discuss the emerging role of vascular cell-specific mineralocorticoid and insulin receptor activation in promoting endothelial stiffness via endothelial Na⁺ channel (EnNaC) activation, and the impact of a decrease in bioavailable nitric oxide (NO) in mediating vascular stiffness in diet-induced obesity (Figure 5).

Arterial stiffness in obesity is associated with structural and functional changes in the intimal, medial, and adventitial layers of the vasculature.²⁴¹ Arterial stiffness is regulated by plasma factors such as aldosterone and insulin, as well as factors derived from the different layers of the vascular wall. Moreover, interactive signaling between different cells of the vascular wall modulates structure and function of cellular and noncellular components. Increased arterial stiffness in obese and insulin resistant states has been related to mechanisms related to both EC and VSMC stiffness, leading to the use of such terms as the stiff EC syndrome^{241–243} and the smooth muscle stiffness syndrome.²⁴² In addition to the role of ECs and VSMCs, vascular adipose and immune cell dysfunction and ECM remodeling contribute to obesity-associated arterial stiffness.^{243,244}

Increased plasma insulin and aldosterone levels lead to heightened activation of vascular MRs and IRs in obesity and insulin resistance states.^{239–243} Further, a downstream mediator of mineralocorticoid and insulin receptor activation, the ion channel EnNaC, has recently been identified as a key molecular determinant of endothelial dysfunction and CV fibrosis and stiffening.^{239,243} Increased activity of EnNaC results in a number of negative consequences including stiffening of the cortical actin cytoskeleton in ECs, impaired endothelial NO release, increased oxidative stress mediated NO destruction, increased vascular permeability and stimulation of an inflammatory environment. Such endothelial alterations impact vascular function and stiffening through increases in vascular constriction

and stimulation of tissue remodeling including fibrosis. In the case of the myocardium, obesity and associated elevations in aldosterone and insulin are associated with coronary vascular endothelial stiffening and related reductions in bioavailable NO leading to heart failure with preserved systolic function.

Recent studies, conducted in female mice fed a diet high in refined carbohydrates and saturated fat showed increased endothelial and aortic stiffness, impaired endothelial-dependent vasorelaxation, aortic fibrosis, aortic oxidative stress, and increased vascular expression of EnNaC.^{239–241} To gain further insight into the vascular role played by EnNaC, we have characterized a mouse model with EC-specific deletion of the α , pore-forming, subunit of EnNaC.²⁴¹ Obesogenic diet-induced abnormalities, along with vascular and cardiac remodeling and fibrosis, were all significantly attenuated in mice with deletion of EnNaC.^{241–243} From a mechanistic standpoint, these studies showed that diet-induced obesity resulted in a heightened inflammatory response that was associated with reduced eNOS activation and NO production and bioavailability. These latter events likely emanated from increased EnNaC activity leading to polymerization of cortical actin fibers, subsequently reducing eNOS activity, and decreasing NO production leading to increased vascular stiffness (Figure 5). This research has further revealed that activation of the endothelial Na⁺ channel by aldosterone and insulin leads to endothelial cortical stiffening, impaired NO production, and subsequent vascular fibrosis and stiffening in diet induced obesity.^{244,245} Additionally, these observations in this obese mouse model also suggest that activation of the endothelial Na⁺ channel in the coronary vasculature promotes myocardial fibrosis, myocardial stiffening, and impaired diastolic relaxation and heart failure with preserved systolic function, a condition that is especially pronounced in obese and insulin resistant females.

Studies performed in epithelial cells have shown that both aldosterone and insulin increase ENaC activity via activation of the ubiquitously expressed SGK-1 (serum and glucocorticoid regulated kinase 1).²⁴⁶ Very recent work has shown that SGK-1 represents a point of convergence for insulin and aldosterone signaling in ECs.²⁴⁴ Consistent with this notion, our preliminary studies have shown that aldosterone and insulin induced increases in EnNaC activity are diminished in isolated ECs from SGK-1 global knock-out mice compared with those of wild-type controls.²⁴⁴ It is also of relevance that evidence exists in humans for SGK-1 playing an important integrative role in the development of the cardiometabolic syndrome. Specifically, an SGK-1 gain of function gene variant that exists in 5% of the population is associated with increased blood pressure and obesity²⁴⁷ and has a particularly strong effect in increasing blood pressure in states of hyperinsulinemia and obesity.²⁴⁷ Further, in rodent models, hyperinsulinism sensitizes the blood

pressure to high fructose and salt intake, an effect involving increased activity of SGK-1.²⁴⁸ Indeed, SGK-1-knock-out mice are protected against salt-induced hypertension in the context of obesity caused by a high-fat and high-fructose diet.²⁴⁸ Finally, increased SGK-1 activity in obesity and hypertension has also been demonstrated in adipocytes²⁴⁹ and immune cells.²⁵⁰ Thus, multiple lines of evidence point toward important contributions of SGK-1 signaling in promoting the cardiometabolic syndrome, vascular stiffness, and associated CVD in obesity.

In summary, obesity is increasing in prevalence, and these increases in obesity are associated with increased consumption of refined carbohydrates and saturated fat and reduced physical activity. These and other environmental factors interact with genetic and epigenetic factors to promote obesity and related CVD (Figure 1). Obesity also negates the CVD protection normally afforded in premenopausal women. The earliest sign of obesity-related CVD is impaired NO-mediated relaxation, which leads to CV stiffness. Recent studies indicate that insulin and mineralocorticoid receptor activation of the ENaC is important in the pathogenesis of CV stiffness, especially in obese females who lose the protection against CVD normally afforded in premenstrual women.

UNANSWERED QUESTIONS AND FUTURE DIRECTIONS

While recent research has highlighted key links between obesity, adipose tissue, and vascular function, a number of important unanswered questions remain. From a basic standpoint, a more complete understanding of the developmental origin and cellular and molecular components of perivascular fat is necessary. Moreover, a comprehensive inventory of the secreted polypeptides and metabolites released by adipose tissues in normal physiology and the obese state will help further illuminate how excess adiposity contributes generally to vascular dysfunction and more specifically to the pathogenesis of hypertension and vascular stiffening. Future studies will also need to uncover the role of environment, genetics, epigenetics, and the microbiome on modulating the interactions between adipose tissues and the vasculature.

ARTICLE INFORMATION

Affiliations

Laboratory of Molecular Metabolism, The Rockefeller University, New York (M.K., P.C.). Dalton Cardiovascular Research Center, University of Missouri, Columbia (M.A.H., J.R.S.). Department of Medical Pharmacology and Physiology (M.A.H., J.R.S.), Diabetes and Cardiovascular Center (J.R.S.), and Department of Medicine (J.R.S.), University of Missouri School of Medicine, Columbia.

Sources of Funding

M. Koenen was supported by the Women and Science Initiative at Rockefeller University; P. Cohen was supported by the Sinsheimer Foundation; M.A. Hill and J.R. Sowers were supported by the National Institutes of Health.

Disclosures

None.

REFERENCES

1. Afshin A, Forouzanfar MH, Reitsma MB, Sur P, Estep K, Lee A, Marczak L, Mokdad AH, Moradi-Lakeh M, Naghavi M, et al; GBD 2015 Obesity Collaborators. Health effects of overweight and obesity in 195 countries over 25 years. *N Engl J Med*. 2017;377:13–27. doi: 10.1056/NEJMoa1614362
2. Gregg EW, Shaw JE. Global health effects of overweight and obesity. *N Engl J Med*. 2017;377:80–81. doi: 10.1056/NEJMe1706095
3. World Health Organization. Obesity and Overweight. Accessed January 11, 2021. <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
4. Ward ZJ, Bleich SN, Cradock AL, Barrett JL, Giles CM, Flax C, Long MW, Gortmaker SL. Projected U.S. state-level prevalence of adult obesity and severe obesity. *N Engl J Med*. 2019;381:2440–2450. doi: 10.1056/NEJMsa1909301
5. Kelly T, Yang W, Chen CS, Reynolds K, He J. Global burden of obesity in 2005 and projections to 2030. *Int J Obes (Lond)*. 2008;32:1431–1437. doi: 10.1038/sj.ijo.2008.102
6. Guanghong J, Hill MA, Sowers JR. Maternal exposure to high fructose and offspring health. *Hypertension*. 2019;74:499–501.
7. Field AE, Coakley EH, Must A, Spadano JL, Laird N, Dietz WH, Rimm E, Colditz GA. Impact of overweight on the risk of developing common chronic diseases during a 10-year period. *Arch Intern Med*. 2001;161:1581–1586. doi: 10.1001/archinte.161.13.1581
8. Hubert HB, Feinleib M, McNamara PM, Castelli WP. Obesity as an independent risk factor for cardiovascular disease: a 26-year follow-up of participants in the Framingham Heart Study. *Circulation*. 1983;67:968–977. doi: 10.1161/01.cir.67.5.968
9. Franks PW, Hanson RL, Knowler WC, Sievers ML, Bennett PH, Looker HC. Childhood obesity, other cardiovascular risk factors, and premature death. *N Engl J Med*. 2010;362:485–493. doi: 10.1056/NEJMoa0904130
10. Juonala M, Magnussen CG, Berenson GS, Venn A, Burns TL, Sabin MA, Srinivasan SR, Daniels SR, Davis PH, Chen W, et al. Childhood adiposity, adult adiposity, and cardiovascular risk factors. *N Engl J Med*. 2011;365:1876–1885. doi: 10.1056/NEJMoa1010112
11. Cote AT, Harris KC, Panagiotopoulos C, Sandor GG, Devlin AM. Childhood obesity and cardiovascular dysfunction. *J Am Coll Cardiol*. 2013;62:1309–1319. doi: 10.1016/j.jacc.2013.07.042
12. Lee MJ, Wu Y, Fried SK. Adipose tissue heterogeneity: implication of depot differences in adipose tissue for obesity complications. *Mol Aspects Med*. 2013;34:1–11. doi: 10.1016/j.mam.2012.10.001
13. Garrison RJ, Kannel WB, Stokes J III, Castelli WP. Incidence and precursors of hypertension in young adults: the Framingham Offspring Study. *Prev Med*. 1987;16:235–251. doi: 10.1016/0091-7435(87)90087-9
14. Bhupathiraju SN, Hu FB. Epidemiology of obesity and diabetes and their cardiovascular complications. *Circ Res*. 2016;118:1723–1735.
15. Jia G, Martinez-Lemus LA, Sowers JR. Interaction of adipogenesis and angiogenesis in dietary-induced obesity. *Diabetes*. 2015;64:2326–2328. doi: 10.2337/db15-0202
16. Aroor AR, McKarns S, Demarco VG, Jia G, Sowers JR. Maladaptive immune and inflammatory pathways lead to cardiovascular insulin resistance. *Metabolism*. 2013;62:1543–1552. doi: 10.1016/j.metabol.2013.07.001
17. Piché M-E, Tchernof A, Després J-P. Obesity phenotypes, diabetes, and cardiovascular diseases. *Circ Res*. 2020;126:1477–1500.
18. DeMarco VG, Aroor AR, Sowers JR. The pathophysiology of hypertension in patients with obesity. *Nat Rev Endocrinol*. 2014;10:364–376. doi: 10.1038/nrendo.2014.44
19. Hall JE, do Carmo JM, da Silva AA, Wang Z, Hall ME. Obesity-induced hypertension. *Circ Res*. 2015;116:991–1006.
20. Virtanen KA, Lidell ME, Orava J, Hegdlin M, Westergren R, Niemi T, Taittonen M, Laine J, Savisto NJ, Enerbäck S, et al. Functional brown adipose tissue in healthy adults. *N Engl J Med*. 2009;360:1518–1525. doi: 10.1056/NEJMoa0808949
21. Leitner BP, Huang S, Brychta RJ, Duckworth CJ, Baskin AS, McGehee S, Tal I, Dieckmann W, Gupta G, Kolodny GM, et al. Mapping of human brown adipose tissue in lean and obese young men. *Proc Natl Acad Sci USA*. 2017;114:8649–8654. doi: 10.1073/pnas.1705287114
22. Zhang F, Hao G, Shao M, Nham K, An Y, Wang Q, Zhu Y, Kusminski CM, Hassan G, Gupta RK, et al. An adipose tissue atlas: an image-guided

- identification of human-like BAT and beige depots in rodents. *Cell Metab*. 2018;27:252.e3–262.e3. doi: 10.1016/j.cmet.2017.12.004
23. Lidell ME, Betz MJ, Dahlqvist Leinhard O, Heglund M, Elander L, Slawik M, Mussack T, Nilsson D, Romu T, Nuutila P, et al. Evidence for two types of brown adipose tissue in humans. *Nat Med*. 2013;19:631–634. doi: 10.1038/nm.3017
 24. Becher T, Palanisamy S, Kramer DJ, Eljalby M, Marx SJ, Wibmer AG, Butler SD, Jiang CS, Vaughan R, Schöder H, Mark A, et al. Brown adipose tissue is associated with cardiometabolic health. *Nat Med*. 2021;27:58–65.
 25. Klaus S, Casteilla L, Bouillaud F, Ricquier D. The uncoupling protein UCP: a membranous mitochondrial ion carrier exclusively expressed in brown adipose tissue. *Int J Biochem*. 1991;23:791–801. doi: 10.1016/0020-711x(91)90062-r
 26. Roesler A, Kazak L. UCP1-independent thermogenesis. *Biochem J*. 2020;477:709–725. doi: 10.1042/BCJ20190463
 27. Wu J, Boström P, Sparks LM, Ye L, Choi JH, Giang AH, Khandekar M, Virtanen KA, Nuutila P, Schaart G, et al. Beige adipocytes are a distinct type of thermogenic fat cell in mouse and human. *Cell*. 2012;150:366–376. doi: 10.1016/j.cell.2012.05.016
 28. Waldén TB, Hansen IR, Timmons JA, Cannon B, Nedergaard J. Recruited vs. nonrecruited molecular signatures of brown, "brite," and white adipose tissues. *Am J Physiol Endocrinol Metab*. 2012;302:E19–E31. doi: 10.1152/ajpendo.00249.2011
 29. Young P, Arch JR, Ashwell M. Brown adipose tissue in the parametrial fat pad of the mouse. *FEBS Lett*. 1984;167:10–14. doi: 10.1016/0014-5793(84)80822-4
 30. Orava J, Nuutila P, Lidell ME, Oikonen V, Noponen T, Viljanen T, Scheinin M, Taittonen M, Niemi T, Enerbäck S, et al. Different metabolic responses of human brown adipose tissue to activation by cold and insulin. *Cell Metab*. 2011;14:272–279. doi: 10.1016/j.cmet.2011.06.012
 31. Wang QA, Tao C, Gupta RK, Scherer PE. Tracking adipogenesis during white adipose tissue development, expansion and regeneration. *Nat Med*. 2013;19:1338–1344. doi: 10.1038/nm.3324
 32. Rosenwald M, Perdikari A, Rülcke T, Wolfrom C. Bi-directional interconversion of brite and white adipocytes. *Nat Cell Biol*. 2013;15:659–667. doi: 10.1038/ncb2740
 33. Roh HC, Tsai LTY, Shao M, Tenen D, Shen Y, Kumari M, Lyubetskaya A, Jacobs C, Dawes B, Gupta RK, et al. Warming induces significant reprogramming of Beige, but not brown, adipocyte cellular identity. *Cell Metab*. 2018;27:1121–1137.e5. doi: 10.1016/j.cmet.2018.03.005
 34. Moulin K, Truel N, André M, Arnauld E, Nibbelink M, Cousin B, Dani C, Pénicaud L, Casteilla L. Emergence during development of the white-adipocyte cell phenotype is independent of the brown-adipocyte cell phenotype. *Biochem J*. 2001;356:659–664. doi: 10.1042/bj3560659
 35. Timmons JA, Wennmalm K, Larsson O, Walden TB, Lassmann T, Petrovic N, Hamilton DL, Gimeno RE, Wahlestedt C, Baar K, et al. Myogenic gene expression signature establishes that brown and white adipocytes originate from distinct cell lineages. *Proc Natl Acad Sci USA*. 2007;104:4401–4406. doi: 10.1073/pnas.0610615104
 36. Xue B, Rim JS, Hogan JC, Coulter AA, Koza RA, Kozak LP. Genetic variability affects the development of brown adipocytes in white fat but not in interscapular brown fat. *J Lipid Res*. 2007;48:41–51. doi: 10.1194/jlr.M600287-JLR200
 37. Seale P, Bjork B, Yang W, Kajimura S, Chin S, Kuang S, Scimè A, Devarakonda S, Conroe HM, Erdjument-Bromage H, et al. PRDM16 controls a brown fat/skeletal muscle switch. *Nature*. 2008;454:961–967. doi: 10.1038/nature07182
 38. Rodeheffer MS, Birsoy K, Friedman JM. Identification of white adipocyte progenitor cells in vivo. *Cell*. 2008;135:240–249. doi: 10.1016/j.cell.2008.09.036
 39. Berry R, Rodeheffer MS. Characterization of the adipocyte cellular lineage in vivo. *Nat Cell Biol*. 2013;15:302–308. doi: 10.1038/ncb2696
 40. Gesta S, Blüher M, Yamamoto Y, Norris AW, Berndt J, Kralisch S, Boucher J, Lewis C, Kahn CR. Evidence for a role of developmental genes in the origin of obesity and body fat distribution. *Proc Natl Acad Sci USA*. 2006;103:6676–6681. doi: 10.1073/pnas.0601752103
 41. Atit R, Sgaier SK, Mohamed OA, Taketo MM, Duford D, Joyner AL, Niswander L, Conlon RA. Beta-catenin activation is necessary and sufficient to specify the dorsal dermal fate in the mouse. *Dev Biol*. 2006;296:164–176. doi: 10.1016/j.ydbio.2006.04.449
 42. Long JZ, Svensson KJ, Tsai L, Zeng X, Roh HC, Kong X, Rao RR, Lou J, Lokurkar I, Baur W, et al. A smooth muscle-like origin for beige adipocytes. *Cell Metab*. 2014;19:810–820. doi: 10.1016/j.cmet.2014.03.025
 43. Sharp LZ, Shinoda K, Ohno H, Scheel DW, Tomoda E, Ruiz L, Hu H, Wang L, Pavlova Z, Gilsanz V, et al. Human BAT possesses molecular signatures that resemble beige/brite cells. *PLoS One*. 2012;7:e49452. doi: 10.1371/journal.pone.0049452
 44. Cannon B, de Jong JMA, Fischer AW, Nedergaard J, Petrovic N. Human brown adipose tissue: classical brown rather than brite/beige? *Exp Physiol*. 2020;105:1191–1200. doi: 10.1113/EP087875
 45. Silva KR, Côrtes I, Liechocki S, Carneiro JR, Souza AA, Borojevic R, Maya-Monteiro CM, Baptista LS. Characterization of stromal vascular fraction and adipose stem cells from subcutaneous, preperitoneal and visceral morbidly obese human adipose tissue depots. *PLoS One*. 2017;12:e0174115. doi: 10.1371/journal.pone.0174115
 46. Kumar RK, Jin Y, Watts SW, Rockwell CE. Naïve, regulatory, activated, and memory immune cells co-exist in PVATs that are comparable in density to non-PVAT fats in health. *Front Physiol*. 2020;11:58.
 47. Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW Jr. Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest*. 2003;112:1796–1808. doi: 10.1172/JCI19246
 48. Lumeng CN, Bodzin JL, Saltiel AR. Obesity induces a phenotypic switch in adipose tissue macrophage polarization. *J Clin Invest*. 2007;117:175–184. doi: 10.1172/JCI29881
 49. Altintas MM, Azad A, Nayer B, Contreras G, Zais J, Faul C, Reiser J, Nayer A. Mast cells, macrophages, and crown-like structures distinguish subcutaneous from visceral fat in mice. *J Lipid Res*. 2011;52:480–488. doi: 10.1194/jlr.M011338
 50. Caspar-Bauguil S, Cousin B, Galinier A, Segafredo C, Nibbelink M, André M, Casteilla L, Pénicaud L. Adipose tissues as an ancestral immune organ: site-specific change in obesity. *FEBS Lett*. 2005;579:3487–3492. doi: 10.1016/j.febslet.2005.05.031
 51. Schipper HS, Rakhshandehroo M, van de Graaf SF, Venken K, Koppen A, Stienstra R, Prop S, Meerding J, Hamers N, Besra G, et al. Natural killer T cells in adipose tissue prevent insulin resistance. *J Clin Invest*. 2012;122:3343–3354. doi: 10.1172/JCI62739
 52. Feuerer M, Herrero L, Cipolletta D, Naaz A, Wong J, Nayer A, Lee J, Goldfine AB, Benoist C, Shoelson S, et al. Lean, but not obese, fat is enriched for a unique population of regulatory T cells that affect metabolic parameters. *Nat Med*. 2009;15:930–939. doi: 10.1038/nm.2002
 53. Nishimura S, Manabe I, Nagasaki M, Eto K, Yamashita H, Ohsugi M, Otsu M, Hara K, Ueki K, Sugiura S, et al. CD8⁺ effector T cells contribute to macrophage recruitment and adipose tissue inflammation in obesity. *Nat Med*. 2009;15:914–920. doi: 10.1038/nm.1964
 54. Wu D, Molofsky AB, Liang HE, Ricardo-Gonzalez RR, Jouihan HA, Bando JK, Chawla A, Locksley RM. Eosinophils sustain adipose alternatively activated macrophages associated with glucose homeostasis. *Science*. 2011;332:243–247. doi: 10.1126/science.1201475
 55. Brigger D, Riether C, van Brummelen R, Mosher KI, Shiu A, Ding Z, Zbären N, Gasser P, Guntern P, Yousef H, et al. Eosinophils regulate adipose tissue inflammation and sustain physical and immunological fitness in old age. *Nat Metab*. 2020;2:688–702. doi: 10.1038/s42255-020-0228-3
 56. Zuk PA, Zhu M, Ashjian P, De Ugarte DA, Huang JL, Mizuno H, Alfonso ZC, Fraser JK, Benhaim P, Hedrick MH. Human adipose tissue is a source of multipotent stem cells. *Mol Biol Cell*. 2002;13:4279–4295. doi: 10.1091/mbc.e02-02-0105
 57. Romijn JA, Fliers E. Sympathetic and parasympathetic innervation of adipose tissue: metabolic implications. *Curr Opin Clin Nutr Metab Care*. 2005;8:440–444. doi: 10.1097/01.mco.0000172586.09762.55
 58. Chi J, Wu Z, Choi CHJ, Nguyen L, Teegene S, Ackerman SE, Crane A, Marchildon F, Tessier-Lavigne M, Cohen P. Three-dimensional adipose tissue imaging reveals regional variation in beige fat biogenesis and PRDM16-dependent sympathetic neurite density. *Cell Metab*. 2018;27:226–236.e3. doi: 10.1016/j.cmet.2017.12.011
 59. Xue Y, Petrovic N, Cao R, Larsson O, Lim S, Chen S, Feldmann HM, Liang Z, Zhu Z, Nedergaard J, et al. Hypoxia-independent angiogenesis in adipose tissues during cold acclimation. *Cell Metab*. 2009;9:99–109. doi: 10.1016/j.cmet.2008.11.009
 60. Villaret A, Galitzky J, Decaunes P, Estève D, Marques MA, Sengenès C, Chiotasso P, Tchonia T, Lafontan M, Kirkland JL, et al. Adipose tissue endothelial cells from obese human subjects: differences among depots in angiogenic, metabolic, and inflammatory gene expression and cellular senescence. *Diabetes*. 2010;59:2755–2763. doi: 10.2337/db10-0398
 61. Elias I, Franckhauser S, Ferré T, Vilà L, Tafuro S, Muñoz S, Roca C, Ramos D, Pujol A, Riu E, et al. Adipose tissue overexpression of vascular endothelial growth factor protects against diet-induced obesity and insulin resistance. *Diabetes*. 2012;61:1801–1813. doi: 10.2337/db11-0832

62. Bagchi M, Kim LA, Boucher J, Walshe TE, Kahn CR, D'Amore PA. Vascular endothelial growth factor is important for brown adipose tissue development and maintenance. *FASEB J*. 2013;27:3257–3271. doi: 10.1096/fj.12-221812
63. Shimizu I, Aghajanian T, Kikuchi R, Shimizu A, Papanicolaou KN, MacLauchlan S, Maruyama S, Walsh K. Vascular rarefaction mediates whitening of brown fat in obesity. *J Clin Invest*. 2014;124:2099–2112. doi: 10.1172/JCI71643
64. Murano I, Barbatelli G, Giordano A, Cinti S. Noradrenergic parenchymal nerve fiber branching after cold acclimation correlates with brown adipocyte density in mouse adipose organ. *J Anat*. 2009;214:171–178. doi: 10.1111/j.1469-7580.2008.01001.x
65. Wang P, Loh KH, Wu M, Morgan DA, Schneeberger M, Yu X, Chi J, Kosse C, Kim D, Rahmouni K, et al. A leptin-BDNF pathway regulating sympathetic innervation of adipose tissue. *Nature*. 2020;583:839–844. doi: 10.1038/s41586-020-2527-y
66. Nguyen KD, Qiu Y, Cui X, Goh YP, Mwangi J, David T, Mukundan L, Brombacher F, Locksley RM, Chawla A. Alternatively activated macrophages produce catecholamines to sustain adaptive thermogenesis. *Nature*. 2011;480:104–108. doi: 10.1038/nature10653
67. Qiu Y, Nguyen KD, Odegaard JI, Cui X, Tian X, Locksley RM, Palmiter RD, Chawla A. Eosinophils and type 2 cytokine signaling in macrophages orchestrate development of functional beige fat. *Cell*. 2014;157:1292–1308. doi: 10.1016/j.cell.2014.03.066
68. Knights AJ, Vohralik EJ, Houweling PJ, Stout ES, Norton LJ, Alexopoulos SJ, Yik JJ, Mat Jusoh H, Olzomer EM, Bell-Anderson KS, et al. Eosinophil function in adipose tissue is regulated by Krüppel-like factor 3 (KLF3). *Nat Commun*. 2020;11:2922. doi: 10.1038/s41467-020-16758-9
69. Brestoff JR, Kim BS, Saenz SA, Stine RR, Monticelli LA, Sonnenberg GF, Thome JJ, Farber DL, Lutfy K, Seale P, et al. Group 2 innate lymphoid cells promote beiging of white adipose tissue and limit obesity. *Nature*. 2015;519:242–246. doi: 10.1038/nature14115
70. Saxton SN, Clark BJ, Withers SB, Eringa EC, Heagerty AM. Mechanistic links between obesity, diabetes, and blood pressure: role of perivascular adipose tissue. *Physiol Rev*. 2019;99:1701–1763. doi: 10.1152/physrev.00034.2018
71. Padilla J, Vieira-Potter VJ, Jia G, Sowers JR. Role of perivascular adipose tissue on vascular reactive oxygen species in type 2 diabetes: a give-and-take relationship. *Diabetes*. 2015;64:1904–1906. doi: 10.2337/db15-0096
72. Jia G, Durante W, Sowers JR. Endothelium-derived hyperpolarizing factors: a potential therapeutic target for vascular dysfunction in obesity and insulin resistance. *Diabetes*. 2016;65:2118–2120. doi: 10.2337/db16-0026
73. Jia G, Arora AR, Sowers JR. The role of mineralocorticoid receptor signaling in the cross-talk between adipose tissue and the vascular wall. *Cardiovasc Res*. 2017;113:1055–1063. doi: 10.1093/cvr/cvx097
74. Watts SW, Flood ED, Garver H, Fink GD, Roccabianca S. A new function for perivascular adipose tissue (PVAT): assistance of arterial stress relaxation. *Sci Rep*. 2020;10:1807. doi: 10.1038/s41598-020-58368-x
75. Lynch FM, Withers SB, Yao Z, Werner ME, Edwards G, Weston AH, Heagerty AM. Perivascular adipose tissue-derived adiponectin activates BKCa channels to induce anticontractile responses. *Am J Physiol Heart Circ Physiol*. 2013;304:H786–H795.
76. Saxton SN, Ryding KE, Aldous RG, Withers SB, Ohanian J, Heagerty AM. Role of sympathetic nerves and adipocyte catecholamine uptake in the vasorelaxant function of perivascular adipose tissue. *Arterioscler Thromb Vasc Biol*. 2018;38:880–891.
77. Abu Bakar H, Robert Dunn W, Daly C, Ralevic V. Sensory innervation of perivascular adipose tissue: a crucial role in artery vasodilatation and leptin release. *Cardiovasc Res*. 2017;113:962–972. doi: 10.1093/cvr/cvx062
78. Police SB, Thatcher SE, Charnigo R, Daugherty A, Cassis LA. Obesity promotes inflammation in periaortic adipose tissue and angiotensin II-induced abdominal aortic aneurysm formation. *Arterioscler Thromb Vasc Biol*. 2009;29:1458–1464.
79. Qi XY, Qu SL, Xiong WH, Rom O, Chang L, Jiang ZS. Perivascular adipose tissue (PVAT) in atherosclerosis: a double-edged sword. *Cardiovasc Diabetol*. 2018;17:134. doi: 10.1186/s12933-018-0777-x
80. Padilla J, Jenkins NT, Vieira-Potter VJ, Laughlin MH. Divergent phenotype of rat thoracic and abdominal perivascular adipose tissues. *Am J Physiol Regul Integr Comp Physiol*. 2013;304:R543–R552. doi: 10.1152/ajpregu.00567.2012
81. Eringa EC, Bakker W, Smulders YM, Serné EH, Yudkin JS, Stehouwer CD. Regulation of vascular function and insulin sensitivity by adipose tissue: focus on perivascular adipose tissue. *Microcirculation*. 2007;14:389–402. doi: 10.1080/10739680701303584
82. Kwok KH, Lam KS, Xu A. Heterogeneity of white adipose tissue: molecular basis and clinical implications. *Exp Mol Med*. 2016;48:e215. doi: 10.1038/emmm.2016.5
83. Gálvez-Prieto B, Bolbrinker J, Stucchi P, de Las Heras AJ, Merino B, Arribas S, Ruiz-Gayo M, Huber M, Wehland M, Kreutz R, et al. Comparative expression analysis of the renin-angiotensin system components between white and brown perivascular adipose tissue. *J Endocrinol*. 2008;197:55–64.
84. Chang L, Villacorta L, Li R, Hamblin M, Xu W, Dou C, Zhang J, Wu J, Zeng R, Chen YE. Loss of perivascular adipose tissue on peroxisome proliferator-activated receptor- γ deletion in smooth muscle cells impairs intravascular thermoregulation and enhances atherosclerosis. *Circulation*. 2012;126:1067–1078. doi: 10.1161/CIRCULATIONAHA.112.104489
85. Henrichot E, Juge-Aubry CE, Pernin A, Pache JC, Velevit V, Dayer JM, Meda P, Carlo Chizzolini C, Meier CA. Production of chemokines by perivascular adipose tissue. *Arterioscler Thromb Vasc Biol*. 2005;25:2594–2599.
86. Fitzgibbons TP, Kogan S, Aouadi M, Hendricks GM, Straubhaar J, Czech MP. Similarity of mouse perivascular and brown adipose tissues and their resistance to diet-induced inflammation. *Am J Physiol Heart Circ Physiol*. 2011;301:H1425–H1437. doi: 10.1152/ajpheart.00376.2011
87. Nedergaard J, Bengtsson T, Cannon B. Unexpected evidence for active brown adipose tissue in adult humans. *Am J Physiol Endocrinol Metab*. 2007;293:E444–E452. doi: 10.1152/ajpendo.00691.2006
88. Efremova A, Senzacqua M, Venema W, Isakov E, Di Vincenzo A, Zingaretti MC, Protasoni M, Thomsen M, Giordano A, Cinti S. A large proportion of mediastinal and perirenal visceral fat of Siberian adult people is formed by UCP1 immunoreactive multilocular and paucilocular adipocytes. *J Physiol Biochem*. 2020;76:185–192. doi: 10.1007/s13105-019-00721-4
89. Kusminski CM, Park J, Scherer PE. MitoNEET-mediated effects on browning of white adipose tissue. *Nat Commun*. 2014;5:3962. doi: 10.1038/ncomms4962
90. Xiong W, Zhao X, Garcia-Barrio MT, Zhang J, Lin J, Chen YE, Jiang Z, Chang L. MitoNEET in perivascular adipose tissue blunts atherosclerosis under mild cold condition in mice. *Front Physiol*. 2017;8:1032. doi: 10.3389/fphys.2017.01032
91. Gao YJ, Zeng ZH, Teoh K, Sharma AM, Abouzahr L, Cybulsky I, Lamy A, Semelhago L, Lee RM. Perivascular adipose tissue modulates vascular function in the human internal thoracic artery. *J Thorac Cardiovasc Surg*. 2005;130:1130–1136. doi: 10.1016/j.jtcvs.2005.05.028
92. Boucher JM, Ryzhova L, Harrington A, Davis-Knowlton J, Turner JE, Cooper E, Maridas D, Ryzhov S, Rosen CJ, Vary CPH, et al. Pathological conversion of mouse perivascular adipose tissue by notch activation. *Arterioscler Thromb Vasc Biol*. 2020;40:2227–2243.
93. Gu P, Xu A. Interplay between adipose tissue and blood vessels in obesity and vascular dysfunction. *Rev Endocr Metab Disord*. 2013;14:49–58. doi: 10.1007/s11154-012-9230-8
94. Greenstein AS, Khavandi K, Withers SB, Sonoyama K, Clancy O, Jeziorska M, Laing I, Yates AP, Pemberton PW, Malik RA, et al. Local inflammation and hypoxia abolish the protective anticontractile properties of perivascular fat in obese patients. *Circulation*. 2009;119:1661–1670.
95. Withers SB, Forman R, Meza-Perez S, Sorobetea D, Sitnik K, Hopwood T, Lawrence CB, Agace WW, Else KJ, Heagerty AM, et al. Eosinophils are key regulators of perivascular adipose tissue and vascular functionality. *Sci Rep*. 2017;7:44571. doi: 10.1038/srep44571
96. Brosnihan KB, Li P, Ferrario CM. Angiotensin-(1–7) dilates canine coronary arteries through kinins and nitric oxide. *Hypertension*. 1996;27:523–528. doi: 10.1161/01.hyp.27.3.523
97. Durand MJ, Zinkevich NS, Riedel M, Gutterman DD, Nasci VL, Salato VK, Hijawi JB, Reuben CF, North PE, Beyer AM. Vascular actions of angiotensin 1–7 in the human microcirculation: novel role for telomerase. *Arterioscler Thromb Vasc Biol*. 2016;36:1254–1262. doi: 10.1161/ATVBAHA.116.307518
98. Bujak-Gizycka B, Madej J, Wolkow PP, Olszanecki R, Drabik L, Rutowski J, Korb R. Measurement of angiotensin metabolites in organ bath and cell culture experiments by liquid chromatography – electrospray ionization – mass spectrometry (LC-ESI-MS). *J Physiol Pharmacol*. 2007;58:529–540.
99. Gao YJ, Lu C, Su LY, Sharma AM, Lee RM. Modulation of vascular function by perivascular adipose tissue: the role of endothelium and hydrogen peroxide. *Br J Pharmacol*. 2007;151:323–331. doi: 10.1038/sj.bjp.0707228
100. Arita Y, Kihara S, Ouchi N, Takahashi M, Maeda K, Miyagawa J, Hotta K, Shimomura I, Nakamura T, Miyaoka K, et al. Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity. *Biochem Biophys Res Commun*. 1999;257:79–83. doi: 10.1006/bbrc.1999.0255
101. Gu W, Nowak WN, Xie Y, Le Bras A, Hu Y, Deng J, Issa Bhaloo S, Lu Y, Yuan H, Fidanis E, et al. Single-cell RNA-sequencing and metabolomics

- analyses reveal the contribution of perivascular adipose tissue stem cells to vascular remodeling. *Arterioscler Thromb Vasc Biol*. 2019;39:2049–2066.
102. Jia G, Sowers JR. Endothelial dysfunction potentially interacts with impaired glucose metabolism to increase cardiovascular risk. *Hypertension*. 2014;64:1192–1193.
 103. Villacorta L, Chang L. The role of perivascular adipose tissue in vasoconstriction, arterial stiffness, and aneurysm. *Horm Mol Biol Clin Invest*. 2015;21:137–147. doi: 10.1515/hmbci-2014-0048
 104. Thanassoulis G, Massaro JM, Corsini E, Rogers I, Schlett CL, Meigs JB, Hoffmann U, O'Donnell CJ, Fox CS. Periaortic adipose tissue and aortic dimensions in the Framingham Heart Study. *J Am Heart Assoc*. 2012;1:e000885. doi: 10.1161/JAHA.112.000885
 105. Li W, Jin D, Takai S, Hayakawa T, Ogata J, Yamanishi K, Yamanishi H, Okamura H. Impaired function of aorta and perivascular adipose tissue in IL-18-deficient mice. *Am J Physiol Heart Circ Physiol*. 2019;317:H1142–H1156. doi: 10.1152/ajpheart.00813.2018
 106. Talman AH, Psaltis PJ, Cameron JD, Meredith IT, Seneviratne SK, Wong DT. Epicardial adipose tissue: far more than a fat depot. *Cardiovasc Diagn Ther*. 2014;4:416–429. doi: 10.3978/j.issn.2223-3652.2014.11.05
 107. Marchington JM, Mattacks CA, Pond CM. Adipose tissue in the mammalian heart and pericardium: structure, foetal development and biochemical properties. *Comp Biochem Physiol B*. 1989;94:225–232. doi: 10.1016/0305-0491(89)90337-4
 108. Sacks HS, Fain JN, Holman B, Cheema P, Chary A, Parks F, Karas J, Optican R, Bahouth SW, Garrett E, et al. Uncoupling protein-1 and related messenger ribonucleic acids in human epicardial and other adipose tissues: epicardial fat functioning as brown fat. *J Clin Endocrinol Metab*. 2009;94:3611–3615. doi: 10.1210/jc.2009-0571
 109. Chatterjee TK, Stoll LL, Denning GM, Harrelson A, Blomkalns AL, Idelman G, Rothenberg FG, Neltner B, Romig-Martin SA, Dickson EW, et al. Proinflammatory phenotype of perivascular adipocytes. *Circ Res*. 2009;104:541–549.
 110. Aldiss P, Davies G, Woods R, Budge H, Sacks HS, Symonds ME. 'Browning' the cardiac and peri-vascular adipose tissues to modulate cardiovascular risk. *Int J Cardiol*. 2017;228:265–274. doi: 10.1016/j.ijcard.2016.11.074
 111. Iacobellis G, Pistilli D, Gucciardo M, Leonetti F, Miraldi F, Brancaccio G, Gallo P, di Gioia CR. Adiponectin expression in human epicardial adipose tissue in vivo is lower in patients with coronary artery disease. *Cytokine*. 2005;29:251–255. doi: 10.1016/j.cyto.2004.11.002
 112. Silaghi A, Achard V, Paulmyer-Lacroix O, Scridon T, Tassistro V, Duncea I, Clément K, Dutour A, Grino M. Expression of adrenomedullin in human epicardial adipose tissue: role of coronary status. *Am J Physiol Endocrinol Metab*. 2007;293:E1443–E1450. doi: 10.1152/ajpendo.00273.2007
 113. Teixeira-Fernandez E, Eiras S, Grigorian-Shamagian L, Fernandez A, Adrio B, Gonzalez-Juanatey JR. Epicardial adipose tissue expression of adiponectin is lower in patients with hypertension. *J Hum Hypertens*. 2008;22:856–863. doi: 10.1038/jhh.2008.75
 114. Le Jemtel TH, Samson R, Ayinapudi K, Singh T, Oparil S. Epicardial adipose tissue and cardiovascular disease. *Curr Hypertens Rep*. 2019;21:36. doi: 10.1007/s11906-019-0939-6
 115. Dicker D, Atar E, Kornowski R, Bachar GN. Increased epicardial adipose tissue thickness as a predictor for hypertension: a cross-sectional observational study. *J Clin Hypertens (Greenwich)*. 2013;15:893–898. doi: 10.1111/jch.12201
 116. Eroğlu S, Sade LE, Yıldırım A, Demir O, Müderrisoğlu H. Association of epicardial adipose tissue thickness by echocardiography and hypertension. *Türk Kardiyol Dern Ars*. 2013;41:115–122. doi: 10.5543/tkda.2013.83479
 117. Austys D, Dobrovolskij A, Jablonskienė V, Dobrovolskij V, Valevičienė N, Stukas R. Epicardial adipose tissue accumulation and essential hypertension in non-obese adults. *Medicina (Kaunas)*. 2019;55:456.
 118. Tomasz M, LiFeng Z, Andrew JD, Mannion JD, Diehl JT, Arafat H, Sarov-Blat L, O'Brien S, Keiper EA, Johnson AG, et al. Human epicardial adipose tissue is a source of inflammatory mediators. *Circulation*. 2003;108:2460–2466.
 119. Iacobellis G, Ribaudo MC, Zappaterreno A, Iannucci CV, Leonetti F. Relation between epicardial adipose tissue and left ventricular mass. *Am J Cardiol*. 2004;94:1084–1087. doi: 10.1016/j.amjcard.2004.06.075
 120. Iacobellis G, Leonetti F, Singh N, M Sharma A. Relationship of epicardial adipose tissue with atrial dimensions and diastolic function in morbidly obese subjects. *Int J Cardiol*. 2007;115:272–273. doi: 10.1016/j.ijcard.2006.04.016
 121. Thanassoulis G, Massaro JM, O'Donnell CJ, Hoffmann U, Levy D, Ellinor PT, Wang TJ, Schnabel RB, Vasan RS, Fox CS, et al. Pericardial fat is associated with prevalent atrial fibrillation: the Framingham Heart Study. *Circ Arrhythm Electrophysiol*. 2010;3:345–350.
 122. Baker AR, Silva NF, Quinn DW, Harte AL, Pagano D, Bonser RS, Kumar S, McTernan PG. Human epicardial adipose tissue expresses a pathogenic profile of adipocytokines in patients with cardiovascular disease. *Cardiovasc Diabetol*. 2006;5:1. doi: 10.1186/1475-2840-5-1
 123. Nguyen Dinh Cat A, Briones AM, Callera GE, Yogi A, He Y, Montezano AC, Touyz RM. Adipocyte-derived factors regulate vascular smooth muscle cells through mineralocorticoid and glucocorticoid receptors. *Hypertension*. 2011;58:479–488. doi: 10.1161/HYPERTENSIONAHA.110.168872
 124. Nguyen Dinh Cat A, Touyz RM. A new look at the renin-angiotensin system—focusing on the vascular system. *Peptides*. 2011;32:2141–2150. doi: 10.1016/j.peptides.2011.09.010
 125. Frederick RC Jr, Kahn BB, Peach MJ, Flier JS. Tissue-specific nutritional regulation of angiotensinogen in adipose tissue. *Hypertension*. 1992;19:339–344. doi: 10.1161/01.hyp.19.4.339
 126. Massiéra F, Bloch-Faure M, Ceiler D, Murakami K, Fukamizu A, Gasc JM, Quignard-Boulange A, Negrel R, Ailhaud G, Seydoux J, et al. Adipose angiotensinogen is involved in adipose tissue growth and blood pressure regulation. *FASEB J*. 2001;15:2727–2729. doi: 10.1096/fj.01-0457fje
 127. Engeli S, Gorzelniak K, Kreutz R, Runkel N, Distler A, Sharma AM. Co-expression of renin-angiotensin system genes in human adipose tissue. *J Hypertens*. 1999;17:555–560. doi: 10.1097/00004872-199917040-00014
 128. Giacchetti G, Faloia E, Sardu C, Camilloni MA, Mariniello B, Gatti C, Garrapa GG, Guerrieri M, Mantero F. Gene expression of angiotensinogen in adipose tissue of obese patients. *Int J Obes Relat Metab Disord*. 2000;24(suppl 2):S142–S143. doi: 10.1038/sj.ijo.0801305
 129. Zhang Y, Somers KR, Becari C, Polonis K, Pfeifer MA, Allen AM, Kellogg TA, Covassin N, Singh P. Comparative expression of renin-angiotensin pathway proteins in visceral versus subcutaneous fat. *Front Physiol*. 2018;9:1370. doi: 10.3389/fphys.2018.01370
 130. Jones BH, Standridge MK, Taylor JW, Moustaid N. Angiotensinogen gene expression in adipose tissue: analysis of obese models and hormonal and nutritional control. *Am J Physiol*. 1997;273:R236–R242. doi: 10.1152/ajpregu.1997.273.1.R236
 131. Liu C, Kraja AT, Smith JA, Brody JA, Franceschini N, Bis JC, Rice K, Morrison AC, Lu Y, Weiss S, et al; CHD Exome Consortium; ExomeBP Consortium; GoT2DGenes Consortium; T2D-GENES Consortium; Myocardial Infarction Genetics and CARDIoGRAM Exome Consortium; CKDGen Consortium. Meta-analysis identifies common and rare variants influencing blood pressure and overlapping with metabolic trait loci. *Nat Genet*. 2016;48:1162–1170. doi: 10.1038/ng.3660
 132. Morimoto H, Mori J, Nakajima H, Kawabe Y, Tsuma Y, Fukuhara S, Kodo K, Ikoma K, Matoba S, Oudit GY, et al. Angiotensin 1–7 stimulates brown adipose tissue and reduces diet-induced obesity. *Am J Physiol Endocrinol Metab*. 2018;314:E131–E138. doi: 10.1152/ajpendo.00192.2017
 133. Kawabe Y, Mori J, Morimoto H, Yamaguchi M, Miyagaki S, Ota T, Tsuma Y, Fukuhara S, Nakajima H, Oudit GY, et al. ACE2 exerts anti-obesity effect via stimulating brown adipose tissue and induction of browning in white adipose tissue. *Am J Physiol Endocrinol Metab*. 2019;317:E1140–E1149. doi: 10.1152/ajpendo.00311.2019
 134. Than A, Xu S, Li R, Leow MK, Sun L, Chen P. Angiotensin type 2 receptor activation promotes browning of white adipose tissue and brown adipogenesis. *Signal Transduct Target Ther*. 2017;2:17022. doi: 10.1038/sigtrans.201722
 135. de Kloet AD, Krause EG, Scott KA, Foster MT, Herman JP, Sakai RR, Seeley RJ, Woods SC. Central angiotensin II has catabolic action at white and brown adipose tissue. *Am J Physiol Endocrinol Metab*. 2011;301:E1081–E1091. doi: 10.1152/ajpendo.00307.2011
 136. Tsukuda K, Mogi M, Iwanami J, Kanno H, Nakaoka H, Wang XL, Bai HY, Shan BS, Kukida M, Higaki A, et al. Enhancement of adipocyte browning by angiotensin II type 1 receptor blockade. *PLoS One*. 2016;11:e0167704. doi: 10.1371/journal.pone.0167704
 137. Hotamisligil GS, Shargill NS, Spiegelman BM. Adipose expression of tumor necrosis factor- α : direct role in obesity-linked insulin resistance. *Science*. 1993;259:87–91. doi: 10.1126/science.7678183
 138. Cildir G, Akincilar SC, Tergaonkar V. Chronic adipose tissue inflammation: all immune cells on the stage. *Trends Mol Med*. 2013;19:487–500. doi: 10.1016/j.molmed.2013.05.001
 139. Spalding KL, Arner E, Westermark PO, Bernard S, Buchholz BA, Bergmann O, Blomqvist L, Hoffstedt J, Näslund E, Britton T, et al. Dynamics of fat cell turnover in humans. *Nature*. 2008;453:783–787. doi: 10.1038/nature06902
 140. Goossens GH. The metabolic phenotype in obesity: fat mass, body fat distribution, and adipose tissue function. *Obes Facts*. 2017;10:207–215. doi: 10.1159/000471488

141. Moreno-Indias I, Tinahones FJ. Impaired adipose tissue expandability and lipogenic capacities as ones of the main causes of metabolic disorders. *J Diabetes Res*. 2015;2015:970375. doi: 10.1155/2015/970375
142. Guzik TJ, Skiba DS, Touyz RM, Harrison DG. The role of infiltrating immune cells in dysfunctional adipose tissue. *Cardiovasc Res*. 2017;113:1009–1023. doi: 10.1093/cvr/cvx108
143. Fried SK, Lee MJ, Karastergiou K. Shaping fat distribution: new insights into the molecular determinants of depot- and sex-dependent adipose biology. *Obesity (Silver Spring)*. 2015;23:1345–1352. doi: 10.1002/oby.21133
144. Palmer BF, Clegg DJ. The sexual dimorphism of obesity. *Mol Cell Endocrinol*. 2015;402:113–119. doi: 10.1016/j.mce.2014.11.029
145. Lee MJ, Fried SK. Sex-dependent depot differences in adipose tissue development and function; role of sex steroids. *J Obes Metab Syndr*. 2017;26:172–180. doi: 10.7570/jomes.2017.26.3.172
146. Haarbo J, Marslew U, Gotfredsen A, Christiansen C. Postmenopausal hormone replacement therapy prevents central distribution of body fat after menopause. *Metabolism*. 1991;40:1323–1326. doi: 10.1016/0026-0495(91)90037-w
147. Chen X, McClusky R, Chen J, Beaven SW, Tontonoz P, Arnold AP, Reue K. The number of x chromosomes causes sex differences in adiposity in mice. *PLoS Genet*. 2012;8:e1002709. doi: 10.1371/journal.pgen.1002709
148. Link JC, Wiese CB, Chen X, Avetisyan R, Ronquillo E, Ma F, Guo X, Yao J, Allison M, Chen YI, et al. X chromosome dosage of histone demethylase KDM5C determines sex differences in adiposity. *J Clin Invest*. 2020;130:5688–5702. doi: 10.1172/JCI140223
149. Liu R, Nikolajczyk BS. Tissue immune cells fuel obesity-associated inflammation in adipose tissue and beyond. *Front Immunol*. 2019;10:1587. doi: 10.3389/fimmu.2019.01587
150. Kanda H, Tateya S, Tamori Y, Kotani K, Hiasa K, Kitazawa R, Kitazawa S, Miyachi H, Maeda S, Egashira K, et al. MCP-1 contributes to macrophage infiltration into adipose tissue, insulin resistance, and hepatic steatosis in obesity. *J Clin Invest*. 2006;116:1494–1505. doi: 10.1172/JCI26498
151. Patsouris D, Li PP, Thapar D, Chapman J, Olefsky JM, Neels JG. Ablation of CD11c-positive cells normalizes insulin sensitivity in obese insulin resistant animals. *Cell Metab*. 2008;8:301–309. doi: 10.1016/j.cmet.2008.08.015
152. Wentworth JM, Naselli G, Brown WA, Doyle L, Phipson B, Smyth GK, Wabitsch M, O'Brien PE, Harrison LC. Pro-inflammatory CD11c+CD206+ adipose tissue macrophages are associated with insulin resistance in human obesity. *Diabetes*. 2010;59:1648–1656. doi: 10.2337/db09-0287
153. Amano SU, Cohen JL, Vangala P, Tencerova M, Nicoloso SM, Yaw JC, Shen Y, Czech MP, Aouadi M. Local proliferation of macrophages contributes to obesity-associated adipose tissue inflammation. *Cell Metab*. 2014;19:162–171. doi: 10.1016/j.cmet.2013.11.017
154. Cancellaro R, Tordjman J, Poitou C, Guilhem G, Bouillot JL, Hugol D, Coussieu C, Basdevant A, Bar Hen A, Bedossa P, et al. Increased infiltration of macrophages in omental adipose tissue is associated with marked hepatic lesions in morbid human obesity. *Diabetes*. 2006;55:1554–1561. doi: 10.2337/db06-0133
155. Murano I, Barbatelli G, Parisani V, Latini C, Muzzonigro G, Castellucci M, Cinti S. Dead adipocytes, detected as crown-like structures, are prevalent in visceral fat depots of genetically obese mice. *J Lipid Res*. 2008;49:1562–1568. doi: 10.1194/jlr.M800019-JLR200
156. Winer S, Chan Y, Paltser G, Truong D, Tsui H, Bahrami J, Dorfman R, Wang Y, Zielinski J, Mastroratti F, et al. Normalization of obesity-associated insulin resistance through immunotherapy: CD4+ T cells control glucose homeostasis. *Nat Med*. 2009;15:921–929.
157. Cipolletta D, Feuerer M, Li A, Kamei N, Lee J, Shoelson SE, Benoist C, Mathis D. PPAR- γ is a major driver of the accumulation and phenotype of adipose tissue Treg cells. *Nature*. 2012;486:549–553. doi: 10.1038/nature11132
158. Fuster JJ, Ouchi N, Gokce N, Walsh K. Obesity-induced changes in adipose tissue microenvironment and their impact on cardiovascular disease. *Circ Res*. 2016;118:1786–1807. doi: 10.1161/CIRCRESAHA.115.306885
159. Osborn O, Olefsky JM. The cellular and signaling networks linking the immune system and metabolism in disease. *Nat Med*. 2012;18:363–374. doi: 10.1038/nm.2627
160. Saito M, Okamatsu-Ogura Y, Matsushita M, Watanabe K, Yoneshiro T, Nio-Kobayashi J, Iwanaga T, Miyagawa M, Kameya T, Nakada K, et al. High incidence of metabolically active brown adipose tissue in healthy adult humans: effects of cold exposure and adiposity. *Diabetes*. 2009;58:1526–1531. doi: 10.2337/db09-0530
161. Yoneshiro T, Aita S, Matsushita M, Kayahara T, Kameya T, Kawai Y, Iwanaga T, Saito M. Recruited brown adipose tissue as an antiobesity agent in humans. *J Clin Invest*. 2013;123:3404–3408. doi: 10.1172/JCI67803
162. Lowell BB, S-Susulic V, Hamann A, Lawitts JA, Himms-Hagen J, Boyer BB, Kozak LP, Flier JS. Development of obesity in transgenic mice after genetic ablation of brown adipose tissue. *Nature*. 1993;366:740–742. doi: 10.1038/366740a0
163. Feldmann HM, Golozoubova V, Cannon B, Nedergaard J. UCP1 ablation induces obesity and abolishes diet-induced thermogenesis in mice exempt from thermal stress by living at thermoneutrality. *Cell Metab*. 2009;9:203–209. doi: 10.1016/j.cmet.2008.12.014
164. Sakamoto T, Nitta T, Maruno K, Yeh YS, Kuwata H, Tomita K, Goto T, Takahashi N, Kawada T. Macrophage infiltration into obese adipose tissues suppresses the induction of UCP1 level in mice. *Am J Physiol Endocrinol Metab*. 2016;310:E676–E687. doi: 10.1152/ajpendo.00028.2015
165. Peterson KR, Flaherty DK, Hasty AH. Obesity alters B cell and macrophage populations in brown adipose tissue. *Obesity (Silver Spring)*. 2017;25:1881–1884. doi: 10.1002/oby.21982
166. Bi P, Shan T, Liu W, Yue F, Yang X, Liang XR, Wang J, Li J, Carlesso N, Liu X, et al. Inhibition of Notch signaling promotes browning of white adipose tissue and ameliorates obesity. *Nat Med*. 2014;20:911–918. doi: 10.1038/nm.3615
167. Lawes CM, Vander Hoorn S, Rodgers A; International Society of Hypertension. Global burden of blood-pressure-related disease, 2001. *Lancet*. 2008;371:1513–1518. doi: 10.1016/S0140-6736(08)60655-8
168. Zhou D, Xi B, Zhao M, Wang L, Veeranki SP. Uncontrolled hypertension increases risk of all-cause and cardiovascular disease mortality in US adults: the NHANES III Linked Mortality Study. *Sci Rep*. 2018;8:9418.
169. Burke GL, Bertoni AG, Shea S, Tracy R, Watson KE, Blumenthal RS, Chung H, Carnethon MR. The impact of obesity on cardiovascular disease risk factors and subclinical vascular disease: the multi-ethnic study of atherosclerosis. *Arch Intern Med*. 2008;168:928–935. doi: 10.1001/archinte.168.9.928
170. Stevens VJ, Obarzanek E, Cook NR, Lee IM, Appel LJ, Smith West D, Milas NC, Mattfeldt-Beman M, Belden L, Bragg C, et al; Trials for the Hypertension Prevention Research Group. Long-term weight loss and changes in blood pressure: results of the Trials of Hypertension Prevention, phase II. *Ann Intern Med*. 2001;134:1–11. doi: 10.7326/0003-4819-134-1-200101020-00007
171. Lee DC, Sui X, Church TS, Lavie CJ, Jackson AS, Blair SN. Changes in fitness and fatness on the development of cardiovascular disease risk factors hypertension, metabolic syndrome, and hypercholesterolemia. *J Am Coll Cardiol*. 2012;59:665–672. doi: 10.1016/j.jacc.2011.11.013
172. Matsuzawa Y, Funahashi T, Nakamura T. The concept of metabolic syndrome: contribution of visceral fat accumulation and its molecular mechanism. *J Atheroscler Thromb*. 2011;18:629–639. doi: 10.5551/jat.7922
173. Huang PL. A comprehensive definition for metabolic syndrome. *Dis Model Mech*. 2009;2:231–237. doi: 10.1242/dmm.001180
174. van Harmelen V, Elizalde M, Ariapart P, Bergstedt-Lindqvist S, Reynisdottir S, Hoffstedt J, Lundkvist I, Bringman S, Arner P. The association of human adipose angiotensinogen gene expression with abdominal fat distribution in obesity. *Int J Obes Relat Metab Disord*. 2000;24:673–678. doi: 10.1038/sj.ijo.0801217
175. Rahmouni K, Mark AL, Haynes WG, Sigmund CD. Adipose depot-specific modulation of angiotensinogen gene expression in diet-induced obesity. *Am J Physiol Endocrinol Metab*. 2004;286:E891–E895. doi: 10.1152/ajpendo.00551.2003
176. Yiannikouris F, Gupta M, Putnam K, Thatcher S, Charnigo R, Rateri DL, Daugherty A, Cassis LA. Adipocyte deficiency of angiotensinogen prevents obesity-induced hypertension in male mice. *Hypertension*. 2012;60:1524–1530.
177. Boustany CM, Brown DR, Randall DC, Cassis LA. AT1-receptor antagonism reverses the blood pressure elevation associated with diet-induced obesity. *Am J Physiol Regul Integr Comp Physiol*. 2005;289:R181–R186. doi: 10.1152/ajpregu.00507.2004
178. Cohen P, Levy JD, Zhang Y, Frontini A, Kolodin DP, Svensson KJ, Lo JC, Zeng X, Ye L, Khandekar MJ, et al. Ablation of PRDM16 and beige adipose causes metabolic dysfunction and a subcutaneous to visceral fat switch. *Cell*. 2014;156:304–316. doi: 10.1016/j.cell.2013.12.021
179. Harms MJ, Lim HW, Ho Y, Shapira SN, Ishibashi J, Rajakumari S, Steger DJ, Lazar MA, Won KJ, Seale P. PRDM16 binds MED1 and controls chromatin architecture to determine a brown fat transcriptional program. *Genes Dev*. 2015;29:298–307. doi: 10.1101/gad.252734.114
180. Cittadini A, Mantzoros CS, Hampton TG, Travers KE, Katz SE, Morgan JP, Flier JS, Douglas PS. Cardiovascular abnormalities in transgenic mice with reduced brown fat: an animal model of human obesity. *Circulation*. 1999;100:2177–2183.

181. Dall'Asta C, Vedani P, Manunta P, Pizzocri P, Marchi M, Paganelli M, Folli F, Pontiroli AE. Effect of weight loss through laparoscopic gastric banding on blood pressure, plasma renin activity and aldosterone levels in morbid obesity. *Nutr Metab Cardiovasc Dis*. 2009;19:110–114. doi: 10.1016/j.numecd.2008.06.001
182. Maffei M, Halaas J, Ravussin E, Pratley RE, Lee GH, Zhang Y, Fei H, Kim S, Lallone R, Ranganathan S. Leptin levels in human and rodent: measurement of plasma leptin and ob RNA in obese and weight-reduced subjects. *Nat Med*. 1995;1:1155–1161. doi: 10.1038/nm1195-1155
183. Considine RV, Sinha MK, Heiman ML, Kriauciunas A, Stephens TW, Nyce MR, Ohannesian JP, Marco CC, McKee LJ, Bauer TL. Serum immunoreactive-leptin concentrations in normal-weight and obese humans. *N Engl J Med*. 1996;334:292–295. doi: 10.1056/NEJM199602013340503
184. Russell CD, Petersen RN, Rao SP, Ricci MR, Prasad A, Zhang Y, Brolin RE, Fried SK. Leptin expression in adipose tissue from obese humans: depot-specific regulation by insulin and dexamethasone. *Am J Physiol*. 1998;275:E507–E515. doi: 10.1152/ajpendo.1998.275.3.E507
185. Van Harmelen V, Reynisdottir S, Eriksson P, Thörne A, Hoffstedt J, Lönnqvist F, Arner P. Leptin secretion from subcutaneous and visceral adipose tissue in women. *Diabetes*. 1998;47:913–917. doi: 10.2337/diabetes.47.6.913
186. Schoof E, Stuppy A, Harig F, Carbon R, Horbach T, Stöhr W, Rascher W, Dötsch J. Comparison of leptin gene expression in different adipose tissues in children and adults. *Eur J Endocrinol*. 2004;150:579–584. doi: 10.1530/eje.0.1500579
187. Simonds SE, Pryor JT, Ravussin E, Greenway FL, Dileone R, Allen AM, Bassi J, Elmquist JK, Keogh JM, Henning E, et al. Leptin mediates the increase in blood pressure associated with obesity. *Cell*. 2014;159:1404–1416. doi: 10.1016/j.cell.2014.10.058
188. Rahmouni K, Morgan DA, Morgan GM, Mark AL, Haynes WG. Role of selective leptin resistance in diet-induced obesity hypertension. *Diabetes*. 2005;54:2012–2018. doi: 10.2337/diabetes.54.7.2012
189. Mark AL, Correia M, Morgan DA, Shaffer RA, Haynes WG. State-of-the-art-lecture: obesity-induced hypertension: new concepts from the emerging biology of obesity. *Hypertension*. 1999;33:537–541. doi: 10.1161/01.hyp.33.1.537
190. Haynes WG. Interaction between leptin and sympathetic nervous system in hypertension. *Curr Hypertens Rep*. 2000;2:311–318. doi: 10.1007/s11906-000-0015-1
191. Mark AL, Shaffer RA, Correia ML, Morgan DA, Sigmund CD, Haynes WG. Contrasting blood pressure effects of obesity in leptin-deficient ob/ob mice and agouti yellow obese mice. *J Hypertens*. 1999;17:1949–1953. doi: 10.1097/00004872-199917121-00026
192. Halaas JL, Boozer C, Blair-West J, Fidathusein N, Denton DA, Friedman JM. Physiological response to long-term peripheral and central leptin infusion in lean and obese mice. *Proc Natl Acad Sci USA*. 1997;94:8878–8883. doi: 10.1073/pnas.94.16.8878
193. Lembo G, Vecchione C, Fratta L, Marino G, Trimarco V, d'Amati G, Trimarco B. Leptin induces direct vasodilation through distinct endothelial mechanisms. *Diabetes*. 2000;49:293–297. doi: 10.2337/diabetes.49.2.293
194. Nakagawa K, Higashi Y, Sasaki S, Oshima T, Matsuura H, Chayama K. Leptin causes vasodilation in humans. *Hypertens Res*. 2002;25:161–165.
195. Bravo PE, Morse S, Borne DM, Aguilar EA, Reisin E. Leptin and hypertension in obesity. *Vasc Health Risk Manag*. 2006;2:163–169. doi: 10.2147/vhrm.2006.2.2.163
196. Mark AL. Selective leptin resistance revisited. *Am J Physiol Regul Integr Comp Physiol*. 2013;305:R566–R581. doi: 10.1152/ajpregu.00180.2013
197. Correia ML, Haynes WG, Rahmouni K, Morgan DA, Sivitz WI, Mark AL. The concept of selective leptin resistance: evidence from agouti yellow obese mice. *Diabetes*. 2002;51:439–442. doi: 10.2337/diabetes.51.2.439
198. Gabriely I, Ma XH, Yang XM, Atzmon G, Rajala MW, Berg AH, Scherer P, Rossetti L, Barzilai N. Removal of visceral fat prevents insulin resistance and glucose intolerance of aging: an adipokine-mediated process? *Diabetes*. 2002;51:2951–2958. doi: 10.2337/diabetes.51.10.2951
199. Langheim S, Dreas L, Veschini L, Maisano F, Foglieni C, Ferrarello S, Sinagra G, Zingone B, Alfieri O, Ferrero E, et al. Increased expression and secretion of resistin in epicardial adipose tissue of patients with acute coronary syndrome. *Am J Physiol Heart Circ Physiol*. 2010;298:H746–H753. doi: 10.1152/ajpheart.00617.2009
200. Park SY, Kim KH, Seo KW, Bae JU, Kim YH, Lee SJ, Lee WS, Kim CD. Resistin derived from diabetic perivascular adipose tissue up-regulates vascular expression of osteopontin via the AP-1 signalling pathway. *J Pathol*. 2014;232:87–97. doi: 10.1002/path.4286
201. Steppan CM, Bailey ST, Bhat S, Brown EJ, Banerjee RR, Wright CM, Patel HR, Ahima RS, Lazar MA. The hormone resistin links obesity to diabetes. *Nature*. 2001;409:307–312. doi: 10.1038/35053000
202. Takata Y, Osawa H, Kurata M, Kurokawa M, Yamauchi J, Ochi M, Nishida W, Okura T, Higaki J, Makino H. Hyperresistinemia is associated with coexistence of hypertension and type 2 diabetes. *Hypertension*. 2008;51:534–539.
203. Jiang Y, Lu L, Hu Y, Li Q, An C, Yu X, Shu L, Chen A, Niu C, Zhou L, et al. Resistin induces hypertension and insulin resistance in mice via a TLR4-dependent pathway. *Sci Rep*. 2016;6:22193. doi: 10.1038/srep22193
204. Calabro P, Samudio I, Willerson JT, Yeh ET. Resistin promotes smooth muscle cell proliferation through activation of extracellular signal-regulated kinase 1/2 and phosphatidylinositol 3-kinase pathways. *Circulation*. 2004;110:3335–3340. doi: 10.1161/01.CIR.0000147825.97879.E7
205. Segawa K, Fukuhara A, Hosogai N, Morita K, Okuno Y, Tanaka M, Nakagawa Y, Kihara S, Funahashi T, Komuro R, et al. Visfatin in adipocytes is upregulated by hypoxia through HIF1 α -dependent mechanism. *Biochem Biophys Res Commun*. 2006;349:875–882. doi: 10.1016/j.bbrc.2006.07.083
206. Gunes F, Akbal E, Cakir E, Akyurek O, Altunbas M, Ozbek M. Visfatin may be a novel marker for identifying stages of essential hypertension in advanced age patients. *Intern Med*. 2012;51:553–557. doi: 10.2169/INTERNALMED.51.6609
207. Dogru T, Sonmez A, Tasci I, Yilmaz MI, Erdem G, Erturk H, Bingol N, Kilic S, Ozgurtas T. Plasma visfatin levels in young male patients with uncomplicated and newly diagnosed hypertension. *J Hum Hypertens*. 2007;21:173–175. doi: 10.1038/sj.jhh.1002114
208. Curat CA, Wegner V, Sengenès C, Miranville A, Tonus C, Busse R, Bouloumié A. Macrophages in human visceral adipose tissue: increased accumulation in obesity and a source of resistin and visfatin. *Diabetologia*. 2006;49:744–747. doi: 10.1007/s00125-006-0173-z
209. Reneau J, Goldblatt M, Gould J, Kindel T, Kastenmeier A, Higgins R, Rengel LR, Schoyer K, James R, Obi B, et al. Effect of adiposity on tissue-specific adiponectin secretion. *PLoS One*. 2018;13:e0198889. doi: 10.1371/journal.pone.0198889
210. Ohashi K, Kihara S, Ouchi N, Kumada M, Fujita K, Hiuge A, Hibuse T, Ryo M, Nishizawa H, Maeda N, et al. Adiponectin replenishment ameliorates obesity-related hypertension. *Hypertension*. 2006;47:1108–1116. doi: 10.1161/01.HYP.0000222368.43759.a1
211. Yilmaz MI, Sonmez A, Caglar K, Celik T, Yenicesu M, Eyileten T, Acikel C, Oguz Y, Yavuz I, Vural A. Effect of antihypertensive agents on plasma adiponectin levels in hypertensive patients with metabolic syndrome. *Nephrology (Carlton)*. 2007;12:147–153. doi: 10.1111/j.1440-1797.2007.00764.x
212. Yin C, Chu H, Li H, Xiao Y. Plasma Sfrp5 and adiponectin levels in relation to blood pressure among obese children. *J Hum Hypertens*. 2017;31:284–291. doi: 10.1038/jhh.2016.76
213. Imatoh T, Miyazaki M, Momose Y, Tanihara S, Une H. Adiponectin levels associated with the development of hypertension: a prospective study. *Hypertens Res*. 2008;31:229–233. doi: 10.1291/hyres.31.229
214. Shatat IF, Freeman KD, Vuguin PM, Dimartino-Nardi JR, Flynn JT. Relationship between adiponectin and ambulatory blood pressure in obese adolescents. *Pediatr Res*. 2009;65:691–695. doi: 10.1203/PDR.0b013e31819ea776
215. Peri-Okonny PA, Ayers C, Maalouf N, Das SR, de Lemos JA, Berry JD, Turer AT, Neeland IJ, Scherer PE, Vongpatanasin W. Adiponectin protects against incident hypertension independent of body fat distribution: observations from the Dallas Heart Study. *Diabetes Metab Res Rev*. 2017;33.
216. Ouchi N, Ohishi M, Kihara S, Funahashi T, Nakamura T, Nagaretani H, Kumada M, Ohashi K, Okamoto Y, Nishizawa H, et al. Association of hypoadiponectinemia with impaired vasoreactivity. *Hypertension*. 2003;42:231–234.
217. Almbrouk TAM, Ugusman AB, Katwan OJ, Salt IP, Kennedy S. Deletion of AMPK α 1 attenuates the anticontractile effect of perivascular adipose tissue (PVAT) and reduces adiponectin release. *Br J Pharmacol*. 2017;174:3398–3410. doi: 10.1111/bph.13633
218. Yang RZ, Lee MJ, Hu H, Pray J, Wu HB, Hansen BC, Shuldiner AR, Fried SK, McLenithan JC, Gong DW. Identification of omentin as a novel depot-specific adipokine in human adipose tissue: possible role in modulating insulin action. *Am J Physiol Endocrinol Metab*. 2006;290:E1253–E1261. doi: 10.1152/ajpendo.00572.2004
219. de Souza Batista CM, Yang RZ, Lee MJ, Glynn NM, Yu DZ, Pray J, Ndubuizu K, Patil S, Schwartz A, Kligen M, et al. Omentin plasma levels and gene expression are decreased in obesity. *Diabetes*. 2007;56:1655–1661. doi: 10.2337/db06-1506

220. Moreno-Navarrete JM, Catalán V, Ortega F, Gómez-Ambrosi J, Ricart W, Frühbeck G, Fernández-Real JM. Circulating omentin concentration increases after weight loss. *Nutr Metab (Lond)*. 2010;7:27. doi: 10.1186/1743-7075-7-27
221. Kazama K, Okada M, Hara Y, Yamawaki H. A novel adipocytokine, omentin, inhibits agonists-induced increases of blood pressure in rats. *J Vet Med Sci*. 2013;75:1029–1034. doi: 10.1292/jvms.12-0537
222. Brunetti L, Leone S, Orlando G, Ferrante C, Recinella L, Chiavaroli A, Di Nisio C, Shohreh R, Manippa F, Ricciuti A, et al. Hypotensive effects of omentin-1 related to increased adiponectin and decreased interleukin-6 in intra-thoracic pericardial adipose tissue. *Pharmacol Rep*. 2014;66:991–995. doi: 10.1016/j.pharep.2014.06.014
223. Ohashi K, Shibata R, Murohara T, Ouchi N. Role of anti-inflammatory adipokines in obesity-related diseases. *Trends Endocrinol Metab*. 2014;25:348–355. doi: 10.1016/j.tem.2014.03.009
224. Zabetian-Targhi F, Mirzaei K, Keshavarz SA, Hossein-Nezhad A. Modulatory role of Omentin-1 in inflammation: cytokines and dietary intake. *J Am Coll Nutr*. 2016;35:670–678. doi: 10.1080/07315724.2015.1126207
225. Sargolzaei J, Chamani E, Kazemi T, Fallah S, Soori H. The role of adiponectin and adipolin as anti-inflammatory adipokines in the formation of macrophage foam cells and their association with cardiovascular diseases. *Clin Biochem*. 2018;54:1–10. doi: 10.1016/j.clinbiochem.2018.02.008
226. Enomoto T, Ohashi K, Shibata R, Higuchi A, Maruyama S, Izumiya I, Walsh K, Murohara T, Ouchi N. Adipolin/C1qdc2/CTRP12 protein functions as an adipokine that improves glucose metabolism*. *J Biol Chem*. 2011;286:34552–34558.
227. Ogawa H, Ohashi K, Ito M, Shibata R, Kanemura N, Yuasa D, Kambara T, Matsuo K, Hayakawa S, Hiramatsu-Ito M, et al. Adipolin/CTRP12 protects against pathological vascular remodelling through suppression of smooth muscle cell growth and macrophage inflammatory response. *Cardiovasc Res*. 2020;116:237–249. doi: 10.1093/cvr/cvz074
228. Saxton SN, Whitley AS, Potter RJ, Withers SB, Grens R, Heagerty AM. Interleukin-33 rescues perivascular adipose tissue anticontractile function in obesity. *Am J Physiol Heart Circ Physiol*. 2020;319:H1387–H1397. doi: 10.1152/ajpheart.00491.2020
229. Titone D, Aroonsakool N, Li J, et al. Increased serum interleukin-33 in patients with pulmonary arterial hypertension: a role for IL-33/ST2 in disease pathogenesis. In: B95. NOVEL THERAPEUTIC TARGETS IN PULMONARY HYPERTENSION: INSIGHTS FROM TRANSLATIONAL AND PRECLINICAL STUDIES. American Thoracic Society International Conference Abstracts. American Thoracic Society; 2014:A3636–A3636.
230. Gutierrez AK. The Role of Interleukin-33 in the Progression of Pulmonary Arterial Hypertension Through an ST2/MyD88 Pathway. 2018. Accessed January 11, 2021. <https://escholarship.org/uc/item/52d491rq>
231. Selcuk A, Bulucu F, Kalafat F, Cakar M, Demirbas S, Karaman M, Ay SA, Saglam K, Balta S, Demirkol S, et al. Skinfold thickness as a predictor of arterial stiffness: obesity and fatness linked to higher stiffness measurements in hypertensive patients. *Clin Exp Hypertens*. 2013;35:459–464. doi: 10.3109/10641963.2012.746357
232. Strasser B, Arvandi M, Pasha EP, Haley AP, Stanforth P, Tanaka H. Abdominal obesity is associated with arterial stiffness in middle-aged adults. *Nutr Metab Cardiovasc Dis*. 2015;25:495–502. doi: 10.1016/j.numecd.2015.01.002
233. Sutton-Tyrrell K, Newman A, Simonsick EM, Havlik R, Pahor M, Lakatta E, Spurgeon H, Vaitkevicius P. Aortic stiffness is associated with visceral adiposity in older adults enrolled in the study of health, aging, and body composition. *Hypertension*. 2001;38:429–433. doi: 10.1161/01.hyp.38.3.429
234. Tounian P, Aggoun Y, Dubern B, Varille V, Guy-Grand B, Sidi D, Girardet JP, Bonnet D. Presence of increased stiffness of the common carotid artery and endothelial dysfunction in severely obese children: a prospective study. *Lancet*. 2001;358:1400–1404. doi: 10.1016/S0140-6736(01)06525-4
235. Cote AT, Phillips AA, Harris KC, Sandor GG, Panagiotopoulos C, Devlin AM. Obesity and arterial stiffness in children: systematic review and meta-analysis. *Arterioscler Thromb Vasc Biol*. 2015;35:1038–1044. doi: 10.1161/ATVBAHA.114.305062
236. Mitchell GF, Hwang SJ, Vasan RS, Larson MG, Pencina MJ, Hamburg NM, Vita JA, Levy D, Benjamin EJ. Arterial stiffness and cardiovascular events: the Framingham Heart Study. *Circulation*. 2010;121:505–511. doi: 10.1161/CIRCULATIONAHA.109.886655
237. Kim HL, Lee JM, Seo JB, Chung WY, Kim SH, Zo JH, Kim MA. The effects of metabolic syndrome and its components on arterial stiffness in relation to gender. *J Cardiol*. 2015;65:243–249. doi: 10.1016/j.jjcc.2014.05.009
238. Cooper LL, Palmisano JN, Benjamin EJ, Larson MG, Vasan RS, Mitchell GF, Hamburg NM. Microvascular function contributes to the relation between aortic stiffness and cardiovascular events: the Framingham Heart Study. *Circ Cardiovasc Imaging*. 2016;9:e004979.
239. Bostick B, Habibi J, DeMarco VG, Jia G, Domeier TL, Lambert MD, Aroor AR, Nistala R, Bender SB, Garro M, et al. Mineralocorticoid receptor blockade prevents Western diet-induced diastolic dysfunction in female mice. *Am J Physiol Heart Circ Physiol*. 2015;308:H1126–H1135.
240. Jia G, Habibi J, Aroor AR, Martinez-Lemus LA, DeMarco VG, Ramirez-Perez FI, Sun Z, Hayden MR, Meininger GA, Mueller KB, et al. Endothelial mineralocorticoid receptor mediates diet-induced aortic stiffness in females. *Circ Res*. 2016;118:935–943. doi: 10.1161/CIRCRESAHA.115.308269
241. Sowers JR, Habibi J, Aroor AR, Yang Y, Lastra G, Hill MA, Whaley-Connell A, Jaisser F, Jia G. Epithelial sodium channels in endothelial cells mediate diet-induced endothelium stiffness and impaired vascular relaxation in obese female mice. *Metabolism*. 2019;99:57–66. doi: 10.1016/j.metabol.2019.153946
242. Sehgel NL, Vatner SF, Meininger GA. "Smooth muscle cell stiffness syndrome"-revisiting the structural basis of arterial stiffness. *Front Physiol*. 2015;6:335. doi: 10.3389/fphys.2015.00335
243. Sowers JR, Habibi J, Jia G, Bostick B, Manrique-Acevedo C, Lastra G, Yang Y, Chen D, Sun Z, Domeier TL, et al. Endothelial sodium channel activation promotes cardiac stiffness and diastolic dysfunction in Western diet fed female mice. *Metabolism*. 2020;109:154223. doi: 10.1016/j.metabol.2020.154223
244. Hill MA, Jaisser F, Sowers JR. Role of the vascular endothelial sodium channel activation in the genesis of pathologically increased cardiovascular stiffness. *Cardiovascular Research*. 2020;cvaa326. doi: 10.1093/cvr/cvaa326
245. Leopold Jane A. Cellular and molecular mechanisms of arterial stiffness associated with obesity. *Hypertension*. 2013;62:1003–1004.
246. Wang J, Barbry P, Maiyar AC, Rozansky DJ, Bhargava A, Leong M, Firestone GL, Pearce D. SGK integrates insulin and mineralocorticoid regulation of epithelial sodium transport. *Am J Physiol Renal Physiol*. 2001;280:F303–F313. doi: 10.1152/ajprenal.2001.280.2.F303
247. von Wörm F, Berglund G, Carlson J, Månsson H, Hedblad B, Melander O. Genetic variance of SGK-1 is associated with blood pressure, blood pressure change over time and strength of the insulin-diastolic blood pressure relationship. *Kidney Int*. 2005;68:2164–2172. doi: 10.1111/j.1523-1755.2005.00672.x
248. Huang DY, Boini KM, Osswald H, Friedrich B, Artunc F, Ullrich S, Rajamanickam J, Palmada M, Wulff P, Kuhl D, et al. Resistance of mice lacking the serum- and glucocorticoid-inducible kinase SGK1 against salt-sensitive hypertension induced by a high-fat diet. *Am J Physiol Renal Physiol*. 2006;291:F1264–F1273. doi: 10.1152/ajprenal.00299.2005
249. Li P, Pan F, Hao Y, Feng W, Song H, Zhu D. SGK1 is regulated by metabolic-related factors in 3T3-L1 adipocytes and overexpressed in the adipose tissue of subjects with obesity and diabetes. *Diabetes Res Clin Pract*. 2013;102:35–42. doi: 10.1016/j.diabetes.2013.08.009
250. Norlander AE, Saleh MA, Pandey AK, Itani HA, Wu J, Xiao L, Kang J, Dale BL, Goleva SB, Laroumanie F, et al. A salt-sensing kinase in T lymphocytes, SGK1, drives hypertension and hypertensive end-organ damage. *JCI Insight*. 2018;2:e92801.