

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

Adipose cell size: Importance in health and disease

Karin G Stenkula¹ and Charlotte Erlanson-Albertsson^{2,*}

¹Glucose Transport and Protein Trafficking, ²Appetite Control, Department of
Experimental Medical Science, Biomedical Center, Lund University, 221 84 Lund,
SWEDEN

Short title: Adipose cell size

*Corresponding author: Charlotte Erlanson-Albertsson

E-mail: charlotte.erlanson-albertsson@med.lu.se

Phone:46-462228589, Fax:46-462224711

Address: BMC, B11, Sölvegatan 19, S-221 84 Lund, Sweden

20	LIST OF CONTENTS
21	Abstract
22	Introduction
23	Adipose cell biology
24	Exploring fat cell function
25	<i>Isolation of viable intact adipocytes</i>
26	<i>Techniques for cell separation according to size</i>
27	<i>Flotation and filtration</i>
28	<i>Flow cytometer</i>
29	<i>Single cell analysis</i>
30	<i>Cell size distribution</i>
31	Adipose cells during disease
32	Hypertrophic cells
33	<i>Increased cell size as a predictor of systemic insulin sensitivity</i>
34	<i>Impaired hormonal response in isolated large adipocytes</i>
35	<i>Inflammation is associated with increased cell size</i>
36	Dysfunctional small fat cells
37	Mechanisms behind impaired cell function
38	<i>Limited cell expansion</i>
39	<i>Impaired differentiation</i>
40	Therapies affecting adipose cell size
41	<i>Calorie-restricted diets with or without exercise</i>

42	<i>Bariatric surgery</i>
43	<i>Weight cycling – body set point</i>
44	<i>Cholesterol-free diet</i>
45	<i>Bioactive compounds</i>
46	<i>Pharmaceutical treatment with PPAR-ligands</i>
47	Perspective and Significance
48	References
49	
50	

Abstract

Adipose tissue is necessary to harbour energy. To handle excess energy, adipose tissue expands by increasing adipocyte size (hypertrophy) and number (hyperplasia). Here, we have summarized the different experimental techniques used to study adipocyte cell size and describe adipocyte size in relation to insulin resistance, type 2 diabetes and diet interventions. Hypertrophic adipocytes have an impaired cellular function and inherent mechanisms restrict their expansion to protect against cell breakage and subsequent inflammation. Reduction of large fat cells by diet restriction, physical activity or bariatric surgery therefore is necessary to improve cellular function and health. Small fat cells may also be dysfunctional and unable to expand. The distribution and function of the entire cell size range of fat cells, from small to very large fat cells, is an important but understudied aspect of adipose tissue biology. To prevent dysmetabolism, therapeutic strategies to expand small fat cells, recruit new fat cells and reduce large fat cells are needed.

Keywords insulin; adipocytes; obesity; cell size distribution; diet

68 **Introduction**

69 Humans have the highest percentage of body fat among mammals. In comparison to
70 other hominids body fat percentage is markedly higher, even in physically active
71 persons as in traditional hunter-gatherer populations. At the same time humans have
72 an elevated metabolic rate compared to other hominids, indicating an increased
73 organ metabolic activity (89). This expansion of energy budget in humans occurred
74 with a dietary shift from low-energy food such as leaves and fruits to high-energy
75 food like tubers and meat. Cooking also increased the availability of nutrients and
76 energy after foraging (35, 39). Increased body fat serves as a buffer against calorie
77 deficiency, especially in women with the need for extra energy during reproduction.
78 The capacity to store fat hence has been a major adaptive factor of our species. In
79 modern society with a surplus of energy-dense food and a minimum of effort to
80 obtain this food, the ability to store excess energy intake becomes problematic. Our
81 great appetite for energy-dense food and the efficiency of the gastrointestinal tract to
82 digest and absorb such food is coupled to an ability to store energy in adipose tissue.
83 With a positive energy balance and an expanding adipose tissue metabolic
84 dysfunction may arise leading to diseases like arteriosclerosis, diabetes, osteoporosis
85 and cancer.

86

87 Adipose tissue is traditionally considered an organ with low resting energy
88 expenditure (37). Adipose tissue is however a highly dynamic tissue, possessing
89 adipocytes of various size, referred to as small and large adipocytes. The properties

of these cells have been extensively explored, focusing on the relationship between cell size and various disease conditions such as inflammation (97, 109, 118), insulin resistance and diabetes (9, 15, 27, 33, 79, 102, 119). Laforest S *et al.* addressed the correlation between adipose cell size and cellular function as well as metabolic disease and concluded that the size of the adipocyte is an important factor to predict pathophysiological conditions (62).

In the present review, we have described the size variation of small and large fat cells during disease conditions, in particular insulin resistance, diabetes, obesity and during diet intervention. To this end we have included a discussion of the different techniques used for adipose cell size measurements. The data discussed in this review support the notion that the proportion of large and small adipocytes rather than a specific cell size *per se* within the adipose tissue depot is important for maintained cellular function and metabolism.

Adipose cell biology

The white adipose tissue (WAT) is an organ that primarily stores energy as triacylglycerol (lipogenesis) during energy surplus, to be mobilized as fatty acids (lipolysis) when needed. These processes are tightly regulated by insulin and adrenalin, which control the activity of lipogenic and lipolytic enzymes. Distinct from WAT, brown adipose tissue (BAT) is specialized for energy expenditure through thermogenesis (16). Browning and beiging of white adipose tissue has gained

attention as a therapeutic strategy to combat obesity (36). However, since to our knowledge there are no studies addressing the size of brown adipocytes or the alteration in cell size during browning of white adipocytes, we will focus on the white adipose tissue in this review.

Most white adipose tissue is distributed in two major depots, the subcutaneous (scWAT) and the visceral white adipose tissue (vWAT) (94). WAT contains both mature adipocytes (roughly 50% total cell number) and a stromavascular fraction, including preadipocytes, endothelial cells, mesenchymal stem cells and macrophages (67). WAT expands both through an increased fat cell volume (hypertrophy) and an increased number of fat cells (hyperplasia) (50). Hyperplasia occurs through differentiation of progenitor cells (adipogenesis) under the influence of various transcriptions factors, among these the peroxisome proliferator-activated receptor gamma (PPAR γ) and CCAAT/enhancer-binding protein alpha (CEBP α) (45). There is a constant turnover of adipocytes, with a life-span of around eight years, as established by measuring the ¹⁴C-incorporation into DNA (112). Both mature adipocytes and the resident inflammatory cells within adipose tissue secrete various hormones and cytokines, named adipokines (56). The most well-studied adipokines are leptin and adiponectin, which regulate appetite and energy metabolism, respectively (55, 127). Others are either pro-inflammatory, such as interleukin-6 (IL-6) (118) and tumor necrosis factor alpha (TNF- α) (42) or anti-inflammatory, such as

apelin (126). The secretion of free fatty acids and adipokines from adipose tissue cross-talks with other organs to affect whole body metabolism.

Exploring fat cell function

Adipocytes vary dramatically in size, with human white adipocytes ranging from <20 to 300 μm in diameter. This extrapolates to a several thousand fold range in cell volume within the same depot (58). This variation is mainly dependent on cellular triglyceride content. A central question is therefore whether the metabolism of adipocytes is size-dependent. To address this, standard methods for isolation of intact adipocytes have been established (93). However, methods to accurately characterize metabolism of isolated adipocytes with respect to their size are far more complex.

Isolation of viable, intact adipocytes

Early in the 1960's, Rodbell established a method for isolation of intact rat adipocytes from epididymal fat pad, a protocol that is still used in the research field of adipocyte biology (93). The method is based on incubating fat tissue in buffer containing collagenase. The fat cells released from the adipose tissue during digestion are then separated from the denser stromavascular cell fraction by flotation. The resultant single cell population contains adipocytes varying between 50 and 100 μm in diameter as assessed by light microscopy. These early studies concluded that isolated adipocytes may serve as a good model for subsequent

investigations of insulin action in adipose tissue. This allows investigation of hormonal responses related to glucose and lipid metabolism.

Subsequently, a Swedish group headed by Per Björntorp, compared the collagenase method, using the protocol of Rodbell (93), with a microscopy method based on fixation of adipose tissue followed by freeze-cutting the tissue (108), to verify that cell size estimation of adipocytes from human tissue had not been distorted during the isolation procedure (111). The cell size distribution was found similar using either method, with cell diameter varying between 40 and 150 μm . The collagenase method introduced by Rodbell *et al* (93) is a gentle and adequate method for isolation of adipocytes and there is minimal risk to rupture large adipocytes as long as centrifugations are avoided. The collagenase method may therefore appear most useful for metabolic and morphological studies of human adipose tissue. However, the microscopy method (108) is better suited for clinical studies. It is less time-consuming and requires a smaller adipose tissue sample for study. The collagenase method was however later reported to trigger an inflammatory response that could interfere with the signal transduction and gene expression of target molecules involved in lipid metabolism, potentially contributing to the relatively short life-span of the isolated cells (98).

Techniques for cell separation according to size

It is difficult to draw conclusions regarding a direct association between function and

size when comparing adipocytes of varying size isolated from different subjects (102, 110, 119). In light of this, a major limitation in characterizing the metabolic function of primary adipocytes in relation to cell size is the technical issue of collecting enough numbers of cells that are well-separated in size from the same subject. Another challenge includes optimization of cell handling, which has to be gentle enough to avoid cell breakage. This is a concern in particular for the very large adipocytes, which contain a huge lipid droplet that makes the cells fragile and prone to breakage. A rapid release of triglycerides will quickly deteriorate the entire cell population. All fat cells irrespective of size have a certain degree of pliability that makes filtration of the cell suspension through filters of different pore size difficult, even though some success has been reported (27, 48, 65, 109) as discussed below.

Flotation and filtration

A strategy to separate isolated cells of different size is to utilize the differences in buoyancy (25, 29). This property was exploited in an early attempt to collect fractions of fat cells of varying sizes from the same fat depot. Meters of dialysis tubing were used (12), resulting in two separated fat cells pools, with a mean diameter of 70 and 95 μm , respectively. Another approach relies on filtration of the cell suspension through nylon mesh of different pore sizes to collect pools of small and large fat cells (27, 48). Through filtration, a successful separation was reported with cells isolated from human subcutaneous fat tissue, producing cell pools with a mean diameter of 58 μm (small cell pool) and 100 μm (large cell pool). However, this technique

requires a substantial amount of starting material and yields poor cell recovery in each fraction. For example, in the study by Jernås *et al* (48), although 4-52 g of fat tissue was used for each subject, a limited amount of material was collected at the end of the separation. In another study, filtration was used to separate human fat cells into two pools starting from a similar amount of material, which resulted in slightly less well-distinct fractions, with a mean cell diameter of 81 and 114 μm , respectively (27). This method allowed a sufficient yield of cells for use in cell signalling studies.

Flow cytometer

A general belief is that isolated adipocytes are too large and fragile for flow cytometer or similar sorting instrumentation. Nonetheless, in a recent method description by Majka *et al* (75), the authors propose that the modern flow systems have the capabilities needed to sort adipocytes up to 250 μm in diameter with relatively low pressure, thereby reducing mechanical stress. The authors have established a protocol to exclude stromal vascular cells, that can potentially interfere with subsequent analysis (76). Indeed, several papers report successful flow cytometer analysis using primary adipocytes isolated from human (26), mouse (82) and rat adipose tissue (8). Thus, with newer sorting instrumental setups, useful methods have been developed that may become more widely adopted for future characterization of adipocytes of various size.

221 *Single Cell Analysis*

222 Methods incorporating use of microscopy have the advantage of functional
223 assessment of single cells of a distinct cell size, though it often includes tedious
224 analysis. Several studies have employed fluorescence microscopy in fixed human
225 adipocytes (27) and live, intact adipocytes isolated from either human (70, 121) or rat
226 adipose tissue (69, 113) for metabolic characterization including hormonal responses.
227 Also, imaging of ex vivo adipose explant from *Rhesus macaques* allowed
228 characterization of fatty acid uptake at a single cell level that included cell size
229 measurements (117). Another technically advanced approach for single cell analysis
230 is patch clamping, which has been established using isolated adipocytes (60, 128).
231 Additionally, the RNA-seq technology has developed rapidly, and is now available
232 at a single cell level (73). A caveat is that even though these methods are, in theory,
233 applicable to adipocytes, these analyses require cell separation, usually carried out
234 by flow sorting, which in itself causes technical problems such as cell breakage. This
235 could explain the lack of published studies using this technique. Considering the vast
236 information that can be gathered by single cell RNA-seq studies, the technical
237 hurdles utilizing this technology for isolated adipocytes will hopefully be resolved.
238 One possibility is the use of a fluid chamber-based system, like Drop seq analysis
239 (74), a method that is applicable for single cell suspensions.

240

241 *Cell size distribution*

In many studies, measurements of cellular and systemic function are correlated with the average fat cell size measured by histological sectioning of intact tissue or isolated cells. This type of size measurement requires only a small amount of tissue/cells and is technically relatively easy to accomplish. However, often this method does not provide causal relationship between cell size and function of individual adipocytes. In addition, it is questionable whether the entire span of cell sizes is detectable using a histological approach, where the very small cell population (<30 μm diameter) easily could be overlooked or underestimated, since the sectioning has to occur at the widest diameter of every cell to obtain a true measurement. Several (semi)-automated image analysis tools have been developed to improve cell size analysis from histologic fat tissue samples (10, 18, 85, 86). This allows faster analysis of an increased cell number compared with manual measurements.

The coulter counter is another commonly used method, where the size of osmium tetroxide-fixed cells is determined by measurement of electric resistance. The method is evaluated to be an accurate method for size measurement in adipose tissue (40), and also for isolated cells from either human or rat adipose tissue (20). A drawback is the use of osmium tetroxide, which is expensive and hazardous. In comparison, the coulter counter measurement has the advantage of measuring a large number of cells (~10000) with a broad size range, 20-300 μm in diameter, whereas histology does not provide the same precision. It also carries a limitation in detecting small fat cells. Caveats raised for the coulter counter method are whether cell debris could

contribute to false-positive measurements of the number of small cells, and the risk of osmium to cause cell swelling (40). Mathematical modelling of data obtained by the coulter counter approach revealed a bimodal cell size distribution (51, 78, 79, 122). It was pointed out that the use of the average cell diameter of the entire cell population could be misleading and that more useful information could be obtained by analysing the mean cell size in each cell population, which was termed a small and a large cell population. Optical analysis on the other hand, usually demonstrates a one-peak Gaussian-like cell size distribution (48, 81), with a reduced frequency of smaller fat ($<30\text{ }\mu\text{m}$) cells. The methods used for analysing adipocytes according to size in both intact tissue and isolated cells are summarized in figure 1. Even though the isolation of fat cells is technically challenging, there is definitely a need for continued research on adipocyte function in relation to cell size.

Fat cells during disease

It is well-known that obesity, which is defined by an excessive white adipose tissue mass, is one of the main risk factors for insulin resistance and type 2 diabetes (92, 104). While the exact mechanisms are not known, it is clear that in obesity the adipocyte itself is unable to regulate excess nutrients, leading to increased circulating levels of fatty acids and glucose, changed adipokine secretion, and dysregulated energy metabolism. In this regard, a limitation in fat tissue expansion, as reviewed by Rutkowski *et al.* (99) and Virtue *et al.* (120), could contribute to systemic insulin resistance and diabetes.

286

287 The fact that adipose tissue contains a mixture of adipose cells, that vastly vary in
288 size, makes it complicated to define if there is a set size threshold that is associated
289 with impaired function. When reviewing original research articles, it is also
290 important to consider: 1) what method that was used to determine cell size; 2) if the
291 biologic function is measured in isolated cells or is merely correlated with cell size;
292 and 3) whether cells of different sizes originated from the same subject. In order to
293 compare different cell size measurements across the literature, we have converted
294 data expressed as μg lipid/cell or cell volume (pl), to cell diameter (μm), using the
295 assumption that the cells are spherical with a density of 0.91 (53). In the text, " μm "
296 implies mean adipocyte diameter unless stated specifically to be otherwise. The
297 described differences regarding cell size and their function attained statistical
298 significance in each individual study.

299

300 **Hypertrophic adipose cells**

301 *Increased cell size as a predictor of systemic insulin sensitivity*

302 In general, large, hypertrophic cells are considered less metabolically favorable and
303 are associated with pathophysiologic conditions. Even in the early studies of human
304 adipocytes, increased fat cell size was shown to correlate with impaired whole body
305 metabolic regulation (9, 11) and systemic insulin resistance (33, 61, 101, 102). In a
306 prospective study of Pima Indians, known for their predisposition for developing
307 insulin resistance and type 2 diabetes, increased adipocyte cell size was shown as an

independent marker of type 2 diabetes (119). The average cell size diameter of the entire cell population, as assayed by coulter counter, positively correlated with systemic glucose tolerance (GTT). It was found that an average cell size of 115 μm correlated with normal GTT; 121 μm with impaired GTT; and 125 μm with diabetes (119). In a comparison between Pima Indians and Caucasian children with similar body weight and body fat mass, Pima Indians had increased average adipose cell size, 101 versus 92 μm , as well as increased circulating levels of glucose and insulin (2). In non-diabetic subjects, an increased average adipose cell size, varying between 75 and 130 μm as determined manually by light microscopy, negatively correlated with both cellular and systemic insulin sensitivity independent of BMI (72). In another study, over-feeding of both insulin resistant (IR) and insulin sensitive (IS) moderately obese subjects normalized for BMI was carried out to test whether IS subjects were protected from cell enlargement (77). At the start of the study, the insulin sensitive subjects had smaller adipocytes, but following a modest weight gain (~3 kg), they displayed an increase in average adipocyte size from 108 to 115 μm as assayed by coulter counter. Such individuals also demonstrated impaired insulin-mediated glucose uptake (IMGU) and increased lipolysis (77). In contrast, the insulin resistant subjects had no significant increase in cell size and only slightly further impairment of IMGU. Thus, it was concluded that an increase in adipose cell size could predict the development of insulin resistance and type 2 diabetes (77). In yet another over-feeding study, the subjects starting with smaller adipocytes, assayed between 57 and 115 μm using coulter counter, turned out to have the most impaired

metabolic profile following weight gain. Thus having small adipocytes *per se* was not sufficient to protect against development of insulin resistance (52). It was thought that the small adipocytes expanded more rapidly, whereas the subjects that initially had large adipocytes instead increased the number of cells (hyperplasia). This supports the notion that any increase in adipocyte size, either from small or normal size cells, could result in an impaired metabolic profile (52). Increased subcutaneous adipocyte cell size was also a predictor of type 2 diabetes incidence in a Swedish cohort of ~1300 women followed over 25 years (71). Together, these studies suggest that increase in subcutaneous adipocyte size is an important factor to predicts type 2 diabetes unrelated to obesity (2, 71, 72, 119).

In contrast, Pima/Papago subjects with type 2 diabetes had similar mean adipose cell size, 120 μm , compared to weight matched healthy individuals (53). However, the glucose metabolism and insulin response in adipocytes isolated from type 2 diabetic subjects were markedly reduced (53). Large subcutaneous fat cells were also observed in obese children with hyperinsulinemia (15), with an average cell size of 115 μm compared with 86 μm in lean controls, as assayed by coulter counter. After body weight reduction through dietary restriction over 8 weeks, the average fat cell size in the obese children decreased to an average cell size diameter of 95 μm , but no direct correlation between cell size and insulin level was observed.

Impaired hormonal response in isolated large adipocytes

352 Direct measurements of cellular hormonal response and metabolism in isolated
353 adipocytes support the concept that enlarged adipocytes are less responsive and less
354 metabolically favorable (1, 33, 61, 102, 122). Insulin responsiveness, measured as %
355 increase of glucose oxidation to CO₂, negatively correlated with cell size, where the
356 mean cell size within each adipose cell tissue sample ranged between 74 and 134 μ m,
357 as assessed by coulter counter (102). No difference in the protein level or degree of
358 activation was found for the insulin signaling down-stream intermediates insulin
359 receptor substrate (IRS)-1 and protein kinase B (PKB), when comparing small versus
360 large human adipocytes isolated by filtration from the same non-diabetic subject (27).
361 Even though the total GLUT4 protein content was similar in the small and large cell
362 populations as assessed by microscopy, average diameter 81 and 114 μ m
363 respectively, image analysis revealed a 2-fold increase in insulin-induced GLUT4
364 translocation to the plasma membrane in the small cell population whereas no
365 increase of GLUT4 translocation was found in the large cell population (27). This
366 supported a reduced insulin sensitivity in the large fat cells. In recent studies using
367 live cell imaging with total internal reflection fluorescence microscopy in human fat
368 cells, impaired GLUT4 dynamics correlated with both increased BMI and insulin
369 resistance of the donor subject (121). In a follow-up analysis, there was however no
370 correlation between GLUT4 trafficking and size of fat cells from 25 to 125 μ m in
371 diameter, isolated from different subjects (70). In non-diabetic subjects an increase in
372 the average adipocyte size, assayed over a range of 75-130 μ m in diameter, correlated

with a decrease in insulin-stimulated glucose uptake (72), again supporting a reduced insulin sensitivity with enlargement of fat cells.

Lipolytic activity, leading to an increased release of circulating FFA is also increased in large human adipose cells (22, 47, 65, 125), an effect regulated by insulin. Using a flotation approach, two cell populations, average diameter of 82 μm of small and 100 μm of large fat cells, were collected from human subcutaneous adipose tissue and characterized for their lipolytic activity (65). Both non-stimulated as well as hormone-induced lipolysis was increased in large adipocytes compared to small adipocytes. These findings were supported by increased protein levels of hormone-sensitive lipase (HSL) and adipose tissue triacylglycerol lipase (ATGL), which are responsible for TAG hydrolysis prior to fatty acid release from fat cells (65). Similar findings were observed in a study by Jacobsson B *et al.* (47), where basal and adrenergically stimulated lipolysis correlated positively with cell size, whereas the insulin-induced anti-lipolytic effect was independent of cell size in the size range studied, 70-115 μm , measured by microscopy.

Inflammation is associated with increased cell size

Altered adipokine release is tightly associated with an increased inflammatory response in adipose tissues, which in turn can deteriorate insulin signaling in mature adipocytes (63, 97, 118). The positive correlation between fat cell size and expression of the inflammatory genes NF- κ B, TNF- γ (6), and expression of TNF receptor (43)

395 suggest that increased adipocyte size is associated with an increased inflammatory
396 response. Adipokine secretion in relation to cell size was also addressed in a study
397 where human adipocytes isolated from the same subject were fractionated via
398 filtration into four different diameter groups - small (73 μm), medium (98 μm), large
399 (113 μm) and very large (127 μm) (109). Increased pro-inflammatory adipokine
400 release, namely IL-6, IL-8, monocyte chemoattractant protein-1, and granulocyte
401 colony-stimulating factor, was found in the very large fat cell group compared with
402 the small fat cell group, confirming large fat cells to be highly inflammatory. By
403 isolation and separation of human adipocytes through filtration into two pools, large
404 and small cells, 58 and 100 μm mean diameter, Jernås *et al* (48) detected a >4-fold
405 increase of several immune-related genes including serum amyloid A in the large fat
406 cell population. In a recent study, non-obese type 2 diabetic subjects had larger
407 adipocytes, average cell diameter of 104 μm , as assessed by light microscopy of
408 isolated cells, than the healthy controls, average cell diameter of 95 μm , but lipolysis
409 and the secretion of inflammatory proteins measured using isolated adipocytes were
410 similar in the two groups (3). Still, increased adipocyte size positively correlated with
411 both lipolysis and systemic insulin resistance, independent of type 2 diabetes (3).
412 This suggests that the appearance of large fat cells precede the development of
413 diabetes. Release of leptin, but not adiponectin, increased with increasing cell size in
414 mature adipocytes isolated from subcutaneous adipose tissue (109). However, the
415 proportion of small cells in human SAT explants, defined as <69 μm mean diameter,
416 and large cells defined as >131 μm mean diameter by histologic sectioning, positively

correlated with adiponectin secretion and circulating serum adiponectin (81). Thus, there seems to be a shift towards an increased pro-inflammatory secretion and decreased anti-inflammatory secretion from the very hypertrophic cells, which could contribute to systemic insulin resistance and type 2 diabetes (119).

Dysfunctional small adipocytes

There are also studies that suggest that impaired expansion of small fat cells is associated with insulin resistance (24, 78, 79, 87). The recent approach of studying not only the average cell size but rather the entire cell size distribution including the very small fat cells and the proportion between different cell sizes could be the reason for the emerging concept of an impaired lipid storage capacity in not only the large cells but also the smaller cells. In morbidly obese subjects, type 2 diabetes was associated with an increased proportion of small adipocytes (<50 μm) in omental, mesenteric and subcutaneous fat depots (24). In a study by McLaughlin *et al.* (79), the coulter counter technique was used to illustrate a bimodal cell size distribution in human subcutaneous adipose tissue, with two cell populations, small and large cells. To verify that the population of so-called small cells represented adipocytes, the analysis was repeated in both intact adipose tissue as well as in isolated adipocytes from the same subject (79). In contrast to expectations, the overall mean adipocyte size was similar for cells from insulin-sensitive and insulin-resistant moderately obese subjects normalized for BMI (~30). However, the proportion of small fat cells was higher in the insulin-resistant group, ratio small/large cells 1.66 in

the insulin sensitive group and 0.94 in the insulin resistant group (79). Presumably, the insulin-resistant subjects had small fat cells that failed to expand appropriately, thus being unable to store sufficient fat, resulting in increased ectopic storage of fat in the liver and the skeletal muscle. This idea was supported in a further study where increased insulin resistance as determined by steady-state plasma glucose concentrations correlated positively with an increased proportion of small fat cells and an increased mean cell size within the large cell population. (78). By coulter counter analysis, type 2 diabetic subjects had a greater mean adipocyte size of the very large cells, 144 versus 171 μm , a lower total cell number in the subcutaneous abdominal fat mass, but a greater proportion of small adipocytes compared with weight-matched non-diabetic subjects (87). The idea of dysfunctional small fat cells is not in conflict with the concept of large adipocytes as less metabolically favourable, but emphasizes the importance of examining the distribution of the entire cell size range. Thus, there seems to be an inability of both small and large fat cells to expand, leading to dysmetabolism and insulin resistance, illustrated in figure 2.

Mechanisms behind impaired cell function

Limited cell expansion

The cellular mechanisms behind impaired function of large adipocytes are not yet resolved but are at least in part due to limitation in further cell expansion. During excess energy, the adipocyte has the ability to maximize its volume several thousand

times. However, with an increased cell size the adipocyte becomes stiff (106) and the membrane signaling disturbed (88, 107). Various intracellular signaling pathways are activated as demonstrated in 3T3-L1 adipocytes, for example the MEK signaling pathway (107). Also, the regulation of fat cell size could potentially be mediated through volume sensitive adipocyte membrane proteins (17), one of them recently being identified and named SWELL 1 (128). The authors demonstrated that gene silencing of SWELL1 led to a decrease in fat cell size and decreased glucose and lipid uptake in both cultured and primary adipocytes. Adipose-specific SWELL1 knock-out caused systemic hyperglycemia and decreased insulin sensitivity (128). These experiments clearly demonstrate that fat cell enlargement is closely linked to glucose metabolism and when this mechanism fails glucose metabolism is deranged. Mice studies have shown that remodeling of the extracellular matrix by knock-down of Collagen VI reduced the local adipose tissue inflammation and allowed increased adipocytes expansion without impaired cellular insulin signaling and unfavorable metabolic consequences (57).

The failure of enzymes promoting triglyceride storage within the fat cell may also contribute to impaired fat cell function. Hence, improvement of the triglyceride storage capacity in adipocytes by adipose tissue-specific overexpression of PEPCK, the rate-limiting step in glyceroneogenesis, leads to increased adipocyte cell size allowing fat mass increase without leading to insulin resistance (28). In humans, increased mRNA expression of the lipid droplet-associated proteins Cidea, FSP27

and perilipin in SAT positively correlated with insulin sensitivity index (HOMA-IR) in weight matched obese subjects (91). Since these proteins promote lipid droplet formation and facilitate triglyceride storage (54), their increased expression is expected to favour increased cell size. A perilipin knock-out mouse model had a 30% reduction of fat tissue mass and was protected against high fat-diet induced obesity (116). Still, these mice were prone to develop impaired glucose tolerance and systemic insulin resistance. Together, these data suggest that a limitation of cell expansion is linked with impaired glucose and impaired fat metabolism both at the cellular and at the systemic level.

Impaired differentiation

Several studies have pinpointed that too few mature adipose cells and impaired differentiation of precursor cells is associated with type 2 diabetes (21) in both non-obese (3) and obese subjects (32, 46). Notably, there was no evidence of fewer precursor cells but rather impaired cell lineage commitment and differentiation (3). Others reported a negative correlation between number of precursors and BMI of the subject (46). The canonical Wnt signaling pathway (96) and bone morphogenetic protein 4 (BMP4) (14) regulate the commitment of mesenchymal stem cells and the differentiation of pre-adipocytes into adipocytes. In a study of non-diabetic insulin-resistant first-degree relatives of type 2 diabetic patients, the average adipose cell size negatively correlated with the expression of Wnt1 signaling genes FZD1, GSK3 β , and LEF1 β -catenin (123). The adipokine Wnt1 inducible signaling pathway

protein 2 (WISP2) was demonstrated to inhibit cell lineage commitment by inhibiting BMP4 (34), and increased expression of WISP2 in human subcutaneous fat tissue positively correlated with increased cell size (34). Thus, this could provide a mechanism for impaired differentiation of new adipocytes, and result in increasing the size of already committed adipocytes.

Therapies affecting adipose cell size

Metabolic improvements are strongly associated with a reduction of fat cell size. These may occur through calorie-restricted diets/exercise, cholesterol-free diets, bariatric surgery, the use of bioactive compounds and pharmacological compounds stimulating PPAR γ . Weight gain seems to be mostly associated with cell expansion (hypertrophy) (112), whereas weight loss leads to a decreased fat cell size independent of the method employed for weight loss; surgical, diet restriction, physical activity or life style intervention (83).

Calorie-restricted diets with or without exercise

Life style intervention usually involves restricted diet, such low-calorie diets (LCD), very-low calorie diets (VLCD) either with or without exercise. In one study, 48 obese subjects were divided into four groups and were fed for six months either 1) a healthy control diet, 2) a 25% calorie restricted diet, 3) a 12.5% restricted diet with 12.5% increase in total energy expenditure by exercise, or 4) a low-calorie diet until a 15% reduction in body weight was achieved (64). Diets were based on American

Heart Association recommendations ($\leq 30\%$ fat). To estimate fat cell size, subcutaneous abdominal needle biopsies were performed and adipocyte size determined by coulter counter technique (38). It was found that in response to 6 months of energy-restricted diet in overweight men and women, body weight, visceral fat mass and fat cell size all decreased, and that addition of exercise further decreased adipose cell size (64). The reduction in fat cell size was most strongly associated with a decrease in body weight and a decrease in percent body fat (64). The intervention also decreased liver fat, but had no effect on muscular fat. Insulin sensitivity was increased to a similar level by diet alone or in conjunction with exercise. The improved insulin sensitivity was linked to body weight reduction, visceral fat reduction and fat cell size reduction, but unrelated to liver fat reduction. It was stated that large fat cell size was associated with impaired adipogenesis and increased TAG content in adipose tissue, liver, muscle and pancreas leading to insulin resistance.

In another 20-week intervention study, overweight women were given a restricted diet alone, together with low-intensity or high-intensity exercise, consisting of 20 min treadmill walking three times per week at 50 respectively 75 % of maximal heart rate (124). The calorie deficiency was adjusted to -2800 kcal/week for all three groups, obtained through diet restriction in the diet group, and through diet- 2400 kcal per week and -400 kcal per week in the exercise groups (124). All three interventions reduced body weight, fat mass and percent fat. To estimate fat cell size,

subcutaneous abdominal and gluteal needle biopsies were performed and adipocyte size determined by microscopic evaluation of isolated fat cells (124). Gluteal adipocyte size decreased similarly in all three groups, whereas abdominal adipocyte size did not change in the diet group but was reduced in the exercise groups. It was concluded that in order to affect fat cell size in the abdomen, exercise needs to be added in a life style intervention program. Hence caloric restriction alone protects against abnormal expansion of fat but may be more efficient in combination with physical activity to prevent dysmetabolism and insulin resistance. The authors did not address whether the reduced fat cell size was due to increased hyperplasia or reduced size of existing fat cells.

Bariatric surgery

Bariatric surgery has been demonstrated to decrease fat cell size in abdominal subcutaneous adipose tissue. Two years after a gastric bypass in 62 obese women, body weight loss amounted to 33% with a simultaneous shift of abdominal subcutaneous fat cell size to lower diameter (5, 41). In these studies, a subcutaneous fat biopsy was obtained from the abdominal wall. Fat cells were isolated through collagenase treatment, and the diameter of 100 cells measured. The reduced average fat cell size correlated strongly with improved insulin sensitivity ($p=0.0057$), whereas body weight loss and fat mass loss did not (41, 100). The hypothesis that visceral adipose tissue induces insulin resistance was dismissed since removal of the major omentum, constituting a substantial portion of this tissue, during bariatric surgery

had no additive effect on insulin sensitivity (5). The mechanism behind normalized fat cell biology after bariatric surgery is unknown, but may be due to reduced inflammatory markers and increased adiponectin levels (41).

Cholesterol-free diet

In the obese state, half of the cholesterol pool in the body is stored in the adipose tissue. With adipocyte hypertrophy there is an increased content of body cholesterol, which is stored up to 90 % in the free form at the surface of the triglyceride droplet, the remainder being part of the plasma membrane (90). Dietary cholesterol has been demonstrated to aggravate adipose tissue inflammation in a mouse model of diet-induced obesity, suggesting that high dietary cholesterol can lead to adipocyte dysfunction (115). In monkeys, a better model for human physiology than rodents, a high-cholesterol diet was demonstrated to induce the enlargement of fat cells (19). In this model, the fat cell size was measured in fixed and embedded samples of omental and subcutaneous adipose tissue, using light microscopy. It was found that in omental fat, the mean diameter of the fat cells were 42.3 (± 2.4) μm with low cholesterol diet, and increased to 49.0 (± 2.7) μm and 58.9 (± 3.1) μm for medium and high cholesterol diets, respectively after 10 weeks (19). At the same time, there was a parallel increase in plasma lipoprotein LDL levels, suggesting that the LDL levels determined the size of the adipocytes. There was no change in adipocyte size of subcutaneous adipose tissue with the medium and high cholesterol diet compared to the low cholesterol diet. Also, a significant increase in adipose tissue expression of

the pro-inflammatory genes IL-6 and IL-8 in the high cholesterol diet group relative to the other groups was observed. There was no difference in glucose and insulin sensitivity, suggesting that 10 weeks of feeding the medium and high cholesterol diets was perhaps not long enough to observe such changes (19). Hypertrophic adipocytes have reduced plasma membrane cholesterol, with a ~40% reduction of membrane cholesterol content (expressed as cholesterol/protein) comparing small (~40 μm) and large (~50 μm) adipocytes, when studied by filtration (66). Since serum amyloid A has been shown to reduce cholesterol uptake from HDL particles, an altered cholesterol metabolism could be one of the mechanisms through which serum amyloid A contributes to the insulin resistance reported in enlarged adipocytes. Dietary cholesterol can thus induce visceral adipocyte enlargement, which may explain metabolic dysfunction with high-cholesterol diets.

Bioactive compounds

The increasing world-wide prevalence of obesity is partially related to the ready availability of highly palatable food, which increases the incidence of hedonic, non-homeostatic eating. Any treatment that suppresses appetite, in particular appetite for palatable food, will reduce the flow of nutrients to the fat cell. Thylakoids are the green leaf membranes present in all chloroplasts, that transform energy from light to ATP. Upon isolation, highly complex membrane proteins, galactolipids, vitamins and antioxidants are the main components (23). These were found to suppress fat digestion in the intestine exerted by pancreatic lipase/colipase, suppress food intake

and reduce body weight as demonstrated both in rodent and in human studies (23). When given to mice fed a high-fat diet, fat cell size was reduced in the thylakoid-fed animals compared to control high-fat fed animals, since some of the fat was retarded in the intestine (114). Thus a reduced flow of nutrients from the intestine to the body reduces fat cell size. Also, an increased energy expenditure leading to loss of energy may reduce fat cell size. A couple of thermogenic compounds are known that lead to body fat loss, without affecting appetite (7). One example of a thermogenic compound is capsaicin, which administrated in combination with exercise led to a decreased fat cell size in mice (84).

Pharmaceutical treatment with PPAR-ligands

Thiazolidinediones (TZDs), also called glitazones, are a class of drugs introduced in the late 1990's that increase insulin sensitivity in muscle, fat and liver (103). TZDs exert their effect by acting as synthetic ligands for PPAR γ , a member of the nuclear receptor superfamily of transcription factors. TZD promote binding of the PPAR γ -retinoid X receptor (RXR) complex to PPAR γ response elements (PPRE) in target genes that are involved in adipogenesis, lipid metabolism and glucose metabolism (68). To date, two PPAR γ isoforms have been identified, PPAR γ 1 and PPAR γ 2. PPAR γ 1 is ubiquitously expressed whereas PPAR γ 2 is mainly expressed in adipocytes and intestine. In adipose tissue, PPAR γ 2 is the master regulator of adipogenesis (95). PPAR γ is not only essential for proper adipocyte differentiation but is also required for function and survival of mature adipocytes in mice (44), even

though PPAR γ silencing in differentiated 3T3-L1 cells only partially led to impaired adipogenesis (105). Besides adipogenesis, PPAR γ regulates gene expression of the GLUT4 transporter that is essential for glucose metabolism, the c-Cbl-associated protein (CAP) involved in insulin signaling, and several adipokines, including adiponectin (31), IL-6 (63) and TNF- α (49). Thus, the insulin sensitizing effect obtained by TZD treatment is multifactorial.

In humans, decreased insulin resistance after 12 weeks of pioglitazone treatment of overweight/obese nondiabetic, insulin-resistant subjects was associated with an increased proportion of small fat cells, and a 25% increase in the absolute number of these cells, assessed by coulter counter measurements (80). In contrast, a significantly increased subcutaneous adipocyte surface area was associated with improved systemic insulin sensitivity (59). A trend towards an increased proportion of small subcutaneous adipocytes (<50 μ m) was shown after 2 months of TZD treatment in type 2 diabetic subjects (13). In the latter study, TZD treatment also improved the anti-lipolytic effect of insulin, and led to a 2.5-fold increase of plasma adiponectin level and a 30% decrease in plasma leptin levels (13). There are, to the best of our knowledge, no studies yet that have addressed the specific effect of TZD's on adipocytes of different sizes. Still, the literature exploring the mechanisms of action of TZD's supports the idea that the insulin sensitizing effect of TZDs, to a large extent, can be explained by increasing the pool of small adipocytes preferentially in the subcutaneous fat depots (4).

659

660 Perspective and Significance

661

662

663 The mechanism by which obesity contributes to dysmetabolism is three-fold: 1) an
664 inability of adipose precursor cells to differentiate into mature fat cells 2) an inability
665 of small fat cells to expand and 3) an inability of large fat cells to expand further.
666 Together, this leads to an impaired ability to store excess energy, whereby increased
667 circulating free fatty acids and adipokines cause disturbed systemic metabolism. The
668 picture of fat cell size should be considered dynamic, where the proportion of
669 various fat cell sizes should be measured in order to gain the greatest degree of
670 insight.

671

672 In agreement with earlier studies, hypertrophic adipocytes are associated with
673 impaired cell function and dysmetabolism. This has been documented through
674 dietary intervention studies, demonstrating that a reduced fat cell size following
675 weight reduction is coupled to improved insulin sensitivity. Whether the
676 hypotrophic fat cell regains its normal function per se or whether other factors
677 inherent in newly recruited fat cells are necessary to provide a healthy metabolism
678 needs to be investigated. Of specific interest will be to characterize the very small fat
679 cells, especially in terms of their capacity to expand, in order to understand their
680 relation to metabolic dysfunction.

681

Acknowledgements

This review was financially supported by the Diabetes foundation, Diabetes Wellness, Crafoord foundation, Royal Physiographic Society, and Runo Svensson foundation.

References

1. **Abbasi F, Brown BW, Jr., Lamendola C, McLaughlin T, and Reaven GM.** Relationship between obesity, insulin resistance, and coronary heart disease risk. *J Am Coll Cardiol* 40: 937-943, 2002.
2. **Abbott WG, and Foley JE.** Comparison of body composition, adipocyte size, and glucose and insulin concentrations in Pima Indian and Caucasian children. *Metabolism* 36: 576-579, 1987.
3. **Acosta JR, Douagi I, Andersson DP, Backdahl J, Ryden M, Arner P, and Laurencikiene J.** Increased fat cell size: a major phenotype of subcutaneous white adipose tissue in non-obese individuals with type 2 diabetes. *Diabetologia* 59: 560-570, 2016.
4. **Adams M, Montague CT, Prins JB, Holder JC, Smith SA, Sanders L, Digby JE, Sewter CP, Lazar MA, Chatterjee VK, and O'Rahilly S.** Activators of peroxisome proliferator-activated receptor gamma have depot-specific effects on human preadipocyte differentiation. *J Clin Invest* 100: 3149-3153, 1997.
5. **Andersson DP, Eriksson-Hogling D, Backdahl J, Thorell A, Lofgren P, Ryden M, Arner P, and Hoffstedt J.** Omentectomy in Addition to Bariatric Surgery-a 5-Year Follow-up. *Obes Surg* 27: 1115-1118, 2017.
6. **Bahceci M, Gokalp D, Bahceci S, Tuzcu A, Atmaca S, and Arikan S.** The correlation between adiposity and adiponectin, tumor necrosis factor alpha, interleukin-6 and high sensitivity C-reactive protein levels. Is adipocyte size associated with inflammation in adults? *J Endocrinol Invest* 30: 210-214, 2007.
7. **Belza A, Frandsen E, and Kondrup J.** Body fat loss achieved by stimulation of thermogenesis by a combination of bioactive food ingredients: a placebo-controlled, double-blind 8-week intervention in obese subjects. *Int J Obes (Lond)* 31: 121-130, 2007.
8. **Bernstein RL, Hyun WC, Davis JH, Fulwyler MJ, and Pershadsingh HA.** Flow cytometric analysis of mature adipocytes. *Cytometry* 10: 469-474, 1989.
9. **Bernstein RS, Grant N, and Kipnis DM.** Hyperinsulinemia and enlarged adipocytes in patients with endogenous hyperlipoproteinemia without obesity or diabetes mellitus. *Diabetes* 24: 207-213, 1975.
10. **Bjornheden T, Jakubowicz B, Levin M, Oden B, Eden S, Sjostrom L, and Lonn M.** Computerized determination of adipocyte size. *Obes Res* 12: 95-105, 2004.
11. **Bjorntorp P, Grimby G, Sanne H, Sjostrom L, Tibblin G, and Wilhelmsen L.** Adipose tissue fat cell size in relation to metabolism in weight-stabile, physically active men. *Horm Metab Res* 4: 182-186, 1972.

- 722 12. **Bjorntorp P, and Sjoström L.** The composition and metabolism in vitro of
723 adipose tissue fat cells of different sizes. *Eur J Clin Invest* 2: 78-84, 1972.
- 724 13. **Boden G, Cheung P, Mozzoli M, and Fried SK.** Effect of thiazolidinediones
725 on glucose and fatty acid metabolism in patients with type 2 diabetes. *Metabolism* 52:
726 753-759, 2003.
- 727 14. **Bowers RR, and Lane MD.** A role for bone morphogenetic protein-4 in
728 adipocyte development. *Cell Cycle* 6: 385-389, 2007.
- 729 15. **Brook CG, and Lloyd JK.** Adipose cell size and glucose tolerance in obese
730 children and effects of diet. *Arch Dis Child* 48: 301-304, 1973.
- 731 16. **Cannon B, and Nedergaard J.** Brown adipose tissue: function and
732 physiological significance. *Physiol Rev* 84: 277-359, 2004.
- 733 17. **Che H, Yue J, Tse HF, and Li GR.** Functional TRPV and TRPM channels in
734 human preadipocytes. *Pflugers Arch* 466: 947-959, 2014.
- 735 18. **Chen HC, and Farese RV, Jr.** Determination of adipocyte size by computer
736 image analysis. *J Lipid Res* 43: 986-989, 2002.
- 737 19. **Chung S, Cuffe H, Marshall SM, McDaniel AL, Ha JH, Kavanagh K, Hong
738 C, Tontonoz P, Temel RE, and Parks JS.** Dietary cholesterol promotes adipocyte
739 hypertrophy and adipose tissue inflammation in visceral, but not in subcutaneous, fat in
740 monkeys. *Arterioscler Thromb Vasc Biol* 34: 1880-1887, 2014.
- 741 20. **Cushman SW, and Salans LB.** Determinations of adipose cell size and
742 number in suspensions of isolated rat and human adipose cells. *J Lipid Res* 19: 269-273,
743 1978.
- 744 21. **Danforth E, Jr.** Failure of adipocyte differentiation causes type II diabetes
745 mellitus? *Nat Genet* 26: 13, 2000.
- 746 22. **Engfeldt P, and Arner P.** Lipolysis in human adipocytes, effects of cell size,
747 age and of regional differences. *Horm Metab Res Suppl* 19: 26-29, 1988.
- 748 23. **Erlanson-Albertsson C, and Albertsson PA.** The Use of Green Leaf
749 Membranes to Promote Appetite Control, Suppress Hedonic Hunger and Loose Body
750 Weight. *Plant foods for human nutrition* 70: 281-290, 2015.
- 751 24. **Fang L, Guo F, Zhou L, Stahl R, and Grams J.** The cell size and distribution
752 of adipocytes from subcutaneous and visceral fat is associated with type 2 diabetes
753 mellitus in humans. *Adipocyte* 4: 273-279, 2015.
- 754 25. **Farnier C, Krief S, Blache M, Diot-Dupuy F, Mory G, Ferre P, and Bazin
755 R.** Adipocyte functions are modulated by cell size change: potential involvement of an
756 integrin/ERK signalling pathway. *Int J Obes Relat Metab Disord* 27: 1178-1186, 2003.
- 757 26. **Festy F, Hoareau L, Bes-Houtmann S, Pequignat AM, Gonthier MP,
758 Munstun A, Hoareau JJ, Cesari M, and Roche R.** Surface protein expression between
759 human adipose tissue-derived stromal cells and mature adipocytes. *Histochem Cell Biol*
760 124: 113-121, 2005.
- 761 27. **Franck N, Stenkula KG, Ost A, Lindstrom T, Stralfors P, and Nystrom
762 FH.** Insulin-induced GLUT4 translocation to the plasma membrane is blunted in large
763 compared with small primary fat cells isolated from the same individual. *Diabetologia*
764 50: 1716-1722, 2007.
- 765 28. **Franckhauser S, Munoz S, Pujol A, Casellas A, Riu E, Otaegui P, Su B,
766 and Bosch F.** Increased fatty acid re-esterification by PEPCK overexpression in adipose
767 tissue leads to obesity without insulin resistance. *Diabetes* 51: 624-630, 2002.
- 768 29. **Guo KY, Halo P, Leibel RL, and Zhang Y.** Effects of obesity on the
769 relationship of leptin mRNA expression and adipocyte size in anatomically distinct fat
770 depots in mice. *Am J Physiol Regul Integr Comp Physiol* 287: R112-119, 2004.

30. **Gustafson B, Hammarstedt A, Hedjazifar S, and Smith U.** Restricted adipogenesis in hypertrophic obesity: the role of WISP2, WNT, and BMP4. *Diabetes* 62: 2997-3004, 2013.
31. **Gustafson B, Jack MM, Cushman SW, and Smith U.** Adiponectin gene activation by thiazolidinediones requires PPAR gamma 2, but not C/EBP alpha-evidence for differential regulation of the aP2 and adiponectin genes. *Biochem Biophys Res Commun* 308: 933-939, 2003.
32. **Gustafson B, and Smith U.** The WNT inhibitor Dickkopf 1 and bone morphogenetic protein 4 rescue adipogenesis in hypertrophic obesity in humans. *Diabetes* 61: 1217-1224, 2012.
33. **Haller H, Leonhardt W, Hanefeld M, and Julius U.** Relationship between adipocyte hypertrophy and metabolic disturbances. *Endokrinologie* 74: 63-72, 1979.
34. **Hammarstedt A, Hedjazifar S, Jenndahl L, Gogg S, Grunberg J, Gustafson B, Klimcakova E, Stich V, Langin D, Laakso M, and Smith U.** WISP2 regulates preadipocyte commitment and PPARgamma activation by BMP4. *Proc Natl Acad Sci U S A* 110: 2563-2568, 2013.
35. **Hardy K, Brand-Miller J, Brown KD, Thomas MG, and Copeland L.** The Importance of Dietary Carbohydrate in Human Evolution. *Q Rev Biol* 90: 251-268, 2015.
36. **Harms M, and Seale P.** Brown and beige fat: development, function and therapeutic potential. *Nature medicine* 19: 1252-1263, 2013.
37. **Hayes M, Chustek M, Wang Z, Gallagher D, Heshka S, Spungen A, Bauman W, and Heymsfield SB.** DXA: potential for creating a metabolic map of organ-tissue resting energy expenditure components. *Obes Res* 10: 969-977, 2002.
38. **Heilbronn LK, Rood J, Janderova L, Albu JB, Kelley DE, Ravussin E, and Smith SR.** Relationship between serum resistin concentrations and insulin resistance in nonobese, obese, and obese diabetic subjects. *J Clin Endocrinol Metab* 89: 1844-1848, 2004.
39. **Heitmann BL, Westerterp KR, Loos RJ, Sorensen TI, O'Dea K, McLean P, Jensen TK, Eisenmann J, Speakman JR, Simpson SJ, Reed DR, and Westerterp-Plantenga MS.** Obesity: lessons from evolution and the environment. *Obes Rev* 13: 910-922, 2012.
40. **Hirsch J, and Gallian E.** Methods for the determination of adipose cell size in man and animals. *J Lipid Res* 9: 110-119, 1968.
41. **Hoffstedt J, Andersson DP, Eriksson Hogling D, Theorell J, Naslund E, Thorell A, Ehrlund A, Ryden M, and Arner P.** Long-term Protective Changes in Adipose Tissue After Gastric Bypass. *Diabetes Care* 40: 77-84, 2017.
42. **Hotamisligil GS, Shargill NS, and Spiegelman BM.** Adipose expression of tumor necrosis factor-alpha: direct role in obesity-linked insulin resistance. *Science* 259: 87-91, 1993.
43. **Hube F, Birgel M, Lee YM, and Hauner H.** Expression pattern of tumour necrosis factor receptors in subcutaneous and omental human adipose tissue: role of obesity and non-insulin-dependent diabetes mellitus. *Eur J Clin Invest* 29: 672-678, 1999.
44. **Imai T, Takakuwa R, Marchand S, Dentz E, Bornert JM, Messaddeq N, Wendling O, Mark M, Desvergne B, Wahli W, Chambon P, and Metzger D.** Peroxisome proliferator-activated receptor gamma is required in mature white and brown adipocytes for their survival in the mouse. *Proc Natl Acad Sci U S A* 101: 4543-4547, 2004.

- 819 45. **Imam MU, Wang Q, Yida Z, and Wang F.** Peroxisome proliferator-
820 activated receptor gamma (PPARgamma) as a target for concurrent management of
821 diabetes and obesity-related cancer. *Current pharmaceutical design* 2017.
- 822 46. **Isakson P, Hammarstedt A, Gustafson B, and Smith U.** Impaired
823 preadipocyte differentiation in human abdominal obesity: role of Wnt, tumor necrosis
824 factor-alpha, and inflammation. *Diabetes* 58: 1550-1557, 2009.
- 825 47. **Jacobsson B, and Smith U.** Effect of cell size on lipolysis and antilipolytic
826 action of insulin in human fat cells. *J Lipid Res* 13: 651-656, 1972.
- 827 48. **Jernas M, Palming J, Sjöholm K, Jennische E, Svensson PA, Gabrielsson
828 BG, Levin M, Sjogren A, Rudemo M, Lystig TC, Carlsson B, Carlsson LM, and Lonn M.**
829 Separation of human adipocytes by size: hypertrophic fat cells display distinct gene
830 expression. *FASEB J* 20: 1540-1542, 2006.
- 831 49. **Jiang C, Ting AT, and Seed B.** PPAR-gamma agonists inhibit production of
832 monocyte inflammatory cytokines. *Nature* 391: 82-86, 1998.
- 833 50. **Jo J, Gavrilova O, Pack S, Jou W, Mullen S, Sumner AE, Cushman SW, and
834 Periwai V.** Hypertrophy and/or Hyperplasia: Dynamics of Adipose Tissue Growth. *PLoS*
835 *computational biology* 5: e1000324, 2009.
- 836 51. **Jo J, Guo J, Liu T, Mullen S, Hall KD, Cushman SW, and Periwai V.**
837 Hypertrophy-driven adipocyte death overwhelms recruitment under prolonged weight
838 gain. *Biophys J* 99: 3535-3544, 2010.
- 839 52. **Johannsen DL, Tchoukalova Y, Tam CS, Covington JD, Xie W, Schwarz
840 JM, Bajpeyi S, and Ravussin E.** Effect of 8 weeks of overfeeding on ectopic fat
841 deposition and insulin sensitivity: testing the "adipose tissue expandability" hypothesis.
842 *Diabetes Care* 37: 2789-2797, 2014.
- 843 53. **Kashiwagi A, Verso MA, Andrews J, Vasquez B, Reaven G, and Foley JE.**
844 In vitro insulin resistance of human adipocytes isolated from subjects with noninsulin-
845 dependent diabetes mellitus. *J Clin Invest* 72: 1246-1254, 1983.
- 846 54. **Keller P, Petrie JT, De Rose P, Gerin I, Wright WS, Chiang SH, Nielsen
847 AR, Fischer CP, Pedersen BK, and MacDougald OA.** Fat-specific protein 27 regulates
848 storage of triacylglycerol. *J Biol Chem* 283: 14355-14365, 2008.
- 849 55. **Kern PA, Di Gregorio GB, Lu T, Rassouli N, and Ranganathan G.**
850 Adiponectin expression from human adipose tissue: relation to obesity, insulin
851 resistance, and tumor necrosis factor-alpha expression. *Diabetes* 52: 1779-1785, 2003.
- 852 56. **Kershaw EE, and Flier JS.** Adipose tissue as an endocrine organ. *J Clin*
853 *Endocrinol Metab* 89: 2548-2556, 2004.
- 854 57. **Khan T, Muise ES, Iyengar P, Wang ZV, Chandalia M, Abate N, Zhang BB,
855 Bonaldo P, Chua S, and Scherer PE.** Metabolic dysregulation and adipose tissue
856 fibrosis: role of collagen VI. *Mol Cell Biol* 29: 1575-1591, 2009.
- 857 58. **Kloting N, Fasshauer M, Dietrich A, Kovacs P, Schon MR, Kern M,
858 Stumvoll M, and Bluher M.** Insulin-sensitive obesity. *Am J Physiol Endocrinol Metab*
859 299: E506-515, 2010.
- 860 59. **Koenen TB, Tack CJ, Kroese JM, Hermus AR, Sweep FC, van der Laak J,
861 Stalenhoef AF, de Graaf J, van Tits LJ, and Stienstra R.** Pioglitazone treatment
862 enlarges subcutaneous adipocytes in insulin-resistant patients. *J Clin Endocrinol Metab*
863 94: 4453-4457, 2009.
- 864 60. **Komai AM, Musovic S, Peris E, Alrifaiy A, El Hachmane MF, Johansson
865 M, Wernstedt Asterholm I, and Olofsson CS.** White Adipocyte Adiponectin Exocytosis
866 Is Stimulated via beta3-Adrenergic Signaling and Activation of Epac1: Catecholamine
867 Resistance in Obesity and Type 2 Diabetes. *Diabetes* 65: 3301-3313, 2016.

61. **Krotkiewski M, Bjorntorp P, Sjostrom L, and Smith U.** Impact of obesity on metabolism in men and women. Importance of regional adipose tissue distribution. *J Clin Invest* 72: 1150-1162, 1983.
62. **Laforest S, Labrecque J, Michaud A, Cianflone K, and Tchernof A.** Adipocyte size as a determinant of metabolic disease and adipose tissue dysfunction. *Crit Rev Clin Lab Sci* 52: 301-313, 2015.
63. **Lagathu C, Bastard JP, Auclair M, Maachi M, Capeau J, and Caron M.** Chronic interleukin-6 (IL-6) treatment increased IL-6 secretion and induced insulin resistance in adipocyte: prevention by rosiglitazone. *Biochem Biophys Res Commun* 311: 372-379, 2003.
64. **Larson-Meyer DE, Heilbronn LK, Redman LM, Newcomer BR, Frisard MI, Anton S, Smith SR, Alfonso A, and Ravussin E.** Effect of calorie restriction with or without exercise on insulin sensitivity, beta-cell function, fat cell size, and ectopic lipid in overweight subjects. *Diabetes Care* 29: 1337-1344, 2006.
65. **Laurencikiene J, Skurk T, Kulyte A, Heden P, Astrom G, Sjolin E, Ryden M, Hauner H, and Arner P.** Regulation of lipolysis in small and large fat cells of the same subject. *J Clin Endocrinol Metab* 96: E2045-2049, 2011.
66. **Le Lay S, Krief S, Farnier C, Lefrere I, Le Liepvre X, Bazin R, Ferre P, and Dugail I.** Cholesterol, a cell size-dependent signal that regulates glucose metabolism and gene expression in adipocytes. *J Biol Chem* 276: 16904-16910, 2001.
67. **Lee MJ, Wu Y, and Fried SK.** Adipose tissue heterogeneity: implication of depot differences in adipose tissue for obesity complications. *Mol Aspects Med* 34: 1-11, 2013.
68. **Lehmann JM, Moore LB, Smith-Oliver TA, Wilkison WO, Willson TM, and Kliewer SA.** An antidiabetic thiazolidinedione is a high affinity ligand for peroxisome proliferator-activated receptor gamma (PPAR gamma). *J Biol Chem* 270: 12953-12956, 1995.
69. **Lizunov VA, Lisinski I, Stenkula K, Zimmerberg J, and Cushman SW.** Insulin regulates fusion of GLUT4 vesicles independent of Exo70-mediated tethering. *J Biol Chem* 284: 7914-7919, 2009.
70. **Lizunov VA, Stenkula KG, Blank PS, Troy A, Lee JP, Skarulis MC, Cushman SW, and Zimmerberg J.** Human adipose cells in vitro are either refractory or responsive to insulin, reflecting host metabolic state. *PLoS One* 10: e0119291, 2015.
71. **Lonn M, Mehlig K, Bengtsson C, and Lissner L.** Adipocyte size predicts incidence of type 2 diabetes in women. *FASEB J* 24: 326-331, 2010.
72. **Lundgren M, Svensson M, Lindmark S, Renstrom F, Ruge T, and Eriksson JW.** Fat cell enlargement is an independent marker of insulin resistance and 'hyperleptinaemia'. *Diabetologia* 50: 625-633, 2007.
73. **Macaulay IC, and Voet T.** Single cell genomics: advances and future perspectives. *PLoS Genet* 10: e1004126, 2014.
74. **Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, Tirosh I, Bialas AR, Kamitaki N, Martersteck EM, Trombetta JJ, Weitz DA, Sanes JR, Shalek AK, Regev A, and McCarroll SA.** Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. *Cell* 161: 1202-1214, 2015.
75. **Majka SM, Miller HL, Helm KM, Acosta AS, Childs CR, Kong R, and Klemm DJ.** Analysis and isolation of adipocytes by flow cytometry. *Methods Enzymol* 537: 281-296, 2014.

- 915 76. **Majka SM, Miller HL, Sullivan T, Erickson PF, Kong R, Weiser-Evans M,**
916 **Nemenoff R, Moldovan R, Morandi SA, Davis JA, and Klemm DJ.** Adipose lineage
917 specification of bone marrow-derived myeloid cells. *Adipocyte* 1: 215-229, 2012.
- 918 77. **McLaughlin T, Craig C, Liu LF, Perelman D, Allister C, Spielman D, and**
919 **Cushman SW.** Adipose Cell Size and Regional Fat Deposition as Predictors of Metabolic
920 Response to Overfeeding in Insulin-Resistant and Insulin-Sensitive Humans. *Diabetes* 65:
921 1245-1254, 2016.
- 922 78. **McLaughlin T, Lamendola C, Coghlan N, Liu TC, Lerner K, Sherman A,**
923 **and Cushman SW.** Subcutaneous adipose cell size and distribution: relationship to
924 insulin resistance and body fat. *Obesity (Silver Spring)* 22: 673-680, 2014.
- 925 79. **McLaughlin T, Sherman A, Tsao P, Gonzalez O, Yee G, Lamendola C,**
926 **Reaven GM, and Cushman SW.** Enhanced proportion of small adipose cells in insulin-
927 resistant vs insulin-sensitive obese individuals implicates impaired adipogenesis.
928 *Diabetologia* 50: 1707-1715, 2007.
- 929 80. **McLaughlin TM, Liu T, Yee G, Abbasi F, Lamendola C, Reaven GM, Tsao**
930 **P, Cushman SW, and Sherman A.** Pioglitazone increases the proportion of small cells in
931 human abdominal subcutaneous adipose tissue. *Obesity (Silver Spring)* 18: 926-931,
932 2010.
- 933 81. **Meyer LK, Ciaraldi TP, Henry RR, Wittgrove AC, and Phillips SA.**
934 Adipose tissue depot and cell size dependency of adiponectin synthesis and secretion in
935 human obesity. *Adipocyte* 2: 217-226, 2013.
- 936 82. **Murakami K, Bujo H, Unoki H, and Saito Y.** High fat intake induces a
937 population of adipocytes to co-express TLR2 and TNFalpha in mice with insulin
938 resistance. *Biochem Biophys Res Commun* 354: 727-734, 2007.
- 939 83. **Murphy J, Moullec G, and Santosa S.** Factors associated with adipocyte
940 size reduction after weight loss interventions for overweight and obesity: a systematic
941 review and meta-regression. *Metabolism* 67: 31-40, 2017.
- 942 84. **Ohyama K, Nogusa Y, Suzuki K, Shinoda K, Kajimura S, and Bannai M.** A
943 combination of exercise and capsinoid supplementation additively suppresses diet-
944 induced obesity by increasing energy expenditure in mice. *Am J Physiol Endocrinol Metab*
945 308: E315-323, 2015.
- 946 85. **Osman OS, Selway JL, Kepczynska MA, Stocker CJ, O'Dowd JF,**
947 **Cawthorne MA, Arch JR, Jassim S, and Langlands K.** A novel automated image analysis
948 method for accurate adipocyte quantification. *Adipocyte* 2: 160-164, 2013.
- 949 86. **Parlee SD, Lentz SI, Mori H, and MacDougald OA.** Quantifying size and
950 number of adipocytes in adipose tissue. *Methods Enzymol* 537: 93-122, 2014.
- 951 87. **Pasarica M, Xie H, Hymel D, Bray G, Greenway F, Ravussin E, and Smith**
952 **SR.** Lower total adipocyte number but no evidence for small adipocyte depletion in
953 patients with type 2 diabetes. *Diabetes Care* 32: 900-902, 2009.
- 954 88. **Pellegrinelli V, Heuvingh J, du Roure O, Rouault C, Devulder A, Klein C,**
955 **Lacasa M, Clement E, Lacasa D, and Clement K.** Human adipocyte function is impacted
956 by mechanical cues. *J Pathol* 233: 183-195, 2014.
- 957 89. **Pontzer H, Brown MH, Raichlen DA, Dunsworth H, Hare B, Walker K,**
958 **Luke A, Dugas LR, Durazo-Arvizu R, Schoeller D, Plange-Rhule J, Bovet P, Forrester**
959 **TE, Lambert EV, Thompson ME, Shumaker RW, and Ross SR.** Metabolic acceleration
960 and the evolution of human brain size and life history. *Nature* 533: 390-392, 2016.
- 961 90. **Prattes S, Horl G, Hammer A, Blaschitz A, Graier WF, Sattler W,**
962 **Zechner R, and Steyrer E.** Intracellular distribution and mobilization of unesterified

- cholesterol in adipocytes: triglyceride droplets are surrounded by cholesterol-rich ER-like surface layer structures. *Journal of cell science* 113 (Pt 17): 2977-2989, 2000.
91. **Puri V, Ranjit S, Konda S, Nicoloro SM, Straubhaar J, Chawla A, Chouinard M, Lin C, Burkart A, Corvera S, Perugini RA, and Czech MP.** Cidea is associated with lipid droplets and insulin sensitivity in humans. *Proc Natl Acad Sci U S A* 105: 7833-7838, 2008.
92. **Randle PJ, Garland PB, Newsholme EA, and Hales CN.** Glucose Fatty-Acid Cycle - Its Role in Insulin Sensitivity and Metabolic Disturbances of Diabetes Mellitus. *Lancet* 1: 785-&, 1963.
93. **Rodbell M.** Metabolism of Isolated Fat Cells. I. Effects of Hormones on Glucose Metabolism and Lipolysis. *J Biol Chem* 239: 375-380, 1964.
94. **Rosen ED, and Spiegelman BM.** What we talk about when we talk about fat. *Cell* 156: 20-44, 2014.
95. **Rosen ED, Walkey CJ, Puigserver P, and Spiegelman BM.** Transcriptional regulation of adipogenesis. *Genes Dev* 14: 1293-1307, 2000.
96. **Ross SE, Hemati N, Longo KA, Bennett CN, Lucas PC, Erickson RL, and MacDougald OA.** Inhibition of adipogenesis by Wnt signaling. *Science* 289: 950-953, 2000.
97. **Rotter V, Nagaev I, and Smith U.** Interleukin-6 (IL-6) induces insulin resistance in 3T3-L1 adipocytes and is, like IL-8 and tumor necrosis factor-alpha, overexpressed in human fat cells from insulin-resistant subjects. *J Biol Chem* 278: 45777-45784, 2003.
98. **Ruan H, Zarnowski MJ, Cushman SW, and Lodish HF.** Standard isolation of primary adipose cells from mouse epididymal fat pads induces inflammatory mediators and down-regulates adipocyte genes. *J Biol Chem* 278: 47585-47593, 2003.
99. **Rutkowski JM, Stern JH, and Scherer PE.** The cell biology of fat expansion. *J Cell Biol* 208: 501-512, 2015.
100. **Ryden M, Andersson DP, Bergstrom IB, and Arner P.** Adipose tissue and metabolic alterations: regional differences in fat cell size and number matter, but differently: a cross-sectional study. *J Clin Endocrinol Metab* 99: E1870-1876, 2014.
101. **Salans LB, Cushman SW, and Weismann RE.** Studies of human adipose tissue. Adipose cell size and number in nonobese and obese patients. *J Clin Invest* 52: 929-941, 1973.
102. **Salans LB, Knittle JL, and Hirsch J.** The role of adipose cell size and adipose tissue insulin sensitivity in the carbohydrate intolerance of human obesity. *J Clin Invest* 47: 153-165, 1968.
103. **Saltiel AR, and Olefsky JM.** Thiazolidinediones in the treatment of insulin resistance and type II diabetes. *Diabetes* 45: 1661-1669, 1996.
104. **Schinner S, Scherbaum WA, Bornstein SR, and Barthel A.** Molecular mechanisms of insulin resistance. *Diabet Med* 22: 674-682, 2005.
105. **Schupp M, Cristancho AG, Lefterova MI, Hanniman EA, Briggs ER, Steger DJ, Qatanani M, Curtin JC, Schug J, Ochsner SA, McKenna NJ, and Lazar MA.** Re-expression of GATA2 cooperates with peroxisome proliferator-activated receptor-gamma depletion to revert the adipocyte phenotype. *J Biol Chem* 284: 9458-9464, 2009.
106. **Shoham N, Girshovitz P, Katzungold R, Shaked NT, Benayahu D, and Gefen A.** Adipocyte stiffness increases with accumulation of lipid droplets. *Biophys J* 106: 1421-1431, 2014.

107. **Shoham N, Gottlieb R, Sharabani-Yosef O, Zaretsky U, Benayahu D, and Gefen A.** Static mechanical stretching accelerates lipid production in 3T3-L1 adipocytes by activating the MEK signaling pathway. *Am J Physiol Cell Physiol* 302: C429-441, 2012.
108. **Sjostrom L, Bjorntorp P, and Vrana J.** Microscopic fat cell size measurements on frozen-cut adipose tissue in comparison with automatic determinations of osmium-fixed fat cells. *J Lipid Res* 12: 521-530, 1971.
109. **Skurk T, Alberti-Huber C, Herder C, and Hauner H.** Relationship between adipocyte size and adipokine expression and secretion. *J Clin Endocrinol Metab* 92: 1023-1033, 2007.
110. **Smith U.** Studies of human adipose tissue in culture. I. Incorporation of glucose and release of glycerol. *Anat Rec* 172: 597-602, 1972.
111. **Smith U, Sjöström L, and Björntorp P.** Comparison of two methods for determining human adipose cell size. *J Lipid Res* 12: 521-530, 1971.
112. **Spalding KL, Arner E, Westermarck PO, Bernard S, Buchholz BA, Bergmann O, Blomqvist L, Hoffstedt J, Naslund E, Britton T, Concha H, Hassan M, Ryden M, Frisen J, and Arner P.** Dynamics of fat cell turnover in humans. *Nature* 453: 783-787, 2008.
113. **Stenkula KG, Lizunov VA, Cushman SW, and Zimmerberg J.** Insulin controls the spatial distribution of GLUT4 on the cell surface through regulation of its postfusion dispersal. *Cell Metab* 12: 250-259, 2010.
114. **Stenkula KG, Stenblom EL, Montelius C, Egecioglu E, and Erlanson-Albertsson C.** Thylakoids reduce body fat and fat cell size by binding to dietary fat making it less available for absorption in high-fat fed mice. *Nutr Metab (Lond)* 14: 4, 2017.
115. **Subramanian S, Han CY, Chiba T, McMillen TS, Wang SA, Haw A, 3rd, Kirk EA, O'Brien KD, and Chait A.** Dietary cholesterol worsens adipose tissue macrophage accumulation and atherosclerosis in obese LDL receptor-deficient mice. *Arterioscler Thromb Vasc Biol* 28: 685-691, 2008.
116. **Tansey JT, Sztalryd C, Gruia-Gray J, Roush DL, Zee JV, Gavrilova O, Reitman ML, Deng CX, Li C, Kimmel AR, and Londos C.** Perilipin ablation results in a lean mouse with aberrant adipocyte lipolysis, enhanced leptin production, and resistance to diet-induced obesity. *Proc Natl Acad Sci U S A* 98: 6494-6499, 2001.
117. **Varlamov O, Somwar R, Cornea A, Kievit P, Grove KL, and Roberts CT, Jr.** Single-cell analysis of insulin-regulated fatty acid uptake in adipocytes. *Am J Physiol Endocrinol Metab* 299: E486-496, 2010.
118. **Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, and Ferrante AW, Jr.** Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest* 112: 1796-1808, 2003.
119. **Weyer C, Foley JE, Bogardus C, Tataranni PA, and Pratley RE.** Enlarged subcutaneous abdominal adipocyte size, but not obesity itself, predicts type II diabetes independent of insulin resistance. *Diabetologia* 43: 1498-1506, 2000.
120. **Virtue S, and Vidal-Puig A.** Adipose tissue expandability, lipotoxicity and the Metabolic Syndrome--an allostatic perspective. *Biochim Biophys Acta* 1801: 338-349, 2010.
121. **Vladimir A, Lizunov J-PL, Monica C. Skarulis, Joshua Zimmerberg,, and Samuel W. Cushman KGS.** Impaired Tethering and Fusion of GLUT4 in Insulin-resistant Human Adipose Cells. In: *Diabetes* 2013.

122. **Yang J, Eliasson B, Smith U, Cushman SW, and Sherman AS.** The size of large adipose cells is a predictor of insulin resistance in first-degree relatives of type 2 diabetic patients. *Obesity (Silver Spring)* 20: 932-938, 2012.
123. **Yang X, Jansson PA, Nagaev I, Jack MM, Carvalho E, Sunnerhagen KS, Cam MC, Cushman SW, and Smith U.** Evidence of impaired adipogenesis in insulin resistance. *Biochem Biophys Res Commun* 317: 1045-1051, 2004.
124. **You T, Murphy KM, Lyles MF, Demons JL, Lenchik L, and Nicklas BJ.** Addition of aerobic exercise to dietary weight loss preferentially reduces abdominal adipocyte size. *Int J Obes (Lond)* 30: 1211-1216, 2006.
125. **Yu J, Yu HC, Kim KA, Kwon KB, Park JW, Kim SZ, Kim SH, and Park BH.** Differences in the amount of lipolysis induced by atrial natriuretic peptide in small and large adipocytes. *J Pept Sci* 14: 972-977, 2008.
126. **Yue P, Jin H, Aillaud M, Deng AC, Azuma J, Asagami T, Kundu RK, Reaven GM, Quertermous T, and Tsao PS.** Apelin is necessary for the maintenance of insulin sensitivity. *Am J Physiol Endocrinol Metab* 298: E59-67, 2010.
127. **Zhang Y, Proenca R, Maffei M, Barone M, Leopold L, and Friedman JM.** Positional cloning of the mouse obese gene and its human homologue. *Nature* 372: 425-432, 1994.
128. **Zhang Y, Xie L, Gunasekar SK, Tong D, Mishra A, Gibson WJ, Wang C, Fidler T, Marthaler B, Klingelutz A, Abel ED, Samuel I, Smith JK, Cao L, and Sah R.** SWELL1 is a regulator of adipocyte size, insulin signalling and glucose homeostasis. *Nat Cell Biol* 19: 504-517, 2017.

Figure Legends

Figure 1.

Different techniques used for characterization of adipose cell function according to size. The method to obtain isolated adipocytes in a single cell suspension from adipose tissue through collagenase digestion was established by Rodbell (93). Cell pools of different sizes can be collected by filtration of the cell suspension through nylon mesh of different pore size (27, 48), or by exploiting their differences in buoyancy (12, 25). The separated cell fractions can be further analyzed (RNA or protein expression, glucose uptake and other functional assays). Protocols to sort adipose cells by flow cytometer have been established (26, 75). Microscopy and patch-clamping techniques allow single cell analysis, even though they usually include tedious experimental handling and data processing (60, 128). Correlative studies often include histological cell size measurements of intact adipose tissue. The coulter counter technique is an alternative method to attain data of cell size distribution within a broad size range (40, 79).

Figure 2.

Precursor adipose cells are differentiated into mature adipocytes, which can expand significantly if needed, having a maximal size of 300 μm in diameter. The large and very large adipocytes are less metabolically favourable, with impaired insulin response (27), increased secretion of free fatty acids (FFA) (65) and pro-inflammatory cytokines (109), and decreased secretion of adiponectin (81); together contributing to

1103 a dysregulated systemic energy metabolism. Also, dysmetabolism could be caused
1104 by an impaired differentiation of the precursor cells (21, 30) in combination with an
1105 impaired ability to expand both small and large mature adipocytes (24, 78). The
1106 distribution of fat cell size hence is dynamic, and the proportion of various fat cell
1107 sizes should be considered in relation to health.

1108

Figure 1




Figure 2

