

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36

Adipose Tissue Inflammation and Metabolic Dysfunction in Obesity

Tatsuo Kawai MD, PhD, Michael V. Autieri PhD, and Rosario Scalia MD, PhD

The Cardiovascular Research Center and The Limole Center for Integrated Lymphatic Research, Lewis Katz School of Medicine at Temple University, Philadelphia, PA

Word count: 6694

Address Correspondence to:
Rosario Scalia MD, PhD
3500 N. Broad Street
MERB Room 1049
Philadelphia, PA 19140
Fax: (215) 707-4003
Phone: (215) 707-3248
Email:rscaliam@temple.edu

Abstract

Several lines of preclinical and clinical research have confirmed that chronic low-grade inflammation of adipose tissue is mechanistically linked to metabolic disease and organ tissue complications in the overweight and obese organism. Despite this widely confirmed paradigm, numerous open questions and knowledge gaps remain to be investigated. This is mainly due to the intricately intertwined crosstalk of various pro- and anti-inflammatory signaling cascades involved in the immune response of expanding adipose depots, particularly the visceral adipose tissue. Adipose Tissue inflammation is initiated and sustained over time by dysfunctional adipocytes that secrete inflammatory adipokines and by infiltration of bone-marrow derived immune cells that signal via production of cytokines and chemokines. Despite its low-grade nature, adipose tissue inflammation negatively impacts remote organ function, a phenomenon that is considered causative of the complications of obesity. The aim of this review is to broadly present an overview of adipose tissue inflammation by highlighting the most recent reports in the scientific literature and summarizing our overall understanding of the field. We also discuss key endogenous antiinflammatory mediators and analyze their mechanistic role(s) in the pathogenesis and treatment of adipose tissue inflammation. In doing so, we hope to stimulate studies to uncover novel physiologic, cellular, and molecular targets for the treatment of obesity.

Immune responses in obesity

59

60

61

62

63

64

Introduction

Obesity is now considered as a worldwide epidemic (171) and a strong risk factor for insulin resistance, type 2 diabetes mellitus (T2DM), cardiovascular diseases (CVD), immune disorders, and non-alcoholic fatty liver disease (NAFLD), in addition to several type of cancers. Overall, obesity is associated with a reduction of quality of life, shortened life span and increased healthcare costs (17, 52, 112, 189). Because of a pathophysiology that recapitulates the more complicated picture of multifactorial chronic disease similar to the aging process, obviously obesity cannot be simply considered the result of an energy imbalance between calorie intake and expenditure. Thus, a host of metabolic abnormalities, oxidative stress, mitochondrial dysfunction, immune dysfunction and chronic low-grade inflammation have been identified in the overweight obese organism (153, 158). While it has been almost universally established that adipose tissue (AT) responds to overnutrition by mounting an immune response, the initial inflammatory trigger remains unfortunately unknown. As a result, the clinical efficacy of drugs targeting the presently discovered inflammatory pathways has been disappointing. Perhaps, as discussed later in this review, little attention has been devoted to endogenous antiinflammatory mediators, other potentially intriguing therapeutic targets in AT inflammation. In this review, we summarize the most recent findings about AT inflammation and obesity, highlighting physiologic, cellular and molecular mechanisms through which AT inflammation contributes to AT dysfunction and related systemic complications.

87 *The inflammatory phenotype of white adipose tissue*

88 White adipose tissue (WAT) is the major fat storing depot and also serves as the
89 largest endocrine organ to secrete adipokines and cytokines systemically. Adipokines
90 are involved in various metabolic and physiologic signaling cascades and regulate
91 insulin signaling, glucose uptake, fatty acid oxidation, and other energy producing and
92 metabolic processes (2). Cytokines regulate inflammation and resolution of
93 inflammation along with adaptive and reparative angiogenesis. Weight gain and
94 obesity cause a phenotypic switch of WAT which is characterized by the appearance of
95 inflamed dysfunctional adipocytes along with infiltration of immune cells in the
96 stromal vascular fraction (80, 81). Inflamed adipocytes secrete, both locally and
97 systemically, pro-inflammatory cytokines which in turn disrupt the normal function of
98 AT itself as well as that of remote organs (79). From this standpoint AT can be
99 considered as an immune and secretory organ, and obesity as an inflammatory immune
100 disease (Figure 1).

101 Both animal and human studies have confirmed the association between
102 increased adiposity and AT inflammation following excessive caloric intake. Using
103 several immunocompromised mouse models, Lee YS et al. demonstrated the essential
104 role of inflammation in the establishment of insulin resistance following long-term
105 consumption of an obesogenic western-type diet (115). A unique feature of the
106 inflammatory responses of expanding WAT is its duration and intensity, i.e., a
107 persistent, low-grade inflammation that fails to resolve. Such obesity-related chronic
108 low-grade inflammation and subsequent altered metabolism has been termed

“metaflammation” (80). Inflammation in general is an energy wasting process that enhances energy expenditure and reduce energy intake in direct and indirect manners. Directly, inflammatory cytokines, such as TNF- α , IL-1, and IL-6, induce energy expenditure by binding to signaling receptors located in the central nervous system or in the tissue of metabolic active organs. Indeed, they are provided with leptin-like properties that promote energy expenditure (9, 217) (151). Induction of leptin expression represents a molecular mechanism of the inflammatory activity. Leptin expression is increased in adipose tissue by inflammation. Leptin transcription is induced by hypoxia (66) and inflammatory mediators (67), all of which is experienced by expanding adipose depots. Furthermore, leptin receptor expression is induced by TNF- α (60), which provides a mechanism by which pro-inflammatory cytokines enhance leptin activity for energy expenditure. Leptin is an adipokine that inhibits appetite and induces energy expenditure (131), thus indirectly increasing energy expenditure.

Surprisingly, the unique features of AT inflammation induced by over nutrition are not associated with a significant increase in energy expenditure, which permits the coexistence of inflammation and weight gain in obese people (31). Nonetheless, inflammation of AT still shares several commonalities with the traditional inflammatory response, that is the infiltration of bone marrow derived immune cells and the secretion of inflammatory mediators including, but not limited to, chemokines and cytokines by adipocytes and resident immune cells. Moreover, because of its conspicuous mass and

distribution, even in the apparently healthy normal weight individual, the inflamed WAT can cause widespread systemic inflammation via release of cytokines (147).

Based on its anatomic location and morphological structure (reviewed in (176)), WAT develops diverse and unique inflammatory phenotypes. Accordingly, it is currently well established that obesity instigates a more complex and intense inflammatory reaction in visceral WAT (VAT) than in subcutaneous WAT (SAT). Indeed, VAT contains more macrophages compared with SAT in obese mice (6, 130, 145) and obese humans (28, 75, 76). With obesity and insulin resistance VAT adipocytes experiences a much higher degree of hypertrophy than SAT in both humans (74, 138) and animal models (94). Furthermore, VAT inflammation in obese humans is associated with decreased expression of lipogenic markers (166), probably due to the fact that more cells are switched to an inflammatory rather than to a lipid storage phenotype, which leads to the development of metabolic complications, such as ectopic lipid deposition in skeletal muscle and liver (166). Since ectopic lipid deposition dampens peripheral insulin signaling, VAT inflammation is considered to play a major impact on obesity-related metabolic disorders such as systemic insulin resistance and development of type 2 diabetes (74, 96, 148). While these lines of research support a higher involvement of VAT inflammation in obesity-related complications, it should be nonetheless noted that many investigators have reported convincing evidence of a role for SAT inflammation in metabolic complications (1, 148, 210). The ongoing controversy on the role that different WAT depots play in metaflammation has been partially addressed in mice. To explain the mechanism underlying the different impact of VAT

and SAT on metabolism, Rytka et al. artificially increased fat mass in mice by transplanting epididymal VAT obtained from littermates C57Bl6/J donor mice into the parietal peritoneum that drains in the caval system or the mesenterium that drains in the liver portal system. The procedure induced AT inflammation in both experimental groups of transplanted mice. However, only the mice with mesenterium transplanted fat experienced elevated IL-6 levels and FFA in the portal vein and developed impaired glucose tolerance (172). Human studies of meal FFA uptake further confirm the heterogeneity in the metabolism of VAT and SAT. Expressed relative to the same mass of adipose tissue, meal FFA uptake is greater in intra-abdominal than abdominal subcutaneous fat in both sexes (87) (124). The direct uptake of plasma FFA is also greater in omental compared to abdominal subcutaneous fat of women (99). These data suggest that the anatomical location and the unique structural feature of VAT enable it to directly feed inflammatory cytokines and metabolites to important organs, such as the liver, which play a role in the regulation of insulin action and systemic metabolism.

In the next sections, we will summarize our current understanding of the cellular and molecular mechanisms implicated in the inflammatory response of the WAT in obesity.

The role of bone marrow adipose tissue (MAT).

In healthy adults, the approximately 10% of adipose tissue located in the bone marrow also regulates whole body energy metabolism, along with local and systemic inflammatory responses. The role of MAT role in energy metabolism and inflammation

has been recently reviewed (16, 21); thus, here we will briefly discuss key relevant aspects of MAT's role in obesity. In terms of metabolic responses, MAT expresses and secretes both adiponectin and leptin. While in vitro studies have reported that primary cultured MAT express lower level of adiponectin (119, 163), studies in mice and humans have found higher levels of adiponectin expression and secretion in MAT over WAT (36). Furthermore, modulation of MAT mass is correlated to serum adiponectin abundance following caloric restriction (36) or treatment with thiazolidinediones (195). MAT also express and secrete leptin (110, 208).

The contribution though of bone marrow adipocytes to the metabolic and inflammatory complications of obesity remains somewhat controversial. Differently than visceral fat adipocytes, bone marrow adipocytes are characterized by lower expression levels of adipose-specific gene such as PPAR γ and FABP4, and higher abundance of inflammatory response signaling pathways. In vitro studies have shown that primary cultured bone marrow adipocytes secrete significant levels of IL-6, but only small levels of IL-1 and TNF alpha (109). Others instead have reported increased expression of inflammatory response genes such as TNF alpha, IL-6 and IL-1 beta (119). With high fat feeding, mice experience increases bone marrow adiposity with adipocytes that surprisingly express decreased inflammatory genes such as TNF alpha and IL-1 beta, (202). It should be noted though that MAT is unquestionably the prevalent source of immune cells for other adipose depots and metabolic organs in obesity. Thus, it is reasonable to conclude that cytokine production by MAT adipocytes might be mainly important to regulate bone marrow kinetics and that MAT

contribution to systemic inflammation in obesity is indirectly supported by production and release of immune cells.

Macrophages, Master Regulators of WAT Inflammation

Infiltration of bone marrow derived inflammatory cells is a key feature of the WAT metaflammation. It is well established that infiltration of inflammatory cells in expanding WAT almost invariably causes adipocyte dysfunction and metabolic dysfunction, such as glucose intolerance and insulin resistance. AT Macrophages (ATMs) were the first immune cell population to be discovered and studied in this process (223), although more recent reports have emphasized the role of earlier neutrophil infiltration (discussed later). During local and systemic inflammatory responses, tissue resident macrophages present antigens to initiate recruitment of other immune cells and secrete cytokines to regulate inflammatory signaling cascades in the host tissue. This physiological role of macrophages appears to be maintained also in the metaflammation of expanding WAT depots (146, 224). The relevance of this phenomenon is confirmed by several clinical reports demonstrating ATMs infiltration in the WAT of obese humans (23, 42, 43, 76, 148, 238). Furthermore, clinical studies have also clarified the relationship between ATMs infiltration in WAT and insulin resistance. WAT obtained from insulin resistant and obese patients contains more macrophages with increased expression of pro-inflammatory mediators (74), and this phenomenon strongly correlates with metabolic dysfunction (96, 226).

Activated ATMs are the main source of pro-inflammatory mediators such as TNF-alpha, iNOS, MCP-1 and IL-6 (224, 232). Hotamisligil et al. initially reported in 1993 that the gene and protein expression of TNF-a are upregulated in AT, especially in ATMs, obtained from obese animal models (82). This first observation was followed by many confirmatory studies in laboratory animals and humans showing the upregulation of proinflammatory adipokines including TNF alpha, IL6, IL18 and MCP-1 in obesity, and their role in the development of insulin resistance and type 2 diabetes (82, 83, 194, 224, 232). The importance of cytokines secreted by ATMs in whole-body metabolisms was mechanistically elucidated by Aouadi M et al. who showed that ATMs-specific silencing of TNF-a and osteopontin using siRNA improves insulin sensitivity and glucose tolerance (10). Osteopontin is a more recently studied inflammatory mediator that appears to favor infiltration, survival, and proliferation of monocytes/macrophages in WAT (201). IL-1 beta produced by the NOD-, LRR- and pyrin domain-containing 3 (NLRP3) inflammasome regulates adipose tissue inflammation during obesity (212) (113) (236). And inhibition of NLRP3 reduces pro-inflammatory cytokines and subsequent macrophage infiltration (240). Proteoglycans are also known to regulate inflammation. Recent report revealed that, in obese condition, adipocytes produce versican and macrophages produce biglycan, and the crosstalk between these proteoglycans affect adipose tissue inflammation and insulin resistance (73).

Recent studies have also emphasized the existence of a complex interplay between WAT signaling and the CNS, an organ that contributes to regulation of

metabolism at multiple steps (reviewed in (85)). A recent line of research demonstrates that factors involved in axonal growth are also implicated in WAT inflammation. Neuroimmune guidance cue netrin-1 and semaphorin 3E (SEMA3E) with its receptor olexinD1 are regulators of the development of neuronal system that are upregulated in the obese WAT where they function as chemoattractant for macrophages (169, 181, 185), also by inhibiting macrophage egression (211). Similarly, the neuronal cytokine fractalkine (CX3CL1) and its receptor modulate monocyte adhesion to adipocyte (179).

The classic accepted paradigm is that tissue resident macrophages originate from bone marrow monocytes that infiltrate tissue during physiologic immunosurveillance or in response to inflammatory events. Interestingly, diet-induced obesity increases the circulating levels of CD11b(+) monocytes, which express the chemoattractant receptor leukotriene B₄ receptor (BLT-1) and BLT-1 has been shown to sustain monocyte trafficking to WAT (191). More recently it has been shown that proinflammatory ATMs express genes involved in myelopoiesis and immune cell recruitment, a process that affects circulating levels of monocytes and neutrophils (19, 132). Thus, obesity generates a self-feeding cycle of monocyte/macrophage infiltration to sustain low-grade chronic inflammation of WAT.

While this classic monocyte/macrophage recruitment remains a confirmed mechanism of inflammatory cell infiltration in expanding WAT, a new emerging concept in the field is that of local macrophage proliferation. Cytokines such as IL-4 are reported to stimulate local macrophage proliferation in AT (86). This local proliferation of ATMs occurs mainly at crown-like structures (CLS) that surround necrotic

adipocytes, resulting in a preferential increase of M2 macrophage in WAT (69). The consequence and impact of local ATMs proliferation, especially of the M2 type, in the pathophysiology of AT dysfunction and metaflammation remains uncharacterized and fertile ground for future research in the field.

The phenotype of obese ATMs is polarized to a pro-inflammatory state, a phenomenon that has been associated to metabolic complications (58, 121, 183, 226). In lean, insulin sensitive mice, ATMs express anti-inflammatory genes such as IL-10 and arginase 1, whereas ATMs in obese mice highly express pro-inflammatory genes such as TNF- α and iNOS (120). Similarly, the major characteristic of WAT obtained from patients with metabolic abnormalities, such as type 2 diabetes patients, is adipocyte hypertrophy with high infiltration of pro-inflammatory macrophages (1). These “metabolically activated” pro-inflammatory ATMs function within the context of low-grade inflammation and therefore appear to have important distinctions compared with classically activated M1 macrophages that occur within the context of a full-fledged inflammatory reaction. In fact while M1 macrophages in acute inflammatory reactions highly express CD38, CD274 and CD319 on their cell surface, the ATMs of the obese organism do not (102); they actually show less inflammatory phenotype with different gene expression profile compared with classically activated M1 macrophages (233). As result, the expression of pro-inflammatory cytokines such as TNF- α and IL-6 in obese ATMs remains lower than classically activated M1 macrophages (102). While these observations are consistent with the occurrence of low-grade inflammation in obesity, the question remains as to how this lower degree of immune cell activation is achieved

and maintained. Current research supports the concept that the concomitant activation of endogenous anti-inflammatory cascades is responsible for dampening macrophage activation and overall WAT inflammation in obesity. Thus, recruitment of anti-inflammatory macrophages via increased IL-10 expression has been observed in WAT after HFD (58), and deletion of IL-10 from the hematopoietic system worsened HFD-induced inflammation and insulin resistance as one may have predicted (101). Similarly, we have observed increased expression levels of the proangiogenic and anti-inflammatory cytokine IL-19 in stromal vascular fraction and visceral adipocytes of mice given a HFD (unpublished observation and Figure 2). Others have implicated a role for IFN-gamma, another key regulator of macrophage activation and phenotypic switches. Thus, IFN-gamma knockout mice exhibit less inflammatory ATMs in conjunction with decreased adipocyte size and increased insulin sensitivity following HFD (141). Furthermore, interferon regulatory factor (IRF)-5 expression is upregulated in obesity, and IRF-5 expression is positively correlated with macrophage infiltration and the secretion of proinflammatory adipokines (188).

Molecular mechanisms operating within macrophages themselves have also been investigated. Thus, it has been reported that suppression of macrophage activation in obesity is partially controlled via adenosine receptors expressed on macrophages, and that these receptors could respond to accumulation of local catabolites (47, 157). In a similar line of research, the inositol-requiring enzyme 1 alpha (IRE1alpha), a key regulator of endoplasmic reticulum stress in metabolic organs, has been recently investigated. Shan B et al. have implicated IRE1alpha in the regulation of WAT

inflammation (180) by showing that myeloid-specific IRE1alpha knockout mice have a reduced inflammatory phenotype and are protected from HFD-induced obesity, insulin resistance, hyperlipidemia and hepatic steatosis (180). More clinically relevant, the peroxisome proliferator-activated receptor (PPAR)-gamma has also been implicated in regulation of insulin resistance as well as inflammation. PPAR-gamma is essential for modulation of metabolically activated macrophages, suggesting that PPAR-gamma may increase macrophages with an anti-inflammatory phenotype and therefore it improves glucose metabolism via an anti-inflammatory mechanism in addition to its direct metabolic actions (142, 167).

Despite this wealth of knowledge, currently anti-inflammatory therapies have unfortunately failed to correct the metaflammation of obesity, which poses both a puzzling research question and a therapeutic obstacle. To help explain these unexpected negative results, Scherer et al. have recently demonstrated that inflammatory cytokines signaling is paradoxically required for appropriate expansion and metabolic flexibility of WAT. They found that congenital silencing of TNF- α or IL1-beta function in diet induced obese mice further exacerbates insulin resistance and glucose intolerance due to loss of compensatory angiogenesis and insufficient WAT remodeling (227). Consistent with this study, clinical data have yielded contrasting evidence on the effect of TNF- α and IL1- beta blocking therapy on glycemic control and insulin sensitivity in obese, diabetic patients. Thus, systemic blockade of TNF- α does not improve insulin resistance in humans and blockade of IL1-beta with Canakinumab was not associated with any reduction in the rate of incident diabetes in prediabetic insulin resistance

patients (48, 56). Overall, how the balance between pro- and anti-inflammatory cytokines ultimately determines whether or not an expanding adipose depot develops a dysfunctional degree of inflammation which impairs its metabolic activity remains insufficiently explored and poorly understood. Further studies are needed to fully understand the link between cytokines and metabolism to identify new therapeutic targets.

Infiltration of Other Immune Cells

While macrophages are the most studied and undoubtedly important immune cell type involved in AT inflammation, other immune cells such as neutrophils and lymphocytes appear to play relevant mechanistic roles (125, 170) (Figure 1). Animal studies demonstrate that administration of an HFD to lean mice causes a very early infiltration of neutrophils in VAT, before the onset of insulin resistance and obesity (51). These infiltrated neutrophils initiate recruitment of macrophages and other antigen-presenting cells. Bi-directional crosstalk between adipocytes and neutrophils has been recently reported to cause WAT inflammation via IL1-beta production with the involvement of infiltrating macrophage (221). This neutrophil-dominated early inflammatory cascade was found causal to the development of hepatic insulin resistance and metabolic disorders (70, 197). Prevention of WAT neutrophil infiltration by inhibiting cytosolic phospholipase A2 alpha (cPLA2a) or the endothelial cell adhesion molecule ICAM-1 reduced TNF- α secretion from adipocytes, preserved hepatic insulin signaling, and prevented abnormal gluconeogenesis (70). Neutrophil

elastase secreted from infiltrated neutrophils was found to be one of the key components affecting metabolic functions. Neutrophil elastase inhibitors or the deletion of neutrophil elastase in mice preserved insulin sensitivity and attenuates ATMs infiltration in HFD-fed mice (197).

The infiltration of T cells into expanding adipose depots is also considered to be another important, although controversial (115), mechanism for macrophage recruitment and WAT inflammation (95, 136). Similar to ATMs, views on the immunomodulation of T-cells in expanding adipose depots remains somewhat controversial, especially for the role of regulatory T cells (Tregs). Some studies reported down regulation of Tregs in obese states, potentially resulting in chronic WAT inflammation and subsequent insulin resistance (46, 57, 152, 229). Others have negated a role for Tregs in obesity (206, 239). The natural killer T (NKT) cells have also been implicated, with more recent reports suggesting that HFD increases invariant NKT-1 cells (iNKT1) along with pro-inflammatory ATMs and decreases invariant NKT cells 10 (iNKT10) along with anti-inflammatory ATMs indicating a role for iNKT cells in the amount of M1 ATMs that invade WAT depots (37, 139). The reader should be aware also of the fact that T-cell infiltration in WAT fluctuates with weight cycling between obese states and weight loss states (8). Interestingly, weight loss induces WAT inflammation, which in turn impairs insulin sensitivity by decreasing PI3K, phosphorylated protein kinase B (PKB) and glucose transporter (GLUT) 4 expression (116). In contrast, long-term ketogenic diet containing extremely high fat causes the depletion of gamma delta T cells, and mice lacking gamma delta T cells show impaired

glucose metabolisms (63). These results suggested that gamma delta T cells had protective effects on adipose tissue inflammation.

Obesity has been also associated with upregulation of B cells (228), which, although based on limited studies, induces insulin resistance via the activation of T-cell and AMTs in obese conditions (50, 228). B cell cytokine production appears to be regulated by activation of the toll like receptors (TLRs) in type 2 diabetes inflammation (84).

Mast cells and dendritic cells also accumulate in obese AT (18, 118). Dendritic cells promote macrophage infiltration into WAT and liver (193), or maintain T cell homeostasis in WAT (165).

Lastly, the role of eosinophils has been also explored by Wu et al., who reported that eosinophils play unique roles in metabolic homeostasis by regulating alternatively activated macrophages (230).

The contribution of Adipokines.

Expanding adipocytes themselves produce various mediators in an autocrine fashion, which are able to promote or attenuate WAT inflammation (Figure 2). These mediators have both immunomodulating and metabolic functions and are collectively termed adipokines. Although the functions of adipokines and their crosstalk are not fully understood, adipokines have emerged as key regulators of WAT metabolic homeostasis, and perhaps systemic inflammation. Adipokines secreted from adipocytes not only contributed to localized WAT inflammation (108), but can also induce systemic

elevation of cytokines responsible for peripheral insulin resistance (209, 213). Adipokines also facilitate infiltration of macrophage in WAT via IRF7-mediated upregulation of MCP-1 (106) and its receptor C-C chemokine receptor-2 (CCR2) (26, 91, 128, 200, 222). An important example of this paradigm is offered by Chemerin, an adipokine associated with obesity, inflammation, angiogenesis, and metabolic syndrome. Chemerin regulates leukocyte recruitment into inflamed organs (24, 25, 34); adipose depots expressing low level of chemerin maintains normal insulin signaling and a non-inflammatory phenotype (32, 196). Calprotectin has been elucidated as another adipokine upregulated in obesity, implicated in WAT inflammation via enhancing adhesion of circulating monocytes or recruitment of macrophages (22, 33, 127).

Along with proinflammatory cytokines, adipocytes also secrete anti-inflammatory cytokines, whose abundance and secretion appear to be reduced with weight gain since obesity definitively skews the balance in favor of inflammatory adipokines. Adiponectin is considered the most important of these mediators with several seminal findings by our laboratory and others elucidating its anti-inflammatory actions (149). Silencing of adiponectin exacerbates insulin sensitivity (39, 105). Of note, the expression of adiponectin in WAT is suppressed in acute inflammatory conditions as well as in chronic obesity (53, 234), while its overexpression reduces WAT inflammation and improves insulin sensitivity (94). Several molecules have been involved in the regulation of adiponectin expression, which may represent targets of antiinflammatory drug therapy. Thus, treatment of adipocytes with glucagon-like

peptide-1 (GLP-1) analogs increase adiponectin expression (156). Estrogen also increases adiponectin expression and attenuates WAT inflammation (144, 168), which protects from insulin resistance. Notably, recent study reported women with metabolic syndrome show lower adiponectin and IL-6 level compared to men with metabolic syndrome (203), suggesting that there were sex-specific pathways regulating obesity-induced inflammation. Fat-specific deletion of Fsp27, a lipid droplet-associated protein expressed in adipocytes in mice causes attenuation of WAT inflammation via upregulation of adiponectin (241). However, Fsp27 null mice model remain insulin resistant, possibly because of hepatic steatosis (241), which underscore the complexity of multi-organ integrated functions such as the metaflammation of obesity, and the challenge of interpretation of genetic mouse models.

Taken together, these data suggested the existence of a vicious cycle between dysfunctional adipocytes and the immune system that helps perpetuate low-grade chronic inflammation in the obese organism. Time-course studies aimed at dissecting the temporal relationship between activation of relevant WAT cell types, including those found in the stromal vascular fraction, during feeding studies may provide important mechanistic information to help resolve the pathophysiology conundrum of WAT inflammation by uncovering potential therapeutic targets and intervention timing.

Molecular Mechanisms Regulating Adipocyte Inflammation

Many molecular mechanisms operating within adipocytes have been suggested as possible regulators of WAT inflammation, including endoplasmic reticulum (ER) stress, hypoxia, and cellular senescence (Figure 2). With long standing knowledge

linking ER stress with the insulin resistance of obesity and diabetes (133, 150), several recent studies have explored this field and uncovered a number of important data. In mice, a high fat diet induces ER stress and chronic inflammation in WAT via FFA-mediated increase of reactive oxygen species (ROS) (92). A more direct causal role for ER stress was demonstrated with genetic manipulation of the ER chaperone GRP78 an essential ER chaperone and a master regulator of ER homeostasis that operates as a repressor of the unfolded protein response (UPR) stress sensors. Macrophage-selective ER chaperone GRP78 knockout mice are protected from HFD-induced AT inflammation and insulin resistance (93). In addition, ER stress preconditioning protects against FFA-induced adipocyte inflammation through inhibition of the IKK/NF- κ B pathway via X-box binding protein 1 (XBP1), a modulator of the UPR (219). Interestingly, obese, insulin resistant humans also experience higher levels of circulating saturated fatty acid (FFAs) compared with lean individuals (44, 77, 134, 184). Furthermore, in obese patients adipocyte markers of ER stress significantly correlate with BMI or percent body fat (182), and a recent clinical study reported that the amount and types of FFAs in cell membrane phospholipids is related to the inflammatory phenotype of ATMs; for instance, the proportion of pro-inflammatory ATMs increased with the proportions of palmitic or palmitoleic acids (162). Conversely, omega-3 fatty acids have been shown to exert protective effects on WAT inflammation via G protein-coupled receptor 120 (GPR120), an omega-3 fatty acids receptor (143).

Metabolism of FFA is mainly regulated by cell surface expressed lipases and the cytosolic fatty acid binding protein 4 (FABP4). In mice, WAT deletion of the hormone-

sensitive lipase (HSL) that regulates mobilization of stored fat induces WAT dysfunction and fatty liver (231), which is consistent with clinical observations showing that null mutation in the HSL gene is associated with WAT inflammation or increased risk of type 2 diabetes (3). FABP4 is secreted from adipocytes where it controls lipolysis and circulating FABP4 is elevated in metabolic disorders. In vitro treatment of adipocytes with FABP4 inhibits adipocyte differentiation and instigate adipocytes inflammation via the p38/NF-kB pathway (49). In a similar line of research, Koliwad SK et al. reported that overexpression of DGAT1 in adipocytes, the enzyme that catalyzes the conversion of diacylglycerol and FFAs to triglyceride (TG), attenuates AT inflammation with improved insulin sensitivity and glucose tolerance in HFD-fed mice, despite the exacerbated accumulation of triglycerides in liver or skeletal muscle (97). Hematopoietic deletion of the fatty acid translocase and the scavenger receptor CD36 that uptakes long chain fatty acids (FA) also attenuates WAT inflammation (135).

Research has focused on the Toll Like Receptor- 4 (TLR4) in the inflammatory action of FFAs in obesity. Earlier reports suggested that FFAs serve as chemokine inducers to initiate macrophage infiltration via IKK beta and JNK pathways, and this effect of FFAs is partially through TLR4 (89). More recently it has been reported that FFAs more directly stimulate TLR4, and initiate TLR4-dependent gene expression of lipid metabolic pathways and membrane lipid composition that promote the inflammatory cascade (54, 111). Furthermore, the expression level of tenascin C, an endogenous activator of TLR4, is also increased in obese WAT (15, 35). Accordingly, TLR4 mutant mice exhibit reduced AT inflammation, preserved adiponectin expression

and increased insulin sensitivity despite higher epididymal AT mass and adipocyte hypertrophy (161). Furthermore, global TLR4 silencing protects mice from AT inflammation, probably via reduced ER stress, augmented autophagy activity, attenuated senescence (62), and reduction of oxidative stress through metabolic reprogramming of mitochondria (117). On the other hand, TLR4 silencing in myeloid cells alone does not preserve insulin sensitivity, indicating the importance of TLR4 expression in metabolic organs in obesity (88, 173). At the mechanistic level, the high mobility group box protein 1 (HMGB1) initiates TLR4-dependent inflammation via NF- κ B and P38 MAPK signaling with subsequent IL-6 secretion (68). TLR4-mediated macrophage activation (7, 29) has also been associated with alterations of gut microbiota due to changes in gut permeability following intake of fat diets (7, 29, 164). Similar to what we have discussed for cytokines, counter regulatory expression of other TLR types appears to take place in obesity. Thus, TLR9 expressed in adipocytes was found to have WAT anti-inflammatory actions, as demonstrated by significantly reduced adiponectin secretion and increased MCP1 secretion in TLR9 knockout mice (204).

Adaptive reactions induced by AT inflammation

Chronic low-grade inflammation mediates not only metabolic dysfunctions, but also the adaptive reactions necessary to maintain physiological conditions in WAT. Interestingly, many of these adaptive reactions are inflammatory in nature. For instance, cellular senescence is another contributor to WAT inflammation, and adipocytes death

is enhanced in obese mice and humans (38, 130). Furthermore, accumulation of senescent T cells and endothelial cells occurs in the obese WAT (177, 187, 216), as well as in the insulin resistant organism (126, 177). Upregulation of senescence markers such as p16, EGFP and beta-galactosidase has been reported in expanding VAT depots (177). Since senescent cells are known to secrete pro-inflammatory cytokines (40), it is quite likely that cellular senescence instigates WAT inflammation in obesity. In support of this view, mice lacking programmed cell death-4 (PDCD4), a selective protein translation inhibitor, are protected from HFD-induced obesity and WAT inflammation (220).

With expansion, AT becomes also hypoxic due to insufficient vascular supply (reviewed in (207)). The reduced oxygenation of obese fat depots remains even in the face of compensatory angiogenesis (64, 72). The ensuing hypoxia contributes to the increased secretion of inflammatory adipokines, decreased expression of adiponectin, and upregulation of hypoxia-inducible factor 1 alpha (HIF-1alpha) (78, 140, 235). HIF-1alpha increases macrophage infiltration and WAT inflammation (155). Relevant to what we discussed in the section above, Snodgrass et al. reported that FFAs exacerbated macrophage-mediated inflammation in response to HIF-1alpha (190). Indeed, HIF-1alpha favors the M1 ATMs switch by increasing saturated fatty acid stimulation of the adenine nucleotide translocase 2 (ANT2) (114), and activation of the interleukin-1 receptor associated kinase M (160). In addition to macrophages, infiltration of other myeloid cells is also mediated by HIF-1alpha (41).

Hypoxia also negatively impacts adipocytes homeostasis. Thus, hypoxia enhances adipocytokine promoter hypomethylation via HIF1-alpha and ten-eleven translocation-1 (TET1) pathway in human adipocytes (5). Taken together, hypoxia in expanded adipocytes is one of the major regulatory mechanisms regulating WAT inflammation and insulin resistance (59).

Adaptive changes have also been described in the lymphatic system of WAT. The lymphatic system is involved in adipose metabolism by regulating lipid absorption and lipid transport and the trafficking of immune cells. HFD-fed obese mice have impaired lymphatic vessel function (20, 225). Clinical studies also found evidence of lymphatic vessel dysfunction in humans (11, 65). Obesity-induced lymphatic dysfunction appears to be related to perilymphatic inflammation (61, 137, 205) and is causative of WAT and fibrosis (175), via increased lymphatic system permeability (104). Recent publications have linked adipokines such as leptin and adiponectin to lymphatic function. Leptin causes lymphedema in obese patients via morphological change of lymphatic ducts (174). Importantly, adiponectin can improve lymphoedema in obese mice by promoting lymphatics formation via AMPK/ALT/eNOS signaling in lymphatic endothelial cells (186).

Inflammation as a therapeutic target in Obesity

Attenuation of WAT inflammation in obesity has been explored in preclinical and clinical studies as a possible treatment strategy to avert the metabolic and vascular complication of obesity. Unfortunately, the data obtained have yielded controversial

results and have also made clear the difficulty of translating results obtained in animal models of obesity to humans.

Clinical reports have investigated anti-cytokine therapy in the fight against insulin resistance, based on evidence of elevated TNF- α in the serum of obese patients. The mechanistic value of this correlative association has been confirmed with studies demonstrating that administration of TNF- α to laboratory animals and humans impairs insulin sensitivity [9, 128]. In line with this results, some human studies have reported reduced glucose intolerance in overweight individuals and reduced progression to T2D in patients with rheumatoid arthritis following TNF- α neutralization [132, 133]. To the contrary, others have reported lack of improved insulin sensitivity in people with T2D patients [129, 130] or rheumatoid arthritis [131] following anti-TNF- α therapy. Discrepant results have been reported also in healthy volunteers in which TNF- α infusion increased glucose uptake in skeletal muscle [134]. Recent evidence in humans have also implicated IL1-beta. However, in the recent CANTOS trial, IL1-beta inhibition with canakinumab failed to reduce HbA1c, glucose or insulin levels in diabetic patients [150]. This dichotomy between WAT inflammation and metabolism is also observed at the cellular level and in response to diet and lifestyle intervention and even bariatric surgery. Although it is well-established that obesity causes infiltration of ATMs in WAT, calorie restriction or rapid weight loss, which improve insulin sensitivity, paradoxically increase the number of ATMs or pro-inflammatory cytokines both in animal models (98, 107, 237) and human obesity (4, 30, 71, 103, 198, 215). Thus, in spite of improvement of systemic metabolic functions, WAT inflammation or the infiltration of resident

macrophages are not changed after weight loss (123, 237). Similarly, the expression of pro-inflammatory genes in WAT is unchanged in the early phase of a very low calorie diet but increases in the later phase (192). Instead, high-intensity exercise improves macrophage polarization toward the anti-inflammatory phenotypes (14). Calorie restriction accompanied with eccentric exercise is also reported to upregulate anti-inflammatory macrophages and downregulate pro-inflammatory macrophages (122). To complicate matters, short-term overfeeding to healthy normal or overweight individuals did not cause upregulation of pro-inflammatory gene expression or macrophage infiltration in SAT, despite the increased body weight and reduction in insulin sensitivity (90, 199). What appears to be important in reducing ATMs infiltration and WT tissue inflammation is the duration of weight loss more than its total reduction (13, 27, 30, 100). Future studies should focus on understanding how chronic adaptation to weight gain and weight cycling links WAT inflammation to insulin resistance. More importantly, integrative physiological studies should be undertaken to fully evaluate the interplay between diet induced WAT inflammation and the function of other metabolic organs such as the cardiovascular system, skeletal muscle and liver.

Many researchers have also investigated the impact of rapid weight loss with bariatric surgery on WAT inflammation, but the results are not consistent. Some studies reported the reduction of WAT inflammation with the improvement of insulin sensitivity after bariatric surgery (12, 27, 214). Conversely other studies show no or only minor changes in AT inflammation after surgery (71, 103, 154). Schmitz J et al.

reported that improved inflammation and insulin action in liver could improve systemic glucose homeostasis such as insulin sensitivity after bariatric surgeries, even if the WAT inflammation parameters including the number of ATMs were not significantly affected (178). Others have reported increased neutrophil infiltration in WAT after bariatric surgery (71, 103).

Several pharmacological and genetic modification strategies have also been investigated in the fight against obesity and metaflammation. In this context, melatonin, a potent antioxidant that improves inflammatory responses and energy metabolism, has been shown to attenuate obesity-associated complications via decreasing inflammatory adipokines including IL-6, MCP-1, leptin and TNF- α (45, 55). Inhibition of glycogen synthase kinase 3 (GSK3), a protein-serine kinase that lies downstream of multiple cell signaling pathways, also reduces WAT inflammation by suppressing migration of monocytes, favoring apoptosis of macrophage, inactivating STAT3, and reducing the levels of FFAs or chemokines secreted by VAT (218).

Genetic modifications have been performed in several mouse models in which the link between WAT inflammation and insulin resistance has been carefully investigated. For example, mice overexpressing calpastatin, the endogenous inhibitor of the calcium-dependent protease calpain, show reduced adipocyte apoptosis and macrophage infiltration in WAT (129). Deletion of C1q/TNF-related protein 7 (CTRP7), a secretory protein of the C1q family, in mice is also improve WAT inflammation, insulin resistance probably via decreased oxidative and ER stress (159).

Conclusion and future direction

In this this review article we have attempted to summarize our current understanding of the role that WAT inflammation plays in the metabolic complications of obesity by highlighting the most recent preclinical and clinical studies. The last decade has clearly demonstrated the existence of a strong relationship between WAT inflammation and insulin resistance, glucose intolerance, type 2 diabetes and obesity. However, numerous knowledge gaps continue to limit our understanding of the complex interactions among adipokines and systemic pathophysiology. Investigation in WAT inflammation have several difficulties to overcome. First, other organs such as brain, liver, and skeletal muscles also contribute to metabolic dysfunction, and an improvement of WAT inflammation is usually associated with an improvement of other organ function that also impact on insulin sensitivity. These integrative metabolic responses make difficult, if not almost impossible, to isolate the mechanistic contribution of WAT inflammation to obesity complications in intact physiologic system. Second, a current limitation of the research in the field is that the fragility of adipocytes makes it difficult to preserve the integrity of the tissue under study. Thus, our current models of the basic mechanisms governing WAT remodeling in obesity are mostly based on extrapolations from *in vitro* or *ex vivo* biochemical approaches or conventional histological analysis. Although these approaches remain useful, they yield limited translational value, as they are inadequate to study integrated physiological responses. Obviously, our understanding of these phenomena would greatly benefit from the availability of tools enabling *in vivo* analysis. Third, there are substantial

differences in characteristics and cell surface markers of immune cells between rodents and humans, and therefore careful interpretation of data obtained in animal models of obesity are warranted. More importantly, genetically modified mice should be validated with the use of wild-type mice in which changes in the target gene byproducts should be observed following induction of obesity. Finally, clinical studies have a limited view of WT tissue inflammation due to the fact that they are largely based on omental fat pad because of accessibility. Since fat depots in different region of the body assume diverse inflammatory phenotypes inclusion of additional adipose depots in future studies may provide a more comprehensive view in the field. We hope the current review will stimulate new interest in this field and will promote obesity research.

647 **Acknowledgments**

648 This work was supported by grant DK096521 to R. Scalia and grants HL141108 and
649 HL117724 to M.V. Autieri.
650

Figure legends

Figure 1. The inflammatory phenotype of expanding adipose tissue. Hypertrophic adipocytes and tissue resident immune cells experience phenotype changes that halt secretion of antiinflammatory, protective cytokines to begin secretion of inflammatory adipokines and cytokines that act both locally and systemically to induce peripheral insulin resistance. The inflammatory reaction is sustained by adipocyte-derived chemoattractants such as CCR2, MCP, and SEMA3A.

Figure 2. With increased energy storage and FFAs cargo, white adipocytes undergo abnormal expansion, which results in hypoxia and remodeling-induced senescence. Hypoxia and senescence initiate and sustain chronic low-grade inflammation. Under these condition, adipocytes experience ER stress and increased ROS production. Dysfunctional adipocytes also secrete inflammatory cytokines at the expenses of production of protective adipokines, such as adiponectin. Noteworthy, inflamed adipose depots also express antiinflammatory mediators, such as IL10 and IL19, whose role in the overall regulation of the immunometabolic response of adipose tissue remains largely unexplored.

References

1. **Acosta JR, Douagi I, Andersson DP, Backdahl J, Ryden M, Arner P, and Laurencikiene J.** Increased fat cell size: a major phenotype of subcutaneous white adipose tissue in non-obese individuals with type 2 diabetes. *Diabetologia* 59: 560-570, 2016.
2. **Ahima RS, and Lazar MA.** Adipokines and the peripheral and neural control of energy balance. *Mol Endocrinol* 22: 1023-1031, 2008.
3. **Albert JS, Yerges-Armstrong LM, Horenstein RB, Pollin TI, Sreenivasan UT, Chai S, Blaner WS, Snitker S, O'Connell JR, Gong DW, Breyer RJ, 3rd, Ryan AS, McLenithan JC, Shuldiner AR, Sztalryd C, and Damcott CM.** Null mutation in hormone-sensitive lipase gene and risk of type 2 diabetes. *N Engl J Med* 370: 2307-2315, 2014.
4. **Aleman JO, Iyengar NM, Walker JM, Milne GL, Da Rosa JC, Liang Y, Giri DD, Zhou XK, Pollak MN, Hudis CA, Breslow JL, Holt PR, and Dannenberg AJ.** Effects of Rapid Weight Loss on Systemic and Adipose Tissue Inflammation and Metabolism in Obese Postmenopausal Women. *J Endocr Soc* 1: 625-637, 2017.
5. **Ali MM, Phillips SA, and Mahmoud AM.** HIF1alpha/TET1 Pathway Mediates Hypoxia-Induced Adipocytokine Promoter Hypomethylation in Human Adipocytes. *Cells* 9: 2020.
6. **Altintas MM, Azad A, Nayer B, Contreras G, Zaias J, Faul C, Reiser J, and Nayer A.** Mast cells, macrophages, and crown-like structures distinguish subcutaneous from visceral fat in mice. *J Lipid Res* 52: 480-488, 2011.
7. **Amar J, Chabo C, Waget A, Klopp P, Vachoux C, Bermudez-Humaran LG, Smirnova N, Berge M, Sulpice T, Lahtinen S, Ouwehand A, Langella P, Rautonen N, Sansonetti PJ, and Burcelin R.** Intestinal mucosal adherence and translocation of commensal bacteria at the early onset of type 2 diabetes: molecular mechanisms and probiotic treatment. *EMBO Mol Med* 3: 559-572, 2011.
8. **Anderson EK, Gutierrez DA, Kennedy A, and Hasty AH.** Weight cycling increases T-cell accumulation in adipose tissue and impairs systemic glucose tolerance. *Diabetes* 62: 3180-3188, 2013.
9. **Anforth HR, Bluth RM, Bristow A, Hopkins S, Lenczowski MJ, Luheshi G, Lundkvist J, Michaud B, Mistry Y, Van Dam AM, Zhen C, Dantzer R, Poole S, Rothwell NJ, Tilders FJ, and Wollman EE.** Biological activity and brain actions of recombinant rat interleukin-1alpha and interleukin-1beta. *Eur Cytokine Netw* 9: 279-288, 1998.

- 705 10. **Aouadi M, Tencerova M, Vangala P, Yawe JC, Nicoloso SM, Amano SU, Cohen**
706 **JL, and Czech MP.** Gene silencing in adipose tissue macrophages regulates
707 whole-body metabolism in obese mice. *Proc Natl Acad Sci U S A* 110: 8278-8283,
708 2013.
- 709 11. **Arnglim N, Simonsen L, Holst JJ, and Bulow J.** Reduced adipose tissue
710 lymphatic drainage of macromolecules in obese subjects: a possible link between
711 obesity and local tissue inflammation? *Int J Obes (Lond)* 37: 748-750, 2013.
- 712 12. **Aron-Wisnewsky J, Tordjman J, Poitou C, Darakhshan F, Hugol D, Basdevant**
713 **A, Aissat A, Guerre-Millo M, and Clement K.** Human adipose tissue
714 macrophages: m1 and m2 cell surface markers in subcutaneous and omental
715 depots and after weight loss. *J Clin Endocrinol Metab* 94: 4619-4623, 2009.
- 716 13. **Auerbach P, Nordby P, Bendtsen LQ, Mehlsen JL, Basnet SK, Vestergaard H,**
717 **Ploug T, and Stallknecht B.** Differential effects of endurance training and weight
718 loss on plasma adiponectin multimers and adipose tissue macrophages in
719 younger, moderately overweight men. *Am J Physiol Regul Integr Comp Physiol* 305:
720 R490-498, 2013.
- 721 14. **Baek KW, Lee DI, Kang SA, and Yu HS.** Differences in macrophage polarization
722 in the adipose tissue of obese mice under various levels of exercise intensity. *J*
723 *Physiol Biochem* 2020.
- 724 15. **Becerril S, Rodriguez A, Catalan V, Mendez-Gimenez L, Ramirez B, Sainz N,**
725 **Llorente M, Unamuno X, Gomez-Ambrosi J, and Fruhbeck G.** Targeted
726 disruption of the iNOS gene improves adipose tissue inflammation and fibrosis
727 in leptin-deficient ob/ob mice: role of tenascin C. *Int J Obes (Lond)* 42: 1458-1470,
728 2018.
- 729 16. **Benova A, and Tencerova M.** Obesity-Induced Changes in Bone Marrow
730 Homeostasis. *Front Endocrinol (Lausanne)* 11: 294, 2020.
- 731 17. **Berrington de Gonzalez A, Hartge P, Cerhan JR, Flint AJ, Hannan L, MacInnis**
732 **RJ, Moore SC, Tobias GS, Anton-Culver H, Freeman LB, Beeson WL, Clipp SL,**
733 **English DR, Folsom AR, Freedman DM, Giles G, Hakansson N, Henderson**
734 **KD, Hoffman-Bolton J, Hoppin JA, Koenig KL, Lee IM, Linet MS, Park Y,**
735 **Pocobelli G, Schatzkin A, Sesso HD, Weiderpass E, Willcox BJ, Wolk A,**
736 **Zeleniuch-Jacquotte A, Willett WC, and Thun MJ.** Body-mass index and
737 mortality among 1.46 million white adults. *N Engl J Med* 363: 2211-2219, 2010.
- 738 18. **Bertola A, Ciucci T, Rousseau D, Bourlier V, Duffaut C, Bonnafous S, Blin-**
739 **Wakkach C, Anty R, Iannelli A, Gugenheim J, Tran A, Bouloumie A, Gual P,**
740 **and Wakkach A.** Identification of adipose tissue dendritic cells correlated with

- 741 obesity-associated insulin-resistance and inducing Th17 responses in mice and
742 patients. *Diabetes* 61: 2238-2247, 2012.
- 743 19. **Bijnen M, Josefs T, Cuijpers I, Maalsen CJ, van de Gaar J, Vroomen M,**
744 **Wijnands E, Rensen SS, Greve JWM, Hofker MH, Biessen EAL, Stehouwer**
745 **CDA, Schalkwijk CG, and Wouters K.** Adipose tissue macrophages induce
746 hepatic neutrophil recruitment and macrophage accumulation in mice. *Gut* 67:
747 1317-1327, 2018.
- 748 20. **Blum KS, Karaman S, Proulx ST, Ochsenbein AM, Luciani P, Leroux JC,**
749 **Wolfrum C, and Detmar M.** Chronic high-fat diet impairs collecting lymphatic
750 vessel function in mice. *PLoS One* 9: e94713, 2014.
- 751 21. **Boroumand P, and Klip A.** Bone marrow adipose cells – cellular interactions and
752 changes with obesity. *Journal of Cell Science* 133: jcs238394, 2020.
- 753 22. **Bouma G, Lam-Tse WK, Wierenga-Wolf AF, Drexhage HA, and Versnel MA.**
754 Increased serum levels of MRP-8/14 in type 1 diabetes induce an increased
755 expression of CD11b and an enhanced adhesion of circulating monocytes to
756 fibronectin. *Diabetes* 53: 1979-1986, 2004.
- 757 23. **Bourlier V, Zakaroff-Girard A, Miranville A, De Barros S, Maumus M,**
758 **Sengenès C, Galitzky J, Lafontan M, Karpe F, Frayn KN, and Bouloumie A.**
759 Remodeling phenotype of human subcutaneous adipose tissue macrophages.
760 *Circulation* 117: 806-815, 2008.
- 761 24. **Bozaoglu K, Bolton K, McMillan J, Zimmet P, Jowett J, Collier G, Walder K,**
762 **and Segal D.** Chemerin is a novel adipokine associated with obesity and
763 metabolic syndrome. *Endocrinology* 148: 4687-4694, 2007.
- 764 25. **Bozaoglu K, Curran JE, Stocker CJ, Zaibi MS, Segal D, Konstantopoulos N,**
765 **Morrison S, Carless M, Dyer TD, Cole SA, Goring HH, Moses EK, Walder K,**
766 **Cawthorne MA, Blangero J, and Jowett JB.** Chemerin, a novel adipokine in the
767 regulation of angiogenesis. *J Clin Endocrinol Metab* 95: 2476-2485, 2010.
- 768 26. **Bruun JM, Lihn AS, Pedersen SB, and Richelsen B.** Monocyte chemoattractant
769 protein-1 release is higher in visceral than subcutaneous human adipose tissue
770 (AT): implication of macrophages resident in the AT. *J Clin Endocrinol Metab* 90:
771 2282-2289, 2005.
- 772 27. **Cancello R, Henegar C, Viguerie N, Taleb S, Poitou C, Rouault C, Coupaye M,**
773 **Pelloux V, Hugol D, Bouillot JL, Bouloumie A, Barbatelli G, Cinti S, Svensson**
774 **PA, Barsh GS, Zucker JD, Basdevant A, Langin D, and Clement K.** Reduction
775 of macrophage infiltration and chemoattractant gene expression changes in

- 776 white adipose tissue of morbidly obese subjects after surgery-induced weight
777 loss. *Diabetes* 54: 2277-2286, 2005.
- 778 28. **Cancello R, Tordjman J, Poitou C, Guilhem G, Bouillot JL, Hugol D, Coussieu**
779 **C, Basdevant A, Bar Hen A, Bedossa P, Guerre-Millo M, and Clement K.**
780 Increased infiltration of macrophages in omental adipose tissue is associated
781 with marked hepatic lesions in morbid human obesity. *Diabetes* 55: 1554-1561,
782 2006.
- 783 29. **Cani PD, Bibiloni R, Knauf C, Waget A, Neyrinck AM, Delzenne NM, and**
784 **Burcelin R.** Changes in gut microbiota control metabolic endotoxemia-induced
785 inflammation in high-fat diet-induced obesity and diabetes in mice. *Diabetes* 57:
786 1470-1481, 2008.
- 787 30. **Capel F, Klimcakova E, Viguerie N, Roussel B, Vitkova M, Kovacikova M,**
788 **Polak J, Kovacova Z, Galitzky J, Maoret JJ, Hanacek J, Pers TH, Bouloumie A,**
789 **Stich V, and Langin D.** Macrophages and adipocytes in human obesity: adipose
790 tissue gene expression and insulin sensitivity during calorie restriction and
791 weight stabilization. *Diabetes* 58: 1558-1567, 2009.
- 792 31. **Carneiro IP, Elliott SA, Siervo M, Padwal R, Bertoli S, Battezzati A, and Prado**
793 **CM.** Is Obesity Associated with Altered Energy Expenditure? *Adv Nutr* 7: 476-
794 487, 2016.
- 795 32. **Cash JL, Hart R, Russ A, Dixon JP, Colledge WH, Doran J, Hendrick AG,**
796 **Carlton MB, and Greaves DR.** Synthetic chemerin-derived peptides suppress
797 inflammation through ChemR23. *J Exp Med* 205: 767-775, 2008.
- 798 33. **Catalan V, Gomez-Ambrosi J, Rodriguez A, Ramirez B, Rotellar F, Valenti V,**
799 **Silva C, Gil MJ, Fernandez-Real JM, Salvador J, and Fruhbeck G.** Increased
800 levels of calprotectin in obesity are related to macrophage content: impact on
801 inflammation and effect of weight loss. *Mol Med* 17: 1157-1167, 2011.
- 802 34. **Catalan V, Gomez-Ambrosi J, Rodriguez A, Ramirez B, Rotellar F, Valenti V,**
803 **Silva C, Gil MJ, Salvador J, and Fruhbeck G.** Increased levels of chemerin and
804 its receptor, chemokine-like receptor-1, in obesity are related to inflammation:
805 tumor necrosis factor-alpha stimulates mRNA levels of chemerin in visceral
806 adipocytes from obese patients. *Surg Obes Relat Dis* 9: 306-314, 2013.
- 807 35. **Catalan V, Gomez-Ambrosi J, Rodriguez A, Ramirez B, Rotellar F, Valenti V,**
808 **Silva C, Gil MJ, Salvador J, and Fruhbeck G.** Increased tenascin C and Toll-like
809 receptor 4 levels in visceral adipose tissue as a link between inflammation and
810 extracellular matrix remodeling in obesity. *J Clin Endocrinol Metab* 97: E1880-1889,
811 2012.

- 812 36. **Cawthorn WP, Scheller EL, Learman BS, Parlee SD, Simon BR, Mori H, Ning X,**
813 **Bree AJ, Schell B, Broome DT, Soliman SS, DelProposto JL, Lumeng CN, Mitra**
814 **A, Pandit SV, Gallagher KA, Miller JD, Krishnan V, Hui SK, Bredella MA,**
815 **Fazeli PK, Klibanski A, Horowitz MC, Rosen CJ, and MacDougald OA.** Bone
816 marrow adipose tissue is an endocrine organ that contributes to increased
817 circulating adiponectin during caloric restriction. *Cell Metab* 20: 368-375, 2014.

- 818 37. **Chen D, Zhao H, Gao X, Chen S, Liu H, Zhang J, Zhang J, and Meng M.**
819 Subcutaneous administration of alpha-GalCer activates iNKT10 cells to promote
820 M2 macrophage polarization and ameliorates chronic inflammation of obese
821 adipose tissue. *Int Immunopharmacol* 77: 105948, 2019.

- 822 38. **Cinti S, Mitchell G, Barbatelli G, Murano I, Ceresi E, Faloia E, Wang S, Fortier**
823 **M, Greenberg AS, and Obin MS.** Adipocyte death defines macrophage
824 localization and function in adipose tissue of obese mice and humans. *J Lipid Res*
825 46: 2347-2355, 2005.

- 826 39. **Combs TP, Pajvani UB, Berg AH, Lin Y, Jelicks LA, Laplante M, Nawrocki AR,**
827 **Rajala MW, Parlow AF, Cheeseboro L, Ding YY, Russell RG, Lindemann D,**
828 **Hartley A, Baker GR, Obici S, Deshaies Y, Ludgate M, Rossetti L, and Scherer**
829 **PE.** A transgenic mouse with a deletion in the collagenous domain of adiponectin
830 displays elevated circulating adiponectin and improved insulin sensitivity.
831 *Endocrinology* 145: 367-383, 2004.

- 832 40. **Coppé J-P, Desprez P-Y, Krtolica A, and Campisi J.** The senescence-associated
833 secretory phenotype: the dark side of tumor suppression. *Annu Rev Pathol* 5: 99-
834 118, 2010.

- 835 41. **Cramer T, Yamanishi Y, Clausen BE, Forster I, Pawlinski R, Mackman N,**
836 **Haase VH, Jaenisch R, Corr M, Nizet V, Firestein GS, Gerber HP, Ferrara N,**
837 **and Johnson RS.** HIF-1alpha is essential for myeloid cell-mediated inflammation.
838 *Cell* 112: 645-657, 2003.

- 839 42. **Curat CA, Miranville A, Sengenès C, Diehl M, Tonus C, Busse R, and**
840 **Bouloumie A.** From blood monocytes to adipose tissue-resident macrophages:
841 induction of diapedesis by human mature adipocytes. *Diabetes* 53: 1285-1292,
842 2004.

- 843 43. **Curat CA, Wegner V, Sengenès C, Miranville A, Tonus C, Busse R, and**
844 **Bouloumie A.** Macrophages in human visceral adipose tissue: increased
845 accumulation in obesity and a source of resistin and visfatin. *Diabetologia* 49: 744-
846 747, 2006.

- 847 44. **Davis JE, Gabler NK, Walker-Daniels J, and Spurlock ME.** Tlr-4 deficiency
848 selectively protects against obesity induced by diets high in saturated fat. *Obesity*
849 (*Silver Spring*) 16: 1248-1255, 2008.
- 850 45. **de Farias T, Cruz MM, de Sa R, Severi I, Perugini J, Senzacqua M, Cerutti SM,**
851 **Giordano A, Cinti S, and Alonso-Vale MIC.** Melatonin Supplementation
852 Decreases Hypertrophic Obesity and Inflammation Induced by High-Fat Diet in
853 Mice. *Front Endocrinol (Lausanne)* 10: 750, 2019.
- 854 46. **Deiuliis J, Shah Z, Shah N, Needleman B, Mikami D, Narula V, Perry K,**
855 **Hazey J, Kampfrath T, Kollengode M, Sun Q, Satoskar AR, Lumeng C,**
856 **Moffatt-Bruce S, and Rajagopalan S.** Visceral adipose inflammation in obesity is
857 associated with critical alterations in regulatory cell numbers. *PLoS One* 6:
858 e16376, 2011.
- 859 47. **DeOliveira CC, Paiva Caria CR, Ferreira Gotardo EM, Ribeiro ML, and**
860 **Gambero A.** Role of A1 and A2A adenosine receptor agonists in adipose tissue
861 inflammation induced by obesity in mice. *Eur J Pharmacol* 799: 154-159, 2017.
- 862 48. **Dhorepatil A, Ball S, Ghosh RK, Kondapaneni M, and Lavie CJ.** Canakinumab:
863 Promises and Future in Cardiometabolic Diseases and Malignancy. *Am J Med*
864 132: 312-324, 2019.
- 865 49. **Dou HX, Wang T, Su HX, Gao DD, Xu YC, Li YX, and Wang HY.** Exogenous
866 FABP4 interferes with differentiation, promotes lipolysis and inflammation in
867 adipocytes. *Endocrine* 67: 587-596, 2020.
- 868 50. **Duffaut C, Galitzky J, Lafontan M, and Bouloumie A.** Unexpected trafficking of
869 immune cells within the adipose tissue during the onset of obesity. *Biochem*
870 *Biophys Res Commun* 384: 482-485, 2009.
- 871 51. **Elgazar-Carmon V, Rudich A, Hadad N, and Levy R.** Neutrophils transiently
872 infiltrate intra-abdominal fat early in the course of high-fat feeding. *J Lipid Res* 49:
873 1894-1903, 2008.
- 874 52. **Emerging Risk Factors C, Wormser D, Kaptoge S, Di Angelantonio E, Wood**
875 **AM, Pennells L, Thompson A, Sarwar N, Kizer JR, Lawlor DA, Nordestgaard**
876 **BG, Ridker P, Salomaa V, Stevens J, Woodward M, Sattar N, Collins R,**
877 **Thompson SG, Whitlock G, and Danesh J.** Separate and combined associations
878 of body-mass index and abdominal adiposity with cardiovascular disease:
879 collaborative analysis of 58 prospective studies. *Lancet* 377: 1085-1095, 2011.
- 880 53. **Engeli S, Feldpausch M, Gorzelniak K, Hartwig F, Heintze U, Janke J, Mohlig**
881 **M, Pfeiffer AF, Luft FC, and Sharma AM.** Association between adiponectin and
882 mediators of inflammation in obese women. *Diabetes* 52: 942-947, 2003.

- 883 54. **Erridge C, and Samani NJ.** Saturated fatty acids do not directly stimulate Toll-
884 like receptor signaling. *Arterioscler Thromb Vasc Biol* 29: 1944-1949, 2009.
- 885 55. **Farias T, Paixao RID, Cruz MM, de Sa R, Simao JJ, Antraco VJ, and Alonso-**
886 **Vale MIC.** Melatonin Supplementation Attenuates the Pro-Inflammatory
887 Adipokines Expression in Visceral Fat from Obese Mice Induced by A High-Fat
888 Diet. *Cells* 8: 2019.
- 889 56. **Ferraz-Amaro I, Arce-Franco M, Muñiz J, López-Fernández J, Hernández-**
890 **Hernández V, Franco A, Quevedo J, Martínez-Martín J, and Díaz-González F.**
891 Systemic blockade of TNF- α does not improve insulin resistance in humans.
892 *Horm Metab Res* 43: 801-808, 2011.
- 893 57. **Feuerer M, Herrero L, Cippolletta D, Naaz A, Wong J, Nayer A, Lee J, Goldfine**
894 **AB, Benoist C, Shoelson S, and Mathis D.** Lean, but not obese, fat is enriched
895 for a unique population of regulatory T cells that affect metabolic parameters.
896 *Nat Med* 15: 930-939, 2009.
- 897 58. **Fujisaka S, Usui I, Bukhari A, Ikutani M, Oya T, Kanatani Y, Tsuneyama K,**
898 **Nagai Y, Takatsu K, Urakaze M, Kobayashi M, and Tobe K.** Regulatory
899 mechanisms for adipose tissue M1 and M2 macrophages in diet-induced obese
900 mice. *Diabetes* 58: 2574-2582, 2009.
- 901 59. **Fujisaka S, Usui I, Ikutani M, Aminuddin A, Takikawa A, Tsuneyama K,**
902 **Mahmood A, Goda N, Nagai Y, Takatsu K, and Tobe K.** Adipose tissue hypoxia
903 induces inflammatory M1 polarity of macrophages in an HIF-1 α -dependent
904 and HIF-1 α -independent manner in obese mice. *Diabetologia* 56: 1403-1412,
905 2013.
- 906 60. **Gan L, Guo K, Cremona ML, McGraw TE, Leibel RL, and Zhang Y.** TNF- α up-
907 regulates protein level and cell surface expression of the leptin receptor by
908 stimulating its export via a PKC-dependent mechanism. *Endocrinology* 153: 5821-
909 5833, 2012.
- 910 61. **Garcia Nores GD, Cuzzzone DA, Albano NJ, Hespe GE, Kataru RP, Torrisi JS,**
911 **Gardenier JC, Savetsky IL, Aschen SZ, Nitti MD, and Mehrara BJ.** Obesity but
912 not high-fat diet impairs lymphatic function. *Int J Obes (Lond)* 40: 1582-1590, 2016.
- 913 62. **Ghosh AK, O'Brien M, Mau T, and Yung R.** Toll-like receptor 4 (TLR4) deficient
914 mice are protected from adipose tissue inflammation in aging. *Aging (Albany NY)*
915 9: 1971-1982, 2017.
- 916 63. **Goldberg EL, Shchukina I, Asher JL, Sidorov S, Artyomov MN, and Dixit VD.**
917 Ketogenesis activates metabolically protective $\gamma\delta$ T cells in visceral adipose tissue.
918 *Nature Metabolism* 2: 50-61, 2020.

- 919 64. **Goossens GH, Bizzarri A, Venteclef N, Essers Y, Cleutjens JP, Konings E,**
920 **Jocken JW, Cajlakovic M, Ribitsch V, Clement K, and Blaak EE.** Increased
921 adipose tissue oxygen tension in obese compared with lean men is accompanied
922 by insulin resistance, impaired adipose tissue capillarization, and inflammation.
923 *Circulation* 124: 67-76, 2011.
- 924 65. **Greene AK, Grant FD, and Slavin SA.** Lower-extremity lymphedema and
925 elevated body-mass index. *N Engl J Med* 366: 2136-2137, 2012.
- 926 66. **Grosfeld A, Andre J, Hauguel-De Mouzon S, Berra E, Pouyssegur J, and**
927 **Guerre-Millo M.** Hypoxia-inducible factor 1 transactivates the human leptin
928 gene promoter. *J Biol Chem* 277: 42953-42957, 2002.
- 929 67. **Grunfeld C, Zhao C, Fuller J, Pollack A, Moser A, Friedman J, and Feingold**
930 **KR.** Endotoxin and cytokines induce expression of leptin, the ob gene product, in
931 hamsters. *J Clin Invest* 97: 2152-2157, 1996.
- 932 68. **Gunasekaran MK, Virama-Latchoumy AL, Girard AC, Planesse C, Guerin-**
933 **Dubourg A, Ottosson L, Andersson U, Cesari M, Roche R, and Hoareau L.**
934 TLR4-dependant pro-inflammatory effects of HMGB1 on human adipocyte.
935 *Adipocyte* 5: 384-388, 2016.
- 936 69. **Haase J, Weyer U, Immig K, Kloting N, Bluher M, Eilers J, Bechmann I, and**
937 **Gericke M.** Local proliferation of macrophages in adipose tissue during obesity-
938 induced inflammation. *Diabetologia* 57: 562-571, 2014.
- 939 70. **Hadad N, Burgazliev O, Elgazar-Carmon V, Solomonov Y, Wueest S, Item F,**
940 **Konrad D, Rudich A, and Levy R.** Induction of cytosolic phospholipase a2alpha
941 is required for adipose neutrophil infiltration and hepatic insulin resistance early
942 in the course of high-fat feeding. *Diabetes* 62: 3053-3063, 2013.
- 943 71. **Hagman DK, Larson I, Kuzma JN, Cromer G, Makar K, Rubinow KB, Foster-**
944 **Schubert KE, van Yserloo B, Billing PS, Landerholm RW, Crouthamel M, Flum**
945 **DR, Cummings DE, and Kratz M.** The short-term and long-term effects of
946 bariatric/metabolic surgery on subcutaneous adipose tissue inflammation in
947 humans. *Metabolism* 70: 12-22, 2017.
- 948 72. **Halberg N, Khan T, Trujillo ME, Wernstedt-Asterholm I, Attie AD, Sherwani**
949 **S, Wang ZV, Landskroner-Eiger S, Dineen S, Magalang UJ, Brekken RA, and**
950 **Scherer PE.** Hypoxia-inducible factor 1alpha induces fibrosis and insulin
951 resistance in white adipose tissue. *Mol Cell Biol* 29: 4467-4483, 2009.
- 952 73. **Han CY, Kang I, Harten IA, Gebe JA, Chan CK, Omer M, Alonge KM, den**
953 **Hartigh LJ, Gomes Kjerulf D, Goodspeed L, Subramanian S, Wang S, Kim F,**
954 **Birk DE, Wight TN, and Chait A.** Adipocyte-Derived Versican and Macrophage-

- 955 Derived Biglycan Control Adipose Tissue Inflammation in Obesity. *Cell Rep* 31:
956 107818-107818, 2020.
- 957 74. **Hardy OT, Perugini RA, Nicoloso SM, Gallagher-Dorval K, Puri V, Straubhaar**
958 **J, and Czech MP.** Body mass index-independent inflammation in omental
959 adipose tissue associated with insulin resistance in morbid obesity. *Surg Obes*
960 *Relat Dis* 7: 60-67, 2011.
- 961 75. **Harlev A, Aricha-Tamir B, Shaco-Levy R, Tarnovski T, Bashan N, Rudich A,**
962 **Sheiner E, Press F, and Wiznitzer A.** Macrophage infiltration and stress-
963 signaling in omental and subcutaneous adipose tissue in diabetic pregnancies. *J*
964 *Matern Fetal Neonatal Med* 27: 1189-1194, 2014.
- 965 76. **Harman-Boehm I, Bluher M, Redel H, Sion-Vardy N, Ovadia S, Avinoach E,**
966 **Shai I, Kloting N, Stumvoll M, Bashan N, and Rudich A.** Macrophage
967 infiltration into omental versus subcutaneous fat across different populations:
968 effect of regional adiposity and the comorbidities of obesity. *J Clin Endocrinol*
969 *Metab* 92: 2240-2247, 2007.
- 970 77. **Holland WL, Bikman BT, Wang LP, Yuguang G, Sargent KM, Bulchand S,**
971 **Knotts TA, Shui G, Clegg DJ, Wenk MR, Pagliassotti MJ, Scherer PE, and**
972 **Summers SA.** Lipid-induced insulin resistance mediated by the
973 proinflammatory receptor TLR4 requires saturated fatty acid-induced ceramide
974 biosynthesis in mice. *J Clin Invest* 121: 1858-1870, 2011.
- 975 78. **Hosogai N, Fukuhara A, Oshima K, Miyata Y, Tanaka S, Segawa K, Furukawa**
976 **S, Tochino Y, Komuro R, Matsuda M, and Shimomura I.** Adipose tissue
977 hypoxia in obesity and its impact on adipocytokine dysregulation. *Diabetes* 56:
978 901-911, 2007.
- 979 79. **Hotamisligil GS.** Foundations of Immunometabolism and Implications for
980 Metabolic Health and Disease. *Immunity* 47: 406-420, 2017.
- 981 80. **Hotamisligil GS.** Inflammation and metabolic disorders. *Nature* 444: 860-867,
982 2006.
- 983 81. **Hotamisligil GS.** Inflammation, metaflammation and immunometabolic
984 disorders. *Nature* 542: 177-185, 2017.
- 985 82. **Hotamisligil GS, Shargill NS, and Spiegelman BM.** Adipose expression of
986 tumor necrosis factor-alpha: direct role in obesity-linked insulin resistance.
987 *Science* 259: 87-91, 1993.
- 988 83. **Huber J, Kiefer FW, Zeyda M, Ludvik B, Silberhumer GR, Prager G, Zlabinger**
989 **GJ, and Stulnig TM.** CC chemokine and CC chemokine receptor profiles in

- visceral and subcutaneous adipose tissue are altered in human obesity. *J Clin Endocrinol Metab* 93: 3215-3221, 2008.
84. **Jagannathan M, McDonnell M, Liang Y, Hasturk H, Hetzel J, Rubin D, Kantarci A, Van Dyke TE, Ganley-Leal LM, and Nikolajczyk BS.** Toll-like receptors regulate B cell cytokine production in patients with diabetes. *Diabetologia* 53: 1461-1471, 2010.
85. **Jais A, and Brüning JC.** Hypothalamic inflammation in obesity and metabolic disease. *The Journal of Clinical Investigation* 127: 24-32, 2017.
86. **Jenkins SJ, Ruckerl D, Cook PC, Jones LH, Finkelman FD, van Rooijen N, MacDonald AS, and Allen JE.** Local macrophage proliferation, rather than recruitment from the blood, is a signature of TH2 inflammation. *Science* 332: 1284-1288, 2011.
87. **Jensen MD, Sarr MG, Dumesic DA, Southorn PA, and Levine JA.** Regional uptake of meal fatty acids in humans. *Am J Physiol Endocrinol Metab* 285: E1282-1288, 2003.
88. **Jia L, Vianna CR, Fukuda M, Berglund ED, Liu C, Tao C, Sun K, Liu T, Harper MJ, Lee CE, Lee S, Scherer PE, and Elmquist JK.** Hepatocyte Toll-like receptor 4 regulates obesity-induced inflammation and insulin resistance. *Nat Commun* 5: 3878, 2014.
89. **Jiao P, Chen Q, Shah S, Du J, Tao B, Tzamelis I, Yan W, and Xu H.** Obesity-related upregulation of monocyte chemotactic factors in adipocytes: involvement of nuclear factor-kappaB and c-Jun NH2-terminal kinase pathways. *Diabetes* 58: 104-115, 2009.
90. **Johannsen DL, Tchoukalova Y, Tam CS, Covington JD, Xie W, Schwarz JM, Bajpeyi S, and Ravussin E.** Effect of 8 weeks of overfeeding on ectopic fat deposition and insulin sensitivity: testing the "adipose tissue expandability" hypothesis. *Diabetes Care* 37: 2789-2797, 2014.
91. **Kanda H, Tateya S, Tamori Y, Kotani K, Hiasa K, Kitazawa R, Kitazawa S, Miyachi H, Maeda S, Egashira K, and Kasuga M.** MCP-1 contributes to macrophage infiltration into adipose tissue, insulin resistance, and hepatic steatosis in obesity. *J Clin Invest* 116: 1494-1505, 2006.
92. **Kawasaki N, Asada R, Saito A, Kanemoto S, and Imaizumi K.** Obesity-induced endoplasmic reticulum stress causes chronic inflammation in adipose tissue. *Sci Rep* 2: 799, 2012.

- 1024 93. **Kim JH, Lee E, Friedline RH, Suk S, Jung DY, Dagdeviren S, Hu X, Inashima K,**
1025 **Noh HL, Kwon JY, Nambu A, Huh JR, Han MS, Davis RJ, Lee AS, Lee KW,**
1026 **and Kim JK.** Endoplasmic reticulum chaperone GRP78 regulates macrophage
1027 function and insulin resistance in diet-induced obesity. *FASEB J* 32: 2292-2304,
1028 2018.
- 1029 94. **Kim JY, van de Wall E, Laplante M, Azzara A, Trujillo ME, Hofmann SM,**
1030 **Schraw T, Durand JL, Li H, Li G, Jelicks LA, Mehler MF, Hui DY, Deshaies Y,**
1031 **Shulman GI, Schwartz GJ, and Scherer PE.** Obesity-associated improvements in
1032 metabolic profile through expansion of adipose tissue. *J Clin Invest* 117: 2621-2637,
1033 2007.
- 1034 95. **Kintscher U, Hartge M, Hess K, Foryst-Ludwig A, Clemenz M, Wabitsch M,**
1035 **Fischer-Posovszky P, Barth TF, Dragun D, Skurk T, Hauner H, Bluher M,**
1036 **Unger T, Wolf AM, Knippschild U, Hombach V, and Marx N.** T-lymphocyte
1037 infiltration in visceral adipose tissue: a primary event in adipose tissue
1038 inflammation and the development of obesity-mediated insulin resistance.
1039 *Arterioscler Thromb Vasc Biol* 28: 1304-1310, 2008.
- 1040 96. **Kloting N, Fasshauer M, Dietrich A, Kovacs P, Schon MR, Kern M, Stumvoll**
1041 **M, and Bluher M.** Insulin-sensitive obesity. *Am J Physiol Endocrinol Metab* 299:
1042 E506-515, 2010.
- 1043 97. **Koliwad SK, Streeper RS, Monetti M, Cornelissen I, Chan L, Terayama K,**
1044 **Naylor S, Rao M, Hubbard B, and Farese RV, Jr.** DGAT1-dependent
1045 triacylglycerol storage by macrophages protects mice from diet-induced insulin
1046 resistance and inflammation. *J Clin Invest* 120: 756-767, 2010.
- 1047 98. **Kosteli A, Sugaru E, Haemmerle G, Martin JF, Lei J, Zechner R, and Ferrante**
1048 **AW, Jr.** Weight loss and lipolysis promote a dynamic immune response in
1049 murine adipose tissue. *J Clin Invest* 120: 3466-3479, 2010.
- 1050 99. **Koutsari C, Ali AH, Mundi MS, and Jensen MD.** Storage of circulating free
1051 fatty acid in adipose tissue of postabsorptive humans: quantitative measures and
1052 implications for body fat distribution. *Diabetes* 60: 2032-2040, 2011.
- 1053 100. **Kovacikova M, Sengenès C, Kovacova Z, Siklova-Vitkova M, Klimcakova E,**
1054 **Polak J, Rossmeislova L, Bajzova M, Hejnova J, Hnevkovska Z, Bouloumie A,**
1055 **Langin D, and Stich V.** Dietary intervention-induced weight loss decreases
1056 macrophage content in adipose tissue of obese women. *Int J Obes (Lond)* 35: 91-98,
1057 2011.
- 1058 101. **Kowalski GM, Nicholls HT, Risis S, Watson NK, Kanellakis P, Bruce CR,**
1059 **Bobik A, Lancaster GI, and Febbraio MA.** Deficiency of haematopoietic-cell-

- 1060 derived IL-10 does not exacerbate high-fat-diet-induced inflammation or insulin
1061 resistance in mice. *Diabetologia* 54: 888-899, 2011.
- 1062 102. **Kratz M, Coats BR, Hisert KB, Hagman D, Mutskov V, Peris E, Schoenfelt KQ,**
1063 **Kuzma JN, Larson I, Billing PS, Landerholm RW, Crouthamel M, Gozal D,**
1064 **Hwang S, Singh PK, and Becker L.** Metabolic dysfunction drives a
1065 mechanistically distinct proinflammatory phenotype in adipose tissue
1066 macrophages. *Cell Metab* 20: 614-625, 2014.
- 1067 103. **Kratz M, Hagman DK, Kuzma JN, Foster-Schubert KE, Chan CP, Stewart S,**
1068 **van Yserloo B, Westbrook EO, Arterburn DE, Flum DR, and Cummings DE.**
1069 Improvements in glycemic control after gastric bypass occur despite persistent
1070 adipose tissue inflammation. *Obesity (Silver Spring)* 24: 1438-1445, 2016.
- 1071 104. **Kuan EL, Ivanov S, Bridenbaugh EA, Victora G, Wang W, Childs EW, Platt**
1072 **AM, Jakubzick CV, Mason RJ, Gashev AA, Nussenzweig M, Swartz MA,**
1073 **Dustin ML, Zawieja DC, and Randolph GJ.** Collecting lymphatic vessel
1074 permeability facilitates adipose tissue inflammation and distribution of antigen
1075 to lymph node-homing adipose tissue dendritic cells. *J Immunol* 194: 5200-5210,
1076 2015.
- 1077 105. **Kubota N, Terauchi Y, Yamauchi T, Kubota T, Moroi M, Matsui J, Eto K,**
1078 **Yamashita T, Kamon J, Satoh H, Yano W, Froguel P, Nagai R, Kimura S,**
1079 **Kadowaki T, and Noda T.** Disruption of adiponectin causes insulin resistance
1080 and neointimal formation. *J Biol Chem* 277: 25863-25866, 2002.
- 1081 106. **Kuroda M, Nishiguchi M, Ugawa N, Ishikawa E, Kawabata Y, Okamoto S,**
1082 **Sasaki W, Miyatake Y, Sebe M, Masumoto S, Tsutsumi R, Harada N, and**
1083 **Sakaue H.** Interferon regulatory factor 7 mediates obesity-associated MCP-1
1084 transcription. *PloS one* 15: e0233390-e0233390, 2020.
- 1085 107. **Lacerda DR, Costa KA, Silveira ALM, Rodrigues DF, Silva AN, Sabino JL,**
1086 **Pinho V, Menezes GB, Soares DD, Teixeira MM, and Ferreira AVM.** Role of
1087 adipose tissue inflammation in fat pad loss induced by fasting in lean and mildly
1088 obese mice. *J Nutr Biochem* 72: 108208, 2019.
- 1089 108. **Lago F, Dieguez C, Gomez-Reino J, and Gualillo O.** Adipokines as emerging
1090 mediators of immune response and inflammation. *Nat Clin Pract Rheumatol* 3:
1091 716-724, 2007.
- 1092 109. **Laharrague P, Fontanilles AM, Tkaczuk J, Corberand JX, Penicaud L, and**
1093 **Casteilla L.** Inflammatory/haematopoietic cytokine production by human bone
1094 marrow adipocytes. *Eur Cytokine Netw* 11: 634-639, 2000.

- 1095 110. **Laharrague P, Truel N, Fontanilles AM, Corberand JX, Penicaud L, and**
1096 **Casteilla L.** Regulation by cytokines of leptin expression in human bone marrow
1097 adipocytes. *Horm Metab Res* 32: 381-385, 2000.
- 1098 111. **Lancaster GI, Langley KG, Berglund NA, Kammoun HL, Reibe S, Estevez E,**
1099 **Weir J, Mellett NA, Pernes G, Conway JRW, Lee MKS, Timpson P, Murphy AJ,**
1100 **Masters SL, Gerondakis S, Bartonicek N, Kaczorowski DC, Dinger ME, Meikle**
1101 **PJ, Bond PJ, and Febbraio MA.** Evidence that TLR4 Is Not a Receptor for
1102 Saturated Fatty Acids but Mediates Lipid-Induced Inflammation by
1103 Reprogramming Macrophage Metabolism. *Cell Metab* 27: 1096-1110 e1095, 2018.
- 1104 112. **Lauby-Secretan B, Scoccianti C, Loomis D, Grosse Y, Bianchini F, Straif K, and**
1105 **International Agency for Research on Cancer Handbook Working G.** Body
1106 Fatness and Cancer--Viewpoint of the IARC Working Group. *N Engl J Med* 375:
1107 794-798, 2016.
- 1108 113. **Lee H-M, Kim J-J, Kim HJ, Shong M, Ku BJ, and Jo E-K.** Upregulated NLRP3
1109 Inflammasome Activation in Patients With Type 2 Diabetes. *Diabetes* 62: 194-204,
1110 2013.
- 1111 114. **Lee YS, Kim JW, Osborne O, Oh DY, Sasik R, Schenk S, Chen A, Chung H,**
1112 **Murphy A, Watkins SM, Quehenberger O, Johnson RS, and Olefsky JM.**
1113 Increased adipocyte O2 consumption triggers HIF-1alpha, causing inflammation
1114 and insulin resistance in obesity. *Cell* 157: 1339-1352, 2014.
- 1115 115. **Lee YS, Li P, Huh JY, Hwang IJ, Lu M, Kim JI, Ham M, Talukdar S, Chen A, Lu**
1116 **WJ, Bandyopadhyay GK, Schwendener R, Olefsky J, and Kim JB.**
1117 Inflammation is necessary for long-term but not short-term high-fat diet-induced
1118 insulin resistance. *Diabetes* 60: 2474-2483, 2011.
- 1119 116. **Li X, Jiang L, Yang M, Wu YW, and Sun JZ.** Impact of weight cycling on CTRP3
1120 expression, adipose tissue inflammation and insulin sensitivity in C57BL/6J mice.
1121 *Exp Ther Med* 16: 2052-2059, 2018.
- 1122 117. **Lin HY, Weng SW, Shen FC, Chang YH, Lian WS, Hsieh CH, Chuang JH, Lin**
1123 **TK, Liou CW, Chang CS, Lin CY, Su YJ, and Wang PW.** Abrogation of Toll-Like
1124 Receptor 4 Mitigates Obesity-Induced Oxidative Stress, Proinflammation, and
1125 Insulin Resistance Through Metabolic Reprogramming of Mitochondria in
1126 Adipose Tissue. *Antioxid Redox Signal* 2020.
- 1127 118. **Liu J, Divoux A, Sun J, Zhang J, Clement K, Glickman JN, Sukhova GK,**
1128 **Wolters PJ, Du J, Gorgun CZ, Doria A, Libby P, Blumberg RS, Kahn BB,**
1129 **Hotamisligil GS, and Shi GP.** Genetic deficiency and pharmacological
1130 stabilization of mast cells reduce diet-induced obesity and diabetes in mice. *Nat*
1131 *Med* 15: 940-945, 2009.

- 1132 119. **Liu LF, Shen WJ, Ueno M, Patel S, and Kraemer FB.** Characterization of age-
1133 related gene expression profiling in bone marrow and epididymal adipocytes.
1134 *BMC Genomics* 12: 212, 2011.
- 1135 120. **Lumeng CN, Bodzin JL, and Saltiel AR.** Obesity induces a phenotypic switch in
1136 adipose tissue macrophage polarization. *J Clin Invest* 117: 175-184, 2007.
- 1137 121. **Lumeng CN, Deyoung SM, Bodzin JL, and Saltiel AR.** Increased inflammatory
1138 properties of adipose tissue macrophages recruited during diet-induced obesity.
1139 *Diabetes* 56: 16-23, 2007.
- 1140 122. **Luo W, Ai L, Wang B, Wang L, Gan Y, Liu C, Jensen J, and Zhou Y.** Eccentric
1141 exercise and dietary restriction inhibits M1 macrophage polarization activated by
1142 high-fat diet-induced obesity. *Life Sci* 243: 117246, 2020.
- 1143 123. **Magkos F, Fraterrigo G, Yoshino J, Luecking C, Kirbach K, Kelly SC, de Las**
1144 **Fuentes L, He S, Okunade AL, Patterson BW, and Klein S.** Effects of Moderate
1145 and Subsequent Progressive Weight Loss on Metabolic Function and Adipose
1146 Tissue Biology in Humans with Obesity. *Cell Metab* 23: 591-601, 2016.
- 1147 124. **Marin P, Lonn L, Andersson B, Oden B, Olbe L, Bengtsson BA, and Bjorntorp**
1148 **P.** Assimilation of triglycerides in subcutaneous and intraabdominal adipose
1149 tissues in vivo in men: effects of testosterone. *The Journal of clinical endocrinology*
1150 *and metabolism* 81: 1018-1022, 1996.
- 1151 125. **McDonnell ME, Ganley-Leal LM, Mehta A, Bigornia SJ, Mott M, Rehman Q,**
1152 **Farb MG, Hess DT, Joseph L, Gokce N, and Apovian CM.** B lymphocytes in
1153 human subcutaneous adipose crown-like structures. *Obesity (Silver Spring)* 20:
1154 1372-1378, 2012.
- 1155 126. **Minamino T, Orimo M, Shimizu I, Kunieda T, Yokoyama M, Ito T, Nojima A,**
1156 **Nabetani A, Oike Y, Matsubara H, Ishikawa F, and Komuro I.** A crucial role for
1157 adipose tissue p53 in the regulation of insulin resistance. *Nat Med* 15: 1082-1087,
1158 2009.
- 1159 127. **Mortensen OH, Nielsen AR, Erikstrup C, Plomgaard P, Fischer CP, Krogh-**
1160 **Madsen R, Lindegaard B, Petersen AM, Taudorf S, and Pedersen BK.**
1161 Calprotectin--a novel marker of obesity. *PLoS One* 4: e7419, 2009.
- 1162 128. **Mulder P, van den Hoek AM, and Kleemann R.** The CCR2 Inhibitor
1163 Propagermanium Attenuates Diet-Induced Insulin Resistance, Adipose Tissue
1164 Inflammation and Non-Alcoholic Steatohepatitis. *PLoS One* 12: e0169740, 2017.
- 1165 129. **Muniappan L, Javidan A, Jiang W, Mohammadmoradi S, Moorlegheh JJ, Katz**
1166 **WS, Balakrishnan A, Howatt DA, and Subramanian V.** Calpain Inhibition

- 1167 Attenuates Adipose Tissue Inflammation and Fibrosis in Diet-induced Obese
1168 Mice. *Sci Rep* 7: 14398, 2017.
- 1169 130. **Murano I, Barbatelli G, Parisani V, Latini C, Muzzonigro G, Castellucci M, and**
1170 **Cinti S.** Dead adipocytes, detected as crown-like structures, are prevalent in
1171 visceral fat depots of genetically obese mice. *J Lipid Res* 49: 1562-1568, 2008.
- 1172 131. **Myers MG, Jr., and Olson DP.** Central nervous system control of metabolism.
1173 *Nature* 491: 357-363, 2012.
- 1174 132. **Nagareddy PR, Kraakman M, Masters SL, Stirzaker RA, Gorman DJ, Grant**
1175 **RW, Dragoljevic D, Hong ES, Abdel-Latif A, Smyth SS, Choi SH, Korner J,**
1176 **Bornfeldt KE, Fisher EA, Dixit VD, Tall AR, Goldberg IJ, and Murphy AJ.**
1177 Adipose tissue macrophages promote myelopoiesis and monocytosis in obesity.
1178 *Cell Metab* 19: 821-835, 2014.
- 1179 133. **Nakatani Y, Kaneto H, Kawamori D, Yoshiuchi K, Hatazaki M, Matsuoka TA,**
1180 **Ozawa K, Ogawa S, Hori M, Yamasaki Y, and Matsuhisa M.** Involvement of
1181 endoplasmic reticulum stress in insulin resistance and diabetes. *J Biol Chem* 280:
1182 847-851, 2005.
- 1183 134. **Nguyen MT, Faveleyukis S, Nguyen AK, Reichart D, Scott PA, Jenn A, Liu-**
1184 **Bryan R, Glass CK, Neels JG, and Olefsky JM.** A subpopulation of
1185 macrophages infiltrates hypertrophic adipose tissue and is activated by free fatty
1186 acids via Toll-like receptors 2 and 4 and JNK-dependent pathways. *J Biol Chem*
1187 282: 35279-35292, 2007.
- 1188 135. **Nicholls HT, Kowalski G, Kennedy DJ, Risis S, Zaffino LA, Watson N,**
1189 **Kanellakis P, Watt MJ, Bobik A, Bonen A, Febbraio M, Lancaster GI, and**
1190 **Febbraio MA.** Hematopoietic cell-restricted deletion of CD36 reduces high-fat
1191 diet-induced macrophage infiltration and improves insulin signaling in adipose
1192 tissue. *Diabetes* 60: 1100-1110, 2011.
- 1193 136. **Nishimura S, Manabe I, Nagasaki M, Eto K, Yamashita H, Ohsugi M, Otsu M,**
1194 **Hara K, Ueki K, Sugiura S, Yoshimura K, Kadowaki T, and Nagai R.** CD8+
1195 effector T cells contribute to macrophage recruitment and adipose tissue
1196 inflammation in obesity. *Nat Med* 15: 914-920, 2009.
- 1197 137. **Nitti MD, Hespe GE, Kataru RP, Garcia Nores GD, Savetsky IL, Torrisi JS,**
1198 **Gardenier JC, Dannenberg AJ, and Mehrara BJ.** Obesity-induced lymphatic
1199 dysfunction is reversible with weight loss. *J Physiol* 594: 7073-7087, 2016.
- 1200 138. **O'Connell J, Lynch L, Cawood TJ, Kwasnik A, Nolan N, Geoghegan J,**
1201 **McCormick A, O'Farrelly C, and O'Shea D.** The relationship of omental and

- 1202 subcutaneous adipocyte size to metabolic disease in severe obesity. *PLoS One* 5:
1203 e9997, 2010.
- 1204 139. **O'Rourke RW, Metcalf MD, White AE, Madala A, Winters BR, Maizlin, II,**
1205 **Jobe BA, Roberts CT, Jr., Slifka MK, and Marks DL.** Depot-specific differences
1206 in inflammatory mediators and a role for NK cells and IFN-gamma in
1207 inflammation in human adipose tissue. *Int J Obes (Lond)* 33: 978-990, 2009.
- 1208 140. **O'Rourke RW, White AE, Metcalf MD, Olivas AS, Mitra P, Larison WG,**
1209 **Cheang EC, Varlamov O, Corless CL, Roberts CT, Jr., and Marks DL.** Hypoxia-
1210 induced inflammatory cytokine secretion in human adipose tissue
1211 stromovascular cells. *Diabetologia* 54: 1480-1490, 2011.
- 1212 141. **O'Rourke RW, White AE, Metcalf MD, Winters BR, Diggs BS, Zhu X, and**
1213 **Marks DL.** Systemic inflammation and insulin sensitivity in obese IFN-gamma
1214 knockout mice. *Metabolism* 61: 1152-1161, 2012.
- 1215 142. **Odegaard JI, Ricardo-Gonzalez RR, Goforth MH, Morel CR, Subramanian V,**
1216 **Mukundan L, Red Eagle A, Vats D, Brombacher F, Ferrante AW, and Chawla A.**
1217 Macrophage-specific PPARgamma controls alternative activation and improves
1218 insulin resistance. *Nature* 447: 1116-1120, 2007.
- 1219 143. **Oh DY, Talukdar S, Bae EJ, Imamura T, Morinaga H, Fan W, Li P, Lu WJ,**
1220 **Watkins SM, and Olefsky JM.** GPR120 is an omega-3 fatty acid receptor
1221 mediating potent anti-inflammatory and insulin-sensitizing effects. *Cell* 142: 687-
1222 698, 2010.
- 1223 144. **Ohlsson C, Hammarstedt A, Vandenput L, Saarinen N, Ryberg H, Windahl SH,**
1224 **Farman HH, Jansson JO, Moverare-Skrtic S, Smith U, Zhang FP, Poutanen M,**
1225 **Hedjazifar S, and Sjogren K.** Increased adipose tissue aromatase activity
1226 improves insulin sensitivity and reduces adipose tissue inflammation in male
1227 mice. *Am J Physiol Endocrinol Metab* 313: E450-E462, 2017.
- 1228 145. **Okamoto Y, Higashiyama H, Rong JX, McVey MJ, Kinoshita M, Asano S, and**
1229 **Hansen MK.** Comparison of mitochondrial and macrophage content between
1230 subcutaneous and visceral fat in db/db mice. *Exp Mol Pathol* 83: 73-83, 2007.
- 1231 146. **Olefsky JM, and Glass CK.** Macrophages, inflammation, and insulin resistance.
1232 *Annu Rev Physiol* 72: 219-246, 2010.
- 1233 147. **Oliveros E, Somers VK, Sochor O, Goel K, and Lopez-Jimenez F.** The concept
1234 of normal weight obesity. *Prog Cardiovasc Dis* 56: 426-433, 2014.
- 1235 148. **Ortega Martinez de Victoria E, Xu X, Koska J, Francisco AM, Scalise M,**
1236 **Ferrante AW, Jr., and Krakoff J.** Macrophage content in subcutaneous adipose

- 1237 tissue: associations with adiposity, age, inflammatory markers, and whole-body
1238 insulin action in healthy Pima Indians. *Diabetes* 58: 385-393, 2009.
- 1239 149. **Ouedraogo R, Gong Y, Berzins B, Wu X, Mahadev K, Hough K, Chan L,**
1240 **Goldstein BJ, and Scalia R.** Adiponectin deficiency increases leukocyte-
1241 endothelium interactions via upregulation of endothelial cell adhesion molecules
1242 in vivo. *The Journal of Clinical Investigation* 117: 1718-1726, 2007.
- 1243 150. **Ozcan U, Cao Q, Yilmaz E, Lee AH, Iwakoshi NN, Ozdelen E, Tuncman G,**
1244 **Gorgun C, Glimcher LH, and Hotamisligil GS.** Endoplasmic reticulum stress
1245 links obesity, insulin action, and type 2 diabetes. *Science* 306: 457-461, 2004.
- 1246 151. **Pamir N, McMillen TS, Kaiyala KJ, Schwartz MW, and LeBoeuf RC.** Receptors
1247 for tumor necrosis factor- α play a protective role against obesity and alter
1248 adipose tissue macrophage status. *Endocrinology* 150: 4124-4134, 2009.
- 1249 152. **Pandolfi JB, Ferraro AA, Sananez I, Gancedo MC, Baz P, Billordo LA,**
1250 **Fainboim L, and Arruvito L.** ATP-Induced Inflammation Drives Tissue-Resident
1251 Th17 Cells in Metabolically Unhealthy Obesity. *J Immunol* 196: 3287-3296, 2016.
- 1252 153. **Pararasa C, Bailey CJ, and Griffiths HR.** Ageing, adipose tissue, fatty acids and
1253 inflammation. *Biogerontology* 16: 235-248, 2015.
- 1254 154. **Pardina E, Ferrer R, Baena-Fustegueras JA, Rivero J, Lecube A, Fort JM, Vargas**
1255 **V, Catalan R, and Peinado-Onsurbe J.** Only C-reactive protein, but not TNF-
1256 α or IL6, reflects the improvement in inflammation after bariatric surgery.
1257 *Obes Surg* 22: 131-139, 2012.
- 1258 155. **Pasarica M, Sereda OR, Redman LM, Albarado DC, Hymel DT, Roan LE, Rood**
1259 **JC, Burk DH, and Smith SR.** Reduced adipose tissue oxygenation in human
1260 obesity: evidence for rarefaction, macrophage chemotaxis, and inflammation
1261 without an angiogenic response. *Diabetes* 58: 718-725, 2009.
- 1262 156. **Pastel E, Joshi S, Knight B, Liversedge N, Ward R, and Kos K.** Effects of
1263 Exendin-4 on human adipose tissue inflammation and ECM remodelling. *Nutr*
1264 *Diabetes* 6: e235, 2016.
- 1265 157. **Pei Y, Li H, Cai Y, Zhou J, Luo X, Ma L, McDaniel K, Zeng T, Chen Y, Qian X,**
1266 **Huo Y, Glaser S, Meng F, Alpini G, Chen L, and Wu C.** Regulation of adipose
1267 tissue inflammation by adenosine 2A receptor in obese mice. *J Endocrinol* 239:
1268 365-376, 2018.
- 1269 158. **Perez LM, Pareja-Galeano H, Sanchis-Gomar F, Emanuele E, Lucia A, and**
1270 **Galvez BG.** 'Adipaging': ageing and obesity share biological hallmarks related to
1271 a dysfunctional adipose tissue. *J Physiol* 594: 3187-3207, 2016.

- 1272 159. **Petersen PS, Lei X, Wolf RM, Rodriguez S, Tan SY, Little HC, Schweitzer MA,**
1273 **Magnuson TH, Steele KE, and Wong GW.** CTRP7 deletion attenuates obesity-
1274 linked glucose intolerance, adipose tissue inflammation, and hepatic stress. *Am J*
1275 *Physiol Endocrinol Metab* 312: E309-E325, 2017.
- 1276 160. **Poblete JMS, Ballinger MN, Bao S, Alghothani M, Nevado JB, Jr., Eubank TD,**
1277 **Christman JW, and Magalang UJ.** Macrophage HIF-1alpha Mediates Obesity-
1278 related Adipose Tissue Dysfunction via Interleukin-1 receptor-associated kinase
1279 M. *Am J Physiol Endocrinol Metab* 2020.
- 1280 161. **Poggi M, Bastelica D, Gual P, Iglesias MA, Gremeaux T, Knauf C, Peiretti F,**
1281 **Verdier M, Juhan-Vague I, Tanti JF, Burcelin R, and Alessi MC.** C3H/HeJ mice
1282 carrying a toll-like receptor 4 mutation are protected against the development of
1283 insulin resistance in white adipose tissue in response to a high-fat diet.
1284 *Diabetologia* 50: 1267-1276, 2007.
- 1285 162. **Poledne R, Malinska H, Kubatova H, Froncek J, Thieme F, Kaueroval S, and**
1286 **Lesna IK.** Polarization of Macrophages in Human Adipose Tissue is Related to
1287 the Fatty Acid Spectrum in Membrane Phospholipids. *Nutrients* 12: 2019.
- 1288 163. **Poloni A, Maurizi G, Serrani F, Mancini S, Zingaretti MC, Frontini A, Cinti S,**
1289 **Olivieri A, and Leoni P.** Molecular and functional characterization of human
1290 bone marrow adipocytes. *Exp Hematol* 41: 558-566 e552, 2013.
- 1291 164. **Poroyko VA, Carreras A, Khalyfa A, Khalyfa AA, Leone V, Peris E, Almendros**
1292 **I, Gileles-Hillel A, Qiao Z, Hubert N, Farre R, Chang EB, and Gozal D.** Chronic
1293 Sleep Disruption Alters Gut Microbiota, Induces Systemic and Adipose Tissue
1294 Inflammation and Insulin Resistance in Mice. *Sci Rep* 6: 35405, 2016.
- 1295 165. **Porsche CE, Delproposto JB, Patrick E, Zamarron BF, and Lumeng CN.**
1296 Adipose tissue dendritic cell signals are required to maintain T cell homeostasis
1297 and obesity-induced expansion. *Mol Cell Endocrinol* 505: 110740, 2020.
- 1298 166. **Poulain-Godefroy O, Lecoer C, Pattou F, Fruhbeck G, and Froguel P.**
1299 Inflammation is associated with a decrease of lipogenic factors in omental fat in
1300 women. *Am J Physiol Regul Integr Comp Physiol* 295: R1-7, 2008.
- 1301 167. **Prieur X, Mok CY, Velagapudi VR, Nunez V, Fuentes L, Montaner D, Ishikawa**
1302 **K, Camacho A, Barbarroja N, O'Rahilly S, Sethi JK, Dopazo J, Oresic M, Ricote**
1303 **M, and Vidal-Puig A.** Differential lipid partitioning between adipocytes and
1304 tissue macrophages modulates macrophage lipotoxicity and M2/M1 polarization
1305 in obese mice. *Diabetes* 60: 797-809, 2011.
- 1306 168. **Qi X, Zhang B, Zhao Y, Li R, Chang HM, Pang Y, and Qiao J.**
1307 Hyperhomocysteinemia Promotes Insulin Resistance and Adipose Tissue

- 1308 Inflammation in PCOS Mice Through Modulating M2 Macrophage Polarization
1309 via Estrogen Suppression. *Endocrinology* 158: 1181-1193, 2017.
- 1310 169. **Ramkhelawon B, Hennessy EJ, Menager M, Ray TD, Sheedy FJ, Hutchison S,**
1311 **Wanschel A, Oldebeken S, Geoffrion M, Spiro W, Miller G, McPherson R,**
1312 **Rayner KJ, and Moore KJ.** Netrin-1 promotes adipose tissue macrophage
1313 retention and insulin resistance in obesity. *Nat Med* 20: 377-384, 2014.
- 1314 170. **Rausch ME, Weisberg S, Vardhana P, and Tortoriello DV.** Obesity in C57BL/6J
1315 mice is characterized by adipose tissue hypoxia and cytotoxic T-cell infiltration.
1316 *Int J Obes (Lond)* 32: 451-463, 2008.
- 1317 171. **Roberto CA, Swinburn B, Hawkes C, Huang TT, Costa SA, Ashe M, Zwicker L,**
1318 **Cawley JH, and Brownell KD.** Patchy progress on obesity prevention: emerging
1319 examples, entrenched barriers, and new thinking. *Lancet* 385: 2400-2409, 2015.
- 1320 172. **Rytka JM, Wueest S, Schoenle EJ, and Konrad D.** The portal theory supported
1321 by venous drainage-selective fat transplantation. *Diabetes* 60: 56-63, 2011.
- 1322 173. **Saberi M, Woods NB, de Luca C, Schenk S, Lu JC, Bandyopadhyay G, Verma**
1323 **IM, and Olefsky JM.** Hematopoietic cell-specific deletion of toll-like receptor 4
1324 ameliorates hepatic and adipose tissue insulin resistance in high-fat-fed mice.
1325 *Cell Metab* 10: 419-429, 2009.
- 1326 174. **Sato A, Kamekura R, Kawata K, Kawada M, Jitsukawa S, Yamashita K, Sato N,**
1327 **Himi T, and Ichimiya S.** Novel Mechanisms of Compromised Lymphatic
1328 Endothelial Cell Homeostasis in Obesity: The Role of Leptin in Lymphatic
1329 Endothelial Cell Tube Formation and Proliferation. *PLoS One* 11: e0158408, 2016.
- 1330 175. **Savetsky IL, Torrisi JS, Cuzzzone DA, Ghanta S, Albano NJ, Gardenier JC,**
1331 **Joseph WJ, and Mehrara BJ.** Obesity increases inflammation and impairs
1332 lymphatic function in a mouse model of lymphedema. *Am J Physiol Heart Circ*
1333 *Physiol* 307: H165-172, 2014.
- 1334 176. **Scalia R.** The microcirculation in adipose tissue inflammation. *Rev Endocr Metab*
1335 *Disord* 14: 69-76, 2013.
- 1336 177. **Schafer MJ, White TA, Evans G, Tonne JM, Verzosa GC, Stout MB, Mazula DL,**
1337 **Palmer AK, Baker DJ, Jensen MD, Torbenson MS, Miller JD, Ikeda Y,**
1338 **Tchkonia T, van Deursen JM, Kirkland JL, and LeBrasseur NK.** Exercise
1339 Prevents Diet-Induced Cellular Senescence in Adipose Tissue. *Diabetes* 65: 1606-
1340 1615, 2016.
- 1341 178. **Schmitz J, Evers N, Awazawa M, Nicholls HT, Bronneke HS, Dietrich A,**
1342 **Mauer J, Bluher M, and Bruning JC.** Obesogenic memory can confer long-term

- 1343 increases in adipose tissue but not liver inflammation and insulin resistance after
1344 weight loss. *Mol Metab* 5: 328-339, 2016.
- 1345 179. **Shah R, Hinkle CC, Ferguson JF, Mehta NN, Li M, Qu L, Lu Y, Putt ME, Ahima**
1346 **RS, and Reilly MP.** Fractalkine is a novel human adipochemokine associated
1347 with type 2 diabetes. *Diabetes* 60: 1512-1518, 2011.
- 1348 180. **Shan B, Wang X, Wu Y, Xu C, Xia Z, Dai J, Shao M, Zhao F, He S, Yang L,**
1349 **Zhang M, Nan F, Li J, Liu J, Liu J, Jia W, Qiu Y, Song B, Han JJ, Rui L, Duan SZ,**
1350 **and Liu Y.** The metabolic ER stress sensor IRE1alpha suppresses alternative
1351 activation of macrophages and impairs energy expenditure in obesity. *Nat*
1352 *Immunol* 18: 519-529, 2017.
- 1353 181. **Sharma M, Schlegel M, Brown EJ, Sansbury BE, Weinstock A, Afonso MS,**
1354 **Corr EM, van Solingen C, Shanley LC, Peled D, Ramasamy R, Schmidt AM,**
1355 **Spite M, Fisher EA, and Moore KJ.** Netrin-1 Alters Adipose Tissue Macrophage
1356 Fate and Function in Obesity. *Immunometabolism* 1: 2019.
- 1357 182. **Sharma NK, Das SK, Mondal AK, Hackney OG, Chu WS, Kern PA, Rasouli N,**
1358 **Spencer HJ, Yao-Borengasser A, and Elbein SC.** Endoplasmic reticulum stress
1359 markers are associated with obesity in nondiabetic subjects. *J Clin Endocrinol*
1360 *Metab* 93: 4532-4541, 2008.
- 1361 183. **Shaul ME, Bennett G, Strissel KJ, Greenberg AS, and Obin MS.** Dynamic, M2-
1362 like remodeling phenotypes of CD11c+ adipose tissue macrophages during high-
1363 fat diet--induced obesity in mice. *Diabetes* 59: 1171-1181, 2010.
- 1364 184. **Shi H, Kokoeva MV, Inouye K, Tzameli I, Yin H, and Flier JS.** TLR4 links
1365 innate immunity and fatty acid-induced insulin resistance. *J Clin Invest* 116: 3015-
1366 3025, 2006.
- 1367 185. **Shimizu I, Yoshida Y, Moriya J, Nojima A, Uemura A, Kobayashi Y, and**
1368 **Minamino T.** Semaphorin3E-induced inflammation contributes to insulin
1369 resistance in dietary obesity. *Cell Metab* 18: 491-504, 2013.
- 1370 186. **Shimizu Y, Shibata R, Ishii M, Ohashi K, Kambara T, Uemura Y, Yuasa D,**
1371 **Kataoka Y, Kihara S, Murohara T, and Ouchi N.** Adiponectin-mediated
1372 modulation of lymphatic vessel formation and lymphedema. *J Am Heart Assoc* 2:
1373 e000438, 2013.
- 1374 187. **Shirakawa K, Yan X, Shinmura K, Endo J, Kataoka M, Katsumata Y, Yamamoto**
1375 **T, Anzai A, Isobe S, Yoshida N, Itoh H, Manabe I, Sekai M, Hamazaki Y,**
1376 **Fukuda K, Minato N, and Sano M.** Obesity accelerates T cell senescence in
1377 murine visceral adipose tissue. *J Clin Invest* 126: 4626-4639, 2016.

- 1378 188. **Sindhu S, Thomas R, Kochumon S, Wilson A, Abu-Farha M, Bennakhi A, Al-**
1379 **Mulla F, and Ahmad R.** Increased Adipose Tissue Expression of Interferon
1380 Regulatory Factor (IRF)-5 in Obesity: Association with Metabolic Inflammation.
1381 *Cells* 8: 2019.
- 1382 189. **Singh GM, Danaei G, Farzadfar F, Stevens GA, Woodward M, Wormser D,**
1383 **Kaptoge S, Whitlock G, Qiao Q, Lewington S, Di Angelantonio E, Vander**
1384 **Hoorn S, Lawes CM, Ali MK, Mozaffarian D, Ezzati M, Global Burden of**
1385 **Metabolic Risk Factors of Chronic Diseases Collaborating G, Asia-Pacific**
1386 **Cohort Studies C, Diabetes Epidemiology: Collaborative analysis of**
1387 **Diagnostic criteria in E, Emerging Risk Factor C, and Prospective Studies C.**
1388 The age-specific quantitative effects of metabolic risk factors on cardiovascular
1389 diseases and diabetes: a pooled analysis. *PLoS One* 8: e65174, 2013.
- 1390 190. **Snodgrass RG, Boss M, Zezina E, Weigert A, Dehne N, Fleming I, Brune B, and**
1391 **Namgaladze D.** Hypoxia Potentiates Palmitate-induced Pro-inflammatory
1392 Activation of Primary Human Macrophages. *J Biol Chem* 291: 413-424, 2016.
- 1393 191. **Spite M, Hellmann J, Tang Y, Mathis SP, Kosuri M, Bhatnagar A, Jala VR, and**
1394 **Haribabu B.** Deficiency of the leukotriene B4 receptor, BLT-1, protects against
1395 systemic insulin resistance in diet-induced obesity. *J Immunol* 187: 1942-1949,
1396 2011.
- 1397 192. **Sramkova V, Rossmeislova L, Krauzova E, Kracmerova J, Koc M, Langin D,**
1398 **Stich V, and Siklova M.** Comparison of Early (2 Days) and Later (28 Days)
1399 Response of Adipose Tissue to Very Low-Calorie Diet in Obese Women. *J Clin*
1400 *Endocrinol Metab* 101: 5021-5029, 2016.
- 1401 193. **Stefanovic-Racic M, Yang X, Turner MS, Mantell BS, Stolz DB, Sumpter TL,**
1402 **Sipula IJ, Dedousis N, Scott DK, Morel PA, Thomson AW, and O'Doherty RM.**
1403 Dendritic cells promote macrophage infiltration and comprise a substantial
1404 proportion of obesity-associated increases in CD11c+ cells in adipose tissue and
1405 liver. *Diabetes* 61: 2330-2339, 2012.
- 1406 194. **Stienstra R, Joosten LA, Koenen T, van Tits B, van Diepen JA, van den Berg SA,**
1407 **Rensen PC, Voshol PJ, Fantuzzi G, Hijmans A, Kersten S, Muller M, van den**
1408 **Berg WB, van Rooijen N, Wabitsch M, Kullberg BJ, van der Meer JW,**
1409 **Kanneganti T, Tack CJ, and Netea MG.** The inflammasome-mediated caspase-1
1410 activation controls adipocyte differentiation and insulin sensitivity. *Cell Metab* 12:
1411 593-605, 2010.
- 1412 195. **Sulston RJ, Learman BS, Zhang B, Scheller EL, Parlee SD, Simon BR, Mori H,**
1413 **Bree AJ, Wallace RJ, Krishnan V, MacDougald OA, and Cawthorn WP.**
1414 Increased Circulating Adiponectin in Response to Thiazolidinediones:

- 1415 Investigating the Role of Bone Marrow Adipose Tissue. *Front Endocrinol*
1416 (*Lausanne*) 7: 128, 2016.
- 1417 196. **Takahashi M, Takahashi Y, Takahashi K, Zolotaryov FN, Hong KS, Kitazawa**
1418 **R, Iida K, Okimura Y, Kaji H, Kitazawa S, Kasuga M, and Chihara K.**
1419 Chemerin enhances insulin signaling and potentiates insulin-stimulated glucose
1420 uptake in 3T3-L1 adipocytes. *FEBS Lett* 582: 573-578, 2008.
- 1421 197. **Talukdar S, Oh DY, Bandyopadhyay G, Li D, Xu J, McNelis J, Lu M, Li P, Yan**
1422 **Q, Zhu Y, Ofrecio J, Lin M, Brenner MB, and Olefsky JM.** Neutrophils mediate
1423 insulin resistance in mice fed a high-fat diet through secreted elastase. *Nat Med*
1424 18: 1407-1412, 2012.
- 1425 198. **Tam CS, Covington JD, Ravussin E, Redman LM, and Pennington CT.** Little
1426 evidence of systemic and adipose tissue inflammation in overweight
1427 individuals(dagger). *Front Genet* 3: 58, 2012.
- 1428 199. **Tam CS, Viardot A, Clement K, Tordjman J, Tonks K, Greenfield JR,**
1429 **Campbell LV, Samocha-Bonet D, and Heilbronn LK.** Short-term overfeeding
1430 may induce peripheral insulin resistance without altering subcutaneous adipose
1431 tissue macrophages in humans. *Diabetes* 59: 2164-2170, 2010.
- 1432 200. **Tamura Y, Sugimoto M, Murayama T, Ueda Y, Kanamori H, Ono K, Ariyasu H,**
1433 **Akamizu T, Kita T, Yokode M, and Arai H.** Inhibition of CCR2 ameliorates
1434 insulin resistance and hepatic steatosis in db/db mice. *Arterioscler Thromb Vasc*
1435 *Biol* 28: 2195-2201, 2008.
- 1436 201. **Tardelli M, Zeyda K, Moreno-Viedma V, Wanko B, Grun NG, Staffler G,**
1437 **Zeyda M, and Stulnig TM.** Osteopontin is a key player for local adipose tissue
1438 macrophage proliferation in obesity. *Mol Metab* 5: 1131-1137, 2016.
- 1439 202. **Tencerova M, Figeac F, Ditzel N, Taipaleenmaki H, Nielsen TK, and Kassem**
1440 **M.** High-Fat Diet-Induced Obesity Promotes Expansion of Bone Marrow
1441 Adipose Tissue and Impairs Skeletal Stem Cell Functions in Mice. *J Bone Miner*
1442 *Res* 33: 1154-1165, 2018.
- 1443 203. **Ter Horst R, van den Munckhof ICL, Schraa K, Aguirre-Gamboa R, Jaeger M,**
1444 **Smeekens SP, Brand T, Lemmers H, Dijkstra H, Galesloot TE, de Graaf J,**
1445 **Xavier RJ, Li Y, Joosten LAB, Rutten JHW, Netea MG, and Riksen NP.** Sex-
1446 Specific Regulation of Inflammation and Metabolic Syndrome in Obesity.
1447 *Arteriosclerosis, thrombosis, and vascular biology* 40: 1787-1800, 2020.
- 1448 204. **Thomalla M, Schmid A, Neumann E, Pfefferle PI, Muller-Ladner U, Schaffler**
1449 **A, and Karrasch T.** Evidence of an anti-inflammatory toll-like receptor 9 (TLR 9)
1450 pathway in adipocytes. *J Endocrinol* 240: 325-343, 2019.

- 1451 205. **Torrissi JS, Hespe GE, Cuzzzone DA, Savetsky IL, Nitti MD, Gardenier JC,**
1452 **Garcia Nores GD, Jowhar D, Kataru RP, and Mehrara BJ.** Inhibition of
1453 Inflammation and iNOS Improves Lymphatic Function in Obesity. *Sci Rep* 6:
1454 19817, 2016.
- 1455 206. **Travers RL, Motta AC, Betts JA, Bouloumie A, and Thompson D.** The impact of
1456 adiposity on adipose tissue-resident lymphocyte activation in humans. *Int J Obes*
1457 *(Lond)* 39: 762-769, 2015.
- 1458 207. **Trayhurn P.** Hypoxia and adipose tissue function and dysfunction in obesity.
1459 *Physiol Rev* 93: 1-21, 2013.
- 1460 208. **Uchihashi K, Aoki S, Shigematsu M, Kamochi N, Sonoda E, Soejima H,**
1461 **Fukudome K, Sugihara H, Hotokebuchi T, and Toda S.** Organotypic culture of
1462 human bone marrow adipose tissue. *Pathol Int* 60: 259-267, 2010.
- 1463 209. **Uysal KT, Wiesbrock SM, Marino MW, and Hotamisligil GS.** Protection from
1464 obesity-induced insulin resistance in mice lacking TNF-alpha function. *Nature*
1465 389: 610-614, 1997.
- 1466 210. **van der Kolk BW, Kalafati M, Adriaens M, van Greevenbroek MMJ,**
1467 **Vogelzangs N, Saris WHM, Astrup A, Valsesia A, Langin D, van der Kallen**
1468 **CJH, Eussen S, Schalkwijk CG, Stehouwer CDA, Goossens GH, Arts ICW,**
1469 **Jocken JWE, Evelo CT, and Blaak EE.** Subcutaneous Adipose Tissue and
1470 Systemic Inflammation Are Associated With Peripheral but Not Hepatic Insulin
1471 Resistance in Humans. *Diabetes* 68: 2247-2258, 2019.
- 1472 211. **van Gils JM, Derby MC, Fernandes LR, Ramkhelawon B, Ray TD, Rayner KJ,**
1473 **Parathath S, Distel E, Feig JL, Alvarez-Leite JI, Rayner AJ, McDonald TO,**
1474 **O'Brien KD, Stuart LM, Fisher EA, Lacy-Hulbert A, and Moore KJ.** The
1475 neuroimmune guidance cue netrin-1 promotes atherosclerosis by inhibiting the
1476 emigration of macrophages from plaques. *Nat Immunol* 13: 136-143, 2012.
- 1477 212. **Vandanmagsar B, Youm Y-H, Ravussin A, Galgani JE, Stadler K, Mynatt RL,**
1478 **Ravussin E, Stephens JM, and Dixit VD.** The NLRP3 inflammasome instigates
1479 obesity-induced inflammation and insulin resistance. *Nature Medicine* 17: 179-188,
1480 2011.
- 1481 213. **Ventre J, Doebber T, Wu M, MacNaul K, Stevens K, Pasparakis M, Kollias G,**
1482 **and Moller DE.** Targeted disruption of the tumor necrosis factor-alpha gene:
1483 metabolic consequences in obese and nonobese mice. *Diabetes* 46: 1526-1531, 1997.
- 1484 214. **Viardot A, Lord RV, and Samaras K.** The effects of weight loss and gastric
1485 banding on the innate and adaptive immune system in type 2 diabetes and
1486 prediabetes. *J Clin Endocrinol Metab* 95: 2845-2850, 2010.


- 1487 215. **Viguerie N, Vidal H, Arner P, Holst C, Verdich C, Avizou S, Astrup A, Saris**
1488 **WH, Macdonald IA, Klimcakova E, Clement K, Martinez A, Hoffstedt J,**
1489 **Sorensen TI, Langin D, and Nutrient-Gene Interactions in Human Obesity--**
1490 **Implications for Dietary Guideline p.** Adipose tissue gene expression in obese
1491 subjects during low-fat and high-fat hypocaloric diets. *Diabetologia* 48: 123-131,
1492 2005.
- 1493 216. **Villaret A, Galitzky J, Decaunes P, Esteve D, Marques MA, Sengenès C,**
1494 **Chiotasso P, Tchkonja T, Lafontan M, Kirkland JL, and Bouloumie A.** Adipose
1495 tissue endothelial cells from obese human subjects: differences among depots in
1496 angiogenic, metabolic, and inflammatory gene expression and cellular
1497 senescence. *Diabetes* 59: 2755-2763, 2010.
- 1498 217. **Wallenius V, Wallenius K, Ahrén B, Rudling M, Carlsten H, Dickson SL,**
1499 **Ohlsson C, and Jansson JO.** Interleukin-6-deficient mice develop mature-onset
1500 obesity. *Nat Med* 8: 75-79, 2002.
- 1501 218. **Wang L, Wang Y, Zhang C, Li J, Meng Y, Dou M, Noguchi CT, and Di L.**
1502 **Inhibiting Glycogen Synthase Kinase 3 Reverses Obesity-Induced White Adipose**
1503 **Tissue Inflammation by Regulating Apoptosis Inhibitor of Macrophage/CD5L-**
1504 **Mediated Macrophage Migration.** *Arterioscler Thromb Vasc Biol* 38: 2103-2116,
1505 2018.
- 1506 219. **Wang M, Chen X, Zheng Z, Yu S, Zhou B, Liu Y, Liu D, Chen Y, and Qian X.**
1507 **Beneficial effect of ER stress preconditioning in protection against FFA-induced**
1508 **adipocyte inflammation via XBP1 in 3T3-L1 adipocytes.** *Mol Cell Biochem* 463: 45-
1509 55, 2020.
- 1510 220. **Wang Q, Dong Z, Liu X, Song X, Song Q, Shang Q, Jiang Y, Guo C, and Zhang**
1511 **L.** Programmed cell death-4 deficiency prevents diet-induced obesity, adipose
1512 tissue inflammation, and insulin resistance. *Diabetes* 62: 4132-4143, 2013.
- 1513 221. **Watanabe Y, Nagai Y, Honda H, Okamoto N, Yanagibashi T, Ogasawara M,**
1514 **Yamamoto S, Imamura R, Takasaki I, Hara H, Sasahara M, Arita M, Hida S,**
1515 **Taniguchi S, Suda T, and Takatsu K.** Bidirectional crosstalk between
1516 neutrophils and adipocytes promotes adipose tissue inflammation. *FASEB J* 33:
1517 11821-11835, 2019.
- 1518 222. **Weisberg SP, Hunter D, Huber R, Lemieux J, Slaymaker S, Vaddi K, Charo I,**
1519 **Leibel RL, and Ferrante AW, Jr.** CCR2 modulates inflammatory and metabolic
1520 effects of high-fat feeding. *J Clin Invest* 116: 115-124, 2006.
- 1521 223. **Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, and Ferrante**
1522 **AW, Jr.** Obesity is associated with macrophage accumulation in adipose tissue.
1523 *The Journal of clinical investigation* 112: 1796-1808, 2003.

- 1524 224. **Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, and Ferrante**
1525 **AW, Jr.** Obesity is associated with macrophage accumulation in adipose tissue. *J*
1526 *Clin Invest* 112: 1796-1808, 2003.
- 1527 225. **Weitman ES, Aschen SZ, Farias-Eisner G, Albano N, Cuzzone DA, Ghanta S,**
1528 **Zampell JC, Thorek D, and Mehrara BJ.** Obesity impairs lymphatic fluid
1529 transport and dendritic cell migration to lymph nodes. *PLoS One* 8: e70703, 2013.
- 1530 226. **Wentworth JM, Naselli G, Brown WA, Doyle L, Phipson B, Smyth GK,**
1531 **Wabitsch M, O'Brien PE, and Harrison LC.** Pro-inflammatory CD11c+CD206+
1532 adipose tissue macrophages are associated with insulin resistance in human
1533 obesity. *Diabetes* 59: 1648-1656, 2010.
- 1534 227. **Wernstedt Asterholm I, Tao C, Morley TS, Wang QA, Delgado-Lopez F, Wang**
1535 **ZV, and Scherer PE.** Adipocyte inflammation is essential for healthy adipose
1536 tissue expansion and remodeling. *Cell Metab* 20: 103-118, 2014.
- 1537 228. **Winer DA, Winer S, Shen L, Wadia PP, Yantha J, Paltser G, Tsui H, Wu P,**
1538 **Davidson MG, Alonso MN, Leong HX, Glassford A, Caimol M, Kenkel JA,**
1539 **Tedder TF, McLaughlin T, Miklos DB, Dosch HM, and Engleman EG.** B cells
1540 promote insulin resistance through modulation of T cells and production of
1541 pathogenic IgG antibodies. *Nat Med* 17: 610-617, 2011.
- 1542 229. **Winer S, Chan Y, Paltser G, Truong D, Tsui H, Bahrami J, Dorfman R, Wang Y,**
1543 **Zielenski J, Mastronardi F, Maezawa Y, Drucker DJ, Engleman E, Winer D,**
1544 **and Dosch HM.** Normalization of obesity-associated insulin resistance through
1545 immunotherapy. *Nat Med* 15: 921-929, 2009.
- 1546 230. **Wu D, Molofsky AB, Liang HE, Ricardo-Gonzalez RR, Jouihan HA, Bando JK,**
1547 **Chawla A, and Locksley RM.** Eosinophils sustain adipose alternatively activated
1548 macrophages associated with glucose homeostasis. *Science* 332: 243-247, 2011.
- 1549 231. **Xia B, Cai GH, Yang H, Wang SP, Mitchell GA, and Wu JW.** Adipose tissue
1550 deficiency of hormone-sensitive lipase causes fatty liver in mice. *PLoS Genet* 13:
1551 e1007110, 2017.
- 1552 232. **Xu H, Barnes GT, Yang Q, Tan G, Yang D, Chou CJ, Sole J, Nichols A, Ross JS,**
1553 **Tartaglia LA, and Chen H.** Chronic inflammation in fat plays a crucial role in the
1554 development of obesity-related insulin resistance. *J Clin Invest* 112: 1821-1830,
1555 2003.
- 1556 233. **Xu X, Grijalva A, Skowronski A, van Eijk M, Serlie MJ, and Ferrante AW, Jr.**
1557 **Obesity activates a program of lysosomal-dependent lipid metabolism in adipose**
1558 **tissue macrophages independently of classic activation.** *Cell Metab* 18: 816-830,
1559 2013.


- 1560 234. **Yamauchi T, Kamon J, Waki H, Terauchi Y, Kubota N, Hara K, Mori Y, Ide T,**
1561 **Murakami K, Tsuboyama-Kasaoka N, Ezaki O, Akanuma Y, Gavrilova O,**
1562 **Vinson C, Reitman ML, Kagechika H, Shudo K, Yoda M, Nakano Y, Tobe K,**
1563 **Nagai R, Kimura S, Tomita M, Froguel P, and Kadowaki T.** The fat-derived
1564 hormone adiponectin reverses insulin resistance associated with both
1565 lipodystrophy and obesity. *Nat Med* 7: 941-946, 2001.
- 1566 235. **Ye J, Gao Z, Yin J, and He Q.** Hypoxia is a potential risk factor for chronic
1567 inflammation and adiponectin reduction in adipose tissue of ob/ob and dietary
1568 obese mice. *Am J Physiol Endocrinol Metab* 293: E1118-1128, 2007.
- 1569 236. **Yin Z, Deng T, Peterson LE, Yu R, Lin J, Hamilton DJ, Reardon PR, Sherman V,**
1570 **Winnier GE, Zhan M, Lyon CJ, Wong STC, and Hsueh WA.** Transcriptome
1571 analysis of human adipocytes implicates the NOD-like receptor pathway in
1572 obesity-induced adipose inflammation. *Molecular and Cellular Endocrinology* 394:
1573 80-87, 2014.
- 1574 237. **Zamarron BF, Mergian TA, Cho KW, Martinez-Santibanez G, Luan D, Singer**
1575 **K, DelProposto JL, Geletka LM, Muir LA, and Lumeng CN.** Macrophage
1576 Proliferation Sustains Adipose Tissue Inflammation in Formerly Obese Mice.
1577 *Diabetes* 66: 392-406, 2017.
- 1578 238. **Zeyda M, Farmer D, Todoric J, Aszmann O, Speiser M, Gyorfi G, Zlabinger GJ,**
1579 **and Stulnig TM.** Human adipose tissue macrophages are of an anti-
1580 inflammatory phenotype but capable of excessive pro-inflammatory mediator
1581 production. *Int J Obes (Lond)* 31: 1420-1428, 2007.
- 1582 239. **Zeyda M, Huber J, Prager G, and Stulnig TM.** Inflammation correlates with
1583 markers of T-cell subsets including regulatory T cells in adipose tissue from
1584 obese patients. *Obesity (Silver Spring)* 19: 743-748, 2011.
- 1585 240. **Zhang T, Fang Z, Linghu K-G, Liu J, Gan L, and Lin L.** Small molecule-driven
1586 SIRT3-autophagy-mediated NLRP3 inflammasome inhibition ameliorates
1587 inflammatory crosstalk between macrophages and adipocytes. *British Journal of*
1588 *Pharmacology* 177: 4645-4665, 2020.
- 1589 241. **Zhou L, Park SY, Xu L, Xia X, Ye J, Su L, Jeong KH, Hur JH, Oh H, Tamori Y,**
1590 **Zingaretti CM, Cinti S, Argente J, Yu M, Wu L, Ju S, Guan F, Yang H, Choi CS,**
1591 **Savage DB, and Li P.** Insulin resistance and white adipose tissue inflammation
1592 are uncoupled in energetically challenged Fsp27-deficient mice. *Nat Commun* 6:
1593 5949, 2015.
- 1594

1595

Lean Adipose Tissue - Anti-inflammatory



Obese Adipose Tissue - Pro-inflammatory



Macrophage

IL-10, Arginase,
Catecholamines




Eosinophil

IL-4, IL-13



Treg cell

IL-10, TGF beta



Th2 cell

IL-4, IL-5, IL-6,
IL-10, IL-13



Macrophage

TNF alpha
IL-1b, IL-6, IL-12



Neutrophil

Elastases,
Cathepsin G
Proteinase-3



Mast cell

TNF alpha
IL-6, IFN gamma



T cell

TNF alpha
IL-2, IFN gamma



NKT cell

IFN gamma



B cell

IgG antibody

Healthy adipocytes



Adipokines

Anti-inflammatory
Adiponectin
Chemerin (low expression)

Inflammation

Dysfunctional adipocytes



Obesity / High fat diet

Saturated Free Fatty Acid

Hypoxia

Cellular
Senescence

Adipokines

Inflammatory
TNF alpha
MCP-1
etc.
Anti-inflammatory
Adiponectin

Anti-inflammatory
Cytokine: IL-19, IL-10

Inflammation

Immune cells
infiltration

Inflammatory
cytokines

Insulin
signaling

Insulin
resistance

