

Theorem 12.1 Rules of Multiplication.

Let a, b , and c belong to a ring R . Then

1. $a0 = 0a = 0$.
2. $a(-b) = (-a)b = -(ab)$.
3. $(-a)(-b) = ab$.
4. $a(b - c) = ab - ac$ and $(b - c)a = ba - ca$.

Furthermore, if R has a unity element 1, then

5. $(-1)a = -a$.
6. $(-1)(-1) = 1$.

Theorem 12.2 Uniqueness of the Unity and Inverses.

If a ring has a unity, it is unique. If a ring element has a multiplicative inverse, it is unique.

Theorem 12.3 Subring Test.

A nonempty subset S of a ring R is a subring if S is closed under subtraction and multiplication – that is, if $a - b$ and ab are in S whenever a and b are in S .

Theorem 13.1 Cancellation.

Let a, b , and c belong to an integral domain. If $a \neq 0$ and $ab = ac$, then $b = c$.

Theorem 13.2 Finite Integral Domains are Fields.

A finite integral domain is a field.

Corollary 13.2.1 \mathbb{Z}_p Is a Field.

For every prime p , \mathbb{Z}_p , the ring of integers modulo p is a field.

Theorem 13.3 Characteristic of a Ring with Unity.

Let R be a ring with unity 1. If 1 has infinite order under addition, then the characteristic of R is 0. If 1 has order n under addition, then the characteristic of R is n .

Theorem 13.4 Characteristic of an Integral Domain.

The characteristic of an integral domain is 0 or prime.

Theorem 14.1 Ideal Test.

A nonempty subset A of a ring R is an ideal of R if

1. $a - b \in A$ whenever $a, b \in A$.
2. ra and ar are in A whenever $a \in A$ and $r \in R$.

Theorem 14.2 Existence of Factor Rings.

Let R be a ring and let A be a subring of R . The set of cosets $\{r + A \mid r \in R\}$ is a ring under the operations $(s + A) + (t + A) = s + t + A$ and $(s + A)(t + A) = st + A$ if and only if A is an ideal of R .

Theorem 14.3 R/A Is an Integral Domain If and Only If A Is Prime.

Let R be a commutative ring with unity and let A be an ideal of R . Then R/A is an integral domain if and only if A is prime.

Theorem 14.4 R/A Is a Field If and Only If A Is Maximal.

Let R be a commutative ring with unity and let A be an ideal of R . Then R/A is a field if and only if A is maximal.

Theorem 15.1 Properties of Ring Homomorphisms.

Let ϕ be a ring homomorphism from a ring R to a ring S . Let A be a subring of R and let B be an ideal of S .

1. For any $r \in R$ and any positive integer n , $\phi(nr) = n\phi(r)$ and $\phi(r^n) = (\phi(r))^n$.

2. $\phi(A) = \{\phi(a) \mid a \in A\}$ is a subring of S .
3. If A is an ideal and ϕ is onto S , then $\phi(A)$ is an ideal.
4. $\phi^{-1}(B) = \{r \in R \mid \phi(r) \in B\}$ is an ideal of R .
5. If R is commutative, then $\phi(R)$ is commutative.
6. If R has a unity 1 , $S \neq \{0\}$, and ϕ is onto, then $\phi(1)$ is the unity of S .
7. ϕ is an isomorphism if and only if ϕ is onto and $\ker \phi = \{r \in R \mid \phi(r) = 0\} = \{0\}$.

Theorem 15.2 Kernels Are Ideals.

Let ϕ be a ring homomorphism from a ring R to a ring S . Then $\ker \phi = \{r \in R \mid \phi(r) = 0\}$ is an ideal of R .

Theorem 15.3 First Isomorphism Theorem for Rings.

Let ϕ be a ring homomorphism from R to S . Then the mapping from $R/\ker \phi$ to $\phi(R)$, given by $r + \ker \phi \rightarrow \phi(r)$, is an isomorphism. In symbols, $R/\ker \phi \approx \phi(R)$. This theorem is often referred to as the *Fundamental Theorem of Ring Homomorphisms*.

Theorem 15.4 Ideals Are Kernels.

Every ideal of a ring R is the kernel of a ring homomorphism of R . In particular, an ideal A is the kernel of the mapping $r \rightarrow r + A$ from R to R/A . This mapping is known as the *natural homomorphism* from R to R/A .

Theorem 15.5 Homomorphism from \mathbb{Z} to a Ring with Unity.

Let R be a ring with unity 1 . The mapping $\phi : \mathbb{Z} \rightarrow R$ given by $n \rightarrow n \cdot 1$ is a ring homomorphism.

Corollary 15.5.1 A Ring with Unity Contains \mathbb{Z}_n or \mathbb{Z} .

If R is a ring with unity and the characteristic of R is $n > 0$, then R contains a subring isomorphic to \mathbb{Z}_n . If the characteristic of R is 0 , then R contains a subring isomorphic to \mathbb{Z} .

Corollary 15.5.2 \mathbb{Z}_m Is a Homomorphic Image of \mathbb{Z} .

For any positive integer m , the mapping of $\phi : \mathbb{Z} \rightarrow \mathbb{Z}_m$ given by $x \rightarrow x \bmod m$ is a ring homomorphism.

Corollary 15.5.3 A Field Contains \mathbb{Z}_p or \mathbb{Q} .

If \mathbb{F} is a field of characteristic p , then \mathbb{F} contains a subfield isomorphic to \mathbb{Z}_p . If \mathbb{F} is a field of characteristic 0 , then \mathbb{F} contains a subfield isomorphic to the rational numbers.

Theorem 15.6 Field of Quotients.

Let D be an integral domain. Then there exists a field \mathbb{F} (called the field of quotients in D) that contains a subring isomorphic to D .

Theorem 16.1 D an Integral Domain Implies $D[x]$ an Integral Domain.

If D is an integral domain, then $D[x]$ is an integral domain.

Theorem 16.2 Division Algorithm for $\mathbb{F}[x]$.

Let \mathbb{F} be a field and let $f(x), g(x) \in \mathbb{F}[x]$ with $g(x) \neq 0$. Then there exist unique polynomials $q(x)$ and $r(x)$ in $\mathbb{F}[x]$ such that $f(x) = g(x)q(x) + r(x)$ and either $r(x) = 0$ or $\deg r(x) < \deg g(x)$.

Corollary 16.2.1 Remainder Theorem.

Let \mathbb{F} be a field, $a \in \mathbb{F}$, and $f(x) \in \mathbb{F}[x]$. Then $f(a)$ is the remainder in the division of $f(x)$ by $x - a$.

Corollary 16.2.2 Factor Theorem.

Let \mathbb{F} be a field, $a \in \mathbb{F}$, and $f(x) \in \mathbb{F}[x]$. Then a is a zero of $f(x)$ if and only if $x - a$ is a factor of $f(x)$.

Corollary 16.2.3 Polynomials of Degree n Have at Most n Zeros.

A polynomial of degree n over a field has at most n zeros, counting multiplicity.

Theorem 16.3 $\mathbb{F}[x]$ Is a PID.

Let \mathbb{F} be a field. Then $\mathbb{F}[x]$ is a principal ideal domain.

Theorem 16.4 Criterion for $I = \langle g(x) \rangle$.

Let \mathbb{F} be a field, I a nonzero ideal in $\mathbb{F}[x]$, and $g(x)$ an element of $\mathbb{F}[x]$. Then, $I = \langle g(x) \rangle$ if and only if $g(x)$ is a nonzero polynomial of minimum degree in I .

Theorem 17.1 Reducibility Test for Degrees 2 and 3.

Let \mathbb{F} be a field. If $f(x) \in \mathbb{F}[x]$ and $\deg f(x)$ is 2 or 3, then $f(x)$ is reducible over \mathbb{F} if and only if $f(x)$ has a zero in \mathbb{F} .

Lemma 17.2 Gauss's Lemma.

The product of two primitive polynomials is primitive.

Theorem 17.3 Reducibility over \mathbb{Q} Implies Reducibility over \mathbb{Z} .

Let $f(x) \in \mathbb{Z}[x]$. If $f(x)$ is reducible over \mathbb{Q} , then it is reducible over \mathbb{Z} .

Theorem 17.4 Mod p Irreducibility Test.

Let p be a prime and suppose that $f(x) \in \mathbb{Z}[x]$ with $\deg f(x) \geq 1$. Let $\bar{f}(x)$ be the polynomial in $\mathbb{Z}_p[x]$ obtained from $f(x)$ by reducing all the coefficients of $f(x)$ modulo p . If $\bar{f}(x)$ is irreducible over \mathbb{Z}_p and $\deg \bar{f}(x) = \deg f(x)$, then $f(x)$ is irreducible over \mathbb{Q} .

Theorem 17.5 Eisenstein's Criterion.

Let

$$f(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_0 \in \mathbb{Z}[x]$$

If there is a prime p such that $p \nmid a_n, p \mid a_{n-1}, \dots, p \mid a_0$ and $p^2 \nmid a_0$, then $f(x)$ is irreducible over \mathbb{Q} .

Corollary 17.5.1 Irreducibility of p th Cyclotomic Polynomial.

For any prime p , the p th cyclotomic polynomial

$$\Phi_p(x) = \frac{x^p - 1}{x - 1} = x^{p-1} + x^{p-2} + \cdots + x + 1$$

is irreducible over \mathbb{Q} .

Theorem 17.6 $\langle p(x) \rangle$ Is Maximal If and Only If $p(x)$ Is Irreducible.

Let \mathbb{F} be a field and let $p(x) \in \mathbb{F}[x]$. Then $\langle p(x) \rangle$ is a maximal ideal in $\mathbb{F}[x]$ if and only if $p(x)$ is irreducible over \mathbb{F} .

Corollary 17.6.1 $\mathbb{F}[x]/\langle p(x) \rangle$ Is a Field.

Let \mathbb{F} be a field and $p(x)$ be an irreducible polynomial over \mathbb{F} . Then $\mathbb{F}[x]/\langle p(x) \rangle$ is a field.

Corollary 17.6.2 $p(x) \mid a(x)b(x)$ Implies $p(x) \mid a(x)$ or $p(x) \mid b(x)$.

Let \mathbb{F} be a field and let $p(x), a(x), b(x) \in \mathbb{F}[x]$. If $p(x)$ is irreducible over \mathbb{F} and $p(x) \mid a(x)b(x)$, then $p(x) \mid a(x)$ or $p(x) \mid b(x)$.

Theorem 17.7 Unique Factorization in $\mathbb{Z}[x]$.

Every polynomial in $\mathbb{Z}[x]$ that is not the zero polynomial or a unit in $\mathbb{Z}[x]$ can be written in the form $b_1 b_2 \cdots b_s p_1(x) p_2(x) \cdots p_m(x)$, where the b_i 's are irreducible polynomials of degree 0 and the $p_i(x)$'s are irreducible polynomials of positive degree. Furthermore, if

$$b_1 b_2 \cdots b_s p_1(x) p_2(x) \cdots p_m(x) = c_1 c_2 \cdots c_t q_1(x) q_2(x) \cdots q_n(x)$$

where the b_i 's and the c_i 's are irreducible polynomials of degree 0 and the $p_i(x)$'s and $q_i(x)$'s are irreducible polynomials of positive degree, then $s = t, m = n$, and, after renumbering the c 's and $q(x)$'s, we have $b_i = \pm c_i$, for $i = 1, \dots, s$, and $p_i(x) = \pm q_i(x)$, for $i = 1, \dots, m$.