

The Primitive Element Thm. (Lecture 15)

Def. Let $L:K$ be a field extension, $\varphi: K \rightarrow L$. Then $L:K$ is a simple extension if $\exists \theta \in L$ s.t. $L = \varphi(K)(\theta)$.

Exm 1) $\mathbb{Q}(\sqrt{2}, \sqrt{3}) = \mathbb{Q}(\sqrt{2} + \sqrt{3})$. Indeed,

$$\frac{1}{\sqrt{2} + \sqrt{3}} = \sqrt{3} - \sqrt{2} \in M \Rightarrow \sqrt{2}, \sqrt{3} \in M.$$

2) Let α, β be alg. over $K \Rightarrow \exists \theta: K(\alpha, \beta) = K(\theta)$
we try to find θ in the following form

$\theta = \alpha + c\beta$, $c \in K$. Clearly, it is enough to prove $\beta \in K(\theta)$. Consider, μ_α^K & μ_β^K . Then

$\mu_\beta^K(\beta) = \mu_\alpha^K(\theta - c\beta) = 0 \Rightarrow \gcd(\mu_\beta^K(x), \mu_\alpha^K(\theta - cx)) \in K(\theta)[x]$. If $g(x) = x - \beta$, then $\beta \in K(\theta)$ and we are done. The 1st polynomial has roots β_1, \dots, β_n , the 2nd: $\alpha_1, \dots, \alpha_m$ and thus we want to find c s.t.

$$\alpha_1 + c\beta_1 - c\beta_i = \theta - c\beta_i = \alpha_j \Leftrightarrow i=j=1$$

Thus, we want $c \neq \frac{\alpha_j - \alpha_1}{\beta_1 - \beta_i}$, $\forall i, j$ (excluding $i=j=1$)

Clearly, if $|K| = \infty$, then such c exists.

If $|K| < \infty$, then K^* is a cyclic group and therefore if $[L:K] < \infty$, then we can put $L = K(g)$, where $K^* = \{1, g, g^2, \dots\}$.

Unfortunately, it can be that $g(x) = (x - \beta)^e$, where $e > 1$ due to the existence of inseparable polynomials in positive characteristic.

Thm (the primitive element theorem)

Let $L:K$ be a finite, separable extension, $K \subseteq L$. Then $L:K$ is a simple extension.

Pf. We assume that $L \subseteq \bar{K}$. If $|K| < \infty$, then $|L| = |K|^{[L:K]} \Rightarrow L^* = \langle g \rangle$ and $L = K(g)$.
Now if $|K| = \infty$, then we use induction on $[L:K]$.
Let $\alpha \in L$ be any element of largest degree over K . If $L = K(\alpha)$, then we are done. Otherwise $\exists \beta \in L \setminus K(\alpha)$.

Suppose that $[K(\alpha, \beta):K] < [L:K] \Rightarrow$ by induction $K(\alpha, \beta) = K(\gamma)$ for some $\gamma \in L$. Then $[K(\gamma):K] = [K(\alpha, \beta):K] > [K(\alpha):K]$ (recall that $\beta \in L \setminus K(\alpha)$) and this contradicts our maximal assumption. Thus $[K(\alpha, \beta):K] = [L:K]$ and hence $L = K(\alpha, \beta)$.

We know that $L:K$ is separable \Rightarrow

we know that \exists ^{see Lecture 14} $[L:K]$ distinct K -hom.
 $\varphi_i : K(\alpha, \beta) \rightarrow \overline{K}$. Put

$$f = \prod_{1 \leq i < j \leq n} ((\varphi_i(\alpha) - \varphi_j(\alpha)) + (\varphi_i(\beta) - \varphi_j(\beta))t)$$

If $f \equiv 0$, then $\varphi_i(\alpha) = \varphi_j(\alpha)$ & $\varphi_i(\beta) = \varphi_j(\beta)$
 $\Rightarrow \varphi_i = \varphi_j$ and this is a contradiction.

We have $|K| = \infty \Rightarrow \exists \delta \in K$ s.t. $f(\delta) \neq 0$.

Put $\theta = \alpha + \delta\beta$. Then $\varphi_i(\theta) \neq \varphi_j(\theta)$, $i \neq j$.

Indeed, otherwise

$$f(\delta) = \prod_{1 \leq i < j \leq n} (\varphi_i(\alpha + \delta\beta) - \varphi_j(\alpha + \delta\beta)) = 0$$

and this is a contradiction. Thus φ_i must
restrict to distinct K -hom. from $K(\theta) \rightarrow \overline{K}$.

As above (see the same Cor. 1 of Lecture
14) we derive that $[K(\theta):K] \geq [L:K]$.

Thus $K(\theta) = L$ as required. \blacksquare

Ex. (Artin) $L:K$ is a finite extension. Then
 $L:K$ is a simple extension \Leftrightarrow the number of
intermediate fields is finite.

Cor. Let $L:K$ is an algebraic, separable
ext. s.t. $\forall \alpha \in L$ one has $\deg m_\alpha^K \leq n$. Then
 $[L:K] \leq n$.

Ex. We want to construct a finite alg (inseparable) extension that is not simple.

First of all, let $K = \mathbb{F}_p(t)$ be our base field. Take any transcendental over K elements. Our first field $M = K(s)$ and L be a splitting field extension for the pol. $(x^p - t)(x^p - s) \in M[x]$. So, we have

$$K \xrightarrow{\infty} M (= K(s)) \xrightarrow{?} L$$

We need to compute $[L:M]$. First consider the pol. $x^p - t$ & M_1 be its splitting field.

L.1. $[M_1:M] = p$

Pf. Let $h(x) = x^p - t \in M[x]$ $\Rightarrow t = \alpha^p \Rightarrow$ so, h is inseparable

$(x - \alpha)^p = x^p - \alpha^p = x^p - t$, so α has multiplicity p .

Let us check that h is irr. over M .

If $h = f \cdot g \Rightarrow f = (x - \alpha)^e, g = (x - \alpha)^{p-e}$ ($M[x]$ is UFD) \Rightarrow since $\gcd(e, p-e) = 1$ we can find $a, b \in \mathbb{Z}$ s.t. $ae + b(p-e) = 1 \Rightarrow x - \alpha = f^a g^b \Rightarrow x - \alpha \in M[x] \Rightarrow \alpha \in M$. It follows that $\exists c, d \in \mathbb{F}_p[s, t]$ with $\alpha = c/d$. Therefore

$t = \frac{c}{d} = \left(\frac{c}{d}\right)^p \Rightarrow c^p = t d^p$. This is a contradiction (compare the degrees of LHS/RHS). Thus h is irreducible over $M[x]$ and hence $[M_1 : M] = \deg h = p$. \square

L.2. $[L : M] = p^2$. In particular, this is an algebraic extension.

Pf. $M \subset M_1 \stackrel{?}{=} F$ \leftarrow the splitting field for $x^p - s$. Let us prove that $[L : M_1] = p$. Indeed, as in the proof of L.1

let $\beta \in L$ s.t. $x^p - s = (x - \beta)^p := H(x)$.

If $H(x)$ is reducible over M_1 , then as above $\beta \in M_1$ and $\exists c, d \in \mathbb{F}_p(\alpha)[s]$ s.t. (recall that $M_1 = M(\alpha)$)

$c^p = s d^p$. Comparing deg we obtain a contradiction again and therefore $H(x)$ is irr. over $M_1 \Rightarrow [F : M_1] = p$ (and $F = M_1(\beta)$)

Clearly $F \supseteq L$ and $L \not\subseteq M_1 \Rightarrow [L : M] = p^2$ (use the tower law) \square

L.3. $L : M$ is not a simple extension.

Pf. Suppose that $\exists \theta \in L$ s.t. $L = M(\theta)$.

Observe that $\forall \theta \in L$ one has $\theta^p \in M$.

Indeed, $\alpha, \beta \in L$ and by construction $\alpha^p = t, \beta^p = s$. Also, $(x^p - t)(x^p - s) = (x - \alpha)^p (x - \beta)^p$

$$\Rightarrow L = M(\alpha, \beta) \Rightarrow \theta = \frac{q(\alpha, \beta)}{r(\alpha, \beta)}, \quad q, r \in M[z_1, z_2]$$

$$\text{Then } \theta^p = \frac{q^p(\alpha, \beta)}{r^p(\alpha, \beta)} = \frac{q(\alpha^p, \beta^p)}{r(\alpha^p, \beta^p)} = \frac{q(t, s)}{r(t, s)} \in M.$$

Thus, we know that $\theta^p = \delta \in M \Rightarrow$
 $\mu_{\theta}^M \mid t^p - \delta \Rightarrow 1 \leq [M(\theta) : M] \leq p.$

On the other hand, $M - M(\theta) - L$

$$\Rightarrow [M(\theta) : M] = \underbrace{p}_p.$$

But $[L : M] = p^2$ and thus $L : M$ cannot be simple. This concludes the proof of L.3 and therefore the construction is complete. $\blacksquare \blacksquare$