

1 extra

Proposition 1. Suppose that K and L are fields and that $\varphi : K \rightarrow L$ is a homomorphism. With t and y denoting indeterminates, extend the homomorphism φ to the mapping $\psi : K[t] \rightarrow L[y]$ by defining

$$\psi(a_0 + a_1t + \cdots + a_nt^n) = \varphi(a_0) + \varphi(a_1)y + \cdots + \varphi(a_n)y^n.$$

Then $\psi : K[t] \rightarrow L[y]$ is an injective homomorphism. Also, when $\varphi : K \rightarrow L$ is surjective, then $\psi : K[t] \rightarrow L[y]$ is surjective and maps irreducible polynomials in $K[t]$ to irreducible polynomials in $L[y]$.

Proposition 2. Suppose $L : K$ is a field extension with $K \subseteq L$, and $\alpha \in L$. Then E_α is a ring homomorphism.

Proposition 3. Let $L : K$ be a field extension with $K \subseteq L$, and suppose that $\alpha \in L$ is algebraic over K . Then

$$I = \ker(E_\alpha) = \{f \in K[t] : f(\alpha) = 0\}$$

is a nonzero ideal of $K[t]$, and there is a unique monic polynomial $\mu_\alpha^K \in K[t]$ that generates I .

Theorem 1.1. Suppose that $L : K$ is a field extension, and that $\alpha \in L$ is algebraic over K . Let g be the minimal polynomial μ_α^K of α over K . Then g is irreducible over K , and $K[t]/(g)$ is a field.

Theorem 1.2. Let K be a field, and suppose that $f \in K[t]$ is irreducible. Then there exists a field extension $L : K$, with associated embedding $\varphi : K[t] \rightarrow L[y]$, having the property that L contains a root of $\varphi(f)$.

Proposition 4. Let $L : K$ be a field extension with $K \subseteq L$. Let $A \subseteq L$ and

$$\mathcal{C} = \{C \subseteq A : C \text{ is a finite set}\}.$$

Then $K(A) = \cup_{C \in \mathcal{C}} K(C)$. Further, when $[K(C) : K] < \infty$ for all $C \in \mathcal{C}$, then $K(A) : K$ is an algebraic extension.

Proposition 5. Let $L : K$ be a field extension with $K \subseteq L$, and suppose that $\alpha \in L$. Then

$$K[\alpha] = \{c_0 + c_1\alpha + \cdots + c_d\alpha^d : d \in \mathbb{Z}_{\geq 0}, c_0, \dots, c_d \in K\}$$

and

$$K(\alpha) = \{f/g : f, g \in K[\alpha], g \neq 0\}.$$

Proposition 6. Let $L : K$ be a field extension with $K \subseteq L$, and suppose that $\alpha \in L$. Then α is algebraic over K if and only if $[K(\alpha) : K] < \infty$.

Definition 1 (Characteristic). Let K be a field with additive identity 0_K and multiplicative identity 1_K . When $n \in \mathbb{N}$, we write $n \cdot 1_K$ to denote $1_K + \cdots + 1_K$ (as an n -fold sum). We define the characteristic of K , denoted by $\text{char}(K)$, to be the smallest positive integer m with the property that $m \cdot 1_K = 0_K$; if no such integer m exists, we define the characteristic of K to be 0.

Proposition 7. Let K be a field with $\text{char}(K) > 0$. Then $\text{char}(K)$ is equal to a prime number p , and then for all $x \in K$ one has $p \cdot x = 0$.

Theorem 1.3 (Localisation principle). Let R be an integral domain, and let I be a prime ideal of R . Define $\varphi : R[X] \rightarrow (R/I)[X]$ by putting

$$\varphi(a_0 + a_1X + \cdots + a_nX^n) = \bar{a}_0 + \bar{a}_1X + \cdots + \bar{a}_nX^n,$$

where $\bar{a}_j = a_j + I$. Then φ is a surjective homomorphism. Moreover, if $f \in R[X]$ is primitive with leading coefficient not in I , then f is irreducible in $R[X]$ whenever $\varphi(f)$ is irreducible in $(R/I)[X]$.

Note: Proposition 3.1 tells us that when $f \in K[t]$ and $\sigma \in \text{Gal}(L : K)$, the mapping σ permutes the roots of f that lie in L .

Theorem 1.4. Suppose that $L : K$ is an algebraic extension, and $\sigma : L \rightarrow L$ is a K -homomorphism. Then σ is an automorphism of L .

Theorem 1.5. *If $L : K$ is a finite extension, then $|\text{Gal}(L : K)| \leq [L : K]$.*

Corollary 1. *Suppose that $L : F$ and $L : F'$ are finite extensions with $F \subseteq L$ and $F' \subseteq L$, and further that $\psi : F \rightarrow F'$ is an isomorphism. Then there are at most $[L : F]$ ways to extend ψ to a homomorphism from L into L .*

Corollary 2. *Let $L : K$ be a finite extension with $K \subseteq L$. Suppose that $\alpha_1, \dots, \alpha_n \in L$ and put $L = K(\alpha_1, \dots, \alpha_n)$. Let $K_0 = K$, and for $1 \leq i \leq n$, let $K_i = K_{i-1}(\alpha_i)$. Then every automorphism $\tau \in \text{Gal}(L : K)$ corresponds to a sequence of homomorphisms $\sigma_1, \dots, \sigma_n$, having the property that $\sigma_0 : K \rightarrow L$ is the inclusion map, one has $\sigma_n = \tau$, and for $1 \leq i \leq n$, the map $\sigma_i : K_i \rightarrow L$ is a homomorphism extending $\sigma_{i-1} : K_{i-1} \rightarrow L$.*

2 Algebraic closures

Corollary 3. *When K is a field, the field \overline{K} is a maximal algebraic extension of K .*

Corollary 4. *Suppose that \overline{K} is an algebraic closure of K , and assume that $K \subseteq \overline{K}$. Take $\alpha \in \overline{K}$ and suppose that $\sigma : K \rightarrow \overline{K}$ is a homomorphism. Then the number of distinct roots of μ_α^K in \overline{K} is equal to the number of distinct roots of $\sigma(\mu_\alpha^K)$ in \overline{K} .*

Proposition 8. *Suppose that L and M are fields having the property that L is algebraically closed, and $\psi : L \rightarrow M$ is a homomorphism. Then $\psi(L)$ is algebraically closed.*

Proposition 9. *If $L : K$ is an algebraic extension, then \overline{L} is an algebraic closure of K , and hence $\overline{L} \cong \overline{K}$. If in addition $K \subseteq L \subseteq \overline{L}$, then we can take $\overline{K} = \overline{L}$.*

3 Splitting field extensions

Definition 2 (Splitting field, splitting field extension). Suppose that $L : K$ is a field extension relative to the embedding $\varphi : K \rightarrow L$, and $f \in K[t] \setminus K$.

- (i) We say that f splits over L if $\varphi(f) = \lambda(t - \alpha_1) \cdots (t - \alpha_n)$, for some $\lambda \in \varphi(K)$ and $\alpha_1, \dots, \alpha_n \in L$.
- (ii) Suppose that f splits over L , and let M be a field with $\varphi(K) \subseteq M \subseteq L$. We say that $M : K$ is a splitting field extension for f if M is the smallest subfield of L containing $\varphi(K)$ over which f splits.
- (iii) More generally, suppose that $S \subseteq K[t] \setminus K$ has the property that every $f \in S$ splits over L . Let M be a field with $\varphi(K) \subseteq M \subseteq L$. We say that $M : K$ is a splitting field extension for S if M is the smallest subfield of L containing $\varphi(K)$ over which every polynomial $f \in S$ splits.

Proposition 10. *Suppose that $L : K$ is a splitting field extension for the polynomial $f \in K[t] \setminus K$ with associated embedding $\varphi : K \rightarrow L$. Let $\alpha_1, \dots, \alpha_n \in L$ be the roots of $\varphi(f)$. Then $L = \varphi(K)(\alpha_1, \dots, \alpha_n)$.*

Proposition 11. *Suppose that $L : K$ is a splitting field extension for the polynomial $f \in K[t] \setminus K$. Then $[L : K] \leq (\deg f)!$*

Proposition 12. *Given $S \subseteq K[t] \setminus K$, there exists a splitting field extension $L : K$ for S , and $L : K$ is an algebraic extension. More explicitly, suppose that \overline{K} is an algebraic closure of K , and that $\overline{K} : K$ is an extension relative to the embedding $\varphi : \overline{K} \rightarrow K$. Let*

$$A = \{\alpha \in \overline{K} : \alpha \text{ is a root of } \varphi(f), \text{ for some } f \in S\}.$$

Put $K' = \varphi(K)$. Then $K'(A) : K$ is a splitting field extension for S .

Theorem 3.1. *Let $f \in K[t] \setminus K$, and suppose that $L : K$ and $M : K$ are splitting field extensions for f . Then $L \cong M$, and thus $[L : K] = [M : K]$.*

Theorem 3.2. *Suppose that $S \subseteq K[t] \setminus K$, and suppose that $L : K$ and $M : K$ are splitting field extensions for S . Then $L \cong M$ and $[L : K] = [M : K]$.*

4 Normal extensions and composita

Proposition 13. *Suppose that $L : M : K$ is a tower of field extensions and $L : K$ is a normal extension. Then $L : M$ is also a normal extension.*

Theorem 4.1. *Suppose that $M : L : K$ is a tower of field extensions having the property that $M : K$ is normal. Assume that $K \subseteq L \subseteq M$. Then the following are equivalent:*

- (i) *the field extension $L : K$ is normal;*
- (ii) *any K -homomorphism of L into M is an automorphism of L ;*
- (iii) *whenever $\sigma : M \rightarrow M$ is a K -automorphism, then $\sigma(L) \subseteq L$.*

Definition 3 (Compositum). Let K_1 and K_2 be fields contained in some field L . The compositum of K_1 and K_2 in L , denoted by K_1K_2 , is the smallest subfield of L containing both K_1 and K_2 .

Proposition 14. *Suppose that $E : K$ and $F : K$ are finite extensions having the property that K , E and F are contained in a field L . Then $EF : K$ is a finite extension.*

Theorem 4.2. *Let $E : K$ and $F : K$ be finite extensions having the property that K , E and F are contained in a field L .*

- (a) *When $E : K$ is normal, then $EF : F$ is normal.*
- (b) *When $E : K$ and $F : K$ are both normal, then $EF : K$ and $E \cap F : K$ are normal.*

5 Separability

Theorem 5.1. *Suppose that $L : M : K$ is a tower of algebraic extensions. Then $L : K$ is separable if and only if $L : M$ and $M : K$ are both separable.*

Theorem 5.2. *Suppose that $E : K$ and $F : K$ are finite extensions with $E \subseteq L$ and $F \subseteq L$, where L is a field.*

- (a) *When $E : K$ is separable, then so too is $EF : F$;*
- (b) *When $E : K$ and $F : K$ are both separable, then so too are $EF : K$ and $E \cap F : K$.*

6 Inseparable polynomials, differentiation, and the Frobenius map

7 The Primitive Element Theorem

8 Fixed fields and Galois extensions

Proposition 15. *Let K , M and L be fields with $K \subseteq L$ and $M \subseteq L$. Suppose that G and H are subgroups of $\text{Aut}(L)$. Then one has the following:*

- (a) *if $K \subseteq M$, then $\text{Gal}(L : K) \geq \text{Gal}(L : M)$;*
- (b) *if $G \leq H$, then $\text{Fix}_L(G) \supseteq \text{Fix}_L(H)$;*
- (c) *one has $K \subseteq \text{Fix}_L(\text{Gal}(L : K))$;*
- (d) *one has $G \leq \text{Gal}(L : \text{Fix}_L(G))$;*
- (e) *one has $\text{Gal}(L : K) = \text{Gal}(L : \text{Fix}_L(\text{Gal}(L : K)))$;*
- (f) *one has $\text{Fix}_L(G) = \text{Fix}_L(\text{Gal}(L : \text{Fix}_L(G)))$.*

Definition 25 (Galois extension). When $L : K$ is a field extension, we say that $L : K$ is a Galois extension if it is an extension that is normal and separable.

Theorem 8.1. *Suppose that $L : K$ is an algebraic extension. Then $L : K$ is Galois if and only if $K = \text{Fix}_L(\text{Gal}(L : K))$.*

Theorem 8.2. *Suppose that L is a field and G is a finite subgroup of $\text{Aut}(L)$, and put $K = \text{Fix}_L(G)$. Then $L : K$ is a finite Galois extension with $[L : K] = |\text{Gal}(L : K)|$, and furthermore $G = \text{Gal}(L : K)$.*

Theorem 8.3. *Suppose that $L : K$ is a finite extension. Then, if $L : K$ is a Galois extension, one has $|\text{Gal}(L : K)| = [L : K]$ and $K = \text{Fix}_L(\text{Gal}(L : K))$. If $L : K$ is not Galois, meanwhile, one has $|\text{Gal}(L : K)| < [L : K]$ and K is a proper subfield of $\text{Fix}_L(\text{Gal}(L : K))$.*

Proposition 16. *Suppose that $L : K$ is a Galois extension, and further that $L : M : K$ is a tower of field extensions. Then $L : M$ is a Galois extension.*

9 The main theorems of Galois theory

Definition 26. Suppose that $L : K$ is a field extension. When G is a subgroup of $\text{Aut}(L)$, we write $\phi(G)$ for $\text{Fix}_L(G)$, and when $L : M : K_0$ is a tower of field extensions with $K_0 = \phi(\text{Gal}(L : K))$, we write $\gamma(M)$ for $\text{Gal}(L : M)$.

Theorem 9.1 (The Fundamental Theorem of Galois Theory). *Suppose that $L : K$ is a finite extension, let $G = \text{Gal}(L : K)$, and put $K_0 = \phi(G)$. Then one has the following:*

- (a) *the map ϕ is a bijection from the set of subgroups of G onto the set of fields M intermediate between L and K_0 , and γ is the inverse map;*
- (b) *if $H \leq G$, then $H \trianglelefteq G$ if and only if $\phi(H) : K_0$ is a normal extension;*
- (c) *if $H \trianglelefteq G$, one has $\text{Gal}(\phi(H) : K_0) \cong G/H$. In particular, if $\sigma \in G$, one has $\sigma|_{\phi(H)} \in \text{Gal}(\phi(H) : K_0)$, and the map $\sigma \mapsto \sigma|_{\phi(H)}$ is a homomorphism of G onto $\text{Gal}(\phi(H) : K_0)$ with kernel H .*

Definition 27 (Galois group of polynomial). When $f \in K[t]$ and $L : K$ is a splitting field extension for f , we define the Galois group of the polynomial f over K to be $\text{Gal}_K(f) = \text{Gal}(L : K)$.

10 Finite fields

11 Solvability and solubility

Definition 28 (Soluble group). A finite group G is soluble if there is a series of groups

$$\{\text{id}\} = G_0 \leq G_1 \leq \cdots \leq G_n = G,$$

with the property that $G_i \trianglelefteq G_{i+1}$ and G_{i+1}/G_i is abelian ($0 \leq i < n$).

Theorem 11.1. *Let K be a field of characteristic 0. Then $f \in K[t]$ is solvable by radicals if and only if $\text{Gal}_K(f)$ is soluble.*

Lemma 1. *Suppose $\text{char}(K) = 0$ and $L : K$ is a radical extension. Then there exists an extension $N : L$ such that $N : K$ is normal and radical.*

Definition 29 (Cyclic extension). The extension $L : K$ is cyclic if $L : K$ is a Galois extension and $\text{Gal}(L : K)$ is a cyclic group.

Lemma 2. *Suppose that $\text{char}(K) = 0$ and let p be a prime number. Also, let $L : K$ be a splitting field extension for $t^p - 1$. Then $\text{Gal}(L : K)$ is cyclic, and hence $L : K$ is a cyclic extension.*

Lemma 3. *Let $\text{char}(K) = 0$ and suppose that n is an integer such that $t^n - 1$ splits over K . Let $L : K$ be a splitting field extension for $t^n - a$, for some $a \in K$. Then $\text{Gal}(L : K)$ is abelian.*

Theorem 11.2. *Let $\text{char}(K) = 0$ and suppose that $L : K$ is Galois. Suppose that there is an extension $M : L$ with the property that $M : K$ is radical. Then $\text{Gal}(L : K)$ is soluble.*

Corollary 5. *Suppose that $\text{char}(K) = 0$. Then $\text{Gal}_K(f)$ is soluble whenever $f \in K[t]$ is soluble by radicals.*

Corollary 6. *There exist quintic polynomials in $\mathbb{Q}[t]$ with insoluble Galois groups, such as $f(t) = t^5 - 4t + 2$, and which are not solvable by radicals.*

Lemma 4. *Let $\text{char}(K) = 0$, and suppose that $L : K$ is a cyclic extension of degree n . Suppose also that K contains a primitive n -th root of 1. Then there exists $\theta \in K$ having the property that $t^n - \theta$ is irreducible over K , and $L : K$ is a splitting field for $t^n - \theta$. Further, if β is a root of $t^n - \theta$ over L , then $L = K(\beta)$.*

Theorem 11.3. *Let $\text{char}(K) = 0$, and suppose that $f \in K[t] \setminus K$. Then f is solvable by radicals whenever $\text{Gal}_K(f)$ is soluble.*