

## GALOIS THEORY: SOLUTIONS TO HOMEWORK 11

1. Suppose that  $L : M : K$  is an algebraic tower of fields. Prove that  $L : K$  is separable if and only if  $L : M$  and  $M : K$  are both separable. [Hint: try using the Primitive Element Theorem].

**Solution:** We showed in Proposition 7.1 that when  $L : K$  is separable, then so too is  $L : M$ . Meanwhile, the separability, in such circumstances, of  $M : K$  is inherited from that of  $L : K$ . Conversely, suppose that  $L : M$  and  $M : K$  are both separable, and suppose that  $\alpha \in L$ . Then since  $L : M$  is separable, one finds that  $\alpha$  is separable over  $M$ . The polynomial  $m_\alpha(M)$  has its coefficients defined in a subfield  $M'$  of  $M$  with  $M' : K$  a finite separable extension. Since  $m_\alpha(M') = m_\alpha(M)$  is separable, we deduce that  $\alpha$  is separable over  $M'$ . Thus, since  $M' : K$  is finite and separable, it follows from the primitive element theorem that there exists  $\beta \in M'$  such that  $M' = K(\beta)$ , whence Theorem 7.4 implies that  $M'(\alpha) : K$ , or equivalently  $K(\alpha, \beta) : K$ , is separable. Consequently, we deduce that  $\alpha \in K(\alpha, \beta)$  is separable over  $K$ . Since this conclusion holds for all  $\alpha \in L$ , we conclude that  $L : K$  is separable.

2. Suppose that  $E : K$  and  $F : K$  are finite extensions with  $K \subseteq E \subseteq L$  and  $K \subseteq F \subseteq L$ , with  $L$  a field.

(a) Show that when  $E : K$  is separable, then so too is  $EF : F$ .

**Solution:** By the primitive element theorem, we may suppose that  $E = K(\alpha)$  for some  $\alpha \in E$  separable over  $K$ . Thus  $EF = F(\alpha)$ . Since  $\alpha$  is separable over  $K$ , it is also separable over  $F$ , and hence it follows from Theorem 7.4 that  $F(\alpha) : F$ , or equivalently  $EF : F$ , is separable.

(b) Show that when  $E : K$  and  $F : K$  are both separable, then so too are  $EF : K$  and  $E \cap F : K$ .

**Solution:** When  $E : K$  and  $F : K$  are both separable, then  $EF : F$  is separable, and hence  $EF : F : K$  is a tower of extensions with  $EF : F$  and  $F : K$  both separable. Then it follows from problem 1 that  $EF : K$  is separable. Likewise, one has the tower  $E : E \cap F : K$  of extensions with  $E : K$  separable. Then it follows from problem 1 that  $E \cap F : K$  is separable.

3. Suppose that  $\text{char}(K) = p > 0$  and that  $L : K$  is a totally inseparable algebraic extension (thus, every element of  $L \setminus K$  is inseparable). Show that whenever  $\alpha \in L$ , then there is a non-negative integer  $n$  and an element  $\theta \in K$  having the property that  $m_\alpha(K) = t^{p^n} - \theta$ .

**Solution:** Suppose that  $\alpha \in L$ . Then  $m_\alpha(K)$  is an irreducible polynomial over  $K$ , so by question 4(a) has the shape  $g(t^{p^n})$  for some non-negative integer  $n$  and an irreducible separable polynomial  $g$ . Suppose that  $g$  has degree 2 or more, and that its distinct roots in  $\overline{K}$  are  $\beta_1, \dots, \beta_d$ . Then for some index  $i$  one has  $\beta_i = \alpha^{p^n}$  and  $m_{\beta_i}(K) = g(t)$ , by the irreducibility of

$g$ . But then  $\beta_i \in L$  is separable, because  $g$  is separable, contradicting the totally inseparable property of the extension  $L : K$ . It follows that  $g$  must have degree 1, and hence  $m_\alpha(K) = t^{p^n} - \theta$ , where  $\theta = \alpha^{p^n} \in K$ .

©Trevor D. Wooley, Purdue University 2024. This material is copyright of Trevor D. Wooley at Purdue University unless explicitly stated otherwise. It is provided exclusively for educational purposes at Purdue University, and is to be downloaded or copied for your private study only. 