

Context-dependent host-microbe interactions in stochastic environments

Joshua C. Fowler^{a,1}, Shaun Ziegler^b, Kenneth D. Whitney^b, Jennifer A. Rudgers^b, and Tom E. X. Miller^a

^aRice University, Department of BioSciences, Houston, TX, 77005; ^bUniversity of New Mexico, Department of Biology, Albuquerque, NM, 87131

This manuscript was compiled on May 27, 2020

Please provide an abstract of no more than 250 words in a single paragraph. Abstracts should explain to the general reader the major contributions of the article. References in the abstract must be cited in full within the abstract itself and cited in the text.

Keyword 1 | Keyword 2 | Keyword 3 | ...

Classic ecological theory predicts that environmental variation will tend to have negative consequences for long-term population growth rates (1, 2). Long-term population growth rates (λ) incur a cost due to temporal variation. Our ability to explore the demographic consequences of environmental variation in nature relies on long-term observational studies and experiments that capture natural climatic variation (cite) plus maybe examples of other studies looking at demographic buffering. While there is appreciation for long-term studies and recognition of the importance of studying both climate mean and variance in ecology, demographic studies that examine demographic buffering are limited due to the need for long term data and the need to account for multiple sources of variation within data (3).

Long-term population growth rates, which are calculated as geometric means, incur a cost due to temporal variation (1, 2). A population will increase over time if the long term growth rate (λ) is > 1 , and can be expected to decrease if $\lambda < 1$. There are two pathways by which population growth can increase: mean λ can increase or variance can decrease (). include math somewhere here. As in the demographic buffering hypothesis, where the fitness consequences of environmental variability may select for buffering in the vital rates that are most consequential for population growth (4), the fitness consequences of species interactions may apply to both the mean and variance of vital rates. Whether variance buffering by species interactions occurs is an underexplored question, but it may come to be of increasing importance under climate change.

Climate projections indicate that environmental variability is expected to increase along with increases in mean climate conditions (5, 6). Contributions from demographic buffering in natural populations may become more important under this scenario and will be important for projecting species' responses to climate change (7). In particular, it is unclear how commonly demographic buffering plays an important role in population dynamics in general, and how species interactions may contribute to demographic buffering (cite). Mutualistic symbioses in particular may have the potential to provide resilience to environmental variability (cite). I don't know about that last sentence, but I need some sort of transition to symbiosis.

In nature, microbial symbionts provide protection from environmental stresses across a broad range of taxa, including stress caused by drought, salinity, and temperature (cite).

Commonly, the benefits from these symbioses are context-dependent where the magnitude of interaction benefit changes depending on environmental conditions (8). This can make it difficult to quantify the net effect of a given interaction, but it also allows for the possibility that interaction strength can vary through time (cite). Symbionts may provide benefits under harsh conditions when they are needed by their hosts, but be neutral or even costly under benign conditions (cite). Over time, this may lead symbiont-associated organisms to experience a reduction in variation in vital rates by reducing the frequency of extreme events (conceptual figure). Variance buffering by symbionts is novel mechanism that may be common across many symbioses that

Using long-term data from experimental grass-fungal endophyte plots, we test the hypothesis that symbionts buffer hosts from the fitness consequences of environmental variability. Specifically, we ask if fungal endophytes buffer demographic variance in their grass hosts, and, if so, what is the relative importance of demographic buffering vs. mean effects in the overall fitness impact of the symbiosis. With 13 years of demographic data, we employ structured, stochastic population models for seven species of cool-season grass hosts that are commonly infected with fungal endophytes (*Lolium arundinaceum*, *Festuca subverticillata*, *Elymus virginicus*, and *Elymus villosus*, *Poa alsodes* and *Poa sylvestris*).

This paragraph is mostly talking off my head about results, but my idea is to include a brief statement of our results. Across species, we find that variance buffering by endophytes contributes (percentage) to population growth rates. While the effect is generally weaker than effects on the mean, we found that buffering was common in the most sensitive vital rates, and was most important for xxx species with xxx life history.

Significance Statement

Authors must submit a 120-word maximum statement about the significance of their research paper written at a level understandable to an undergraduate educated scientist outside their field of speciality. The primary goal of the Significance Statement is to explain the relevance of the work in broad context to a broad readership. The Significance Statement appears in the paper itself and is required for all research papers.

Please provide details of author contributions here.

Please declare any conflict of interest here.

¹To whom correspondence should be addressed. E-mail: jcf3rice.edu

79 Results

80 Discussion

81 Guide to using this template on Overleaf

82 Please note that whilst this template provides a preview of the
83 typeset manuscript for submission, to help in this preparation,
84 it will not necessarily be the final publication layout. For
85 more detailed information please see the [PNAS Information](#)
86 [for Authors](#).

87 If you have a question while using this template on Overleaf,
88 please use the help menu (“?”) on the top bar to search for [help](#)
89 [and tutorials](#). You can also [contact the Overleaf support team](#)
90 at any time with specific questions about your manuscript or
91 feedback on the template.

92 **Author Affiliations.** Include department, institution, and com-
93 plete address, with the ZIP/postal code, for each author. Use
94 lower case letters to match authors with institutions, as shown
95 in the example. Authors with an ORCID ID may supply this
96 information at submission.

97 **Submitting Manuscripts.** All authors must submit their arti-
98 cles at [PNAScentral](#). If you are using Overleaf to write your
99 article, you can use the “Submit to PNAS” option in the top
100 bar of the editor window.

101 **Format.** Many authors find it useful to organize their
102 manuscripts with the following order of sections; Title, Author
103 Affiliation, Keywords, Abstract, Significance Statement, Re-
104 sults, Discussion, Materials and methods, Acknowledgments,
105 and References. Other orders and headings are permitted.

106 **Manuscript Length.** PNAS generally uses a two-column for-
107 mat averaging 67 characters, including spaces, per line. The
108 maximum length of a Direct Submission research article is
109 six pages and a Direct Submission Plus research article is ten
110 pages including all text, spaces, and the number of characters
111 displaced by figures, tables, and equations. When submitting
112 tables, figures, and/or equations in addition to text, keep the
113 text for your manuscript under 39,000 characters (including
114 spaces) for Direct Submissions and 72,000 characters (including
115 spaces) for Direct Submission Plus.

116 **References.** References should be cited in numerical order as
117 they appear in text; this will be done automatically via bibtex,
118 e.g. . All references should be included in the main manuscript
119 file.

120 **Data Archival.** PNAS must be able to archive the data es-
121 sential to a published article. Where such archiving is not
122 possible, deposition of data in public databases, such as Gen-
123 Bank, ArrayExpress, Protein Data Bank, Unidata, and others
124 outlined in the Information for Authors, is acceptable.

125 **Language-Editing Services.** Prior to submission, authors who
126 believe their manuscripts would benefit from professional edit-
127 ing are encouraged to use a language-editing service (see list at
128 [www.pnas.org/site/authors/language-editing.xhtml](#)). PNAS
129 does not take responsibility for or endorse these services, and
130 their use has no bearing on acceptance of a manuscript for
131 publication.

Table 1. Comparison of the fitted potential energy surfaces and ab initio benchmark electronic energy calculations

Species	CBS	CV	G3
1. Acetaldehyde	0.0	0.0	0.0
2. Vinyl alcohol	9.1	9.6	13.5
3. Hydroxyethylidene	50.8	51.2	54.0

nomenclature for the TSs refers to the numbered species in the table.

Digital Figures. Only TIFF, EPS, and high-resolution PDF
for Mac or PC are allowed for figures that will appear in the
main text, and images must be final size. Authors may submit
U3D or PRC files for 3D images; these must be accompanied
by 2D representations in TIFF, EPS, or high-resolution PDF
format. Color images must be in RGB (red, green, blue) mode.
Include the font files for any text.

Figures and Tables should be labelled and referenced in the
standard way using the `\label{}` and `\ref{}` commands.

Figure ?? shows an example of how to insert a column-wide
figure. To insert a figure wider than one column, please use
the `\begin{figure*}...\end{figure*}` environment. Fig-
ures wider than one column should be sized to 11.4 cm or 17.8
cm wide. Use `\begin{SCfigure*}...\end{SCfigure*}` for a
wide figure with side captions.

Tables. In addition to including your tables within this
manuscript file, PNAS requires that each table be uploaded to
the submission separately as a Table file. Please ensure that
each table .tex file contains a preamble, the `\begin{document}`
command, and the `\end{document}` command. This is neces-
sary so that the submission system can convert each file to
PDF.

Single column equations. Authors may use 1- or 2-column
equations in their article, according to their preference.

To allow an equation to span both columns, use the
`\begin{figure*}...\end{figure*}` environment mentioned
above for figures.

Note that the use of the `widetext` environment for equa-
tions is not recommended, and should not be used.

Supporting Information (SI). Authors should submit SI as a
single separate PDF file, combining all text, figures, tables,
movie legends, and SI references. PNAS will publish SI un-
composed, as the authors have provided it. Additional details
can be found here: [policy on SI](#). For SI formatting instructions
click [here](#). The PNAS Overleaf SI template can be found [here](#).
Refer to the SI Appendix in the manuscript at an appropri-
ate point in the text. Number supporting figures and tables
starting with S1, S2, etc.

Authors who place detailed materials and methods in an
SI Appendix must provide sufficient detail in the main text
methods to enable a reader to follow the logic of the procedures
and results and also must reference the SI methods. If a paper
is fundamentally a study of a new method or technique, then
the methods must be described completely in the main text.

SI Datasets. Supply Excel (.xls), RTF, or PDF files. This file
type will be published in raw format and will not be edited or
composed.

$$\begin{aligned}
 (x+y)^3 &= (x+y)(x+y)^2 \\
 &= (x+y)(x^2+2xy+y^2) \\
 &= x^3+3x^2y+3xy^2+x^3.
 \end{aligned}
 \tag{1}$$

SI Movies. Supply Audio Video Interleave (avi), Quicktime (mov), Windows Media (wmv), animated GIF (gif), or MPEG files and submit a brief legend for each movie in a Word or RTF file. All movies should be submitted at the desired reproduction size and length. Movies should be no more than 10 MB in size.

3D Figures. Supply a composable U3D or PRC file so that it may be edited and composed. Authors may submit a PDF file but please note it will be published in raw format and will not be edited or composed.

Materials and Methods

Plant propagation and endophyte treatment. Seeds from naturally infected populations of seven species of cool-season grasses (*Agrostis perennans*, *Elymus villosus*, *Elymus virginicus*, *Festuca subverticillata*, *Lolium arundinaceum*, *Poa alsodes*, and *Poa sylvestris*) were collected in the Spring of 2006 for Lilly Dickie Woods and Bayles Road in Brown. Co. IN. Seeds with shared maternal ancestry were either experimentally disinfected by heat treatments or left naturally infected to reduce confounding genotype effects. Seeds were surface sterilized with XXXX and cold stratified for XXXX weeks, then germinated in the XXXX for XXXX weeks. They were then grown in the greenhouse at Indiana University for XXXX weeks.

Experimental design. We collected long-term demographic data from experimental plots established in 2007. We established 10 plots for *Lolium arundinaceum*, *Festuca subverticillata*, *Elymus virginicus*, and *Elymus villosus* and 18 plots for *Poa alsodes* and *Poa sylvestris* with 25 individuals.

Data analysis. We used statistics

ACKNOWLEDGMENTS. Please include your acknowledgments here, set in a single paragraph. Please do not include any acknowledgments in the Supporting Information, or anywhere else in the manuscript.

1. R. C. Lewontin and D. Cohen. On Population Growth in a Randomly Varying Environment. *Proceedings of the National Academy of Sciences*, 62(4):1056–1060, April 1969. ISSN 0027-8424, 1091-6490. . URL <https://www.pnas.org/content/62/4/1056>. Publisher: National Academy of Sciences Section: Biological Sciences: Zoology.
2. Shripad D. Tuljapurkar. Population dynamics in variable environments. III. Evolutionary dynamics of *r*-selection. *Theoretical Population Biology*, 21(1):141–165, February 1982. ISSN 0040-5809. . URL <http://www.sciencedirect.com/science/article/pii/0040580982900107>.
3. Christoffer H. Hilde, Marlène Gamelon, Bernt-Erik Sæther, Jean-Michel Gaillard, Nigel G. Yoccoz, and Christophe Pélabon. The Demographic Buffering Hypothesis: Evidence and Challenges. *Trends in Ecology & Evolution*, 0(0), March 2020. ISSN 0169-5347. . URL [https://www.cell.com/trends/ecology-evolution/abstract/S0169-5347\(20\)30050-1](https://www.cell.com/trends/ecology-evolution/abstract/S0169-5347(20)30050-1). Publisher: Elsevier.
4. Catherine A Pfister. Patterns of variance in stage-structured populations: evolutionary predictions and ecological implications. *Proceedings of the National Academy of Sciences*, 95(1):213–218, 1998.
5. IPCC (Intergovernmental Panel on Climate Change). *Managing the risks of extreme events and disasters to advance climate change adaptation: special report of the intergovernmental panel on climate change. Special Report*. Cambridge University Press, 2012.
6. David T Iles, Robert F Rockwell, and David N Koons. Shifting vital rate correlations alter predicted population responses to increasingly variable environments. *The American Naturalist*, 193(3):E57–E64, 2019.
7. Daniel F Doak, William F Morris, Cathy Pfister, Bruce E Kendall, and Emilio M Bruna. Correctly estimating how environmental stochasticity influences fitness and population growth. *The American Naturalist*, 166(1):E14–E21, 2005.

8. Scott A Chamberlain, Judith L Bronstein, and Jennifer A Rudgers. How context dependent are species interactions? *Ecology letters*, 17(7):881–890, 2014.

239
240