

Context-dependent host-microbe interactions in stochastic environments

Joshua C. Fowler^{a,1}, Shaun Ziegler^b, Kenneth D. Whitney^b, Jennifer A. Rudgers^b, and Tom E. X. Miller^a

^aRice University, Department of BioSciences, Houston, TX, 77005; ^bUniversity of New Mexico, Department of Biology, Albuquerque, NM, 87131

This manuscript was compiled on October 21, 2022

Microbial symbioses are ubiquitous in nature, yet our ability to predict how these interactions affect species' responses to climate change has been limited by interaction outcomes that vary with environmental context. Increased environmental variability is a key prediction of climate change. Here we present and test a novel hypothesis: microbial symbionts may buffer hosts from environmental variability by being beneficial in harsh years while being neutral or costly in good years. Long-term data reveal that, while weaker than effects on the mean of population growth, variance buffering is an important contribution to hosts long-term population growth. Additionally, the symbiosis becomes more mutualistic under simulated increases in variability, driven by an increased contribution from variance buffering. Please provide an abstract of no more than 250 words in a single paragraph. Abstracts should explain to the general reader the major contributions of the article. References in the abstract must be cited in full within the abstract itself and cited in the text.

Keyword 1 | Keyword 2 | Keyword 3 | ...

Along with increases in average temperatures, global climate change is driving increases in the variability of precipitation events, temperature extremes, and droughts (1–3). Discerning the effects of environmental variability on population dynamics and species interactions are therefore key objectives of global change ecology.

*Classic theory predicts that long-term population growth rates (equivalently, population mean fitness) will decline under increased environmental variability due to negative effects of bad years that outweigh positive effects of good years (a consequence of nonlinear averaging) (4, 5). For example, in unstructured populations, the long-term stochastic growth rate in fluctuating environments (λ_s) will always be less than the growth rate averaged across environments ($\bar{\lambda}$) by an amount proportional to the environmental variance (σ^2): $\log(\lambda_s) \approx \log(\bar{\lambda}) - \frac{\sigma^2}{2\bar{\lambda}^2}$. Populations structured by size or stage are expected to similarly experience negative effects of variability (6, 7). Thus, there are two pathways to increase population viability in a stochastic environment λ_s : increasing the mean growth rate and/or dampening temporal fluctuation in growth rates, also called variance “buffering”.

Given the potential for negative fitness consequences of increasing environmental stochasticity under global change, there is growing interest in factors that can amplify or dampen demographic fluctuations. Much attention has focused on the properties of species or their environment as modifiers of fluctuations in fitness, including variation in vital rates (survival, growth and reproduction)(9), correlations between vital rates (11), transient shifts in stage structure (12), and the degree of environmental autocorrelation (13, 14). In contrast, little is known about how biotic interactions influence patterns

of demographic variability (15), due in part to the paucity of long-term interaction data that are required to detect such effects.

Our work focused on the potential for microbial symbionts to buffer host plants against negative effects of environmental variability. Microbial symbioses are ubiquitous across plant and animal host taxa and can be crucial determinants of host fitness (16, 17). Yet we know little about their potential influence on responses to climate change and environmental variation (18). Across a broad range of taxa, mutualistic host-associated microbes provide protection from environmental stresses including drought, extreme temperatures, and enemies (19–23).[†] The role they play may be under-appreciated, and it can be difficult to quantify the net outcome of a given interaction because they are often viewed as being context-dependent where the magnitude of benefit depends on environmental conditions (24). Rather than considering context-dependence as some unexplainable intricacy of species interactions, environmental variation opens up the possibility for interaction strength to vary through time (25, 26) and to influence the variance of population growth rates.[‡]

We hypothesize that symbionts may provide benefits under harsh conditions when they are needed by their hosts, but be neutral or even costly under benign conditions. Over time, this would lead symbiotic hosts to experience a reduction in variation in vital rates by reducing the frequency of extreme years. Incorporating context-dependence in this way, we highlight a novel mechanism by which symbionts can act as mutualists that may come to be of increasing importance in a more variable future.[§]

Using data from a unique, long-term symbiont-removal

[†] This sentence seems to contradict the previous sentence.

[‡] I agree that this is the place to introduce context-dependence and connect climate variability to variability in interaction outcome. But I don't think what you have here works.

[§] Similar comment as above. I think this is the place to lay out the hypotheses of symbionts elevating mean and/or reducing variance. But I don't think this is sufficiently explanatory or provides sufficient context or rationale for the ideas to really hit home.

Significance Statement

Authors must submit a 120-word maximum statement about the significance of their research paper written at a level understandable to an undergraduate educated scientist outside their field of speciality. The primary goal of the Significance Statement is to explain the relevance of the work in broad context to a broad readership. The Significance Statement appears in the paper itself and is required for all research papers.

Please provide details of author contributions here.

Please declare any conflict of interest here.

¹To whom correspondence should be addressed. E-mail: jcf3rice.edu

* It would be nice to start paragraph 2 here, but that probably leaves P1 too skimpy.

experiment, we test the hypothesis that context-dependent benefits of microbial symbionts buffer hosts from the fitness consequences of environmental variability[†]. The experiment consists of annually censused plots initiated in 2007 with seven grass species that host Epichloë fungal endophytes. These specialized fungal endophytes are unique to grasses, occurring in at least 30%[‡] of cool-season (C3) species, and are primarily vertically transmitted from maternal plant to seed (27). While they have been associated with contributing to drought tolerance for their hosts^{**}, these benefits are commonly context-dependent^{††} (28–30). And so^{‡‡}, we ask first how fungal endophytes influence the mean and interannual variance of their hosts' vital rates; next, we ask if these vital rate effects buffer variance in fitness and, if so, what is the relative contribution of variance buffering vs. mean effects to the overall effect of the symbiosis on long-term growth rates. To answer these questions, we build structured, stochastic population models for these seven grass host species (*Agrostis perennans*, *Elymus villosus*, *Elymus virginicus*, *Festuca subverticillata*, *Lolium arundinaceum*, *Poa alsodes* and *Poa sylvestris*).^{§§} These long-term plots contain either naturally symbiotic plants (E+) or those which have had their symbionts experimentally removed (E-).^{¶¶} Each annual census is a sample of climatic variation. Across 14 years, the data contain 31,216 individual-transition years.^{***} After quantifying endophyte effects on mean and variance in population growth rates, we use simulations to explore the consequences of variance buffering under increased variance^{†††} and construct climate-explicit population models^{§§§} to evaluate the role of climate drivers as explanations for this buffering.

Intriguingly, we find that the symbiosis contributes positively to long-term population growth rates through both mean and variance buffering effects. Integrating across diverse effects on vital rates, contributions to long-term growth rates from effects on the mean are 4.17 times greater on average than contributions from variance buffering. However, these effects varied between species; for example, two species (*A. perennans* and *P. sylvestris*) have contributions from variance buffering which are greater than mean effects. Additionally, the effect of mutualism increases under simulations with increased variance driven by greater contributions of variance buffering. In the most extreme scenario, we find that variance buffering contributions across species are 1.5 times greater than effects on the mean on average, and that variance buffering contributions are greater than mean effects for five out of seven species.^{§§§}

[†] In addition to this hypothesis, I think it is important to emphasize the question of relative importance of mean vs variance effects. Like Volker always says, yes/no hypotheses are often a little dull.

[‡] I think the 30% number comes from a Leuchtmann paper.

^{**} Awkward phrasing.

^{††} This is a little confusing because it sounds like you are saying that the drought benefit is context-dependent, but I think drought is the context, so the benefits vary with presence or absence of drought.

^{‡‡} too colloquial

^{§§} Mention the models somewhere in this paragraph but don't list the species. Species list should definitely go somewhere and should include the endophyte species to the best of our known – maybe this is an appendix table.

^{¶¶} Should come earlier when you first introduce the "unique" experiment.

^{***} Put in results.

^{†††} Should be emphasized earlier. It's cool and novel.

^{§§§} I would not mention this here since it is not central to the story of the paper.

^{§§§} I would eliminate this paragraph entirely. Not needed, since you launch directly into results from here.

Results

Endophyte effects on the mean and variance of vital rates.

Across eight host species, seven vital rates, 14 years, and *large number* individual host-plants, we found that Epichloë fungal endophyte symbiosis generally had positive effects on host demographic performance and negative effects on inter-annual demographic variance. The mean endophyte effect was positive for *number* out of 56 host species–vital rate combinations, and was particularly strong for host survival and growth. Endophytes also consistently reduced inter-annual variance for the majority of host species and vital rates (*number* out of 56 host species–vital rate combinations; Fig. 1), consistent with the hypothesis of variance buffering. Variance buffering effects were particularly strong for reproductive vital rates (e.g. likelihood of flowering and panicle production); This might align with predictions from life-history theory suggesting that reproductive vital rates are likely to experience greater year-to-year variance (¹⁷).

The¹⁸ relative magnitude of symbiont effects on means and variances was idiosyncratic across species and across vital rates. For example, endophytes increased mean adult survival and caused a modest variance reduction for *P. alsodes*, while for *F. subverticillata*, effects from variance buffering were stronger with a relatively weaker mean effect. In other vital rates, such as in seedling growth, *P. alsodes* experiences stronger buffering than *F. subverticillata*. Interestingly, there were also certain vital rates that showed costs¹⁹ of endophyte symbiosis, such as *A. perennans* and *F. subverticillata* which had lower mean germination rates when partnered with endophytes.²⁰

Endophyte effects on the mean and variance of population growth rates.

Not all vital rates contribute equally to fitness, so we used stochastic matrix models (where tiller number was the integer-valued state variable) to integrate the diverse effects described above into comprehensive measures for the mean and variance of host fitness. We found that, on average across host species, endophyte-symbiotic populations experienced a 9.2 % increase in mean fitness $\bar{\lambda}$. Our hierarchical Bayesian analysis, which propagates uncertainty from the underlying data through model predictions, indicates 92.8% confidence that endophytes increased $\bar{\lambda}$. The standard deviation²¹, reflecting inter-annual variability in fitness, was 6.6 % lower in endophyte-symbiotic populations than endophyte-free populations, averaging across host species (with 66% certainty²² that the endophyte effect was negative) (Fig. 2). For some host species, the standard deviation was reduced by as much as 44.3% (*L. arundinaceum*) and 28.5% (*F. subverticillata*), while for others, variance effects were much smaller, or even

^{¶¶¶} I changed all your tenses to past tense.

¹⁷ This is an interesting and possibly useful observation here, but it implies that negative effects on variance were proportional to the absolute value of variance. This seems plausible - is that the case? Also, the best citation would be Pfister 1998.

¹⁸ We talked about updating the figure so that individual examples would pop out of the matrix. Presumably you would reference that figure here where you talk about specific examples.

¹⁹ If you are going to highlight costs then it would be good to show both reductions in mean and increases in variance.

²⁰ Not sure how much to talk about demographic compensation (i.e. do we see less variance in vital rates that are more important). We don't really assess this. And at least looking at the actual sigma values for each species, it seems like none of the vital rates have hugely different baseline standard deviations. TOM: I would not talk about demographic compensation.

²¹ The figure in this draft shows CV. We should be consistent with what is shown and described.

²² The CV figure I am looking at is more than 66% certainty.



Fig. 1. needs to be updated pdf version, but putting this here for now. Effect of endophytes on vital rates across species




Fig. 2. needs to be updated pdf version, but putting this here for now. Effect of endophytes on mean and CV of population growth

slightly positive (*E. villosus* and *E. virginicus*).²³ Here is where I think the biplot figure makes an interesting point. For some species the benefit is through the mean, for some it is through variance, for some it is both – but we don't see any negative-negative. It's really compelling evidence for mutualism being pervasive here, but taking different forms.²⁴

Contribution of mean and variance effects to long-term growth rates. Having documented positive fitness effects of endophyte symbiosis via mean benefits and / or variance buffering, we next asked about the relative importance of those two pathways of host-symbiont mutualism for the stochastic growth rate λ_S . We decomposed the overall effect of the symbiosis on λ_S into

contributions from mean and variance buffering effects (32)²⁵. We found that variance buffering provided a small benefit overall (XXX%)²⁶ (Fig. 3) but note that even small changes in fitness can have profound effects over long time periods²⁷.

Role of symbiotic buffering under increased variance. In simulations with increased variance, we find that the contribution of variance buffering increases.²⁸

Endophytes are buffering their hosts from environmental variability. Symbiotic and non-symbiotic populations responded

²³ I am thinking it might be valuable to show that biplot

²⁴ This comment is neither here nor there, but we can probably expect reviewers to ask was the realized population growth patterns were, i.e., did E+ plots grow faster than E- plots. Not sure how much we should go there yet (because the populations mostly shrank) but I do think we need to better communicate that we are talking about asymptotic growth rates.

²⁵ I don't know if there's a better citation for this, this is more about IPM's— Tom: I don't think we need a citation here, but we should briefly describe how we did this, since readers will get here before the methods. I've tried to add a little of this throughout the Results.

²⁶ Seems like this section is still under construction, but be sure to emphasize that across species there is really strong evidence for mutualism (positive effect on λ_S) and in the current regime of environmental variability that is dominated by the mean benefits.

²⁷ probably would not say this

²⁸ Skipping this for now, but it might make more sense to talk about the ambient and increased variance scenarios in the same results section, since they are in the same figure.




Fig. 3. needs to be updated pdf version, but putting this here for now. Stochastic lambda contributions with simulations of increased variance

distinctly to environmental drivers. Our climate-explicit analysis shows that on average, the population growth of non-endophytic plants was more sensitive to our drought index, SPEI, than endophytic populations (FIGURE). This pattern was particularly pronounced for species which showed strong buffering in our earlier analysis (SPECIES), suggesting that endophytes are buffering their hosts from this aspect of inter-annual variation, although a large amount of variation remained unexplained.²⁹

Discussion

Guide to using this template on Overleaf

Please note that whilst this template provides a preview of the typeset manuscript for submission, to help in this preparation, it will not necessarily be the final publication layout. For more detailed information please see the [PNAS Information for Authors](#).

If you have a question while using this template on Overleaf, please use the help menu (“?”) on the top bar to search for [help and tutorials](#). You can also [contact the Overleaf support team](#) at any time with specific questions about your manuscript or feedback on the template.

Author Affiliations. Include department, institution, and complete address, with the ZIP/postal code, for each author. Use lower case letters to match authors with institutions, as shown in the example. Authors with an ORCID ID may supply this information at submission.

Submitting Manuscripts. All authors must submit their articles at [PNAScentral](#). If you are using Overleaf to write your article, you can use the “Submit to PNAS” option in the top bar of the editor window.

Format. Many authors find it useful to organize their manuscripts with the following order of sections; Title, Author Affiliation, Keywords, Abstract, Significance Statement, Results, Discussion, Materials and methods, Acknowledgments, and References. Other orders and headings are permitted.

²⁹ I am wondering if it might be better to incorporate this material into the ealier section that describes variance buffering, so really what we are doing here is just providing a more mechanistic explanation for that phenomenological result.

Manuscript Length. PNAS generally uses a two-column format averaging 67 characters, including spaces, per line. The maximum length of a Direct Submission research article is six pages and a Direct Submission Plus research article is ten pages including all text, spaces, and the number of characters displaced by figures, tables, and equations. When submitting tables, figures, and/or equations in addition to text, keep the text for your manuscript under 39,000 characters (including spaces) for Direct Submissions and 72,000 characters (including spaces) for Direct Submission Plus.

References. References should be cited in numerical order as they appear in text; this will be done automatically via bibtex, e.g. . All references should be included in the main manuscript file.

Data Archival. PNAS must be able to archive the data essential to a published article. Where such archiving is not possible, deposition of data in public databases, such as GenBank, ArrayExpress, Protein Data Bank, Unidata, and others outlined in the Information for Authors, is acceptable.

Language-Editing Services. Prior to submission, authors who believe their manuscripts would benefit from professional editing are encouraged to use a language-editing service (see list at [www.pnas.org/site/authors/language-editing.xhtml](#)). PNAS does not take responsibility for or endorse these services, and their use has no bearing on acceptance of a manuscript for publication.

Digital Figures. Only TIFF, EPS, and high-resolution PDF for Mac or PC are allowed for figures that will appear in the main text, and images must be final size. Authors may submit U3D or PRC files for 3D images; these must be accompanied by 2D representations in TIFF, EPS, or high-resolution PDF format. Color images must be in RGB (red, green, blue) mode. Include the font files for any text.

Figures and Tables should be labelled and referenced in the standard way using the `\label{}` and `\ref{}` commands.

Figure ?? shows an example of how to insert a column-wide figure. To insert a figure wider than one column, please use the `\begin{figure*}...\end{figure*}` environment. Figures wider than one column should be sized to 11.4 cm or 17.8

$$\begin{aligned}
 (x + y)^3 &= (x + y)(x + y)^2 \\
 &= (x + y)(x^2 + 2xy + y^2) \\
 &= x^3 + 3x^2y + 3xy^2 + y^3.
 \end{aligned}
 \tag{1}$$

Table 1. Comparison of the fitted potential energy surfaces and ab initio benchmark electronic energy calculations

Species	CBS	CV	G3
1. Acetaldehyde	0.0	0.0	0.0
2. Vinyl alcohol	9.1	9.6	13.5
3. Hydroxyethylidene	50.8	51.2	54.0

nomenclature for the TSs refers to the numbered species in the table.

cm wide. Use `\begin{SCfigure*}...\end{SCfigure*}` for a wide figure with side captions.

Tables. In addition to including your tables within this manuscript file, PNAS requires that each table be uploaded to the submission separately as a Table file. Please ensure that each table .tex file contains a preamble, the `\begin{document}` command, and the `\end{document}` command. This is necessary so that the submission system can convert each file to PDF.

Single column equations. Authors may use 1- or 2-column equations in their article, according to their preference.

To allow an equation to span both columns, use the `\begin{figure*}...\end{figure*}` environment mentioned above for figures.

Note that the use of the `widetext` environment for equations is not recommended, and should not be used.

Supporting Information (SI). Authors should submit SI as a single separate PDF file, combining all text, figures, tables, movie legends, and SI references. PNAS will publish SI uncomposed, as the authors have provided it. Additional details can be found here: [policy on SI](#). For SI formatting instructions click [here](#). The PNAS Overleaf SI template can be found [here](#). Refer to the SI Appendix in the manuscript at an appropriate point in the text. Number supporting figures and tables starting with S1, S2, etc.

Authors who place detailed materials and methods in an SI Appendix must provide sufficient detail in the main text methods to enable a reader to follow the logic of the procedures and results and also must reference the SI methods. If a paper is fundamentally a study of a new method or technique, then the methods must be described completely in the main text.

SI Datasets. Supply Excel (.xls), RTF, or PDF files. This file type will be published in raw format and will not be edited or composed.

SI Movies. Supply Audio Video Interleave (avi), Quicktime (mov), Windows Media (wmv), animated GIF (gif), or MPEG files and submit a brief legend for each movie in a Word or RTF file. All movies should be submitted at the desired reproduction size and length. Movies should be no more than 10 MB in size.

3D Figures. Supply a composable U3D or PRC file so that it may be edited and composed. Authors may submit a PDF file but please note it will be published in raw format and will not be edited or composed.

Materials and Methods

Natural history of grass-endophyte symbiosis. To study the effects of context-dependent microbial symbiosis, we focused on *Epichloë* fungal endophytes, which live in the aboveground tissue of many species of cool-season grasses and grow into their hosts' seeds where they can be transmitted vertically from mother to offspring plants. This vertical transmission couples host and symbiont fitness and leads to the expectation that the interaction be mutualistic, else the fungi cause their host to be selected out of the population (33–35). While there are demonstrated benefits against herbivory(36) and under drought stress (37) for some host species, these interactions outcomes are commonly context-dependent (28, 29).

Plant propagation and endophyte removal. Seeds from naturally infected populations of seven species of cool-season grasses (*Agrostis perennans*, *Elymus villosus*, *Elymus virginicus*, *Festuca subverticillata*, *Lolium arundinaceum*, *Poa alsodes*, and *Poa sylvestris*) were collected during the 2006 growing season from Lilly Dickie Woods (39.238533, -86.218150) and the Bayles Road Teaching and Research Preserve (39.220167, -86.542683) in Brown Co. IN. To reduce confounding genotype effects, seeds with shared maternal ancestry were disinfected with heat treatments (6d in a drying oven at 60° C for *E. villosus*, *E. virginicus*, *F. subverticillata*, and *L. arundinaceum*; 7d in a drying oven at 60° C for *P. alsodes*, and *P. sylvestris*; and 12 min. in a hot water bath at 60° C for *A. perennans*)³⁰ or left naturally infected. Seeds were surface sterilized with bleach and cold stratified for ??? weeks, then germinated in a growth chamber before being transferred to the greenhouse at Indiana University and allowed to grow for XXXX weeks. We confirmed the endophyte status of these plants using leaf peels, where tissue from the leaf sheath is stained with aniline blue dye and examined for the presence of fungal hyphae (38). Then, we established the experimental plots with **vegetatively propagated clones of similar sizes from the plants**³¹ to reduce the potential for negative side effects of heat treatments (39).

Experimental design and data collection. During the spring of 2007, we established 10 3x3 plots for *A. perennans*, *E. villosus*, *E. virginicus*, *F. subverticillata*, and *L. arundinaceum* and 18 plots for *P. alsodes* and *P. sylvestris*. For each species, an equal number of plots were randomly assigned to each endophyte status, E+ or E-. Each plot was planted with 20 evenly spaced symbiotic or symbiont-free individuals respectively and each plant marked with aluminum tags.

Each summer starting in 2007, we censused all individuals in each plot for survival, growth and reproduction, garnering a dataset covering 14 years that contains 31,216 individual transition years. After clearing out leaf litter, for each plant alive in the previous year, we marked its survival and measured its size as a count of the number of tillers. Further, we collected reproductive data by counting the number of reproductive tillers, and then counting the number of seed-bearing spikelets on up to three of those reproductive tillers. In 2009, we took additional counts of seeds per inflorescence. Together, we use these measurements to estimate seed production. In each plot, we also survey for and tag any unmarked individuals.

³⁰ need to double check methods for temp, duration, etc.

³¹ not sure this happened

New recruits typically have one tiller and are non-reproductive, but we also find and tag any individuals that may have been missed in previous censuses.

We typically expect plots of each endophyte status to maintain their status because the fungi are almost entirely vertically transmitted and plots are spaced at least 5 m apart, limiting the possibility for unwanted dispersal between plots or horizontal transmission of the fungus. Seeds from reproductive individuals are opportunistically taken and scored for their endophyte status. Overall, these scores reflect a 97.5% faithfulness of recruits to their expected endophyte status across species and plots (Supplement data).

Demographic modeling. Armed with this demographic data, we next constructed size-structured, stochastic population models. These models describe transitions between sizes (measured as a count of tillers) from one year to the next. For all species, we include a 1 year reproductive delay in the population model following the observation that these newly recruited plants are rarely observed flowering in their first year. Our population model can be expressed as:

$$\mathbf{n}_{t+1} = \mathbf{A}\mathbf{n}_t \quad [2]$$

where \mathbf{n}_{t+1} is a vector of abundances across sizes in year $t+1$ for each species and endophyte status.

$$\mathbf{n}_{t+1} = \begin{bmatrix} size^{sdlg} \\ size_i \\ \vdots \\ size_N \end{bmatrix} \quad [3]$$

and \mathbf{A} is expressed as a $N+1 \times N+1$ matrix:

$$\mathbf{A} = \begin{bmatrix} 0 & F_i & \cdot & \cdot & F_N \\ T^{sdlg} & T_i & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ T^{sdlg} & \cdot & \cdot & \cdot & T_N \end{bmatrix} \quad [4]$$

in which T and F are size-transition (i.e. survival and growth) and reproduction kernels drawn from our vital rate estimates for each species and endophyte status.

Statistical analysis of vital rates. We modeled the effect of endophyte symbiosis on the mean and variance of vital rates by fitting generalized linear mixed models (GLMM) to the long-term data with year and plot random effects. We fit all vital rate models in a hierarchical Bayesian framework using Rstan (?), allowing us to propagate uncertainty from the vital rate estimates to our population model (40).

The probabilities of survival and flowering are recorded as successes or failures and consequently are modeled as Bernoulli processes. We modeled growth (measured as the number of tillers in year $t+1$), and the number of flowering tillers with the zero-truncated Poisson-Inverse Gaussian distribution, and the number of spikelets per inflorescence with the Negative Binomial distribution. Each of these size-dependent vital rates are modeled with the same structure of linear predictor (μ)

For example, growth ($G_{i,t1}$) of a given individual (i) in year $t+1$ is modeled as:

$$G_{i,t1} \sim P(IG(\mu_{s,e}, \lambda_{s,e})) \quad [5]$$

Similarly, survival $S_{i,t1}$ in year $t+1$ is modeled as:

$$S_{i,t1} \sim Bernoulli(\mu_{s,e}) \quad [6]$$

Where μ , for each species (s), is a linear function of the logarithm of plant size in year t (t), the plot level endophyte status (e), whether the plant was part of the initial transplanting or naturally recruited into the plot (r), along with random effects to account for plot(p), and year variation specific to each species and endophyte status. Thus μ can be written:

$$\begin{aligned} \mu_{s,e} &= \beta_s^1 + \beta_s^2 \log(size_t) + \beta_{s,e}^3 + \beta_r^4 \\ &\quad + \tau + \rho \end{aligned} \quad [7]$$

$$\begin{aligned} \tau &\sim N(0, \sigma_{s,e}) \\ \rho &\sim N(0, \sigma_p) \end{aligned}$$

For all species, we account for a reproductive delay by modeling seedling growth and survival separately from adult growth and survival. Seedlings are those plants that are recruited into the plot in a given year, and typically have only one tiller. So, for seedlings, growth ($G_{1,t1}^{sdlg}$) is modelled as:

$$G_{1,t1}^{sdlg} \sim P(IG(\mu_{s,e}^{sdlg}, \lambda_{s,e})) \quad [8]$$

Similarly, survival ($S_{i,t1}^{sdlg}$) in year $t+1$ is modeled as:

$$S_{1,t1}^{sdlg} \sim Bernoulli(\mu_{s,e}^{sdlg}) \quad [9]$$

Here, $\mu_{s,e}^{sdlg}$ is the linear function for these seedling specific vital rates. It does not include size-dependence or an effect to account for the initial benefits of greenhouse rearing.

$$\begin{aligned} \mu_{s,e}^{sdlg} &= \beta_s^1 + \beta_{s,e}^3 \\ &\quad + \tau + \rho \\ \tau &\sim N(0, \sigma_{s,e}) \\ \rho &\sim N(0, \sigma_p) \end{aligned} \quad [10]$$

The final element of fecundity comes from recruitment of plant into the plot. For this, we modeled the probability of germination as a proportion of seeds produced from the preceding year with a Binomial regression.

$$R_{1,t1} \sim Binomial(\mu_{s,e}^{sdlg}) \quad [11]$$

$$\begin{aligned} \mu_{s,e}^{rec} &= \beta_s^1 + \beta_{s,e}^3 \\ &\quad + \tau + \rho \\ \tau &\sim N(0, \sigma_{s,e}) \\ \rho &\sim N(0, \sigma_p) \end{aligned} \quad [12]$$

We ran each vital rate model for 2500 warm-up and 2500 MCMC sampling iterations with three chains using rStan. We assessed model convergence with trace plots of posterior chains and checked for \hat{R} values less than 1.01, indicating low within and between chain variation (41, 42). For those models that show poor convergence, we extended the MCMC sampling to include 5000 warm-up and 5000 sampling iterations, which was only necessary for seedling growth. For each of these vital rate models, we graphically check model fit with posterior predictive checks comparing simulated data from 500 posterior draws with the observed data (See supplement?). These checks provide evidence that our models are accurately recreating size-specific growth, survival, and reproduction patterns in our data across endophyte treatments.

We calculate the effect of endophytes on mean and variance in population growth rates by assembling matrix models with and without endophyte symbiosis following equation [2]. To do this, we calculated annual population growth rates for symbiotic and non-symbiotic populations for the 13 observed transition years by adjusting our vital rate estimates according to the year random effects, incorporating model uncertainty by sampling 500 posterior draws from our vital rate models. By using observed years, we maintain the structure of vital rate correlations as observed in the field. We then calculated the mean and variance of these annual growth rates.

Stochastic growth rate simulation experiment. We decomposed the contribution of endophyte symbiosis to long-term stochastic growth rates by calculating the geometric mean of 500 randomly sampled annual population growth rates, for simulated populations including mean-only, variance-only, and mean and variance endophyte effects compared to those without endophyte effects (13).can add better citations for this

$$\log(\lambda_s) E[\log(\Sigma(\mathbf{n}_{t+1})/\Sigma(\mathbf{n}_t))] \quad [13]$$

To explore the potential effects of future increased climate variability, we calculated the stochastic growth rates as above for simulations which sample the most extreme observed years more frequently. To do this, we randomly sampled the six annual population growth rates that were furthest from the mean growth rate across E+ and E- populations. Additionally, we tested the influence of temporal autocorrelation by sampling either alternating

- 461 **Estimating climate drivers of environmental context-dependence.** To
462 ask whether the variance buffering effects of symbiosis are driven by
463 environmental variation, and to forecast population dynamics under
464 future climate change scenarios, we built climate-explicit population
465 models. These models are built from vital rates as described above
466 with the inclusion of a parameter describing the effect, for each
467 endophyte status and species, of 12-month SPEI, a drought index
468 intended to incorporate precipitation and evapotranspiration (). We
469 calculated SPEI with weather station data from Bloomington, IN via
470 the 'rnoaa' and 'spei' R packages (). This weather station has the
471 most complete climate record over the study period and the historic
472 record when compared to other nearby weather stations and is
473 comparable to downscaled climate data (could include correlations)
474 (See Supplement?). Preliminary model selection using the climwin
475 package () with simpler vital rate models suggested that a 12-
476 month SPEI would reasonably capture the relevant climate signal
477 (see appendix?).
- 478 **Forecasting under alternative climate forcings.** We performed model
479 experiments to explore the effect of altered mean and variance in
480 climate as expected under climate change. First, we generated a
481 50 year climate time series by randomly drawing observed SPEI
482 values,
- 483 **ACKNOWLEDGMENTS.** Please include your acknowledgments
484 here, set in a single paragraph. Please do not include any acknowl-
485 edgments in the Supporting Information, or anywhere else in the
486 manuscript.
- 487 1. IPCC (Intergovernmental Panel on Climate Change). *Managing the risks of extreme events*
488 *and disasters to advance climate change adaptation: special report of the intergovernmental*
489 *panel on climate change. Special Report.* Cambridge University Press, 2012.
 - 490 2. Sonia Seneviratne, Neville Nicholls, David Easterling, Clare Goodess, Shinjiro Kanae, James
491 Kossin, Yali Luo, Jose Marengo, Kathleen McInnes, Mohammad Rahimi, et al. Changes in
492 climate extremes and their impacts on the natural physical environment. 2012.
 - 493 3. Thomas F Stocker, Dahe Qin, G-K Plattner, Lisa V Alexander, Simon K Allen, Nathaniel L
494 Bindoff, F-M Bréon, John A Church, Ulrich Cubasch, Seita Emori, et al. Technical summary.
495 In *Climate change 2013: the physical science basis. Contribution of Working Group I to the*
496 *Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, pages 33–115.
497 Cambridge University Press, 2013.
 - 498 4. R. C. Lewontin and D. Cohen. On Population Growth in a Randomly Varying Environment.
499 *Proceedings of the National Academy of Sciences*, 62(4):1056–1060, April 1969. ISSN
500 0027-8424, 1091-6490. . URL <https://www.pnas.org/content/62/4/1056>. Publisher: National
501 Academy of Sciences Section: Biological Sciences: Zoology.
 - 502 5. Shripad D. Tuljapurkar. Population dynamics in variable environments. III. Evolutionary dy-
503 namics of r-selection. *Theoretical Population Biology*, 21(1):141–165, February 1982. ISSN
504 0040-5809. . URL <http://www.sciencedirect.com/science/article/pii/0040580982900107>.
 - 505 6. Joel E Cohen. Comparative statics and stochastic dynamics of age-structured populations.
506 *Theoretical population biology*, 16(2):159–171, 1979.
 - 507 7. Shripad Tuljapurkar. *Population dynamics in variable environments*, volume 85. Springer
508 Science & Business Media, 2013.
 - 509 8. Catherine A Pfister. Patterns of variance in stage-structured populations: evolutionary predic-
510 tions and ecological implications. *Proceedings of the National Academy of Sciences*, 95(1):
511 213–218, 1998.
 - 512 9. William F Morris, Catherine A Pfister, Shripad Tuljapurkar, Chirakal V Haridas, Carol L Boggs,
513 Mark S Boyce, Emilio M Bruna, Don R Church, Tim Coulson, Daniel F Doak, et al. Longevity
514 can buffer plant and animal populations against changing climatic variability. *Ecology*, 89(1):
515 19–25, 2008.
 - 516 10. Eric S Menges. Population viability analysis for an endangered plant. *Conservation biology*, 4
517 (1):52–62, 1990.
 - 518 11. Aldo Compagnoni, Andrew J Bibian, Brad M Ochocki, Haldre S Rogers, Emily L Schultz,
519 Michelle E Sneek, Bret D Elder, Amy M Iler, David W Inouye, Hans Jacquemyn, et al. The
520 effect of demographic correlations on the stochastic population dynamics of perennial plants.
521 *Ecological Monographs*, 86(4):480–494, 2016.
 - 522 12. Martha M Ellis and Elizabeth E Crone. The role of transient dynamics in stochastic population
523 growth for nine perennial plants. *Ecology*, 94(8):1681–1686, 2013.
 - 524 13. SD Tuljapurkar and Steven Hecht Orzack. Population dynamics in variable environments i.
525 long-run growth rates and extinction. *Theoretical Population Biology*, 18(3):314–342, 1980.
 - 526 14. John Fieberg and Stephen P Ellner. Stochastic matrix models for conservation and manage-
527 ment: a comparative review of methods. *Ecology letters*, 4(3):244–266, 2001.
 - 528 15. Christoffer H. Hilde, Marlène Gamelon, Bernt-Erik Sæther, Jean-Michel Gaillard, Nigel G.
529 Yoccoz, and Christophe Pélabon. The Demographic Buffering Hypothesis: Evidence and
530 Challenges. *Trends in Ecology & Evolution*, 0(0), March 2020. ISSN 0169-5347. . URL
531 [https://www.cell.com/trends/ecology-evolution/abstract/S0169-5347\(20\)30050-1](https://www.cell.com/trends/ecology-evolution/abstract/S0169-5347(20)30050-1). Publisher:
532 Elsevier.
 - 533 16. RJ Rodriguez, JF White Jr, Anne E Arnold, and a RS and Redman. Fungal endophytes:
534 diversity and functional roles. *New phytologist*, 182(2):314–330, 2009.
 - 535 17. Margaret McFall-Ngai, Michael G Hadfield, Thomas CG Bosch, Hannah V Carey, Tomislav
536 Domazet-Lošo, Angela E Douglas, Nicole Dubilier, Gerard Eberl, Tadashi Fukami, Scott F
537 Gilbert, et al. Animals in a bacterial world, a new imperative for the life sciences. *Proceedings*
538 *of the National Academy of Sciences*, 110(9):3229–3236, 2013.
 - 539 18. Jennifer A Rudgers, Michelle E Afkhami, Lukas Bell-Dereske, Y Anny Chung, Kerri M Crawford,
540 Stephanie N Kivlin, Michael A Mann, and Martin A Nuñez. Climate disruption of plant-microbe
541 interactions. *Annual review of ecology, evolution, and systematics*, 51(1), 2020.
 - 542 19. Jacob A Russell and Nancy A Moran. Costs and benefits of symbiont infection in aphids:
543 variation among symbionts and across temperatures. *Proceedings of the Royal Society B:*
544 *Biological Sciences*, 273(1586):603–610, 2006.
 - 545 20. Jeremy C Brownlie and Karyn N Johnson. Symbiont-mediated protection in insect hosts.
546 *Trends in microbiology*, 17(8):348–354, 2009.
 - 547 21. Stephanie N Kivlin, Sarah M Emery, and Jennifer A Rudgers. Fungal symbionts alter plant
548 responses to global change. *American Journal of Botany*, 100(7):1445–1457, 2013.
 - 549 22. Chris Corbin, Eleanor Rose Heyworth, Julia Ferrari, and Gregory DD Hurst. Heritable sym-
550 bionts in a world of varying temperature. *Heredity*, 118(1):10–20, 2017.
 - 551 23. Kenneth D Hoadley, Allison M Lewis, Drew C Wham, D Tye Pettay, Chris Grasso, Robin Smith,
552 Dustin W Kemp, Todd C LaJeunesse, and Mark E Warner. Host–symbiont combinations
553 dictate the photo-physiological response of reef-building corals to thermal stress. *Scientific*
554 *Reports*, 9(1):1–15, 2019.
 - 555 24. Scott A Chamberlain, Judith L Bronstein, and Jennifer A Rudgers. How context dependent are
556 species interactions? *Ecology letters*, 17(7):881–890, 2014.
 - 557 25. Pedro Jordano. Spatial and temporal variation in the avian-frugivore assemblage of prunus
558 mahaleb: patterns and consequences. *Oikos*, pages 479–491, 1994.
 - 559 26. Ian Billick and Kirk Tonkel. The relative importance of spatial vs. temporal variability in
560 generating a conditional mutualism. *Ecology*, 84(2):289–295, 2003.
 - 561 27. Gregory P Cheplick, Stanley Faeth, and Stanley H Faeth. *Ecology and evolution of the*
562 *grass-endophyte symbiosis*. OUP USA, 2009.
 - 563 28. Gregory P Cheplick. Recovery from drought stress in lolium perenne (poaceae): are fungal
564 endophytes detrimental? *American Journal of Botany*, 91(12):1960–1968, 2004.
 - 565 29. S Kannadan and JA Rudgers. Endophyte symbiosis benefits a rare grass under low water
566 availability. *Functional Ecology*, 22(4):706–713, 2008.
 - 567 30. Facundo A Decunta, Luis I Pérez, Dariusz P Malinowski, Marco A Molina-Montenegro, and
568 Pedro E Gundel. A systematic review on the effects of epichloë fungal endophytes on drought
569 tolerance in cool-season grasses. *Frontiers in plant science*, 12:644731, 2021.
 - 570 31. Jennifer A Rudgers, Tom EX Miller, Shaun M Ziegler, and Kelly D Craven. There are many
571 ways to be a mutualist: endophytic fungus reduces plant survival but increases population
572 growth. *Ecology*, 93(3):565–574, 2012.
 - 573 32. Mark Rees and Stephen P Ellner. Integral projection models for populations in temporally
574 varying environments. *Ecological Monographs*, 79(4):575–594, 2009.
 - 575 33. Paul EM Fine. Vectors and vertical transmission: an epidemiologic perspective. *Annals of the*
576 *New York Academy of Sciences*, 266(1):173–194, 1975.
 - 577 34. AE Douglas. Host benefit and the evolution of specialization in symbiosis. *Heredity*, 81(6):
578 599–603, 1998.
 - 579 35. Jennifer A Rudgers, Michelle E Afkhami, Megan A Rúa, Andrew J Davitt, Samantha Hammer,
580 and Valérie M Huguet. A fungus among us: broad patterns of endophyte distribution in the
581 grasses. *Ecology*, 90(6):1531–1539, 2009.
 - 582 36. D Brem and A Leuchtman. Epichloë grass endophytes increase herbivore resistance in the
583 woodland grass brachypodium sylvaticum. *Oecologia*, 126(4):522–530, 2001.
 - 584 37. Cyd E Hamilton and Taryn L Bauerle. A new currency for mutualism? fungal endophytes alter
585 antioxidant activity in hosts responding to drought. *Fungal Diversity*, 54(1):39–49, 2012.
 - 586 38. Charles W Bacon and James F White. Stains, media, and procedures for analyzing endophytes.
587 In *Biotechnology of endophytic fungi of grasses*, pages 47–56. CRC Press, 2018.
 - 588 39. Jennifer A Rudgers and Angela L Swafford. Benefits of a fungal endophyte in elymus virginicus
589 decline under drought stress. *Basic and Applied Ecology*, 10(1):43–51, 2009.
 - 590 40. Bret D Elder and Tom EX Miller. Quantifying demographic uncertainty: Bayesian methods for
591 integral projection models. *Ecological Monographs*, 86(1):125–144, 2016.
 - 592 41. Stephen P Brooks and Andrew Gelman. General methods for monitoring convergence of
593 iterative simulations. *Journal of computational and graphical statistics*, 7(4):434–455, 1998.
 - 594 42. Andrew Gelman and Jennifer Hill. *Data analysis using regression and multilevel/hierarchical*
595 *models*. Cambridge university press, 2006.