

# Context-dependent host-microbe interactions in stochastic environments

Joshua C. Fowler<sup>a,1</sup>, Shaun Ziegler<sup>b</sup>, Kenneth D. Whitney<sup>b</sup>, Jennifer A. Rudgers<sup>b</sup>, and Tom E. X. Miller<sup>a</sup>

<sup>a</sup>Rice University, Department of BioSciences, Houston, TX, 77005; <sup>b</sup>University of New Mexico, Department of Biology, Albuquerque, NM, 87131

This manuscript was compiled on October 18, 2022

Microbial symbioses are ubiquitous in nature, yet our ability to predict how these interactions affect species' responses to climate change has been limited by interaction outcomes that vary with environmental context. Increased environmental variability is a key prediction of climate change. Here we present and test a novel hypothesis: microbial symbionts may buffer hosts from environmental variability by being beneficial in harsh years while being neutral or costly in good years. Long-term data reveal that, while weaker than effects on the mean of population growth, variance buffering is an important contribution to hosts long-term population growth. Additionally, the symbiosis becomes more mutualistic under simulated increases in variability, driven by an increased contribution from variance buffering.

Please provide an abstract of no more than 250 words in a single paragraph. Abstracts should explain to the general reader the major contributions of the article. References in the abstract must be cited in full within the abstract itself and cited in the text.

Keyword 1 | Keyword 2 | Keyword 3 | ...

Along with increases in average temperatures, global climate change is driving increases in the variability of precipitation events, temperature extremes, and droughts (1–3). Discerning the effects of environmental variability on population dynamics and species interactions are therefore key objectives of global change ecology.

\*Classic theory predicts that long-term population growth rates (equivalently, population mean fitness) will decline under increased environmental variability due to negative effects of bad years that outweigh positive effects of good years (a consequence of nonlinear averaging) (4, 5). For example, in unstructured populations, the long-term stochastic growth rate in fluctuating environments ( $\lambda_s$ ) will always be less than the growth rate averaged across environments ( $\bar{\lambda}$ ) by an amount proportional to the environmental variance ( $\sigma^2$ ):  $\log(\lambda_s) \approx \log(\bar{\lambda}) - \frac{\sigma^2}{2\bar{\lambda}^2}$ . Populations structured by size or stage are expected to similarly experience negative effects of variability (6, 7). Thus, there are two pathways to increase population viability in a stochastic environment  $\lambda_s$ : increasing the mean growth rate and/or dampening temporal fluctuation in growth rates, also called variance “buffering”.

Given the potential for negative fitness consequences of increasing environmental stochasticity under global change, there is growing interest in factors that can amplify or dampen demographic fluctuations. Much attention has focused on the properties of species or their environment as modifiers of fluctuations in fitness, including variation in vital rates (survival, growth and reproduction)(8), correlations between vital rates (9), transient shifts in stage structure (10), and the degree of environmental autocorrelation (11, 12). In contrast,

little is known about how biotic interactions influence patterns of demographic variability (13), due in part to the paucity of long-term interaction data that are required to detect such effects.

Our work focused on the potential for microbial symbionts to buffer host plants against negative effects of environmental variability. Microbial symbioses are ubiquitous across plant and animal host taxa and can be crucial determinants of host fitness (14, 15). Yet we know little about their potential influence on responses to climate change and environmental variation (16). Across a broad range of taxa, mutualistic host-associated microbes provide protection from environmental stresses including drought, extreme temperatures, and enemies (17–21).<sup>†</sup> The role they play may be under-appreciated, and it can be difficult to quantify the net outcome of a given interaction because they are often viewed as being context-dependent where the magnitude of benefit depends on environmental conditions (22). Rather than considering context-dependence as some unexplainable intricacy of species interactions, environmental variation opens up the possibility for interaction strength to vary through time (23, 24) and to influence the variance of population growth rates.<sup>‡</sup>

We hypothesize that symbionts may provide benefits under harsh conditions when they are needed by their hosts, but be neutral or even costly under benign conditions. Over time, this would lead symbiotic hosts to experience a reduction in variation in vital rates by reducing the frequency of extreme years. Incorporating context-dependence in this way, we highlight a novel mechanism by which symbionts can act as mutualists that may come to be of increasing importance in a more variable future.<sup>§</sup>

<sup>†</sup> This sentence seems to contradict the previous sentence.

<sup>‡</sup> I agree that this is the place to introduce context-dependence and connect climate variability to variability in interaction outcome. But I don't think what you have here works.

<sup>§</sup> Similar comment as above. I think this is the place to lay out the hypotheses of symbionts elevating mean and/or reducing variance. But I don't think this is sufficiently explanatory or provides sufficient context or rationale for the ideas to really hit home.

## Significance Statement

Authors must submit a 120-word maximum statement about the significance of their research paper written at a level understandable to an undergraduate educated scientist outside their field of speciality. The primary goal of the Significance Statement is to explain the relevance of the work in broad context to a broad readership. The Significance Statement appears in the paper itself and is required for all research papers.

Please provide details of author contributions here.

Please declare any conflict of interest here.

<sup>1</sup>To whom correspondence should be addressed. E-mail: jcf3rice.edu

\* It would be nice to start paragraph 2 here, but that probably leaves P1 too skimpy.

Using data from a unique, long-term symbiont-removal experiment, we test the hypothesis that context-dependent benefits of microbial symbionts buffer hosts from the fitness consequences of environmental variability<sup>¶</sup>. The experiment consists of annually censused plots initiated in 2007 with seven grass species that host Epichloë fungal endophytes. These specialized fungal endophytes are unique to grasses, occurring in at least 30%<sup>||</sup> of cool-season (C3) species, and are primarily vertically transmitted from maternal plant to seed (25). While they have been associated with contributing to drought tolerance for their hosts<sup>\*\*</sup>, these benefits are commonly context-dependent<sup>††</sup>(26–28). And so<sup>††</sup>, we ask first how fungal endophytes influence the mean and interannual variance of their hosts' vital rates; next, we ask if these vital rate effects buffer variance in fitness and, if so, what is the relative contribution of variance buffering vs. mean effects to the overall effect of the symbiosis on long-term growth rates. To answer these questions, we build structured, stochastic population models for these seven grass host species (*Agrostis perennans*, *Elymus villosus*, *Elymus virginicus*, *Festuca subverticillata*, *Lolium arundinaceum*, *Poa alsodes* and *Poa sylvestris*).<sup>§§</sup> These long-term plots contain either naturally symbiotic plants (E+) or those which have had their symbionts experimentally removed (E-).<sup>¶¶</sup> Each annual census is a sample of climatic variation. Across 14 years, the data contain 31,216 individual-transition years.<sup>\*\*\*</sup> After quantifying endophyte effects on mean and variance in population growth rates, we use simulations to explore the consequences of variance buffering under increased variance<sup>†††</sup> and construct climate-explicit population models<sup>†††</sup> to evaluate the role of climate drivers as explanations for this buffering.

Intriguingly, we find that the symbiosis contributes positively to long-term population growth rates through both mean and variance buffering effects. Integrating across diverse effects on vital rates, contributions to long-term growth rates from effects on the mean are 4.17 times greater on average than contributions from variance buffering. However, these effects varied between species; for example, two species (*A. perennans* and *P. sylvestris*) have contributions from variance buffering which are greater than mean effects. Additionally, the effect of mutualism increases under simulations with increased variance driven by greater contributions of variance buffering. In the most extreme scenario, we find that variance buffering contributions across species are 1.5 times greater than effects on the mean on average, and that variance buffering contributions are greater than mean effects for five out of seven species.<sup>§§§</sup>

<sup>¶</sup> In addition to this hypothesis, I think it is important to emphasize the question of relative importance of mean vs variance effects. Like Volker always says, yes/no hypotheses are often a little dull.

<sup>||</sup> I think the 30% number comes from a Leuchtmann paper.

<sup>\*\*</sup> Awkward phrasing.

<sup>††</sup> This is a little confusing because it sounds like you are saying that the drought benefit is context-dependent, but I think drought is the context, so the benefits vary with presence or absence of drought.

<sup>†††</sup> too colloquial

<sup>§§</sup> Mention the models somewhere in this paragraph but don't list the species.

<sup>¶¶</sup> Should come earlier when you first introduce the "unique" experiment.

<sup>\*\*\*</sup> Put in results.

<sup>†††</sup> Should be emphasized earlier. It's cool and novel.

<sup>†††</sup> I would not mention this here since it is not central to the story of the paper.

<sup>§§§</sup> I would eliminate this paragraph entirely. Not needed, since you launch directly into results from here.

## Results

### Endophyte effects on the mean and variance of vital rates.

Vital rate models reveal that Epichloë fungal endophyte symbiosis has positive effects on vital rate means, particularly for host survival and growth across species. At the same time, endophytes consistently buffer variance in the majority of host vital rates (Fig. 1). The magnitude of these effects vary across species and across vital rates. For example, endophytes increase mean adult survival and buffer variance slightly for *P. alsodes*, while for *F. subverticillata*, effects from variance buffering are stronger with a relatively weaker mean effect. In other vital rates, such as in seedling growth, *P. alsodes* experiences stronger buffering than *F. subverticillata*. Variance buffering effects are notably present across species' reproductive vital rates (e.g. likelihood of flowering and panicle production); This might align with predictions from life-history theory suggesting that reproductive vital rates are likely to experience greater year-to-year variance (). Interestingly, there are also certain vital rates that indicate potential costs of endophyte symbiosis, such as *A. perennans* and *F. subverticillata* which have lower average germination rates when partnered with endophytes.<sup>¶¶¶</sup>

### Endophyte effects on the mean and variance of population growth rates.

To assess the total impact of endophyte symbiosis on host population growth rates, we integrate the diverse effects of endophytes on the mean and variance of host vital rates into structured demographic models (29). We find that, on average across species, populations with endophytes experience a mean population growth rate that is 9.2 % higher (with a 92.8% posterior probability that effects are greater than zero) and a standard deviation that is 6.6 % smaller than non-symbiotic populations (with a 66% posterior probability that effects are less than zero). (Fig. 2). For some species, the standard deviation is reduced on average by as much as 44.3 % for *L. arundinaceum* and 28.5 % for *F. subverticillata*, while for others, variance buffering effects are much smaller, or even slightly positive for *E. villosus* and *E. virginicus*.

### Contribution of mean and variance effects to long-term growth rates.

We decomposed the overall effect of the symbiosis into contributions from mean and variance buffering effects on long-term population growth rates (30)<sup>17</sup>. We found that variance buffering provided a small benefit overall (XXX%) (Fig. 3) but note that even small changes in fitness can have profound effects over long time periods.

**Role of symbiotic buffering under increased variance.** In simulations with increased variance, we find that the contribution of variance buffering increases.

### Endophytes are buffering their hosts from environmental variability.

Symbiotic and non-symbiotic populations responded distinctly to environmental drivers. Our climate-explicit analysis shows that on average, the population growth of non-endophytic plants was more sensitive to our drought index, SPEI, than endophytic populations (FIGURE). This pattern

<sup>¶¶¶</sup> Not sure how much to talk about demographic compensation (i.e. do we see less variance in vital rates that are more important). We don't really assess this. And at least looking at the actual sigma values for each species, it seems like none of the vital rates have hugely different baseline standard deviations.

<sup>17</sup> I don't know if there's a better citation for this, this is more about IPM's




Fig. 1. needs to be updated pdf version, but putting this here for now. Effect of endophytes on vital rates across species




Fig. 2. needs to be updated pdf version, but putting this here for now. Effect of endophytes on mean and CV of population growth




Fig. 3. needs to be updated pdf version, but putting this here for now. Stochastic lambda contributions with simulations of increased variance

was particularly pronounced for species which showed strong buffering in our earlier analysis (SPECIES), suggesting that endophytes are buffering their hosts from this aspect of inter-annual variation, although a large amount of variation remained unexplained.

## Discussion

### Guide to using this template on Overleaf

Please note that whilst this template provides a preview of the typeset manuscript for submission, to help in this preparation, it will not necessarily be the final publication layout. For more detailed information please see the [PNAS Information for Authors](#).

If you have a question while using this template on Overleaf, please use the help menu (“?”) on the top bar to search for [help and tutorials](#). You can also [contact the Overleaf support team](#) at any time with specific questions about your manuscript or feedback on the template.

**Author Affiliations.** Include department, institution, and complete address, with the ZIP/postal code, for each author. Use lower case letters to match authors with institutions, as shown in the example. Authors with an ORCID ID may supply this information at submission.

**Submitting Manuscripts.** All authors must submit their articles at [PNAScentral](#). If you are using Overleaf to write your article, you can use the “Submit to PNAS” option in the top bar of the editor window.

**Format.** Many authors find it useful to organize their manuscripts with the following order of sections; Title, Author Affiliation, Keywords, Abstract, Significance Statement, Results, Discussion, Materials and methods, Acknowledgments, and References. Other orders and headings are permitted.

**Manuscript Length.** PNAS generally uses a two-column format averaging 67 characters, including spaces, per line. The maximum length of a Direct Submission research article is six pages and a Direct Submission Plus research article is ten pages including all text, spaces, and the number of characters displaced by figures, tables, and equations. When submitting tables, figures, and/or equations in addition to text, keep the text for your manuscript under 39,000 characters (including spaces) for Direct Submissions and 72,000 characters (including spaces) for Direct Submission Plus.

**References.** References should be cited in numerical order as they appear in text; this will be done automatically via bibtex, e.g. . All references should be included in the main manuscript file.

**Data Archival.** PNAS must be able to archive the data essential to a published article. Where such archiving is not possible, deposition of data in public databases, such as GenBank, ArrayExpress, Protein Data Bank, Unidata, and others outlined in the Information for Authors, is acceptable.

**Language-Editing Services.** Prior to submission, authors who believe their manuscripts would benefit from professional editing are encouraged to use a language-editing service (see list at

**Table 1. Comparison of the fitted potential energy surfaces and ab initio benchmark electronic energy calculations**

Species	CBS	CV	G3
1. Acetaldehyde	0.0	0.0	0.0
2. Vinyl alcohol	9.1	9.6	13.5
3. Hydroxyethylidene	50.8	51.2	54.0

nomenclature for the TSs refers to the numbered species in the table.

[www.pnas.org/site/authors/language-editing.xhtml](http://www.pnas.org/site/authors/language-editing.xhtml)). PNAS does not take responsibility for or endorse these services, and their use has no bearing on acceptance of a manuscript for publication.

**Digital Figures.** Only TIFF, EPS, and high-resolution PDF for Mac or PC are allowed for figures that will appear in the main text, and images must be final size. Authors may submit U3D or PRC files for 3D images; these must be accompanied by 2D representations in TIFF, EPS, or high-resolution PDF format. Color images must be in RGB (red, green, blue) mode. Include the font files for any text.

Figures and Tables should be labelled and referenced in the standard way using the `\label{}` and `\ref{}` commands.

Figure ?? shows an example of how to insert a column-wide figure. To insert a figure wider than one column, please use the `\begin{figure*}...\end{figure*}` environment. Figures wider than one column should be sized to 11.4 cm or 17.8 cm wide. Use `\begin{SCfigure*}...\end{SCfigure*}` for a wide figure with side captions.

**Tables.** In addition to including your tables within this manuscript file, PNAS requires that each table be uploaded to the submission separately as a Table file. Please ensure that each table .tex file contains a preamble, the `\begin{document}` command, and the `\end{document}` command. This is necessary so that the submission system can convert each file to PDF.

**Single column equations.** Authors may use 1- or 2-column equations in their article, according to their preference.

To allow an equation to span both columns, use the `\begin{figure*}...\end{figure*}` environment mentioned above for figures.

Note that the use of the `widetext` environment for equations is not recommended, and should not be used.

**Supporting Information (SI).** Authors should submit SI as a single separate PDF file, combining all text, figures, tables, movie legends, and SI references. PNAS will publish SI uncomposed, as the authors have provided it. Additional details can be found here: [policy on SI](#). For SI formatting instructions click [here](#). The PNAS Overleaf SI template can be found [here](#). Refer to the SI Appendix in the manuscript at an appropriate point in the text. Number supporting figures and tables starting with S1, S2, etc.

Authors who place detailed materials and methods in an SI Appendix must provide sufficient detail in the main text methods to enable a reader to follow the logic of the procedures and results and also must reference the SI methods. If a paper is fundamentally a study of a new method or technique, then the methods must be described completely in the main text.



$$\begin{aligned}
(x+y)^3 &= (x+y)(x+y)^2 \\
&= (x+y)(x^2+2xy+y^2) \\
&= x^3+3x^2y+3xy^2+y^3.
\end{aligned}
\tag{1}$$

**SI Datasets.** Supply Excel (.xls), RTF, or PDF files. This file type will be published in raw format and will not be edited or composed.

**SI Movies.** Supply Audio Video Interleave (avi), Quicktime (mov), Windows Media (wmv), animated GIF (gif), or MPEG files and submit a brief legend for each movie in a Word or RTF file. All movies should be submitted at the desired reproduction size and length. Movies should be no more than 10 MB in size.

**3D Figures.** Supply a composable U3D or PRC file so that it may be edited and composed. Authors may submit a PDF file but please note it will be published in raw format and will not be edited or composed.

## Materials and Methods

**Natural history of grass-endophyte symbiosis.** To study the effects of context-dependent microbial symbiosis, we focused on *Epichloë* fungal endophytes, which live in the aboveground tissue of many species of cool-season grasses and grow into their hosts' seeds where they can be transmitted vertically from mother to offspring plants. This vertical transmission couples host and symbiont fitness and leads to the expectation that the interaction be mutualistic, else the fungi cause their host to be selected out of the population (31–33). While there are demonstrated benefits against herbivory (34) and under drought stress (35) for some host species, these interactions outcomes are commonly context-dependent (26, 27).

**Plant propagation and endophyte removal.** Seeds from naturally infected populations of seven species of cool-season grasses (*Agrostis perennans*, *Elymus villosus*, *Elymus virginicus*, *Festuca subverticillata*, *Lolium arundinaceum*, *Poa alsodes*, and *Poa sylvestris*) were collected during the 2006 growing season from Lilly Dickie Woods (39.238533, -86.218150) and the Bayles Road Teaching and Research Preserve (39.220167, -86.542683) in Brown Co. IN. To reduce confounding genotype effects, seeds with shared maternal ancestry were disinfected with heat treatments (6d in a drying oven at 60° C for *E. villosus*, *E. virginicus*, *F. subverticillata*, and *L. arundinaceum*; 7d in a drying oven at 60° C for *P. alsodes*, and *P. sylvestris*; and 12 min. in a hot water bath at 60° C for *A. perennans*)<sup>18</sup> or left naturally infected. Seeds were surface sterilized with bleach and cold stratified for ??? weeks, then germinated in a growth chamber before being transferred to the greenhouse at Indiana University and allowed to grow for XXXX weeks. We confirmed the endophyte status of these plants using leaf peels, where tissue from the leaf sheath is stained with aniline blue dye and examined for the presence of fungal hyphae (36). Then, we established the experimental plots with vegetatively propagated clones of similar sizes from the plants<sup>19</sup> to reduce the potential for negative side effects of heat treatments (37).

**Experimental design and data collection.** During the spring of 2007, we established 10 3x3 plots for *A. perennans*, *E. villosus*, *E. virginicus*, *F. subverticillata*, and *L. arundinaceum* and 18 plots for *P. alsodes* and *P. sylvestris*. For each species, an equal number of plots were randomly assigned to each endophyte status, E+ or E-. Each

plot was planted with 20 evenly spaced symbiotic or symbiont-free individuals respectively and each plant marked with aluminum tags.

Each summer starting in 2007, we censused all individuals in each plot for survival, growth and reproduction, garnering a dataset covering 14 years that contains 31,216 individual transition years. After clearing out leaf litter, for each plant alive in the previous year, we marked its survival and measured its size as a count of the number of tillers. Further, we collected reproductive data by counting the number of reproductive tillers, and then counting the number of seed-bearing spikelets on up to three of those reproductive tillers. In 2009, we took additional counts of seeds per inflorescence. Together, we use these measurements to estimate seed production. In each plot, we also survey for and tag any unmarked individuals. New recruits typically have one tiller and are non-reproductive, but we also find and tag any individuals that may have been missed in previous censuses.

We typically expect plots of each endophyte status to maintain their status because the fungi are almost entirely vertically transmitted and plots are spaced at least 5 m apart, limiting the possibility for unwanted dispersal between plots or horizontal transmission of the fungus. Seeds from reproductive individuals are opportunistically taken and scored for their endophyte status. Overall, these scores reflect a 97.5% faithfulness of recruits to their expected endophyte status across species and plots (Supplement data).

**Demographic modeling.** Armed with this demographic data, we next constructed size-structured, stochastic population models. These models describe transitions between sizes (measured as a count of tillers) from one year to the next. For all species, we include a 1 year reproductive delay in the population model following the observation that these newly recruited plants are rarely observed flowering in their first year. Our population model can be expressed as:

$$\mathbf{n}_{t+1} = \mathbf{A}\mathbf{n}_t \tag{2}$$

where  $\mathbf{n}_{t+1}$  is a vector of abundances across sizes in year  $t+1$  for each species and endophyte status.

$$\mathbf{n}_{t+1} = \begin{bmatrix} size^{sdlg} \\ size_i \\ \vdots \\ size_N \end{bmatrix} \tag{3}$$

and  $\mathbf{A}$  is expressed as a  $N+1 \times N+1$  matrix:

$$\mathbf{A} = \begin{bmatrix} 0 & F_i & \dots & F_N \\ T^{sdlg} & T_i & \dots & \vdots \\ \vdots & \vdots & \ddots & \vdots \\ T^{sdlg} & \vdots & \dots & T_N \end{bmatrix} \tag{4}$$

in which  $T$  and  $F$  are size-transition (i.e. survival and growth) and reproduction kernels drawn from our vital rate estimates for each species and endophyte status.

**Statistical analysis of vital rates.** We modeled the effect of endophyte symbiosis on the mean and variance of vital rates by fitting generalized linear mixed models (GLMM) to the long-term data with year and plot random effects. We fit all vital rate models in a hierarchical Bayesian framework using Rstan (?), allowing us to propagate uncertainty from the vital rate estimates to our population model (38).

The probabilities of survival and flowering are recorded as successes or failures and consequently are modeled as Bernoulli processes. We modeled growth (measured as the number of tillers

<sup>18</sup> need to double check methods for temp, duration, etc.

<sup>19</sup> not sure this happened

in year  $t+1$ ), and the number of flowering tillers with the zero-truncated Poisson-Inverse Gaussian distribution, and the number of spikelets per inflorescence with the Negative Binomial distribution. Each of these size-dependent vital rates are modeled with the same structure of linear predictor ( $\mu$ )

For example, growth ( $G_{i,t1}$ ) of a given individual (i) in year  $t+1$  is modeled as:

$$G_{i,t1} \sim P(IG(\mu_{s,e}, \lambda_{s,e})) \quad [5]$$

Similarly, survival  $S_{i,t1}$  in year  $t+1$  is modeled as:

$$S_{i,t1} \sim \text{Bernoulli}(\mu_{s,e}) \quad [6]$$

Where  $\mu$ , for each species (s), is a linear function of the logarithm of plant size in year  $t$  ( $t$ ), the plot level endophyte status (e), whether the plant was part of the initial transplanting or naturally recruited into the plot (r), along with random effects to account for plot(p), and year variation specific to each species and endophyte status. Thus  $\mu$  can be written:

$$\begin{aligned} \mu_{s,e} &= \beta_s^1 + \beta_s^2 \log(\text{size}_t) + \beta_s^3 + \beta_r^4 \\ &\quad + \tau + \rho \\ \tau &\sim N(0, \sigma_{s,e}) \\ \rho &\sim N(0, \sigma_p) \end{aligned} \quad [7]$$

For all species, we account for a reproductive delay by modeling seedling growth and survival separately from adult growth and survival. Seedlings are those plants that are recruited into the plot in a given year, and typically have only one tiller. So, for seedlings, growth ( $G_{1,t1}^{sdlg}$ ) is modelled as:

$$G_{1,t1}^{sdlg} \sim P(IG(\mu_{s,e}^{sdlg}, \lambda_{s,e})) \quad [8]$$

Similarly, survival ( $S_{i,t1}^{sdlg}$ ) in year  $t+1$  is modeled as:

$$S_{1,t1}^{sdlg} \sim \text{Bernoulli}(\mu_{s,e}^{sdlg}) \quad [9]$$

Here,  $\mu_{s,e}^{sdlg}$  is the linear function for these seedling specific vital rates. It does not include size-dependence or an effect to account for the initial benefits of greenhouse rearing.

$$\begin{aligned} \mu_{s,e}^{sdlg} &= \beta_s^1 + \beta_{s,e}^3 \\ &\quad + \tau + \rho \\ \tau &\sim N(0, \sigma_{s,e}) \\ \rho &\sim N(0, \sigma_p) \end{aligned} \quad [10]$$

The final element of fecundity comes from recruitment of plant into the plot. For this, we modeled the probability of germination as a proportion of seeds produced from the preceding year with a Binomial regression.

$$R_{1,t1} \sim \text{Binomial}(\mu_{s,e}^{sdlg}) \quad [11]$$

$$\begin{aligned} \mu_{s,e}^{rec} &= \beta_s^1 + \beta_{s,e}^3 \\ &\quad + \tau + \rho \\ \tau &\sim N(0, \sigma_{s,e}) \\ \rho &\sim N(0, \sigma_p) \end{aligned} \quad [12]$$

We ran each vital rate model for 2500 warm-up and 2500 MCMC sampling iterations with three chains using rStan. We assessed model convergence with trace plots of posterior chains and checked for  $\hat{R}$  values less than 1.01, indicating low within and between chain variation (39, 40). For those models that show poor convergence, we extended the MCMC sampling to include 5000 warm-up and 5000 sampling iterations, which was only necessary for seedling growth. For each of these vital rate models, we graphically check model fit with posterior predictive checks comparing simulated data from 500 posterior draws with the observed data (See supplement?). These checks provide evidence that our models are accurately recreating size-specific growth, survival, and reproduction patterns in our data across endophyte treatments.

We calculate the effect of endophytes on mean and variance in population growth rates by assembling matrix models with and without endophyte symbiosis following equation [2]. To do this, we calculated annual population growth rates for symbiotic and

non-symbiotic populations for the 13 observed transition years by adjusting our vital rate estimates according to the year random effects, incorporating model uncertainty by sampling 500 posterior draws from our vital rate models. By using observed years, we maintain the structure of vital rate correlations as observed in the field. We then calculated the mean and variance of these annual growth rates.

**Stochastic growth rate simulation experiment.** We decomposed the contribution of endophyte symbiosis to long-term stochastic growth rates by calculating the geometric mean of 500 randomly sampled annual population growth rates, for simulated populations including mean-only, variance-only, and mean and variance endophyte effects compared to those without endophyte effects (11).can add better citations for this

$$\log(\lambda_s) \ E[\log(\Sigma(\mathbf{n}_{t+1})/\Sigma(\mathbf{n}_t))] \quad [13]$$

To explore the potential effects of future increased climate variability, we calculated the stochastic growth rates as above for simulations which sample the most extreme observed years more frequently. To do this, we randomly sampled the six annual population growth rates that were furthest from the mean growth rate across E+ and E- populations. Additionally, we tested the influence of temporal autocorrelation by sampling either alternating

**Estimating climate drivers of environmental context-dependence.** To ask whether the variance buffering effects of symbiosis are driven by environmental variation, and to forecast population dynamics under future climate change scenarios, we built climate-explicit population models. These models are built from vital rates as described above with the inclusion of a parameter describing the effect, for each endophyte status and species, of 12-month SPEI, a drought index intended to incorporate precipitation and evapotranspiration (). We calculated SPEI with weather station data from Bloomington, IN via the 'rnoaa' and 'spei' R packages (). This weather station has the most complete climate record over the study period and the historic record when compared to other nearby weather stations and is comparable to downscaled climate data (could include correlations) (See Supplement?). Preliminary model selection using the climwin package () with simpler vital rate models suggested that a 12-month SPEI would reasonably capture the relevant climate signal (see appendix?).

**Forecasting under alternative climate forcings.** We performed model experiments to explore the effect of altered mean and variance in climate as expected under climate change. First, we generated a 50 year climate time series by randomly drawing observed SPEI values,

**ACKNOWLEDGMENTS.** Please include your acknowledgments here, set in a single paragraph. Please do not include any acknowledgments in the Supporting Information, or anywhere else in the manuscript.

1. IPCC (Intergovernmental Panel on Climate Change). *Managing the risks of extreme events and disasters to advance climate change adaptation: special report of the intergovernmental panel on climate change. Special Report.* Cambridge University Press, 2012.
2. Sonia Seneviratne, Neville Nicholls, David Easterling, Clare Goodess, Shinjiro Kanae, James Kossin, Yali Luo, Jose Marengo, Kathleen McInnes, Mohammad Rahimi, et al. Changes in climate extremes and their impacts on the natural physical environment. 2012.
3. Thomas F Stocker, Dahe Qin, G-K Plattner, Lisa V Alexander, Simon K Allen, Nathaniel L Bindoff, F-M Br  n, John A Church, Ulrich Cubasch, Seita Emori, et al. Technical summary. In *Climate change 2013: the physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, pages 33–115. Cambridge University Press, 2013.
4. R. C. Lewontin and D. Cohen. On Population Growth in a Randomly Varying Environment. *Proceedings of the National Academy of Sciences*, 62(4):1056–1060, April 1969. ISSN 0027-8424, 1091-6490. . URL <https://www.pnas.org/content/62/4/1056>. Publisher: National Academy of Sciences Section: Biological Sciences: Zoology.
5. Shripad D. Tuljapurkar. Population dynamics in variable environments. III. Evolutionary dynamics of r-selection. *Theoretical Population Biology*, 21(1):141–165, February 1982. ISSN 0040-5809. . URL <http://www.sciencedirect.com/science/article/pii/0040580982900107>.
6. Joel E Cohen. Comparative statics and stochastic dynamics of age-structured populations. *Theoretical population biology*, 16(2):159–171, 1979.
7. Shripad Tuljapurkar. *Population dynamics in variable environments*, volume 85. Springer Science & Business Media, 2013.

8. William F Morris, Catherine A Pfister, Shripad Tuljapurkar, Chirrakal V Haridas, Carol L Boggs, Mark S Boyce, Emilio M Bruna, Don R Church, Tim Coulson, Daniel F Doak, et al. Longevity can buffer plant and animal populations against changing climatic variability. *Ecology*, 89(1): 19–25, 2008.
9. Aldo Compagnoni, Andrew J Bibian, Brad M Ochocki, Haldre S Rogers, Emily L Schultz, Michelle E Sneek, Bret D Elder, Amy M Iler, David W Inouye, Hans Jacquemyn, et al. The effect of demographic correlations on the stochastic population dynamics of perennial plants. *Ecological Monographs*, 86(4):480–494, 2016.
10. Martha M Ellis and Elizabeth E Crone. The role of transient dynamics in stochastic population growth for nine perennial plants. *Ecology*, 94(8):1681–1686, 2013.
11. SD Tuljapurkar and Steven Hecht Orzack. Population dynamics in variable environments i. long-run growth rates and extinction. *Theoretical Population Biology*, 18(3):314–342, 1980.
12. John Fieberg and Stephen P Ellner. Stochastic matrix models for conservation and management: a comparative review of methods. *Ecology letters*, 4(3):244–266, 2001.
13. Christoffer H. Hilde, Marlène Gamelon, Bernt-Erik Sæther, Jean-Michel Gaillard, Nigel G. Yoccoz, and Christophe Pélabon. The Demographic Buffering Hypothesis: Evidence and Challenges. *Trends in Ecology & Evolution*, 0(0), March 2020. ISSN 0169-5347. . URL [https://www.cell.com/trends/ecology-evolution/abstract/S0169-5347\(20\)30050-1](https://www.cell.com/trends/ecology-evolution/abstract/S0169-5347(20)30050-1). Publisher: Elsevier.
14. RJ Rodriguez, JF White Jr, Anne E Arnold, and a RS and Redman. Fungal endophytes: diversity and functional roles. *New phytologist*, 182(2):314–330, 2009.
15. Margaret McFall-Ngai, Michael G Hadfield, Thomas CG Bosch, Hannah V Carey, Tomislav Domazet-Lošo, Angela E Douglas, Nicole Dubilier, Gerard Eberl, Tadashi Fukami, Scott F Gilbert, et al. Animals in a bacterial world, a new imperative for the life sciences. *Proceedings of the National Academy of Sciences*, 110(9):3229–3236, 2013.
16. Jennifer A Rudgers, Michelle E Afkhami, Lukas Bell-Dereske, Y Anny Chung, Kerri M Crawford, Stephanie N Kivlin, Michael A Mann, and Martin A Nuñez. Climate disruption of plant-microbe interactions. *Annual review of ecology, evolution, and systematics*, 51(1), 2020.
17. Jacob A Russell and Nancy A Moran. Costs and benefits of symbiont infection in aphids: variation among symbionts and across temperatures. *Proceedings of the Royal Society B: Biological Sciences*, 273(1586):603–610, 2006.
18. Jeremy C Brownlie and Karyn N Johnson. Symbiont-mediated protection in insect hosts. *Trends in microbiology*, 17(8):348–354, 2009.
19. Stephanie N Kivlin, Sarah M Emery, and Jennifer A Rudgers. Fungal symbionts alter plant responses to global change. *American Journal of Botany*, 100(7):1445–1457, 2013.
20. Chris Corbin, Eleanor Rose Heyworth, Julia Ferrari, and Gregory DD Hurst. Heritable symbionts in a world of varying temperature. *Heredity*, 118(1):10–20, 2017.
21. Kenneth D Hoadley, Allison M Lewis, Drew C Wham, D Tye Pettay, Chris Grasso, Robin Smith, Dustin W Kemp, Todd C LaJeunesse, and Mark E Warner. Host–symbiont combinations dictate the photo-physiological response of reef-building corals to thermal stress. *Scientific Reports*, 9(1):1–15, 2019.
22. Scott A Chamberlain, Judith L Bronstein, and Jennifer A Rudgers. How context dependent are species interactions? *Ecology letters*, 17(7):881–890, 2014.
23. Pedro Jordano. Spatial and temporal variation in the avian-frugivore assemblage of prunus mahaleb: patterns and consequences. *Oikos*, pages 479–491, 1994.
24. Ian Billick and Kirk Tonkel. The relative importance of spatial vs. temporal variability in generating a conditional mutualism. *Ecology*, 84(2):289–295, 2003.
25. Gregory P Cheplick, Stanley Faeth, and Stanley H Faeth. *Ecology and evolution of the grass-endophyte symbiosis*. OUP USA, 2009.
26. Gregory P Cheplick. Recovery from drought stress in *lilium perenne* (poaceae): are fungal endophytes detrimental? *American Journal of Botany*, 91(12):1960–1968, 2004.
27. S Kannadan and JA Rudgers. Endophyte symbiosis benefits a rare grass under low water availability. *Functional Ecology*, 22(4):706–713, 2008.
28. Facundo A Decunta, Luis I Pérez, Dariusz P Malinowski, Marco A Molina-Montenegro, and Pedro E Gundel. A systematic review on the effects of epichloë fungal endophytes on drought tolerance in cool-season grasses. *Frontiers in plant science*, 12:644731, 2021.
29. Jennifer A Rudgers, Tom EX Miller, Shaun M Ziegler, and Kelly D Craven. There are many ways to be a mutualist: endophytic fungus reduces plant survival but increases population growth. *Ecology*, 93(3):565–574, 2012.
30. Mark Rees and Stephen P Ellner. Integral projection models for populations in temporally varying environments. *Ecological Monographs*, 79(4):575–594, 2009.
31. Paul EM Fine. Vectors and vertical transmission: an epidemiologic perspective. *Annals of the New York Academy of Sciences*, 266(1):173–194, 1975.
32. AE Douglas. Host benefit and the evolution of specialization in symbiosis. *Heredity*, 81(6): 599–603, 1998.
33. Jennifer A Rudgers, Michelle E Afkhami, Megan A Rúa, Andrew J Davitt, Samantha Hammer, and Valérie M Hugué. A fungus among us: broad patterns of endophyte distribution in the grasses. *Ecology*, 90(6):1531–1539, 2009.
34. D Brem and A Leuchtmann. Epichloë grass endophytes increase herbivore resistance in the woodland grass *brachypodium sylvaticum*. *Oecologia*, 126(4):522–530, 2001.
35. Cyd E Hamilton and Taryn L Bauerle. A new currency for mutualism? fungal endophytes alter antioxidant activity in hosts responding to drought. *Fungal Diversity*, 54(1):39–49, 2012.
36. Charles W Bacon and James F White. Stains, media, and procedures for analyzing endophytes. In *Biotechnology of endophytic fungi of grasses*, pages 47–56. CRC Press, 2018.
37. Jennifer A Rudgers and Angela L Swafford. Benefits of a fungal endophyte in *elymus virginicus* decline under drought stress. *Basic and Applied Ecology*, 10(1):43–51, 2009.
38. Bret D Elder and Tom EX Miller. Quantifying demographic uncertainty: Bayesian methods for integral projection models. *Ecological Monographs*, 86(1):125–144, 2016.
39. Stephen P Brooks and Andrew Gelman. General methods for monitoring convergence of iterative simulations. *Journal of computational and graphical statistics*, 7(4):434–455, 1998.
40. Andrew Gelman and Jennifer Hill. *Data analysis using regression and multilevel/hierarchical models*. Cambridge university press, 2006.