

Context-dependent host-microbe interactions in stochastic environments

Joshua C. Fowler^{a,1}, Shaun Ziegler^b, Kenneth D. Whitney^b, Jennifer A. Rudgers^b, and Tom E. X. Miller^a

^aRice University, Department of BioSciences, Houston, TX, 77005; ^bUniversity of New Mexico, Department of Biology, Albuquerque, NM, 87131

This manuscript was compiled on April 25, 2022

Please provide an abstract of no more than 250 words in a single paragraph. Abstracts should explain to the general reader the major contributions of the article. References in the abstract must be cited in full within the abstract itself and cited in the text.

Keyword 1 | Keyword 2 | Keyword 3 | ...

Along with increases in average temperatures, global climate change is driving increases in the variability of precipitation events, temperature extremes, and droughts (1–3). Thus discerning the effects of variability on population dynamics and interactions between species is pivotal to forecasting the future of ecological systems. Classic ecological theory predicts that long-term population growth rates will be reduced by environmental variability (4, 5). This stochastic variability means that populations can expect to experience good years and bad years. The long-term stochastic growth rate (λ_s), which is the long-run geometric mean of annual growth rates, captures this variability; the geometric mean will always be less than expected from the mean growth rate alone.

Following Lewontin and Cohen (4), λ_s can be approximated as:

$$\log(\lambda_s) \approx \log(\bar{\lambda}) - \frac{\sigma^2}{2\bar{\lambda}^2} \quad [1]$$

Where $\bar{\lambda}$ is the mean of annual population growth rates (λ_t) and σ^2 is the variance (4). Populations will increase over time if λ_s is greater than 1, and can be expected to decrease if λ_s is less than 1. Here, there are two pathways to increase λ_s : (1) increasing the mean growth rate, and/or (2) reducing the variance in growth rates. That both mean and variance can determine fitness underlies understanding of which aspects of a species' life history influence its success (6) and has important implications for population viability analysis (7). Anything that limits the negative effects of bad years, while being neutral or costly in good years has the potential to decrease the impact of interannual environmental variability on population dynamics because it would limit variance. * Whether species interactions contribute to variance buffering† is an underexplored question (8).

Host-associated microbes are ubiquitous in nature. Across a broad range of taxa‡, microbial symbionts provide their hosts with protection from environmental stresses including drought, temperature§, and enemies (9). Commonly, the benefits from these symbioses are context-dependent where the magnitude of interaction benefit depends on environmental conditions (10).

* Not sure this is helpful. Obviously, "Anything" is mutualism, but if you don't know that yet then "Anything" will be confusing. But I do think you need some bridge to symbiosis.

† This should be defined.

‡ You only cite one paper on plant-fungal interactions – so not very convincing as a broad range of taxa

§ Temperature per se is not a stress.

This can make it difficult to quantify the net effect of a given interaction, but it also allows for the possibility that interaction strength can vary through time (cite). Symbionts may provide benefits under harsh conditions when they are needed by their hosts, but be neutral or even costly under benign conditions (cite). Over time, this may lead symbiont-associated organisms to experience a reduction in variation in vital rates by reducing the frequency of extreme years (conceptual figure). Embracing context-dependence in this way, we reveal a novel mechanism by which symbionts can act as mutualists that may come to be of increasing importance in a more variable future.

Using long-term data from experimental grass-fungal endophyte plots¶, we test the hypothesis that context-dependent benefits of microbial symbionts buffer hosts from the fitness consequences of environmental variability. Specifically, we ask first how fungal endophytes influence the mean and interannual variance of their hosts' vital rates; next, we ask if these vital rate effects buffer variance in fitness and, if so, what is the relative importance of variance buffering vs. mean effects in the overall fitness impact of the symbiosis. With 14 years of demographic data, we employ structured, stochastic population models for seven species of cool-season grass hosts that are commonly infected with fungal endophytes|| (*Lolium arundinaceum*, *Festuca subverticillata*, *Elymus virginicus*, and *Elymus villosus*, *Poa alsodes* and *Poa sylvestris*). These long-term data, in which each annual census is a sample of weather variation, allow us to construct a climate-explicit population models, which we use to evaluate the importance of buffering under forecasted changes in the mean and variance of climate drivers.

This paragraph is mostly talking off my head about results, but my idea is to include a brief statement of our results.**

¶ This needs context. No one will know what this means.

|| I think we need a more thorough description of the experiment here - it's a novel experiment, at least in temporal scale, so we will want to sell it.

** Agree we will want a punchy summary that leaves readers wanting to continue into the Results.

Significance Statement

Authors must submit a 120-word maximum statement about the significance of their research paper written at a level understandable to an undergraduate educated scientist outside their field of speciality. The primary goal of the Significance Statement is to explain the relevance of the work in broad context to a broad readership. The Significance Statement appears in the paper itself and is required for all research papers.

Please provide details of author contributions here.

Please declare any conflict of interest here.

¹To whom correspondence should be addressed. E-mail: jcf3rice.edu

67 Across species, we find that variance buffering by endophytes
68 contributes (percentage) to population growth rates. While
69 the effect is generally weaker than effects on the mean, we
70 found that buffering was common in the most sensitive vital
71 rates, and was most important for xxx species with xxx life
72 history.

73 Results

74 Discussion

75 Guide to using this template on Overleaf

76 Please note that whilst this template provides a preview of the
77 typeset manuscript for submission, to help in this preparation,
78 it will not necessarily be the final publication layout. For
79 more detailed information please see the [PNAS Information](#)
80 [for Authors](#).

81 If you have a question while using this template on Overleaf,
82 please use the help menu (“?”) on the top bar to search for [help](#)
83 [and tutorials](#). You can also [contact the Overleaf support team](#)
84 at any time with specific questions about your manuscript or
85 feedback on the template.

86 **Author Affiliations.** Include department, institution, and com-
87 plete address, with the ZIP/postal code, for each author. Use
88 lower case letters to match authors with institutions, as shown
89 in the example. Authors with an ORCID ID may supply this
90 information at submission.

91 **Submitting Manuscripts.** All authors must submit their arti-
92 cles at [PNAScentral](#). If you are using Overleaf to write your
93 article, you can use the “Submit to PNAS” option in the top
94 bar of the editor window.

95 **Format.** Many authors find it useful to organize their
96 manuscripts with the following order of sections; Title, Author
97 Affiliation, Keywords, Abstract, Significance Statement, Re-
98 sults, Discussion, Materials and methods, Acknowledgments,
99 and References. Other orders and headings are permitted.

100 **Manuscript Length.** PNAS generally uses a two-column for-
101 mat averaging 67 characters, including spaces, per line. The
102 maximum length of a Direct Submission research article is
103 six pages and a Direct Submission Plus research article is ten
104 pages including all text, spaces, and the number of characters
105 displaced by figures, tables, and equations. When submitting
106 tables, figures, and/or equations in addition to text, keep the
107 text for your manuscript under 39,000 characters (including
108 spaces) for Direct Submissions and 72,000 characters (including
109 spaces) for Direct Submission Plus.

110 **References.** References should be cited in numerical order as
111 they appear in text; this will be done automatically via bibtex,
112 e.g. . All references should be included in the main manuscript
113 file.

114 **Data Archival.** PNAS must be able to archive the data es-
115 sential to a published article. Where such archiving is not
116 possible, deposition of data in public databases, such as Gen-
117 Bank, ArrayExpress, Protein Data Bank, Unidata, and others
118 outlined in the Information for Authors, is acceptable.

Table 1. Comparison of the fitted potential energy surfaces and ab initio benchmark electronic energy calculations

Species	CBS	CV	G3
1. Acetaldehyde	0.0	0.0	0.0
2. Vinyl alcohol	9.1	9.6	13.5
3. Hydroxyethylidene	50.8	51.2	54.0

nomenclature for the TSs refers to the numbered species in the table.

Language-Editing Services. Prior to submission, authors who believe their manuscripts would benefit from professional editing are encouraged to use a language-editing service (see list at [www.pnas.org/site/authors/language-editing.xhtml](#)). PNAS does not take responsibility for or endorse these services, and their use has no bearing on acceptance of a manuscript for publication.

Digital Figures. Only TIFF, EPS, and high-resolution PDF for Mac or PC are allowed for figures that will appear in the main text, and images must be final size. Authors may submit U3D or PRC files for 3D images; these must be accompanied by 2D representations in TIFF, EPS, or high-resolution PDF format. Color images must be in RGB (red, green, blue) mode. Include the font files for any text.

Figures and Tables should be labelled and referenced in the standard way using the `\label{}` and `\ref{}` commands.

Figure ?? shows an example of how to insert a column-wide figure. To insert a figure wider than one column, please use the `\begin{figure*}...\end{figure*}` environment. Figures wider than one column should be sized to 11.4 cm or 17.8 cm wide. Use `\begin{SCfigure*}...\end{SCfigure*}` for a wide figure with side captions.

Tables. In addition to including your tables within this manuscript file, PNAS requires that each table be uploaded to the submission separately as a Table file. Please ensure that each table .tex file contains a preamble, the `\begin{document}` command, and the `\end{document}` command. This is necessary so that the submission system can convert each file to PDF.

Single column equations. Authors may use 1- or 2-column equations in their article, according to their preference.

To allow an equation to span both columns, use the `\begin{figure*}...\end{figure*}` environment mentioned above for figures.

Note that the use of the `widetext` environment for equations is not recommended, and should not be used.

Supporting Information (SI). Authors should submit SI as a single separate PDF file, combining all text, figures, tables, movie legends, and SI references. PNAS will publish SI uncomposed, as the authors have provided it. Additional details can be found here: [policy on SI](#). For SI formatting instructions click [here](#). The PNAS Overleaf SI template can be found [here](#). Refer to the SI Appendix in the manuscript at an appropriate point in the text. Number supporting figures and tables starting with S1, S2, etc.

Authors who place detailed materials and methods in an SI Appendix must provide sufficient detail in the main text methods to enable a reader to follow the logic of the procedures

$$\begin{aligned}
(x+y)^3 &= (x+y)(x+y)^2 \\
&= (x+y)(x^2 + 2xy + y^2) \\
&= x^3 + 3x^2y + 3xy^2 + y^3.
\end{aligned}
\tag{2}$$

and results and also must reference the SI methods. If a paper is fundamentally a study of a new method or technique, then the methods must be described completely in the main text.

SI Datasets. Supply Excel (.xls), RTF, or PDF files. This file type will be published in raw format and will not be edited or composed.

SI Movies. Supply Audio Video Interleave (avi), Quicktime (mov), Windows Media (wmv), animated GIF (gif), or MPEG files and submit a brief legend for each movie in a Word or RTF file. All movies should be submitted at the desired reproduction size and length. Movies should be no more than 10 MB in size.

3D Figures. Supply a composable U3D or PRC file so that it may be edited and composed. Authors may submit a PDF file but please note it will be published in raw format and will not be edited or composed.

Materials and Methods

Natural history of grass-endophyte symbiosis. ^{††}

Plant propagation and endophyte removal. Seeds from naturally infected populations of seven species of cool-season grasses (*Agrostis perennans*, *Elymus villosus*, *Elymus virginicus*, *Festuca subverticillata*, *Lolium arundinaceum*, *Poa alsodes*, and *Poa sylvestris*^{††}) were collected in the Spring of 2006^{§§§} from Lilly Dickie Woods (Lat,Lon) and Bayles Road (Lat,Lon) in Brown Co. IN. Seeds with shared maternal ancestry were either experimentally disinfected by heat treatments^{¶¶} or left naturally infected to reduce confounding genotype effects. Seeds were surface sterilized with XXXX and cold stratified for XXXX weeks, then germinated in a growth chamber for XXXX weeks. Seedlings were then transferred to the greenhouse at Indiana University and allowed to grow for XXXX weeks. We confirmed the endophyte status of these plants using leaf peels^{***} (cite), and vegetatively propagated clones of similar sizes from the plants^{†††}. These clones were used to establish the experimental plots, and cloning reduces the potential for negative effects of heat treatments (cite).

Experimental design and data collection. In 2007^{†††}, we established 10 3x3 plots for *Lolium arundinaceum*, *Festuca subverticillata*, *Elymus virginicus*, and *Elymus villosus* and 18 plots for *Poa alsodes* and *Poa sylvestris*. For each species, an equal number of plots were randomly assigned to each endophyte status, E+ or E-, and was planted with only symbiotic or symbiont-free plants respectively. Each plot was planted with 25^{§§§} evenly spaced individuals and each plant marked with aluminum tags.

In each Summer starting in 2007, we censused all original transplants and any recruits for survival, growth and reproduction. After

clearing out leaf litter, for each plant alive in the previous year, we marked its survival. We measured the size of each plant as a count of the number of tillers. Further, we collected reproductive data by counting the number of reproductive tillers, and then counting the number of seed-bearing spikelets on up to three of those reproductive tillers. In XXXX year, we took additional counts of seeds per inflorescence (list of species) or seeds per spikelet (list of species). Together, we use these measurements to estimate seed production. In each plot, we also survey for and mark any unmarked individuals. New recruits are typically a size of one tiller and non-reproductive, but we also find and mark any individuals who may have been missed in previous censuses.

We typically expect plots of each endophyte status to maintain their status as the fungus is almost entirely vertically transmitted, and plots are spaced at least XX m apart, limiting the possibility for unwanted dispersal between plots or horizontal transmission of the fungus. Seeds from reproductive individuals are opportunistically taken and scored for their endophyte status. These scores reflect a XXXX% faithfulness of recruits to their expected endophyte status (Supplement data)^{¶¶¶}

In sum, this individual-level demographic dataset covers 14 years and contains 30,XXXX individual transition-years.¹⁷

Demographic modeling. Armed with this demographic data, we next constructed size-structured, stochastic population models. This demographic model describes transitions between sizes (measured as a count of tillers) from one year to the next.

$$\mathbf{n}_{t+1} = \mathbf{A}\mathbf{n}_t \tag{3}$$

Model description and estimation. We modeled the mean and variance effects on each vital rate by fitting generalized linear mixed models (GLMM) to the long-term data with year and plot random effects. We fit all vital rate models in a Bayesian framework using Rstan, allowing us to propagate uncertainty from the vital rate estimates to our population model (11).

For all species, the population model includes a 1 year reproductive delay. We account for this by modeling seedling growth and survival separately from adult growth and survival. Seedlings are those plants that are recruited into the plot in a given year, and typically have only one tiller.

The probability of survival and flowering are recorded as successes or failures and consequently are modeled as Bernoulli processes. We modeled growth (measured as the number of tillers in year t+1), the number of flowering tillers, and the number of spikelets per inflorescence with the zero-truncated Poisson-Inverse Gaussian distribution. Where possible, each of these size dependent vital rates are modeled with the same structure of linear predictor (μ)

For example, growth of a given individual in year t+1 is modeled as:

$$\begin{aligned}
G_{i,t+1} &\sim P(IG(\mu_{s,e}, \lambda_{s,e})) \\
\mu_{s,e} &= \beta_s^1 + \beta_s^2 \log(\text{size}_t) + \beta_{s,e}^3 + \beta_r^4 \\
&\quad + \tau + \rho \\
\tau &\sim N(0, \sigma_{s,e,t}) \\
\rho &\sim N(0, \sigma_p)
\end{aligned}
\tag{4}$$

^{¶¶¶} We had these data in the original LTREB proposal.

¹⁷ Move this earlier and also mention at the end of intro. The total number of plant-years is impressive.

^{††} I think we need a brief section like this.

^{†††} italicize species names.

^{§§} check with Jenn

^{¶¶} need methods for temp, duration, etc.

^{***} need to be described

^{†††} not sure this happened

^{†††} dates?

^{§§§} I think 20, check data.

Similarly, survival in year $t+1$ is modeled as:

$$\begin{aligned} S_{i,t+1} &\sim \text{Bernoulli}(\mu_{s,e}) \\ \mu_{s,e} &= \beta_s^1 + \beta_s^2 \log(\text{size}_t) + \beta_{s,e}^3 + \beta_r^4 \\ &\quad + \tau + \rho \\ \tau &\sim N(0, \sigma_{s,e}) \\ \rho &\sim N(0, \sigma_p) \end{aligned} \quad [5]$$

Where μ , for each species (s), is a linear function of the logarithm of plant size in year t , the plot level endophyte status (e), the status as an initial experimental transplant plant (r), along with random effects to account for plot (p), and year random effects specific to each species and endophyte status. Thus μ can be written:

$$\begin{aligned} \mu_{s,e} &= \beta_s^1 + \beta_s^2 \log(\text{size}_t) + \beta_{s,e}^3 + \beta_r^4 \\ &\quad + \tau + \rho \\ \tau &\sim N(0, \sigma_{s,e}) \\ \rho &\sim N(0, \sigma_p) \end{aligned} \quad [6]$$

For survival

Model assessment.

Life table response experiment.

Estimating climate drivers of environmental context-dependence.

Climate data.

Climate-explicit Model description and estimation.

Climate-explicit Model assessment.

Forecasting under alternative climate forcings. We used statistics

ACKNOWLEDGMENTS. Please include your acknowledgments here, set in a single paragraph. Please do not include any acknowledgments in the Supporting Information, or anywhere else in the manuscript.

1. IPCC (Intergovernmental Panel on Climate Change). *Managing the risks of extreme events and disasters to advance climate change adaptation: special report of the intergovernmental panel on climate change. Special Report.* Cambridge University Press, 2012.
2. Sonia Seneviratne, Neville Nicholls, David Easterling, Clare Goodess, Shinjiro Kanae, James Kossin, Yali Luo, Jose Marengo, Kathleen McInnes, Mohammad Rahimi, et al. Changes in climate extremes and their impacts on the natural physical environment. 2012.
3. Thomas F Stocker, Dahe Qin, G-K Plattner, Lisa V Alexander, Simon K Allen, Nathaniel L Bindoff, F-M Bréon, John A Church, Ulrich Cubasch, Seita Emori, et al. Technical summary. In *Climate change 2013: the physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, pages 33–115. Cambridge University Press, 2013.
4. R. C. Lewontin and D. Cohen. On Population Growth in a Randomly Varying Environment. *Proceedings of the National Academy of Sciences*, 62(4):1056–1060, April 1969. ISSN 0027-8424, 1091-6490. . URL <https://www.pnas.org/content/62/4/1056>. Publisher: National Academy of Sciences Section: Biological Sciences: Zoology.
5. Shripad D. Tuljapurkar. Population dynamics in variable environments. III. Evolutionary dynamics of r -selection. *Theoretical Population Biology*, 21(1):141–165, February 1982. ISSN 0040-5809. . URL <http://www.sciencedirect.com/science/article/pii/0040580982900107>.
6. Catherine A Pfister. Patterns of variance in stage-structured populations: evolutionary predictions and ecological implications. *Proceedings of the National Academy of Sciences*, 95(1):213–218, 1998.
7. Eric S Menges. Population viability analysis for an endangered plant. *Conservation biology*, 4(1):52–62, 1990.
8. Christoffer H. Hilde, Marlène Gamelon, Bernt-Erik Sæther, Jean-Michel Gaillard, Nigel G. Yoccoz, and Christophe Pélabon. The Demographic Buffering Hypothesis: Evidence and Challenges. *Trends in Ecology & Evolution*, 0(0), March 2020. ISSN 0169-5347. . URL [https://www.cell.com/trends/ecology-evolution/abstract/S0169-5347\(20\)30050-1](https://www.cell.com/trends/ecology-evolution/abstract/S0169-5347(20)30050-1). Publisher: Elsevier.
9. Stephanie N Kivlin, Sarah M Emery, and Jennifer A Rudgers. Fungal symbionts alter plant responses to global change. *American Journal of Botany*, 100(7):1445–1457, 2013.
10. Scott A Chamberlain, Judith L Bronstein, and Jennifer A Rudgers. How context dependent are species interactions? *Ecology letters*, 17(7):881–890, 2014.
11. Bret D Elder and Tom EX Miller. Quantifying demographic uncertainty: Bayesian methods for integral projection models. *Ecological Monographs*, 86(1):125–144, 2016.