

Effect of probiotics on the nutritional status of severe stroke patients with nasal feeding that receive enteral nutrition

A protocol for systematic review and meta-analysis of randomized controlled trials

Xiaomin Liu, MS^a, Yasu Zhang, PhD^{a,*}, Jiahao Chu, BMS^b, Jie Zheng, PhD^a, Xue Cheng, PhD^a, Xinmin Li, PhD^c, Junzi Long, MS^a

Abstract

Background: Malnutrition is commonly observed after stroke and is closely associated with poor clinical outcomes. So, early nutrition support is particularly crucial for severe stroke patients. However, a significant number of critically ill patients are intolerant to enteral nutrition (EN). Probiotics have been widely used in malnutrition by various diseases and have a low incidence of enteral intolerance. So, we aim to elucidate the efficacy of probiotics in EN in improving the nutritional status and clinical prognosis of severe stroke patients with nasal feeding.

Method: Embase, PubMed, Sinomed, Web of Science, Cochrane Library, China National Knowledge Infrastructure, Wanfang database, and Vip Journal Integration Platform were searched from inception to March 31, 2021. Randomized controlled trials that applied probiotics in patients with severe stroke were included. The data were extracted and the risk of bias was assessed independently by 2 evaluators.

Results: Twenty-four studies comprising 2003 participants of randomized controlled trials were included. The result of pooled analyses showed that probiotics in EN were associated with better outcomes than EN alone on Glasgow Coma Scale score (mean difference [MD] = 1.03, 95% confidence intervals [CI]: 0.78–1.27; $P < .00001$), infection events (odds ratio [OR] = 0.25, 95% CI: 0.15–0.43; $P < .00001$), rate of intestinal flora dysbiosis (OR = 0.24, 95% CI: 0.12–0.48; $P < .0001$), gastrointestinal complications (OR = 0.25, 95% CI: 0.16–0.37, $P < .00001$), time to reach target nutrition (MD = −1.80, 95% CI: −2.42 to 1.18, $P < .00001$), prealbumin content (MD = 25.83, 95% CI: 13.68–37.99, $P < .0001$).

Conclusion: Our results demonstrated that probiotics supplementation might be an effective intervention for improving the clinical prognosis in severe stroke patients with nasal feeding, but no significant effect on increasing muscle circumference.

Abbreviations: CI = confidence interval, MD = mean difference, OR = odds ratio.

Keywords: enteral nutrition, nutrition statue, probiotics, severe stroke

1. Introduction

Malnutrition is commonly observed after stroke and is strongly related to poor clinical outcomes.^[1] It is mostly attributed to dysphagia, disorders of consciousness, mobility impairments,

and gastrointestinal dysfunction, among which dysphagia is the main risk factor for malnutrition in stroke patients.^[2] The prevalence of malnutrition after stroke ranges from 6.1% to 62%.^[3,4] The evidence showed that early nutritional status after

Editor: Manal Kamel Youssef.

Supplemental Digital Content is available for this article.

The authors report no conflicts of interest.

Funding Sources: The study supported by National Natural Science Foundation of China (ID: 81704135), Foundation of He'nan Educational Committee (ID: 18A360012), and the key scientific and technological project of Henan Province (ID: 192102310424).

The datasets generated during and/or analyzed during the current study are publicly available.

^a School of Rehabilitation Medicine, ^b First School of Clinical Medicine, ^c School of Basic Medical Science, Henan University of Chinese Medicine, Zhengzhou, China.

* Correspondence: Yasu Zhang, School of Rehabilitation Medicine, Henan University of Chinese Medicine, Zhengzhou 450046, China (e-mail: zhangyasu16@163.com).

Copyright © 2021 the Author(s). Published by Wolters Kluwer Health, Inc.

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial License 4.0 (CCBY-NC), where it is permissible to download, share, remix, transform, and buildup the work provided it is properly cited. The work cannot be used commercially without permission from the journal.

How to cite this article: Liu X, Zhang Y, Chu J, Zheng J, Cheng X, Li X, Long J. Effect of probiotics on the nutritional status of severe stroke patients with nasal feeding that receive enteral nutrition: a protocol for systematic review and meta-analysis of randomized controlled trials. *Medicine* 2021;100:17(e25657).

Received: 18 January 2021 / Received in final form: 30 March 2021 / Accepted: 3 April 2021

<http://dx.doi.org/10.1097/MD.00000000000025657>

However, the effectiveness of probiotics in severe stroke patients with nasal feeding has not been systematically evaluated. So the present study is necessary to fill this gap. The aim of this review is to investigate the effect of the probiotics on nutrition status and clinical efficacy in severe stroke patients with nasal feeding.

This study was undertaken according the PRISMA guidelines, and had registered this review on PROSPERO: CRD42020173643. <https://www.crd.york.ac.uk/prospero/>.

We searched Embase (from 1974 to March 31, 2021), PubMed (before March 31, 2021), Sinomed (before March 31, 2021), Web of Science (from 1950 to March 31, 2021), Cochrane Library (before March 31, 2021), China National Knowledge Infrastructure (from 1999 to March 31, 2021), VIP Journal Integration Platform (from 1989 to March 31, 2021), and Wanfang database (before March 31, 2021) for randomized controlled trials that assessed the effect of probiotics in EN on severe stroke patients published. The search was conducted using the following terms: (stroke OR apoplexy OR cerebral hemorrhage OR cerebrovascular accident OR cerebral infarction OR brain infarction OR cerebral haemorrhage OR cerebral apoplexy OR cerebrovascular disease OR brain vascular accident) AND (probiotics OR prebiotics OR symbiotic OR lactobacillus OR synbiotics OR lactobacterium OR bifidobacterium OR lactobacilli OR lactic acid bacteria).

2.2.1. Inclusion criteria.

1. The study design was randomized controlled trials.
2. All subjects were under a clinical diagnosis of severe stroke or Glasgow Coma Scale score ≤ 9 .
3. Treatment course ≥ 14 days.

1. Patients with heart, liver, kidney, and other severe organ failure, or malignancy, metabolic diseases.

- The study selection process was shown in Figure 1. A total of 1225 potentially eligible articles were found through databases. Among them, 1162 publications were excluded for duplication or inconsistent with inclusion criteria by reading the title and abstract. Sixty-three full-texts of articles were reviewed, and eventually 24 articles were included in the final selection.




Figure 1. Flowchart of study selection process.

3.2. Study characteristics

The main characteristics of the included studies were summarized in Table 1. The 24 studies were published between 2013 and 2021. A total of 2003 participants were included in this review, of which 57% were males. A total of 1007 patients were allocated to probiotics supplement with EN. The sample size of each study ranged from 56 to 124, and the participants mostly were 60 to 80 years' old. The treatment course ranged from 14 to 60 days.

3.3. Quality assessment

Among these 24 studies, 23 studies^[13–22,24–35] described detailed random grouping methods. The rest of the researches^[23] did not mention specific grouping methods. In incomplete outcome data, selective reporting, and other bias domain, all studies were considered to have a low risk of bias. Also, the included studies in our meta-analysis had unclear risk in terms of the allocation concealment and blinding for outcome assessment. The quality of these articles was presented in Supplementary Table 2, <http://links.lww.com/MD/G40>.

4. Meta-analysis results

4.1. Glasgow Coma Scale score

Eight trials^[16,21,24,27,30–32,35] reported the effect of probiotics on Glasgow Coma Scale score, and the results revealed that

probiotics in EN were associated with a significant improvement on Glasgow Coma Scale score (MD=1.03, 95% CI: 0.78–1.27; $P < .00001$) without significant heterogeneity ($I^2 = 46\%$) (Fig. 2).

4.2. Infection rate

Six trials^[13,17,19,22,24,25] showed the effect of probiotics on infection rate; the result suggested that probiotics in EN were associated with lower infection events (OR=0.25, 95% CI: 0.15–0.43; $P < .00001$). There was no significant heterogeneity ($I^2 = 40\%$) (Fig. 3)

4.3. Rate of intestinal flora dysbiosis

Four trials^[14,19,22,35] reported the effect of probiotics on intestinal flora dysbiosis. The pooled result showed a lower incidence of intestinal flora dysbiosis in probiotics group (OR=0.24, 95% CI: 0.12–0.48; $P < .0001$) without significant heterogeneity ($I^2 = 0\%$) (Fig. 4).

4.4. Gastrointestinal complication

Twelve trials^[14,15,18,20,22,23–26,28,31,33] reported the effect of probiotics on gastrointestinal complications, and the pooled analysis revealed that the probiotics in EN decreased the gastrointestinal complications (OR=0.25, 95% CI: 0.16–0.37; $P < .00001$) without significant heterogeneity ($I^2 = 0\%$) (Fig. 5).

Table 1**Trials characteristics.**

First author (year)	Sample size (M/F)		Age, y ($\bar{X} \pm S$)		Duration of illness	Intervention		Duration of treatment	Result
	Treatment	Control	Treatment	Control		Treatment	Control		
Wang et al (2021) ^[13]	51 (28/23)	51 (26/25)	68.23 \pm 8.72	67.62 \pm 8.21	\leq 48 h	Probiotics + EN + basic treatment	Basic treatment + EN	14 days	F G
Dong (2020) ^[14]	36 (—/—)	36 (—/—)	61.1 \pm 1.9	61.1 \pm 1.9	—	Probiotics + EN + basic treatment	Basic treatment + EN	14 days	B C
Liang et al (2020) ^[15]	30 (—/—)	30 (—/—)	60.19 \pm 18.65	62.13 \pm 13.97	—	Probiotics + EN + basic treatment	Basic treatment + EN	60 days	C
Ma (2020) ^[16]	47 (25/22)	46 (26/20)	52 \pm 6	52 \pm 6	\leq 72 h	Probiotics + EN + basic treatment	Basic treatment + EN	20 days	A F
Wang (2020) ^[17]	30 (20/10)	30 (18/12)	66.18 \pm 5.42	65.07 \pm 2.46	\leq 24 h	Probiotics + EN + basic treatment	Basic treatment + EN	16 days	G
Xie (2020) ^[18]	30 (15/15)	30 (16/14)	61.9 \pm 4.5	61.8 \pm 4.7	\leq 72 h	Probiotics + EN + basic treatment	Basic treatment + EN	14 days	C F
Chen (2019) ^[19]	35 (20/15)	35 (19/16)	75.47 \pm 4.59	75.23 \pm 4.52	—	Probiotics + EN + basic treatment	Basic treatment + EN	14 days	B D G
Chen (2019) ^[20]	34 (18/16)	34 (19/15)	72.06 \pm 6.43	72.06 \pm 6.43	—	Probiotics + EN + basic treatment	Basic treatment + EN	14 days	C
Jin et al (2019) ^[21]	28 (13/15)	28 (17/11)	62.18 \pm 11.12	62.07 \pm 10.94	\leq 48 h	Probiotics + EN + basic treatment	Basic treatment + EN	14 days	A D F
Li et al (2019) ^[22]	43 (24/19)	43 (27/16)	60.9 \pm 8.6	61.66 \pm 10.64	—	Probiotics + EN + basic treatment	Basic treatment + EN	14 days	B C G
Li and Li (2019) ^[23]	40 (24/16)	40 (25/15)	70.21 \pm 0.62	69.74 \pm 0.44	\leq 72 h	Probiotics + EN + basic treatment	Basic treatment + EN	14 days	C E
Dong (2018) ^[24]	41 (24/17)	41 (23/18)	62.4 \pm 4.1	63.4 \pm 4.3	—	Probiotics + EN + basic treatment	Basic treatment + EN	28 days	A C G
Gao (2018) ^[25]	40 (21/19)	40 (15/25)	58.2 \pm 2.1	51.1 \pm 2.3	—	Probiotics + EN + basic treatment	Basic treatment + EN	14 days	C G
Sun et al (2018) ^[26]	50 (26/24)	50 (27/23)	71.52 \pm 7.08	72.17 \pm 7.22	—	Probiotics + EN + basic treatment	Basic treatment + EN	14 days	C E
Wu (2018) ^[27]	31 (18/13)	32 (16/16)	58.48 \pm 8.09	58.59 \pm 9.97	—	Probiotics + EN + basic treatment	Basic treatment + EN	21 days	A D F
Zhou (2018) ^[28]	45 (28/17)	45 (27/18)	70.17 \pm 5.54	69.78 \pm 4.97	—	Probiotics + EN + basic treatment	Basic treatment + EN	14 days	C
Zhang et al (2017) ^[29]	44 (24/20)	44 (23/21)	—	—	\leq 6 h	Probiotics + EN + basic treatment	Basic treatment + EN	14 days	F
Chen and Chen (2016) ^[30]	45 (25/20)	45 (23/22)	69.9 \pm 7.2	70.3 \pm 6.7	\leq 3 days	Probiotics + EN + basic treatment	Basic treatment + EN	14 days	A
Hou (2016) ^[31]	38 (20/18)	38 (22/16)	72.15 \pm 7.56	71.89 \pm 7.42	\leq 3 days	Probiotics + EN + basic treatment	Basic treatment + EN	14 days	A C E
Hou et al (2016) ^[32]	38 (20/18)	38 (22/16)	72.2 \pm 7.6	71.9 \pm 7.4	\leq 3 days	Probiotics + EN + basic treatment	Basic treatment + EN	14 days	A
Wu et al (2016) ^[33]	62 (33/29)	62 (34/28)	54.22 \pm 4.29	54.13 \pm 4.56	—	Probiotics + EN + basic treatment	Basic treatment + EN	14 days	C F
Bai (2014) ^[34]	61 (44/17)	59 (47/12)	64.55 \pm 10.45	64.37 \pm 10.63	\leq 72 h	Probiotics + EN + basic treatment	Basic treatment + EN	14 days	F
Wu et al (2014) ^[35]	47 (25/22)	40 (21/19)	72.5 \pm 8.4	72.2 \pm 9.1	\leq 72 h	Probiotics + EN + basic treatment	Basic treatment + EN	21 days	A B
Bai et al (2013) ^[36]	61 (—/—)	59 (—/—)	—	—	\leq 72 h	Probiotics + EN + basic treatment	Basic treatment + EN	28 days	D

X = mean value, A = Glasgow Coma Scale score, B = rate of intestinal flora dysbiosis, C = gastrointestinal complication, D = time to reach target nutrition. E: mid arm muscle circumference, EN = enteral nutrition, EN = enteral nutrition, F = female, F = prealbumin content, G = infection rate, M = male, S = standard deviation.

4.5. Time to reach target nutrition

Four trials^[19,21,27,36] reported the effect of probiotics on time to reach target nutrition. The pooled result showed that compared to EN alone, probiotics in EN were associated with shorter time to reach target nutrition (MD = -1.80, 95% CI: -2.42 to 1.18 $P < .00001$). There was significant heterogeneity ($I^2 = 95\%$) (Fig. 6), but the main outcomes were not affected by excluding any particular study, so the random-effect model was selected.

4.6. Mid arm muscle circumference

Three trials^[23,26,31] reported the effect of probiotics on mid arm muscle circumference, and result revealed that probiotics in EN

exerted no positive effect on mid arm muscle circumference (MD = 1.76, 95% CI: 0.37–3.89; $P = .1$). There was significant heterogeneity ($I^2 = 91\%$) (Fig. 7).

4.7. Prealbumin content

Eight trials^[13,16,18,21,27,29,33,34] reported the effect of probiotics on prealbumin content, and the pooled result showed that probiotics were associated with improvement on prealbumin content (MD = 25.83, 95% CI: 13.68–37.99, $P < .0001$). There was significant heterogeneity ($I^2 = 93\%$) (Fig. 8), but the main outcomes were not affected by excluding any particular study, so the random-effect model was selected.




Figure 2. Forest plots of effect of probiotics on the Glasgow Coma Scale.




Figure 3. Forest plots of effect of probiotics on the infection rate.




Figure 4. Forest plots of effect of probiotics on the rate of intestinal flora dysbiosis.




Figure 5. Forest plots of effect of probiotics on the gastrointestinal complications.




Figure 6. Forest plots of effect of probiotics on the time to reach target nutrition.




Figure 7. Forest plots of effect of probiotics on the mid arm muscle circumference.




Figure 8. Forest plots of effect of probiotics on the prealbumin content.

4.8. Publication bias

The funnel plot was used to evaluate the publication bias of the 12 randomized controlled trials. Figure 9 showed the publication bias of the whole study was small.

5. Discussion

Our analysis suggested that probiotics had a positive impact on balancing the intestinal flora. The application of probiotics ameliorated gastrointestinal function, nutritional status, and immunity, and what's more improved Glasgow Coma Scale score, which indicated that probiotics were beneficial in improving the state of consciousness.

The results showed that the time to reach target nutrition was decreased, and prealbumin content was improved in the group of probiotics in EN. This meant that compared to EN alone,




Figure 9. Funnel plot of publication bias.

probiotics in EN could improve nutrition status. The studies included in this meta-analysis suggested that probiotics in EN reduced the incidence of gut microbiome dysbiosis, protected intestinal mucosal integrity, and alleviated gastrointestinal complications.^[14,15,18–20,22–26,28,31,33,35] The mechanism of improving the nutritional status of probiotics was that probiotics could stabilize intestinal homeostasis, accelerate gastrointestinal peristalsis, and inhibit disuse atrophy of intestinal mucosal.^[37,38] With the restoration of gastrointestinal function, probiotics ultimately facilitated the absorption of nutrients.^[39] However, the effect on muscle circumference needs to be further validated.

Present researches confirmed that probiotics had positive effect on reducing the incidence of infection and state of consciousness in patients with severe stroke. Infection after stroke, accompanied by inflammatory responses, facilitated the aggregation of proinflammatory cytokines into the region of brain injury and aggravated neurotoxicity.^[40–43] Research had found that probiotics therapy may be an efficient strategy to induce persistent immune regulation of the central nervous system and reduce neuroinflammation.^[44] The researches performed by Jin et al^[45] and Wan et al^[46] found that probiotics could improve the immunity of critically ill patients, which were consistent with the results of our study. It was important that the reduction of infection rate strongly correlates with improvement of nutritional status.^[47–49] Besides, our result indicated the improvement in Glasgow Coma Scale score. However, it was not enough to prove the recovery of neurological function. Therefore, further studies are needed to confirm whether probiotics could improve neurological function in patients with severe stroke.

Based on our finding, we found that probiotics supplementation was carried out in the early stage or within 72 hours after stroke. Probiotic formula was also analyzed in this meta-analysis, and it showed that eight studies used Bifidobacterium triple viable, ten studies used Bifidobacterium tetravaccine, and 1 study

added xylooligosaccharides (XOS) to probiotics. Researches showed that XOS could increase Bifidobacteria and fortify the integrity of intestinal epithelium barrier and gastrointestinal absorption function.^[50,51] Based on comprehensive analysis, it is recommended to set the course of treatment to ≥ 14 days. Marzorati M et al^[52] found 14-day probiotic intervention had a positive impact on the gut microbiome dysbiosis, as evidenced by a rapid decline in Enterobacteriaceae bacteria and increase of SCFA-producing gut flora. Wu et al^[53] also discovered a positive effect of 14-day probiotic intervention on antibiotic-induced intestinal flora dysbiosis. Therefore, we expect our findings to inform the treatment of stroke recovery.

There are several limitations in this meta-analysis. First, all the studies were performed in China. The reason is that probiotics and intestinal flora are a hot topic of research in China. In summary, it was found that the number of articles published on probiotics and intestinal flora from 2015 to 2021 is twice the number published from 2010 to 2015. Secondly, there was significant heterogeneity in the analysis of the 4 indexes, and the heterogeneity might be explained by the difference of the trial design, type of probiotic strains, characteristics of patients, and course of treatment, and the heterogeneity may be a barrier to be accepted widely of our findings. Thirdly, the Glasgow Coma Scale score outcome needs to be interpreted carefully for the scores of the 2 studies not improve,^[21,32,35] although the difference was statistically significant after combined analysis. In the future, high-quality randomized controlled trials will be required to determine the efficacy of various probiotic formulas with different treatment duration. Also, high-quality trials based on large human cohorts should be designed to clarify the underlying mechanisms of various probiotic formulas in ameliorating nutrition status of severe stroke.

6. Conclusion

In conclusion, our results show that probiotics in EN are useful for improving nutritional status, contributing to the clinical prognosis of severe stroke patients with nasal feeding, and providing clinical evidence for probiotics in the treatment of severe stroke patients. Considering the limitations of our meta-analysis, further research is needed to ensure a high level of evidence demonstrating the beneficial effects of probiotics in enteral nutrition.

Acknowledgments

The authors thank Weijia Lin, Yingying Li, and Man Yuan for their suggestions in article design and writing.

Author contributions

Xiaomin Liu drafted the manuscript. Jiahao Chu and Junzi Long searched related literatures, extracted data and assessed the quality of literature. Jie Zheng, Xue Cheng, and Xinmin Li revised and reviewed the article manuscript. Yasu Zhang had primary responsibility for final content. All authors approved the final manuscript.

Data curation: Jiahao Chu, Junzi Long.

Methodology: Jiahao Chu, Junzi Long.

Software: Jie Zheng, Xue Cheng.

Supervision: Yasu Zhang.

Visualization: Xue Cheng.

Writing – original draft: Xiaomin Liu.

Writing – review & editing: Xinmin Li.

References

- [1] FOOD Trial Collaboration. Poor nutritional status on admission predicts poor outcomes after stroke: observational data from the FOOD trial. *Stroke* 2003;34:1450–6.
- [2] Gomes F, Emery PW, Weekes CE. Risk of malnutrition is an independent predictor of mortality, length of hospital stay, and hospitalization costs in stroke patients. *J Stroke Cerebrovasc Dis* 2016;25:799–806.
- [3] Foley NC, Salter KL, Robertson J, et al. Which reported estimate of the prevalence of malnutrition after stroke is valid? *Stroke* 2009;40:e66–74.
- [4] Mosselman MJ, Kruitwagen CL, Schuurmans MJ, et al. Malnutrition and risk of malnutrition in patients with stroke: prevalence during hospital stay. *J Neurosci Nurs* 2013;45:194–204.
- [5] Wierzbicki K, Horynieccki M, Mamak D, et al. Does the nutritional status of acute stroke patients affect the neurological status in the early post-stroke period? *Neurol Res* 2020;4:1–7.
- [6] Pinto A, Tuttolomondo A, Di Raimondo D, et al. Risk factors profile and clinical outcome of ischemic stroke patients admitted in a Department of Internal Medicine and classified by TOAST classification. *Int Angiol* 2006;25:261–7.
- [7] Tuttolomondo A, Pedone C, Pinto A, et al. Gruppo Italiano di Farmacoepidemiologia dell'Anziano (GIFA) researchers. Predictors of outcome in acute ischemic cerebrovascular syndromes: The GIFA study. *Int J Cardiol* 2008;125:391–6.
- [8] Tuttolomondo A, Di Sciacca R, Di Raimondo D, et al. Effects of clinical and laboratory variables and of pretreatment with cardiovascular drugs in acute ischaemic stroke: a retrospective chart review from the GIFA study. *Int J Cardiol* 2011;151:318–22.
- [9] Gungabissoon U, Hacquoil K, Bains C, et al. Prevalence, risk factors, clinical consequences, and treatment of enteral feed intolerance during critical illness. *J Parenter Enteral Nutr* 2015;39:441–8.
- [10] Robertson C, Savva GM, Clapuci R, et al. Incidence of necrotising enterocolitis before and after introducing routine prophylactic Lactobacillus and Bifidobacterium probiotics. *Arch Dis Child Fetal Neonatal Ed* 2020;105:380–6.
- [11] Pavelescu D, Mirea L, Grintescu I. Could selected probiotics have beneficial effects on clinical outcome of severe traumatic brain injury patients? *Crit Care* 2014;18:472.
- [12] Higgins JP, Altman DG, Gotzsche PC. The Cochrane collaboration's tool for assessing risk of bias in randomized trials. *BMJ* 2011;343:d5928.
- [13] Wang J, Ru R, Sun J, et al. Effects of enteral nutrition combined with probiotics on nutritional status in patients with acute ischemic stroke after neural intervention therapy. *Journal of Chengdu Medical College* 2021;16:29–32.
- [14] Dong GH. Clinical effect of probiotics combined with enteral nutrition on patients with severe stroke. *Chinese Remedies & Clinics* 2020;20:1687–8.
- [15] Liang XQ, Mou D, Lei WJ, et al. Effect of probiotics-combined early enteral nutrition treatment in treatment of patients with severe cerebral hemorrhage. *Modern Hospitals* 2020;20:1227–9.
- [16] Ma YH. Effect of Bifidobacterium tetani combined with enteral nutrition on gastrointestinal and neurological functions in patients with severe cerebral hemorrhage. *Chinese Remedies & Clinics* 2020;20:1474–6.
- [17] Wang SS. Early enteral nutrition combined with symbiogenic preparations on the improvement of cellular immune function in elderly hypertensive patients with cerebral hemorrhage. *J Mod Med Health* 2020;20:1474–6.
- [18] Xie YM. Effects of early enteral nutrition combined with probiotics on nutritional status and gastrointestinal function in nasogastric stroke patients. *Journal of North Pharmacy* 2020;17: 44–45, 52.
- [19] Chen K. Analysis of the effect of probiotics combined with early indwelling gastric tube nasal feeding enteral nutrition support in elderly patients with severe stroke. *Yiyao Qianyan* 2019;9:155–6.
- [20] Chen JQ. Effect of Bifidobacterium tetra vaccine tablets combined with enteral nutrition on gastrointestinal complications and intestinal mucosal barrier function in patients with severe stroke. *Med Forum* 2019;23: 2397–8.
- [21] Ji X, Shi Y, Yuan B, et al. The effect of early enteral nutrition supplemented with probiotics on ventilator-associated pneumonia in patients with severe stroke. *Chin J Microecol* 2019;31:174–8.

- [22] Li YY, Yan RY, Li DC, et al. Effect of probiotic enteral nutrition intervention on intestinal flora and inflammatory state in patients with severe cerebral hemorrhage after operation. *Hebei Med* 2019;25:572–7.
- [23] Li JH, Li JY. Effect of Bifidobacterium tetravaccine tablets combined with enteral nutrition on gastrointestinal complications and intestinal mucosal barrier function in patients with severe stroke. *Journal of China Prescription Drug* 2019;17:125–6.
- [24] Dong WY, Chen H, Zhu FL. Effects of Bifidobacterium tetravaccine tablets on neurological function and nutritional status in patients with severe ischemic stroke. *Chinese Journal of Rural Medicine and Pharmacy* 2018;25:6–7.
- [25] Gao SJ. Effect of early enteral nutrition combined with probiotics in patients with acute severe stroke. *China Medical Engineering* 2018;26:34–7.
- [26] Sun HC, Zhang XG, Liu Z, et al. Effect of bifidobacterium tetravaccine Tablets on intestinal function in patients with severe stroke. *Chin J Microecol* 2018;30:1292–6.
- [27] Wu D. Effect of Early Enteral Nutrition With Bifidobacterium on Nutritional Status in Patients With Severe Cerebral Hemorrhage. Hebei: Hebei Medical University; 2018.
- [28] Zhou Q. Effect of bifid triple viable capsules combined with early enteral nutrition on nutritional status and complications in patients with severe stroke. *Diet Health* 2018;5:66–7.
- [29] Zhang Y, Wang J, Zhu HL, et al. Effects of Bifidobacterium-containing enteral nutrition intervention on the nutritional status and intestinal flora disturbance in patients with the severe cerebral infarction. *Journal of Hainan Medical University* 2017;23:1723–5.
- [30] Chen DY, Chen J. Influence of bifid triple viable capsules combined with early enteral nutrition on nutriture and complication of patients with severe stroke. *Journal of Brain and Nervous Diseases* 2016;79–81.
- [31] Hou JH. Influence of Bifidobacterium tetravaccine tablets combined with enteral nutrition on complications of digestive canal and intestinal mucous membrane barrier function of patients with severe stroke. *Chinese Journal of Pharmaco epidemiology* 2016;25: 11–13, 42.
- [32] Hou JH, Zhuo JM, Zhong WG, et al. Effects of Bifidobacterium tetravaccine tablets on nutritional status and immune indexes in patients with severe stroke. *Chinese Journal of Rural Medicine and Pharmacy* 2016;23:6–7.
- [33] Wu LJ, Li JP, Guo BW. Additional Bifidobacteria in severe stroke patients receiving enteral nutrition. *Parenteral & Enteral Nutrition* 2016;23: 220–2.
- [34] Bai GY. Clinical Research of Probiotics Enteral Nutrition for the Treatment of Critically Ill Patients With Cerebral Apoplexy. Xinjiang: Xinjiang Medical University; 2014.
- [35] Wu Y, Yu Y, Liu JF, et al. Preliminary study on application of early enteral nutrition combined with Probiotics for acute severe stroke Patients. *Clinical Medicine of China* 2014;30:1–4.
- [36] Bai GY, Fan M, Tao YL, et al. Probiotics with enteral nutrition therapy clinical observation of patients with severe stroke. *Guangdong Trace Elements Science* 2013;20:10–4.
- [37] d'Ettorre G, Rossi G, Scagnolari C, et al. Probiotic supplementation promotes a reduction in T-cell activation, an increase in Th17 frequencies, and a recovery of intestinal epithelium integrity and mitochondrial morphology in ART-treated HIV-1-positive patients. *Immun Inflamm Dis* 2017;5:244–60.
- [38] Wen Y, Li J, Long Q, et al. The efficacy and safety of probiotics for patients with constipation-predominant irritable bowel syndrome: a systematic review and meta-analysis based on seventeen randomized controlled trials. *Int J Surg* 2020;79:111–9.
- [39] Yu XY, Yin HH, Zhu JC. Increased gut absorptive capacity in rats with severe head injury after feeding with probiotics. *Nutrition* 2011;27: 100–7.
- [40] Tuttolomondo A, Maida C, Pinto A. Diabetic foot syndrome: immune-inflammatory features as possible cardiovascular markers in diabetes. *World J Orthop* 2015;6:62–76.
- [41] Petta S, Valenti L, Tuttolomondo A, et al. Interferon lambda 4 rs368234815 TT> (G variant is associated with liver damage in patients with nonalcoholic fatty liver disease. *Hepatology* 2017;66:1885–93.
- [42] Di Raimondo D, Tuttolomondo A, Buttà C, et al. Metabolic and anti-inflammatory effects of a home-based programme of aerobic physical exercise. *Int J Clin Pract* 2013;67:1247–53.
- [43] Tuttolomondo A, Maida C, Pinto A. Diabetic foot syndrome as a possible cardiovascular marker in diabetic patients. *J Diabetes Res* 2015;2015:268390.
- [44] Frank MG, Fonken LK, Watkins LR, et al. Could probiotics be used to mitigate neuroinflammation? *ACS Chem Neurosci* 2019;10:13–5.
- [45] Jin Y, Xu H, Chen Y, et al. Therapeutic effect of Bifidobacterium combined with early enteral nutrition in the treatment of severe acute pancreatitis: a pilot study. *Eur Rev Med Pharmacol Sci* 2018;22: 4018–24.
- [46] Wan G, Wang L, Zhang G, et al. Effects of probiotics combined with early enteral nutrition on endothelin-1 and C-reactive protein levels and prognosis in patients with severe traumatic brain injury. *J Int Med Res* 2019;48:300060519888112.
- [47] Kokura Y, Wakabayashi H, Nishioka S, et al. Nutritional intake is associated with activities of daily living and complications in older inpatients with stroke. *Geriatr Gerontol Int* 2018;18:1334–9.
- [48] Nii M, Maeda K, Wakabayashi H, et al. Nutritional improvement and energy intake are associated with functional recovery in patients after cerebrovascular disorders. *J Stroke Cerebrovasc Dis* 2016;25:57–62.
- [49] Alwarawrah Y, Kiernan K, MacIver NJ. Changes in nutritional status impact immune cell metabolism and function. *Front Immunol* 2018;9: 1055.
- [50] Sheng K, He S, Sun M, et al. Synbiotic supplementation containing Bifidobacterium infantis and xylooligosaccharides alleviates dextran sulfate sodium-induced ulcerative colitis. *Food Funct* 2020;11:3964–74.
- [51] Finegold SM, Li Z, Summanen PH, et al. Xylooligosaccharide increases bifidobacteria but not lactobacilli in human gut microbiota. *Food Funct* 2014;5:436–45.
- [52] Marzorati M, Abbeele PVD, Bubeck SS, et al. Bacillus subtilis HU58 and Bacillus coagulans SC208 probiotics reduced the effects of antibiotic-induced gut microbiome dysbiosis in an M-SHIME® model. *Microorganisms* 2020;8:1028.
- [53] Wu J, Gan T, Zhang Y, et al. The prophylactic effects of BIFICO on the antibiotic-induced gut dysbiosis and gut microbiota. *Gut Pathog* 2020;12:41.