

[Home](#) > [Translational Stroke Research](#) > [Article](#)Original Article | [Published: 25 June 2020](#)

# Optimizing Dosing of Vagus Nerve Stimulation for Stroke Recovery

[David T. Pruitt](#) , [Tanya T. Danaphongse](#), [Megan Lutchman](#), [Nishi Patel](#), [Priyanka Reddy](#), [Vanesse Wang](#), [Anjana Parashar](#), [Robert L. Rennaker II](#), [Michael P. Kilgard](#) & [Seth A. Hays](#)

[Translational Stroke Research](#) **12**, 65–71 (2021)

**805** Accesses | **24** Citations | [Metrics](#)

## Abstract

Vagus nerve stimulation (VNS) paired with rehabilitative training enhances recovery of function in models of stroke and is currently under investigation for use in chronic stroke patients. Dosing is critical in translation of pharmacological therapies, but electrical stimulation therapies often fail to comprehensively explore dosing parameters in preclinical studies. Varying VNS parameters has non-monotonic effects on plasticity in the central nervous system, which may directly impact efficacy for stroke. We sought to optimize stimulation intensity to maximize recovery of motor function in a model of ischemic stroke. The study design was preregistered prior to beginning data collection (DOI:

<https://doi.org/10.17605/OSF.IO/BMJJEK>). After training on an automated assessment of forelimb function and receiving an ischemic lesion in motor cortex, rats were separated into groups that received rehabilitative training paired with VNS at distinct stimulation intensities (sham, 0.4 mA, 0.8 mA, or 1.6 mA). Moderate-intensity VNS at 0.8 mA enhanced recovery of function compared with all other groups. Neither 0.4 mA nor 1.6 mA VNS was sufficient to improve functional recovery compared with equivalent rehabilitation without VNS. These results demonstrate that moderate-intensity VNS delivered during rehabilitation improves recovery and defines an optimized intensity paradigm for clinical implementation of VNS therapy.

---

This is a preview of subscription content, [access via your institution](#).

---

Access options
<p>Buy article PDF</p> <p><b>USD 39.95</b></p> <p>Price excludes VAT (USA) Tax calculation will be finalised during checkout.</p> <p>Instant access to the full article PDF.</p>

[Rent this article via DeepDyve.](#)

[Learn more about Institutional subscriptions](#)

## Data and Code Availability

---

All data and analysis code from this study have been made available on Github

(<https://github.com/davepruitt/OptimalDosingIntensityVNS>).

## References

---

1. Benjamin EJ, Blaha MJ, Chiuve SE, Cushman M, Das SR, Deo R, et al. Heart disease and stroke statistics-2017 update: a report from the American Heart Association. *Circulation*. 2017;135:e146–603.
  2. Hays SA. Improving stroke rehabilitation with vagus nerve stimulation. Springer, Singapore, pp. 503–515.
  3. Meyers EC, Solorzano BR, James J, et al. Vagus nerve stimulation enhances stable plasticity and generalization of stroke recovery. *Stroke* 2018; STROKEAHA.117.019202.
-

4. Hays SA, Khodaparast N, Hulsey DR, Ruiz A, Sloan AM, Rennaker RL II, et al. Vagus nerve stimulation during rehabilitative training improves functional recovery after intracerebral hemorrhage. *Stroke*. 2014;45:3097–100.

---

5. Dawson J, Pierce D, Dixit A, et al. Safety, feasibility, and efficacy of vagus nerve stimulation paired with upper-limb rehabilitation after ischemic stroke. *Stroke* 2015; STROKEAHA.115.010477-.

---

6. Kimberley TJ, Pierce D, Prudente CN, et al. Vagus nerve stimulation paired with upper limb rehabilitation after chronic stroke. *Stroke*; 49. 2018.  
<https://doi.org/10.1161/STROKEAHA.118.022279>.

---

7. Stroke Therapy Academic Industry Roundtable (STAIR). Recommendations for standards regarding preclinical neuroprotective and restorative drug development. *Stroke*. 1999;30:2752–8.

---

8. Borland MS, Vrana WA, Moreno NA, et al. Cortical map plasticity as a function of vagus nerve stimulation intensity. *Brain Stimul*. 2015.  
<https://doi.org/10.1016/j.brs.2015.08.018>.

---

9. Morrison RA, Hulseley DR, Adcock KS, et al. Vagus nerve stimulation intensity influences motor cortex plasticity. *Brain Stimul.* .2018  
<https://doi.org/10.1016/J.BRS.2018.10.017>.

---

10. Hulseley DR, Riley JR, Loerwald KW, Rennaker RL II, Kilgard MP, Hays SA. Parametric characterization of neural activity in the locus coeruleus in response to vagus nerve stimulation. *Exp Neurol.* 2017;289:21–30.

---

11. Hulseley DR, Shedd CM, Sarker SF, et al. Norepinephrine and serotonin are required for vagus nerve stimulation directed cortical plasticity. *Exp Neurol*; 320. 2019.  
<https://doi.org/10.1016/j.expneurol.2019.112975>.

---

12. Meyers E, Sindhurakar A, Choi R, Solorzano R, Martinez T, Sloan A, et al. The supination assessment task: an automated method for quantifying forelimb rotational function in rats. *J Neurosci Methods.* 2016.  
<https://doi.org/10.1016/j.jneumeth.2016.03.007>.

---

13. Engineer ND, Riley JR, Seale JD, Vrana WA, Shetake JA, Sudanagunta SP, et al. Reversing

pathological neural activity using targeted plasticity. *Nature*. 2011;470:101–4.

---

14. Rios M, Bucksot J, Rahebi K, Engineer C, Kilgard M, Hays S. Protocol for construction of rat nerve stimulation cuff electrodes. *Methods Protoc*. 2019;2:19.

---

15. Bucksot JE, Castelan KM, Skipton SK, et al. Parametric characterization of the rat Hering-Breuer reflex evoked with implanted and non-invasive vagus nerve stimulation. *Exp Neurol*. 2020;327:113220.

---

16. Bucksot JE, Wells AJ, Rahebi KC, Sivaji V, Romero-Ortega M, Kilgard MP, et al. Flat electrode contacts for vagus nerve stimulation. *PLoS One*. 2019;14:e0215191.

---

17. Hammer N, Löffler S, Cakmak YO, et al. Cervical vagus nerve morphometry and vascularity in the context of nerve stimulation - a cadaveric study. *Sci Rep*. 2018;8:1–9.

---

18. Planitzer U, Hammer N, Bechmann I, Glätzner J, Löffler S, Möbius R, et al. Positional relations of the cervical vagus nerve revisited. *Neuromodulation*. 2017;20:361–8.

---

19. Salgado H, Köhr G, Treviño M. Noradrenergic tone determines dichotomous control of cortical spike-timing-dependent plasticity. *Sci Rep*; 2012. <https://doi.org/10.1038/srep00417>.

---

20. Engineer ND, Kimberley TJ, Prudente CN, et al. Targeted vagus nerve stimulation for rehabilitation after stroke. *Frontiers in Neuroscience*; 13 2019; <https://doi.org/10.3389/fnins.2019.00280>.

---

21. Claflin ES, Krishnan C, Khot SP. Emerging treatments for motor rehabilitation after stroke. *The Neurohospitalist*. 2015;5:77–88.

---

22. Elias GJB, Namasivayam AA, Lozano AM. Deep brain stimulation for stroke: current uses and future directions. *Brain Stimulation*. 2018;11:3–28.

---

23. Levy RM, Harvey RL, Kissela BM, Winstein CJ, Lutsep HL, Parrish TB, et al. Epidural electrical stimulation for stroke rehabilitation: results of the prospective, multicenter, randomized, single-blinded everest trial. *Neurorehabil Neural Repair*. 2016;30:107–19.

---

24. Harvey RL, Edwards D, Dunning K, et al.  
Randomized sham-controlled trial of navigated  
repetitive transcranial magnetic stimulation for  
motor recovery in stroke. *Stroke*.  
2018;49:2138–46.

---

25. Buell EP, Loerwald KW, Engineer CT, et al.  
Cortical map plasticity as a function of vagus  
nerve stimulation rate. *Brain Stimul*. 2018.  
<https://doi.org/10.1016/J.BRS.2018.07.045>.

---

26. Borland MS, Engineer CT, Vrana WA, Moreno  
NA, Engineer ND, Vanneste S, et al. The interval  
between VNS-tone pairings determines the  
extent of cortical map plasticity. *Neuroscience*.  
2018;369:76–86.

---

27. Loerwald KW, Buell EP, Borland MS, Rennaker  
RL II, Hays SA, Kilgard MP. Varying stimulation  
parameters to improve cortical plasticity  
generated by VNS-tone pairing. *Neuroscience*.  
2018;388:239–47.

---

28. Loerwald KW, Borland MS, Rennaker RL, et al.  
The interaction of pulse width and current  
intensity on the extent of cortical plasticity  
evoked by vagus nerve stimulation. *Brain  
Stimul*. 2017.  
<https://doi.org/10.1016/J.BRS.2017.11.007>.



29. Buell EP, Borland MS, Loerwald KW, Chandler C, Hays SA, Engineer CT, et al. Vagus nerve stimulation rate and duration determine whether sensory pairing produces neural plasticity. *Neuroscience*. 2019;406:290–9.

---

## Acknowledgments

---

We thank Jaimee Nguyen, Ardalan Naghian, Veda Nashi, Kishan Thomala, Ankita Shankar, Joel Wright, Sangavi Manalan, Priyanka Vayalali, Anya Ali, Emmanuel Aykara, Medhi Zaidi, Jennifer Le, Nidhi Desai, Riley Dickson, Tony Castillo, Jenitta Kunjammattil, Tyler Short, Amber Ho, Shashi Obulasetty, Emad Sidiqi, and Ahmed Alshaikhsalama who assisted with rodent behavioral training.

---

## Funding

---

This project was supported by the National Institute of Health R01 NS085167 (MPK) and R01 NS094384 (SAH).

---

## Author information

---

### Authors and Affiliations

**Texas Biomedical Device Center, BSB11 800  
W Campbell Rd, Richardson, TX, 75080, USA**

David T. Pruitt, Tanya T. Danaphongse, Megan Lutchman, Nishi Patel, Priyanka Reddy, Vanesse

Wang, Anjana Parashar, Robert L. Rennaker

II, Michael P. Kilgard & Seth A. Hays

**Erik Jonsson School of Engineering and  
Computer Science, Richardson, TX, USA**

Robert L. Rennaker II & Seth A. Hays

**School of Behavioral and Brain Sciences, The  
University of Texas at Dallas, Richardson, TX,  
USA**

Michael P. Kilgard

### Contributions

DTP participated in study design, surgeries, data acquisition, data analysis, data interpretation, and software development and wrote the manuscript.

TTD participated in study design, surgeries, and data acquisition. ML/NP/PR/VW/AP participated in data acquisition, surgeries, tissue processing, and histological analysis. RLR participated in study design. MPK participated in study design and data interpretation. SAH participated in study design, data interpretation, and writing of the manuscript.

### Corresponding author

Correspondence to [David T. Pruitt](#).

### Ethics declarations

---

### Conflict of Interest

MPK is a consultant for MicroTransponder which develops VNS-related technologies.

## Ethical Approval

All applicable international, national, and/or institutional guidelines for the care and use of animals were followed. All experimental procedures were approved by the University of Texas Institutional Animal Care and Use Committee.

## Additional information

---

### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Rights and permissions

---

### [Reprints and Permissions](#)

## About this article

---

### Cite this article

Pruitt, D.T., Danaphongse, T.T., Lutchman, M. *et al.* Optimizing Dosing of Vagus Nerve Stimulation for Stroke Recovery. *Transl. Stroke Res.* **12**, 65–71 (2021).  
<https://doi.org/10.1007/s12975-020-00829-6>

Received	Revised	Accepted
10 January 2020	27 May 2020	14 June 2020

Published	Issue Date
25 June 2020	February 2021

DOI

<https://doi.org/10.1007/s12975-020-00829-6>

## Keywords

**Vagus**   **Stroke**   **Dosing**   **Motor**

**Stimulation**

Not logged in - 73.171.108.13

Not affiliated

**SPRINGER NATURE**

© 2023 Springer Nature Switzerland AG. Part of [Springer Nature](#).