

FORMULAS FROM ELEMENTARY MATHEMATICS

A. Algebra.

1. Laws of Exponents.

$$a^m a^n = a^{m+n}, \quad (ab)^m = a^m b^m, \quad (a^m)^n = a^{mn}, \quad a^{m/n} = \sqrt[n]{a^m}.$$

$$\text{If } a \neq 0 : \frac{a^m}{a^n} = a^{m-n}, \quad a^0 = 1, \quad a^{-m} = \frac{1}{a^m}.$$

2. Zero.

$a \cdot 0 = 0 \cdot a = 0$ for any finite number a .

$$\text{If } a \neq 0, \quad \frac{0}{a} = 0, \quad 0^a = 0, \quad a^0 = 1.$$

Division by zero is not defined.

3. Fractions.

$$\frac{a}{b} + \frac{c}{d} = \frac{ad + bc}{bd}, \quad \frac{a}{b} \cdot \frac{c}{d} = \frac{ac}{bd}, \quad \frac{a/b}{c/d} = \frac{a}{b} \cdot \frac{d}{c}, \quad \frac{-a}{b} = -\frac{a}{b} = \frac{a}{-b}.$$

$$\frac{(a/b) + (c/d)}{(e/f) + (g/h)} = \frac{(a/b) + (c/d)}{(e/f) + (g/h)} \cdot \frac{bdfh}{bdfh} = \frac{(ad + bc)fh}{(eh + fg)bd}.$$

4. Binomial Theorem, for $n = \text{positive integer}$.

$$(a+b)^n = a^n + na^{n-1}b + \frac{n(n-1)}{1 \cdot 2}a^{n-2}b^2 + \frac{n(n-1)(n-2)}{1 \cdot 2 \cdot 3}a^{n-3}b^3 + \dots + nab^{n-1} + b^n.$$

5. Proportionality factor. If y is proportional to x , or y varies directly as x , then $y = kx$ for some constant k , called the *proportionality factor*.

6. Remainder Theorem and Factor Theorem. If the polynomial $f(x)$ is divided by $x - r$ until a remainder R independent of x is obtained, then $R = f(r)$. In particular, $x - r$ is a *factor of $f(x)$* if and only if r is a *root* of the equation $f(x) = 0$.

7. *Quadratic formula.* If $a \neq 0$, the roots of the equation $ax^2 + bx + c = 0$ are given by the formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

B. Geometry. (A = area, B = area of base, C = circumference, S = lateral area or surface area, V = volume.)

1. *Triangle.* $A = \frac{1}{2}bh.$

2. *Similar Triangles.* $\frac{a'}{a} = \frac{b'}{b} = \frac{c'}{c}.$

3. *Theorem of Pythagoras.* $c^2 = a^2 + b^2.$

4. *Parallelogram.* $A = bh.$

5. *Trapezoid.* $A = \frac{1}{2}(a + b)h.$

6. *Circle.* $A = \pi r^2,$ $C = 2\pi r.$

7. *Any cylinder or prism with parallel bases.* $V = Bh.$

8. *Right circular cylinder.* $V = \pi r^2 h,$ $S = \pi r s.$

9. *Any cone or pyramid.* $V = \frac{1}{3}Bh.$

10. *Right circular cone.* $V = \frac{1}{3}\pi r^2 h,$ $S = \pi r s.$

11. *Sphere.* $V = \frac{4}{3}\pi r^3,$ $A = 4\pi r^2.$

C. Trigonometry.

1. Definitions and fundamental identities.

$$\sin(\theta) = \frac{y}{r} = \frac{1}{\csc(\theta)}.$$

$$\cos(\theta) = \frac{x}{r} = \frac{1}{\sec(\theta)}.$$

$$\tan(\theta) = \frac{y}{x} = \frac{1}{\cot(\theta)}.$$

$$\sin(-\theta) = -\sin(\theta), \quad \cos(-\theta) = \cos(\theta).$$

$$\sin^2 \theta + \cos^2 \theta = 1.$$

$$\sin(2\theta) = 2(\sin \theta \cos \theta), \quad \cos(2\theta) = \cos^2 \theta - \sin^2 \theta.$$

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta.$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta.$$

2. Angles and sides of triangle.

Law of cosines: $a^2 = b^2 + c^2 - 2bc \cos A$.

$$\text{Law of sines: } \frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}.$$

$$\text{Area} = \frac{1}{2}bc \sin A = \frac{1}{2}ac \sin B = \frac{1}{2}ab \sin C.$$