## OBSERVATIONS OF PWNE WITH THE FERMI GAMMA-RAY SPACE TELESCOPE

# A DISSERTATION SUBMITTED TO THE DEPARTMENT OF PHYSICS AND THE COMMITTEE ON GRADUATE STUDIES OF STANFORD UNIVERSITY IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE DEGREE OF DOCTOR OF PHILOSOPHY

Joshua Jeremy Lande January 2013

© Copyright by Joshua Jeremy Lande 2013 All Rights Reserved I certify that I have read this dissertation and that, in my opinion, it is fully adequate in scope and quality as a dissertation for the degree of Doctor of Philosophy.

(Stefan Funk) Principal Adviser

I certify that I have read this dissertation and that, in my opinion, it is fully adequate in scope and quality as a dissertation for the degree of Doctor of Philosophy.

(Elliott Bloom)

I certify that I have read this dissertation and that, in my opinion, it is fully adequate in scope and quality as a dissertation for the degree of Doctor of Philosophy.

(Roger Romani)

Approved for the University Committee on Graduate Studies

#### Abstract

Two things fill the mind with ever-increasing wonder and awe, the more often and the more intensely the mind of thought is drawn to them: the starry heavens above me and the moral law within me." – Immanuel Kant

The launch of the *Fermi* Gamma-ray space telescope in 2008 offered an unprecedented view into the  $\gamma$ -ray sky.

All the things we can learn with the LAT

Development of a new analysis method for studying spatially-extended PWNe using pointlike.

A monte-carlo validation of the analysis method.

Search for new spatially-extended sources with the LAT.

Observations of PWNe in the off-peak region of LAT detected pulsars.

Search for PWNe counterparts to TeV sources.

Using the population of PWNe to understand the radiation mechanism of PWNe.

#### Acknowledgement

Acknowkege the educational institutes which taught me physics: My high school HB Woodlawn, my undergraduate institution Marlboro College, and my Stanford University.

First, I would like to acknowlege those mentors who inspired me to get a PhD.

- Mark Dodge, my high school physics teacher.
- Ron Turner, my internship adviser at Analytic Services (ANSER) during the GWU Science and Engineering Apprentice Program (SEAP)
- Anthony Tyson at UC Davis for my SULI Internship
- Apurva Mehta and Sam Webb sam Web at SLAC SULI Internship.

During my PhD I was helped by an almost overwhelming large number of people in the LAT collaboration.

People at Stanford/SLAC: Stefan Funk, Elliott Bloom, Markus Ackermann, Tobias Jogler, Junichiro Katsuta, Yasunobu Uchiyama

Pointlike collaborators: Matthew Kerr, Toby Burnett, Eric Wallace, Marshall Roth

Pulsar Collaborators: David Smith, Matthew Kerr, Peter den Hartog, Tyrel Johnson, Damien Parent, Ozlem Celik

Careful review of text: Jean Ballet, Johann Cohen-Tanugi

I would like to thank the PWN Thank the people in Bordeaux: Marianne Lemoine-Goumard, Romain Rousseau, and Marie-Hélène Grondin

Fermi SLAC Grad Students: Keith Bechtol, Alex Drlica-Wagner, Alice Allafort, Herman Lee Yvonne Edmonds, Bijan Berenji, Ping Wang, Warit Mitthumsiri

Additional Astro Stanford Graduate Students: Helen Craig, Michael Shaw, Adam Van Etten, Kyle Watters

Additonal Graduate Students at Stanford: Dan Riley, Joel Frederico, Ahmed Ismail, Joshua Cogan, Kunal Sahasrabuddhe,

#### Contents

Abstract  Acknowledgement						
						1
	1.1	Gamn	na-ray Detectors	1		
		1.1.1	The Fermi Gamma-ray Space Telescope	1		
		1.1.2	H.E.S.S	1		
	1.2	Galac	tic Gamma-ray Astrophysics	1		
		1.2.1	Pulsars	1		
		1.2.2	Pulsar Wind Nebulae	1		
		1.2.3	Supernova Remenants	1		
	1.3	Radia	tion Processes	1		
		1.3.1	Synchrotron	2		
		1.3.2	Inverse Compton	2		
		1.3.3	Bremsstrahlung	2		
		1.3.4	Pi0 Decay	2		
	1.4	Source	es Detected by the Fermi LAT	2		
		1.4.1	Modeling the galactic background	2		
		1.4.2	The Second Fermi-LAT Source Catalog	2		
		1.4.3	The Second Fermi Pulsar Catalog	2		
		1.4.4	PWN Detected by the LAT	3		

<b>2</b>	Ma	ximum-likelihood analysis of LAT data	4			
	2.1	Motivations for Maximum-Likelihood Analysis of Gamma-ray Data .	4			
	2.2	Defining a Model of the Sources in the Sky	4			
	2.3	The LAT Instrument Response Functions	4			
	2.4	Application of Binned Maximum-Likelihood to LAT Data with the				
		Science Tools	4			
	2.5	pointlike	5			
	2.6	Extended Source Analysis in pointlike	Ę			
3	Search for Spatially-extended Sources					
	3.1	Analysis Method	7			
	3.2	Validation of the TS Distribution	7			
	3.3	Extended Source Detection Threshold	7			
	3.4	Testing Against Source Confusion	7			
	3.5	Test of 2LAC Sources	7			
	3.6	Systematic Errors on Extension	7			
	3.7	Extended Source Search Method	7			
	3.8	New Extended Sources	7			
	3.9	Discussion	7			
4	Sea	rch for PWNe associated with Gamma-loud Pulsars	8			
5	Search for PWNe associated with TeV Pulsars					
	5.1	List of Candidates	Ę.			
	5.2	Analysis Method	Ć			
	5.3	Sources Detected	Ć			
6	Sea	Search for PWNe associated with High Edot Pulsars				
7	Por	Population Study of LAT-detected PWNe				

#### List of Tables

## List of Figures

#### Introduction

- 1.1 Gamma-ray Detectors
- 1.1.1 The Fermi Gamma-ray Space Telescope
- 1.1.2 H.E.S.S.
- 1.2 Galactic Gamma-ray Astrophysics
- 1.2.1 Pulsars
- 1.2.2 Pulsar Wind Nebulae
- 1.2.3 Supernova Remenants
- 1.3 Radiation Processes
  - The non-thermal radiation processes typical in astrophysics are most comonly

- 1.3.1 Synchrotron
- 1.3.2 Inverse Compton
- 1.3.3 Bremsstrahlung
- 1.3.4 Pi0 Decay

#### 1.4 Sources Detected by the Fermi LAT

#### 1.4.1 Modeling the galactic background

- Galactic diffuse emission is primarily composed of . . .
- Something about how great galprop is.
- Something about

#### 1.4.2 The Second Fermi-LAT Source Catalog

- Citation is Nolan et al. (2012)
- Source classification method
- Number of sources detected by the LAT
- Forward reference Chapter 2, which does a more thorough analysis.

#### 1.4.3 The Second Fermi Pulsar Catalog

- Process of detecting Pulsars with the LAT
- Number of pulsars detected by the LAT

#### 1.4.4 PWN Detected by the LAT

 $\operatorname{Crab}$ 

 $Vela\ X$ 

MSH 15-52

 ${\rm HESS~J1857}$ 

2FGL J1857 + 026

1. http://arxiv.org/pdf/1206.3324v1.pdf

## Maximum-likelihood analysis of LAT data

- 2.1 Motivations for Maximum-Likelihood Analysis of Gamma-ray Data
  - 1. Why maximum likelihod analysis is important for LAT data
- 2.2 Defining a Model of the Sources in the Sky
- 2.3 The LAT Instrument Response Functions
- 2.4 Application of Binned Maximum-Likelihood to LAT Data with the Science Tools
  - 1. Bin the LAT data
  - 2. Convert a model of the sky into model predicted counts
  - 3. poisson likelihood

- 4. Particular implemenation of maximum likelihood anlaysis
- 5. Describe gtbin, gtselect, gtlike

#### 2.5 pointlike

- 1. Developed for Speed
- 2. Sparce Matricies,

#### 2.6 Extended Source Analysis in pointlike

# Search for Spatially-extended Sources

- 3.1 Analysis Method
- 3.2 Validation of the TS Distribution
- 3.3 Extended Source Detection Threshold
- 3.4 Testing Against Source Confusion
- 3.5 Test of 2LAC Sources
- 3.6 Systematic Errors on Extension
- 3.7 Extended Source Search Method
- 3.8 New Extended Sources
- 3.9 Discussion

Search for PWNe associated with Gamma-loud Pulsars

# Search for PWNe associated with TeV Pulsars

- 5.1 List of Candidates
- 5.2 Analysis Method
- 5.3 Sources Detected

Search for PWNe associated with High Edot Pulsars

Population Study of LAT-detected PWNe

## Bibliography

Nolan, P. L., Abdo, A. A., Ackermann, M., et al. 2012, ApJS, 199, 31