

Todo list

Get a picture of EGRET to include	11
Short description of the history of TeV astronomy	12
What is the LAT effective area?	12
Make note of “Air force had early warning of pulsars” paper	13
First gamma-ray detection	14
When was the PSR, PWN connection made	14
XX	
Get image of Synchrotron radiation (from R&L to include in discussion. . . .	18
Why no Bremsstrahlung radiation from PWN. Maybe a back-of-the-envelope estimate	21
Describe the characteristic π^0 cutoff energy	21
Include discussion of modeling, if time permitting	21
Describe Catalog	22
Dig up HESS reference of HESS J1514-59.	23
Where are sites of acceleratoin	30
Discuss pulsar evolution “The Evolution and Structure of Pulsar Wind Nebulae” – Bryan M. Gaensler and Patrick O. Slane	34
Describe Mattana’s work on pulsar wind nebulae (PWNe): “On the evolution of the Gamma- and X-ray luminosities of Pulsar Wind Nebulae”	34
Describe SNR Reverse Shock	35
Include discussion of cooling in Matanna et al 2009 (equation 12	36
Look up scaling relationships for IC and Sync radiation from Adam Van Etten’s thesis	36

what section discusses energy dependent psf?	38
What are the benefits of maximum likelihood	39
Describe Wilk's Therorem and it's application to parameter error estimation .	39
WHAT SECTION DESCRIBES EXTENDED SOURCE PDFs	41
FINISH DISCUSSION	41
Discuss how diffuse background is more complciated and requires a mapcube.	41
LINK TO arXiv:1206.1896 for MORE THOUROUGH DISCUSSION OF EF- FECTIVE AREA	42
DISCUSS HOW EFFECTIVE AREA IS A FUNCTION OF DIFFERENT THINGS	42
What is the range of the integrals	42
BETTER DISCUSSION OF PSF OF THE LAT, WHAT ITS SCALE IS... .	43
Why discard time dispersion	43
WRITE ENERGY DISPERSION AS A DELTA FUNCTION	43
FINISH	44
Figure out how the θ depedence of the IRFs factors into this calcualtion . . .	44
Write Section or Perform simple MC Simulation to demonstrate significance of detection	45
Mention that Acero et al. (in prep.). has a better discussion of the sources . .	129
Mention that Acero et al. (in prep.) has a better discussion of the sources detected for the new PWN	130
Forward reference population study in	131
What would make good future work. Something about CTA population study, something about improved modeling liek HESS J1825, something about better PSF	138

OBSERVATIONS OF PWNE WITH THE FERMI GAMMA-RAY
SPACE TELESCOPE

A DISSERTATION
SUBMITTED TO THE DEPARTMENT OF PHYSICS
AND THE COMMITTEE ON GRADUATE STUDIES
OF STANFORD UNIVERSITY
IN PARTIAL FULFILLMENT OF THE REQUIREMENTS
FOR THE DEGREE OF
DOCTOR OF PHILOSOPHY

Joshua Jeremy Lande

April 2013

© Copyright by Joshua Jeremy Lande 2013
All Rights Reserved

I certify that I have read this dissertation and that, in my opinion, it is fully adequate in scope and quality as a dissertation for the degree of Doctor of Philosophy.

(Stefan Funk) Principal Adviser

I certify that I have read this dissertation and that, in my opinion, it is fully adequate in scope and quality as a dissertation for the degree of Doctor of Philosophy.

(Elliott Bloom)

I certify that I have read this dissertation and that, in my opinion, it is fully adequate in scope and quality as a dissertation for the degree of Doctor of Philosophy.

(Roger Romani)

Approved for the University Committee on Graduate Studies

Contents

Abstract	iv
Acknowledgement	v
1 Overview	3
2 Gamma-ray Astrophysics	4
2.1 Astronomy and the Atmosphere	4
2.2 The History of Gamma-ray Astrophysics	5
2.3 The <i>Fermi</i> Gamma-ray Space Telescope	12
2.3.1 The Tracker	12
2.3.2 The Calorimeter	12
2.3.3 Anti-Coincidence Detector	12
2.3.4 Gamma-ray Burst Monitor	12
2.4 Astrophysical Sources of Gamma-rays	12
2.4.1 Pulsars	12
2.4.2 Pulsar Wind Nebulae	14
2.5 Radiation Processes in Gamma-ray Astrophysics	17
2.5.1 Synchrotron	18
2.5.2 Inverse Compton	19
2.5.3 Bremsstrahlung	20
2.5.4 Pion Decay	21
2.6 The Galactic Diffuse and Isotropic Gamma-ray Background	21
2.7 Sources Detected by the Fermi the Large Area Telescope	22

2.7.1	The Second Fermi Catalog	22
2.7.2	The Second Fermi Pulsar Catalog	23
2.7.3	Pulsar Wind Nebulae Detected by The Large Area Telescope .	23
3	The Pulsar/Pulsar Wind Nebula System	25
3.1	Neutron Star Formation	25
3.2	Pulsar Evolution	26
3.3	Pulsar Magnetosphere	30
3.4	Pulsar Wind Nebulae Structure	31
3.5	Pulsar Wind Nebula Emission	35
4	Maximum-likelihood analysis of LAT data	37
4.1	Motivations for Maximum-Likelihood Analysis of Gamma-ray Data .	38
4.2	Description of Maximum-Likelihood Analysis	39
4.3	Defining a Model of the Sources in the Sky	39
4.4	The LAT Instrument Response Functions	42
4.5	Binned Maximum-Likelihood of LAT Data with the Science Tools . .	44
4.6	The Alternate Maximum-Likelihood Package <code>pointlike</code>	46
5	Analysis of Spatially Extended LAT Sources	47
5.1	Introduction	47
5.2	Analysis Method	50
5.2.1	Modeling Extended Sources in the <code>pointlike</code> Package	50
5.2.2	Extension Fitting	52
5.2.3	<code>gtlike</code> Analysis Validation	55
5.2.4	Comparing Source Sizes	55
5.3	Validation of the TS Distribution	57
5.3.1	Point-like Source Simulations Over a Uniform Background . .	57
5.3.2	Point-like Source Simulations Over a Structured Background .	59
5.3.3	Extended Source Simulations Over a Structured Background .	63
5.4	Extended Source Detection Threshold	68
5.5	Testing Against Source Confusion	70

5.6	Test of 2LAC Sources	79
6	Search for Spatially-extended LAT Sources	83
6.1	Analysis of Extended Sources Identified in the 2FGL Catalog	84
6.2	Systematic Errors on Extension	85
6.3	Extended Source Search Method	88
6.4	New Extended Sources	94
6.4.1	2FGL J0823.0–4246	97
6.4.2	2FGL J0851.7–4635	99
6.4.3	2FGL J1615.0–5051	99
6.4.4	2FGL J1615.2–5138	103
6.4.5	2FGL J1627.0–2425c	104
6.4.6	2FGL J1632.4–4753c	105
6.4.7	2FGL J1712.4–3941	108
6.4.8	2FGL J1837.3–0700c	110
6.4.9	2FGL J2021.5+4026	112
6.5	Discussion	115
7	Search for PWNe associated with Gamma-loud Pulsars	121
7.1	Off-peak Phase Selection	121
7.2	Off-peak Analysis Method	121
7.3	Off-peak Results	121
7.4	Off-Peak Individual Source Discussion	121
8	Search for PWNe associated with TeV Pulsars	122
8.1	Introduction	123
8.2	List of very high energy (VHE) PWN Candidates	123
8.3	Analysis Method	126
8.4	Sources Detected	128
8.4.1	“PWN” and “PWNe” Sources	129
8.4.2	“O” Sources	131
8.4.3	“PSR” Sources	131

9	Population Study of LATs-detected PWNe	132
9.1	Summary of the PWNe detected by the LATs	133
9.2	The Evolution of γ -ray Emitting PWNe with the Properties of their Pulsars	133
10	Future Work (or Outlook)??	138

List of Tables

5.1	Monte Carlo Spectral Parameters	62
5.2	Extension Detection Threshold	71
6.1	Analysis of the twelve extended sources included in the 2FGL catalog	86
6.2	Nearby Residual-induced Sources	93
6.3	Extension fit for the nine additional extended sources	95
6.4	Dual localization, alternative PSF, and alternative approach to modeling the diffuse emission	96
8.1	List of analyzed VHE sources	124
8.1	List of analyzed VHE sources	125
8.2	Spatial and spectral results for detected VHE sources	128
9.1	Spatial and spectral results for detected VHE sources	134
9.1	Spatial and spectral results for detected VHE sources	135

List of Figures

2.1	Transparency of the atmosphere of the earth to photons of varying wavelenthts. This figure is from Carroll & Ostlie (2006)	5
2.2	The experimental design of Explorer XI. This figure is from Kraushaar et al. (1965).	7
2.3	The position of all 621 cosmic γ -rays detected by the Third Orbiting Solar Observatory (OSO-3). This figure is from Kraushaar et al. (1972).	8
2.4	A map of the sources observed by COS-B. The filled circles represent brighter sources. The unshaded region corresponds to the parts of the sky observed by COS-B. This figure is from Swanenburg et al. (1981).	10
2.5	The position of the Energetic Gamma Ray Experiment Telescope (EGRET) sources in the sky in galactic coordinates. The size of the source markers corresponds to the overall source intensity. This figure is from (Hartman et al. 1999).	11
2.6	The Orion plate from Bevis' book <i>Uranographia Britannica</i> . The Crab nebula can be found on the horn of Taurus the Bull on the top of the figure and the source is marked by a cloudy symbol. This figure was reproduced from Ashworth (1981).	15
3.1	The rotating dipole model of a puslar. This figure is taken from (Carroll & Ostlie 2006).	27
3.2	The magnetosphere for a rotating pulsar. The pulsar is on the bottom left of the plot. This figure is from Goldreich & Julian (1969).	30

3.3	The regions of emission in a pulsar/PWN system. This figure shows (top) the pulsar's magnetosphere, (middle), the unshocked pulsar wind and (bottom) the shocked pulsar wind which can be observed as the PWN. "R", "O", "X", and " γ " describe sites of radio, optical, X-ray, and γ -ray emission respectively. "CR", "Sy", and "IC" refer to regions of curvature, inverse Compton, and synchrotron emission. Figure is taken from Aharonian & Bogovalov (2003).	33
5.1	Counts maps and TS profiles for the SNR IC 443. (a) TS vs. extension of the source. (b) TS_{ext} for individual energy bands. (c) observed radial profile of counts in comparison to the expected profiles for a spatially extended source (solid and colored red in the online version) and for a point-like source (dashed and colored blue in the online version). (d) smoothed counts map after subtraction of the diffuse emission compared to the smoothed LAT PSF (inset). Both were smoothed by a 0.1° 2D Gaussian kernel. Plots (a), (c), and (d) use only photons with energies between 1 GeV and 100 GeV. Plots (c) and (d) include only photons which converted in the front part of the tracker and have an improved angular resolution (Atwood et al. 2009).	54
5.2	A comparison of a 2D Gaussian and uniform disk spatial model of extended sources before and after convolving with the PSF for two energy ranges. The solid black line is the PSF that would be observed for a power-law source of spectral index 2. The dashed line and the dash-dotted lines are the brightness profile of a Gaussian with $r_{68} = 0.5$ and the convolution of this profile with the LAT PSF respectively (colored red in the online version). The dash-dot-dotted and the dot-dotted lines are the brightness profile of a uniform disk with $r_{68} = 0.5$ and the convolution of this profile with the LAT PSF respectively (colored blue in the online version).	56

5.3	Cumulative distribution of the TS for the extension test when fitting simulated point-like sources in the 1 GeV to 100 GeV energy range. The four plots represent simulated sources of different spectral indices and the different lines (colored in the online version) represent point-like sources with different 100 MeV to 100 GeV integral fluxes. The dashed line (colored red) is the cumulative density function of Equation 5.11.	60
5.4	The same plot as Figure 5.3 but fitting in the 10 GeV to 100 GeV energy range.	61
5.5	Cumulative distribution of TS_{ext} for sources simulated on top of the Galactic diffuse and isotropic background.	64
5.6	The distribution of TS values when fitting 985 statistically independent simulations of W44. (a) is the distribution of TS values when fitting W44 as a point-like source and (b) is the distribution of TS_{ext} when fitting the source with a uniform disk or a radially-symmetric Gaussian spatial model. (c) is the distribution of the change in TS when fitting the source with an elliptical disk spatial model compared to fitting it with a radially-symmetric disk spatial model and (d) when fitting the source with an elliptical ring spatial model compared to an elliptical disk spatial model.	66
5.7	The distribution of fit parameters for the Monte Carlo simulations of W44. The plots show the distribution of best fit (a) flux (b) spectral index and (c) 68% containment radius. The dashed vertical lines represent the simulated values of the parameters.	67

5.8	The detection threshold to resolve an extended source with a uniform disk model for a two-year exposure. All sources have an assumed power-law spectrum and the different line styles (colors in the electronic version) correspond to different simulated spectral indices. The lines with no markers correspond to the detection threshold using photons with energies between 100 MeV and 100 GeV, while the lines with star-shaped markers correspond to the threshold using photons with energies between 1 GeV and 100 GeV.	69
5.9	The LAT detection threshold for four spectral indices and three backgrounds ($1\times$, $10\times$, and $100\times$ the Sreekumar-like isotropic background) for a two-year exposure. The left-hand plots are the detection threshold when using photons with energies between 1 GeV and 100 GeV and the right-hand plots are the detection threshold when using photons with energies between 10 GeV and 100 GeV. The flux is integrated only in the selected energy range. Overlaid on this plot are the LAT-detected extended sources placed by the magnitude of the nearby Galactic diffuse emission and the energy range they were analyzed with. The star-shaped markers (colored red in the electronic version) are sources with a spectral index closer to 1.5, the triangular markers (colored blue) an index closer to 2, and the circular markers (colored green) an index closer to 2.5. The triangular marker in plot (d) below the sensitivity line is MSH 15–52.	72

5.10	The projected detection threshold of the LAT to extension after 10 years for a power-law source of spectral index 2 against 10 times the isotropic background in the energy range from 1 GeV to 100 GeV (solid line colored red in the electronic version) and 10 GeV to 100 GeV (dashed line colored blue). The shaded gray regions represent the detection threshold assuming the sensitivity improves from 2 to 10 years by the square root of the exposure (top edge) and linearly with exposure (bottom edge). The lower plot shows the factor increase in sensitivity. For small extended sources, the detection threshold of the LAT to the extension of a source will improve by a factor larger than the square root of the exposure.	73
5.11	(a) and (b) are the distribution of TS_{ext} and of $TS_{2\text{pts}}$ when fitting simulated spatially extended sources of varying sizes as both an extended source and as two point-like sources. (c) and (d) are the distribution of $TS_{\text{ext}} - TS_{2\text{pts}}$ for the same simulated sources. (a) and (c) represent sources fit in the 1 GeV to 100 GeV energy range and (b) and (d) represent sources fit in the 10 GeV to 100 GeV energy range. In (c) and (d), the plus-shaped markers (colored red in the electronic version) are fits where $TS_{\text{ext}} \geq 16$	76
5.12	The distribution of $TS_{\text{ext}} - TS_{2\text{pts}}$ when fitting two simulated point-like sources of varying separations as both an extended source and as two point-like sources. (a), and (b) represent simulations of two point-like sources with the same spectral index and (c) and (d) represent simulations of two point-like sources with different spectral indices. (a) and (c) fit the simulated sources in the 1 GeV to 100 GeV energy range and (b) and (d) fit in the 10 GeV to 100 GeV energy range. The plus-shaped markers (colored red in the electronic version) are fits where $TS_{\text{ext}} \geq 16$	77

5.13	The cumulative density of TS_{ext} for the 733 clean AGN in 2LAC that were significant above 1 GeV calculated with <code>pointlike</code> (dashed line colored blue in the electronic version) and with <code>gtlike</code> (solid line colored black). AGN are too far and too small to be resolved by the LAT. Therefore, the cumulative density of TS_{ext} is expected to follow a $\chi^2_1/2$ distribution (Equation 5.11, the dash-dotted line colored red).	81
6.1	A TS map generated for the region around the SNR IC 443 using photons with energies between 1 GeV and 100 GeV. (a) TS map after subtracting IC 443 modeled as a point-like source. (b) same as (a), but IC 443 modeled as an extended source. The cross represents the best fit position of IC 443.	90
6.2	A diffuse-emission-subtracted 1 GeV to 100 GeV counts map of the region around 2FGL J1856.2+0450c smoothed by a $0^\circ.1$ 2D Gaussian kernel. The plus-shaped marker and circle (colored red in the online version) represent the center and size of the source fit with a radially-symmetric uniform disk spatial model. The black crosses represent the positions of other 2FGL sources. The extension is statistically significant, but the extension encompasses many 2FGL sources and the emission does not look to be uniform. Although the fit is statistically significant, it likely corresponds to residual features of inaccurately modeled diffuse emission picked up by the fit.	92

- 6.3 A diffuse-emission-subtracted 1 GeV to 100 GeV counts map of 2FGL J0823.0–4246 smoothed by a $0^\circ.1$ 2D Gaussian kernel. The triangular marker (colored red in the online version) represents the 2FGL position of this source. The plus-shaped marker and the circle (colored red) represent the best fit position and extension of this source assuming a radially-symmetric uniform disk model. The two star-shaped markers (colored green) represent 2FGL sources that were removed from the background model. From left to right, these sources are 2FGL J0823.4–4305 and 2FGL J0821.0–4254. The lower right inset is the model predicted emission from a point-like source with the same spectrum as 2FGL J0823.4–4305 smoothed by the same kernel. This source is spatially coincident with the Puppis A SNR. The light blue contours correspond to the X-ray image of Puppis A observed by *ROSAT* (Petre et al. 1996). 98
- 6.4 A diffuse-emission-subtracted 10 GeV to 100 GeV counts map of 2FGL J0851.7–4635 smoothed by a $0^\circ.25$ 2D Gaussian kernel. The triangular marker (colored red in the electronic version) represents the 2FGL position of this source. The plus-shaped marker and the circle (colored red) are the best fit position and extension of this source assuming a radially-symmetric uniform disk model. The three black crosses represent background 2FGL sources. The three star-shaped markers (colored green) represent other 2FGL sources that were removed from the background model. They are (from left to right) 2FGL J0853.5–4711, 2FGL J0855.4–4625, and 2FGL J0848.5–4535. The circular and square-shaped marker (colored blue) represents the 2FGL and relocalized position of another 2FGL source. This extended source is spatially coincident with the Vela Jr. SNR. The contours (colored light blue) correspond to the TeV image of Vela Jr. (Aharonian et al. 2007b). . . 100

- 6.5 A diffuse-emission-subtracted 10 GeV to 100 GeV counts map of 2FGL J1615.0–5051 (upper left) and 2FGL J1615.2–5138 (lower right) smoothed by a $0^\circ.1$ 2D Gaussian kernel. The triangular markers (colored red in the electronic version) represent the 2FGL positions of these sources. The cross-shaped markers and the circles (colored red) represent the best fit positions and extensions of these sources assuming a radially symmetric uniform disk model. The two black crosses represent background 2FGL sources and the star-shaped marker (colored green) represents 2FGL J1614.9-5212, another 2FGL source that was removed from the background model. The contours (colored light blue) correspond to the TeV image of HESS J1616–508 (left) and HESS J1614–518 (right) (Aharonian et al. 2006e). 101
- 6.6 A diffuse-emission-subtracted 1 GeV to 100 GeV counts map of (a) the region around 2FGL J1627.0–2425 smoothed by a $0^\circ.1$ 2D Gaussian kernel and (b) with the emission from 2FGL J1625.7–2526 subtracted. The triangular marker (colored red in the online version) represents the 2FGL position of this source. The plus-shaped marker and the circle (colored red) represent the best fit position and extension of this source assuming a radially-symmetric uniform disk model and the black cross represents a background 2FGL source. The contours in (a) correspond to the $100\ \mu\text{m}$ image observed by IRAS (Young et al. 1986). The contours in (b) correspond to CO ($J = 1 \rightarrow 0$) emission integrated from $-8\ \text{km s}^{-1}$ to $20\ \text{km s}^{-1}$. They are from de Geus et al. (1990), were cleaned using the moment-masking technique (Dame 2011), and have been smoothed by a $0^\circ.25$ 2D Gaussian kernel. 104

- 6.7 A diffuse-emission-subtracted 10 GeV to 100 GeV counts map of 2FGL J1632.4–4753c (a) smoothed by a $0\text{.}1^\circ$ 2D Gaussian kernel and (b) with the emission from the background sources subtracted. The triangular marker (colored red in the electronic version) represents the 2FGL position of this source. The plus-shaped marker and the circle (colored red) are the best fit position and extension of 2FGL J1632.4–4753c assuming a radially-symmetric uniform disk model. The four black crosses represent background 2FGL sources subtracted in (b). The circular and square-shaped markers (colored blue) represent the 2FGL and relocalized positions respectively of two additional background 2FGL sources subtracted in (b). The star-shaped marker (colored green) represents 2FGL J1634.4–4743c, another 2FGL source that was removed from the background model. The contours (colored light blue) correspond to the TeV image of HESS J1632–478 (Aharonian et al. 2006e). . . . 106
- 6.8 The spectral energy distribution of four extended sources associated with unidentified extended TeV sources. The black points with circular markers are obtained by the LAT. The points with plus-shaped markers (colored red in the electronic version) are for the associated H.E.S.S. sources. (a) the LAT SED of 2FGL J1615.0–5051 together with the H.E.S.S. SED of HESS J1616–508. (b) 2FGL J1615.2–5138 and HESS J1614–518. (c) 2FGL J1632.4–4753c and HESS J1632–478. (d) 2FGL J1837.3–0700c and HESS J1837–069. The H.E.S.S. data points are from (Aharonian et al. 2006e). Both LAT and H.E.S.S. spectral errors are statistical only. 107

6.9	A diffuse-emission-subtracted 10 GeV to 100 GeV counts map of 2FGL J1712.4–3941 (a) smoothed by a $0^\circ.15$ 2D Gaussian kernel and (b) with the emission from the background sources subtracted. This source is spatially coincident with RX J1713.7–3946 and was recently studied in Abdo et al. (2011). The triangular marker (colored red in the online version) represents the 2FGL position of this source. The plus-shaped marker and the circle (colored red) are the best fit position and extension of this source assuming a radially symmetric uniform disk model. The two black crosses represent background 2FGL sources subtracted in (b). The contours (colored light blue) correspond to the TeV image (Aharonian et al. 2007c).	109
6.10	A diffuse-emission-subtracted 10 GeV to 100 GeV counts map of the region around 2FGL J1837.3–0700c (a) smoothed by a $0^\circ.15$ 2D Gaussian kernel and (b) with the emission from the background sources subtracted. The triangular marker (colored red in the online version) represents the 2FGL position of this source. The plus-shaped marker and the circle (colored red) represent the best fit position and extension of 2FGL J1837.3–0700c assuming a radially-symmetric uniform disk model. The circular and square-shaped markers (colored blue) represent the 2FGL and the relocalized positions respectively of two background 2FGL sources subtracted in (b). The star-shaped marker (colored green) represents 2FGL J1835.5–0649, another 2FGL source that was removed from the background model. The contours (colored light blue) correspond to the TeV image of HESS J1837–069 (Aharonian et al. 2006e). The diamond-shaped marker (colored orange) represents the position of PSR J1838–0655 and the hexagonal-shaped marker (colored purple) represents the position AX J1837.3–0652 (Gotthelf & Halpern 2008).	111

6.11	A diffuse-emission-subtracted 10 GeV to 100 GeV counts map of the region around 2FGL J2021.5+4026 smoothed by a 0.1° 2D Gaussian kernel. The triangular marker (colored red in the online version) represents the 2FGL position of this source. The plus-shaped marker and the circle (colored red) represent the best fit position and extension of 2FGL J2021.5+4026 assuming a radially-symmetric uniform disk model. The star-shaped marker (colored green) represents 2FGL J2019.1+4040, a 2FGL source that was removed from the background model. 2FGL J2021.5+4026 is spatially coincident with the γ -Cygni SNR. The contours (colored light blue) correspond to the 408MHz image of γ -Cygni observed by the Canadian Galactic Plane Survey (Taylor et al. 2003).	113
6.12	The spectral energy distribution of the extended sources Puppis A (2FGL J0823.0–4246) and γ -Cygni (2FGL J2021.5+4026). The lines (colored red in the online version) are the best fit power-law spectral models of these sources. Puppis A has a spectral index of 2.21 ± 0.09 and γ -Cygni has an index of 2.42 ± 0.19 . The spectral errors are statistical only. The upper limit is at the 95% confidence level. . . .	114
6.13	The 21 spatially extended sources detected by the LAT at GeV energies with 2 years of data. The twelve extended sources included in 2FGL are represented by the circular markers (colored red in the online version). The nine new extended sources are represented by the triangular markers (colored orange). The source positions are overlaid on a 100 MeV to 100 GeV Aitoff projection sky map of the LAT data in Galactic coordinates.	116

6.14	A comparison of the sizes of extended sources detected at both GeV and TeV energies. The TeV sizes of W30, 2FGL J1837.3–0700c, 2FGL J1632.4–4753c, 2FGL J1615.0–5051, and 2FGL J1615.2–5138 are from Aharonian et al. (2006e). The TeV sizes of MSH 15–52, HESS J1825–137, Vela X, Vela Jr., RX J1713.7–3946 and W28 are from Aharonian et al. (2005a, 2006c,d, 2007b,c, 2008a). The TeV size of IC 443 is from Acciari et al. (2009) and W51C is from Krause et al. (2011). The TeV sizes of MSH 15–52, HESS J1614–518, HESS J1632–478, and HESS J1837–069 have only been reported with an elliptical 2D Gaussian fit and so the plotted sizes are the geometric mean of the semi-major and semi-minor axis. The LAT extension of Vela X is from Abdo et al. (2010). The TeV sources were fit assuming a 2D Gaussian surface brightness profile so the plotted GeV and TeV extensions were first converted to r_{68} (see Section 5.2.4). Because of their large sizes, the shape of RX J1713.7–3946 and Vela Jr. were not directly fit at TeV energies and so are not included in this comparison. On the other hand, dedicated publications by the LAT collaboration on these sources showed that their morphologies are consistent (Abdo et al. 2011; Tanaka et al. 2011). The LAT extension errors are the statistical and systematic errors added in quadrature.	117
6.15	The distributions of the sizes of 18 extended LAT sources at GeV energies (colored blue in the electronic version) and the sizes of the 40 extended H.E.S.S. sources at TeV energies (colored red). The H.E.S.S. sources were fit with a 2D Gaussian surface brightness profile so the LAT and H.E.S.S. sizes were first converted to r_{68} . The GeV size of Vela X is taken from Abdo et al. (2010). Because of their large sizes, the shape of RX J1713.7–3946 and Vela Jr. were not directly fit at TeV energies and are not included in this comparison. Centaurus A is not included because of its large size.	118

6.16	The distribution of spectral indices of the 1873 2FGL sources (colored red in the electronic version) and the 21 spatially extended sources (colored blue). The index of Centaurus A is taken from Nolan et al. (2012) and the index of Vela X is taken from Abdo et al. (2010).	120
9.1	...	135
9.2	...	136
9.3	...	137

List of Acronyms

1FHL the first *Fermi* hard-source list. 126

2CG the second COS-B catalog. 9

2FGL the second *Fermi* catalog. 22, 38, 122, 127

2PC the second *Fermi* pulsar catalog. 23, 124, 125

3EG the Third EGRET Catalog. 11

ACD Anti-Coincidence Detector. vii, 12

arcsec second of arc. 32

BPL broken-power law. 40

CGRO the Compton Gamma Ray Observatory. 10

CGS the Centimetre-Gram-Second System of Units. 40, 41

ECPL exponentially-cutoff power law. 40, 41

EGRET the Energetic Gamma Ray Experiment Telescope. xii, 9–11

ESA the European Space Agency. 9

FWHM full width at half maximum. 8

GBM Gamma-ray Burst Monitor. vii, 12

IACT Imaging air Cherenkov detector. 48, 119, 122, 126

IC inverse Compton. 6, 19, 20, 31, 32, 35, 126

LAT the Large Area Telescope. iv, v, vii, viii, 12, 22, 23, 122, 123, 125–128, 132

MIT the Massachusetts Institute of Technology. 6, 13

MSC massive star cluster. 125

MSP millisecond pulsar. 27

NASA the National Aeronautics and Space Administration. 9, 10

NRL the Naval Research Laboratory. 13

NS neutron star. 13, 25, 26, 30

OSO-3 the Third Orbiting Solar Observatory. xii, 8, 9, 22

PL power law. 40, 41

PSF point spread function. 126

PWN pulsar wind nebula. iv, v, xiii, 1, 6, 14, 25, 29, 31–35, 122, 123, 125–129, 132

SA solid angle. 42, 43

SAS-2 the second Small Astronomy Satellite. 9, 10

SNR supernova remnant. 29, 32, 41

UNID unidentified source. 125, 126

VHE very high energy. ix, xi, 122–130, 134

Chapter 8

Search for PWNe associated with TeV Pulsars

This chapter is based the first part of the the paper “Constraints on the Galactic Population of TeV Pulsar Wind Nebulae using Fermi Large Area Telescope Observations” by Acero et al which is currently in prep.

In Chapter 6, we searched for spatially-extended sources in the 2FGL catalog. This search showed that the spatial analysis of *Fermi* sources is important in identifying γ -ray emitting pulsar wind nebulae (PWNe). In that work, we analyzed the γ -ray emitting PWNs HESS J1825–137 and MSH 15–52 which had previously been detected in the second *Fermi* catalog (2FGL). In addition, this analysis discovered that there were three additional spatially-extended *Fermi* sources coincident with PWNs candidates (HESS J1616–508, HESS J1632–478, HESS J1837–069).

In Chapter 7, we then searched for γ -ray emitting PWNs by looking in the off-peak emission of the Large Area Telescope (LAT)-detected pulsars. In that analysis, we detected four γ -ray emitting PWNs (Vela–X, the Crab Nebula, MSH 15–52, and 3C 58).

In this chapter, we continue our search for γ -ray emitting PWNs by searching for PWNs which had previously been detected at very high energy (VHE) energies by Imaging air Cherenkov detectors (IACTs). We note that the work presented here is a very condensed version of the results presented in the accompanying work (Acero

et al. in prep.). We refer to that publication for a more detailed discussion of the analysis.

8.1 Introduction

We took all sources detected at VHE energies and potentially associated with PWNs and performed a search at GeV energies for γ -ray emission. As was seen in previous chapters, many PWNs have been detected both at GeV and VHE energies. Therefore, we suspect that by searching LAT data in the regions of VHE PWNs might yield the discovery of new LAT-detecting PWN.

In addition, there are several PWN which have been detected at VHE energies which do not have an associated γ -ray pulsar (such as HESS J1825–137 and HESS J1837–069). We therefore suspect that this search strategy may discovery new PWN which were not previously discovered either in the off-peak search discussed in Chapter 7 or in other dedicated analyses.

8.2 List of VHE PWN Candidates

Table 8.1. List of analyzed VHE sources

Name	Class	l (deg.)	b (deg.)	Pulsar	2PC	Reference
VER J0006+727	PWN	119.58	10.20	PSR J0007+7303	Y	McArthur (2011)
MGRO J0631+105	PWN	201.30	0.51	PSR J0631+1036	Y	Abdo et al. (2009c)
MGRO J0632+17	PWN	195.34	3.78	PSR J0633+1746	Y	Abdo et al. (2009c)
HESS J1018–589	UNID	284.23	–1.72	PSR J1016–5857	Y	H. E. S. S. Collaboration et al. (2012)
HESS J1023–575	MSC	284.22	–0.40	PSR J1023–5746	Y	H.E.S.S. Collaboration et al. (2011e)
HESS J1026–582	PWN	284.80	–0.52	PSR J1028–5819	Y	H.E.S.S. Collaboration et al. (2011e)
HESS J1119–614	PWN	292.10	–0.49	PSR J1119–6127	Y	Presentation ^a
HESS J1303–631	PWN	304.24	–0.36	PSR J1301–6305	N	Aharonian et al. (2005b)
HESS J1356–645	PWN	309.81	–2.49	PSR J1357–6429	Y	H.E.S.S. Collaboration et al. (2011d)
HESS J1418–609	PWN	313.25	0.15	PSR J1418–6058	Y	Aharonian et al. (2006b)
HESS J1420–607	PWN	313.56	0.27	PSR J1420–6048	Y	Aharonian et al. (2006b)
HESS J1427–608	UNID	314.41	–0.14	...	N	Aharonian et al. (2008c)
HESS J1458–608	PWN	317.75	–1.70	PSR J1459–6053	Y	de los Reyes et al. (2012)
HESS J1503–582	UNID	319.62	0.29	...	N	Renaud et al. (2008)
HESS J1507–622	UNID	317.95	–3.49	...	N	H.E.S.S. Collaboration et al. (2011c)
HESS J1514–591	PWN	320.33	–1.19	PSR J1513–5908	Y	Aharonian et al. (2005a)
HESS J1554–550	PWN	327.16	–1.07	...	N	Acero et al. (2012)
HESS J1616–508	PWN	332.39	–0.14	PSR J1617–5055	N	Aharonian et al. (2006e)
HESS J1626–490	UNID	334.77	0.05	...	N	Aharonian et al. (2008c)
HESS J1632–478	PWN	336.38	0.19	...	N	Aharonian et al. (2006e)
HESS J1634–472	UNID	337.11	0.22	...	N	Aharonian et al. (2006e)
HESS J1640–465	PWN	338.32	–0.02	...	N	Aharonian et al. (2006e)
HESS J1702–420	UNID	344.30	–0.18	PSR J1702–4128	Y	Aharonian et al. (2006e)
HESS J1708–443	PWN	343.06	–2.38	PSR J1709–4429	Y	H.E.S.S. Collaboration et al. (2011b)
HESS J1718–385	PWN	348.83	–0.49	PSR J1718–3825	Y	Aharonian et al. (2007a)
HESS J1729–345	UNID	353.44	–0.13	...	N	H.E.S.S. Collaboration et al. (2011a)
HESS J1804–216	UNID	8.40	–0.03	PSR J1803–2149	Y	Aharonian et al. (2006e)
HESS J1809–193	PWN	11.18	–0.09	PSR J1809–1917	N	Aharonian et al. (2007a)
HESS J1813–178	PWN	12.81	–0.03	PSR J1813–1749	N	Aharonian et al. (2006e)
HESS J1818–154	PWN	15.41	0.17	PSR J1818–1541	N	Hofverberg (2011)
HESS J1825–137	PWN	17.71	–0.70	PSR J1826–1334	N	Aharonian et al. (2006c)
HESS J1831–098	PWN	21.85	–0.11	PSR J1831–0952	N	Sheidaei (2011)
HESS J1833–105	PWN	21.51	–0.88	PSR J1833–1034	Y	Djannati-Ataï et al. (2008)
HESS J1834–087	UNID	23.24	–0.31	...	N	Aharonian et al. (2006e)
HESS J1837–069	UNID	25.18	–0.12	PSR J1836–0655	N	Aharonian et al. (2006e)
HESS J1841–055	UNID	26.80	–0.20	PSR J1838–0537	Y	Aharonian et al. (2008c)
HESS J1843–033	UNID	29.30	0.51	...	N	Hoppe (2008)
MGRO J1844–035	UNID	28.91	–0.02	...	N	Abdo et al. (2009c)

Table 8.1 (cont'd)

Name	Class	l (deg.)	b (deg.)	Pulsar	2PC	Reference
HESS J1846–029	PWN	29.70	–0.24	PSR J1846–0258	N	Djannati-Ataï et al. (2008)
HESS J1848–018	UNID	31.00	–0.16	...	N	Chaves et al. (2008)
HESS J1849–000	PWN	32.64	0.53	PSR J1849–001	N	Terrier et al. (2008)
HESS J1857+026	UNID	35.96	–0.06	PSR J1856+0245	N	Aharonian et al. (2008c)
HESS J1858+020	UNID	35.58	–0.58	...	N	Aharonian et al. (2008c)
MGRO J1900+039	UNID	37.42	–0.11	...	N	Abdo et al. (2009c)
MGRO J1908+06	UNID	40.39	–0.79	PSR J1907+0602	Y	Aharonian et al. (2009)
HESS J1912+101	PWN	44.39	–0.07	PSR J1913+1011	N	Aharonian et al. (2008b)
VER J1930+188	PWN	54.10	0.26	PSR J1930+1852	N	Acciari et al. (2010)
MGRO J1958+2848	PWN	65.85	–0.23	PSR J1958+2846	Y	Abdo et al. (2009c)
VER J1959+208	PSR	59.20	–4.70	PSR J1959+2048	Y	Hall et al. (2003)
VER J2016+372	UNID	74.94	1.15	...	N	Aliu (2011)
MGRO J2019+37	PWN	75.00	0.39	PSR J2021+3651	Y	Abdo et al. (2007)
MGRO J2031+41A	UNID	79.53	0.64	...	N	Abdo et al. (2007)
MGRO J2031+41B	UNID	80.25	1.07	PSR J2032+4127	Y	Bartoli et al. (2012)
MGRO J2228+61	PWN	106.57	2.91	PSR J2229+6114	Y	Abdo et al. (2009c)

^aThis source was presented at the "Supernova Remnants and Pulsar Wind Nebulae in the Chandra Era", 2009. See http://cxc.harvard.edu/cdo/snr09/pres/DjannatiAtai_Arache_v2.pdf.

Note. — The VHE sources that we searched for using LAT observations. The classifications come from TeVCat and are PWN for pulsar wind nebula, unidentified source (UNID) for unidentified source, and MSC for massive star cluster. We include HESS J1023–575 because it is potentially a PWN. For sources with an associated pulsar, column 4 includes the pulsar's name. Column 5 describes if the pulsar has been detected by the LAT and included in the second *Fermi* pulsar catalog (2PC) (See Chapter 7).

We used TeVCat to define the our target list of VHE sources. TeVCat is a catalog of sources detected at VHE energies by IACTs.¹ We selected all sources from this catalog where the emission was classified as being due to a PWN. In addition, we included all UNID sources within 5° of the galactic plane since they could be due to a PWN. Finally, we included HESS J1023–575. Although this source is classified as a massive star cluster in the TeVCat, de Naurois & H.E.S.S. Collaboration (2013) suggested that the emission could be due to PWN. The list of all sources included in our analysis as well as their classification in TeVCat can be found in Table 8.1.

8.3 Analysis Method

In this search, our analysis method was very similar to the analysis in Chapter 7. We used the same hybrid `pointlike`/`gtlike` approach for studying the spatial and spectral character of each source and modeled the region using the same standard background models.

The major difference was that this analysis was performed only for $E > 10$ GeV. As can be seen in Chapter 7, for energies much lower than 10 GeV, source analysis becomes strongly influenced by Galactic-diffuse emission and systematic errors associated with incorrectly modeling the emission. On the other hand, the γ -ray emission from PWN is expected to be the rising component of an inverse Compton (IC) peak which falls at VHE energies. Therefore, the emission observed by the LAT is expected to be hard and most significant at higher energies. Therefore, we expect that starting the analysis at 10 GeV will significantly reduce systematics associated with this analysis while preserving most of the space for discovery.

Because the analysis was performed only in this high energy range where the point spread function (PSF) of the LAT is much improved, we were able to use a smaller region of interest (a radius of 5° in `pointlike` and a square of size $7^\circ \times 7^\circ$ in `gtlike`).

Another differences is that we used an event class with less background contamination (Pass 7 Clean instead of Pass 7 Source) and modeled nearby background sources using the first *Fermi* hard-source list (1FHL) (Ackermann et al. in prep).

¹TeVCat can be found at <http://tevcat.uchicago.edu>.

For our analysis, we assume the GeV emission to have a power-law spectral model and to have whatever was the published spatial model observed at VHE energies. We define TS_{TeV} as the likelihood-ratio test for the significance of the source assuming it to have the power-law spectral model and the published VHE spatial model. We claim a detection when $TS_{\text{TeV}} > 16$. Our significance test has only two degrees of freedom: the flux and spectral index. Therefore, following Wilk's Theorem (see Section 5.3.1), this corresponds to a 3.6σ detection threshold. When the source is significantly-detected, we quote the best-fit spectral parameters and otherwise we derive a upper limit on the flux of any potential emission.

We note that Acero et al. (in prep.) performs a more detailed morphological analysis which also fits the positions of the sources assuming their emissions to be point-like and spatially-extended. That work then uses TS_{ext} (See Section 5.2.2) to test if the emission is spatially-extended and otherwise computes an upper limit on any potential spatial extension. The more detailed analysis in Acero et al. (in prep.) is needed to compare the spatial overlap between the GeV and VHE emission. But for brevity, we omit the details and simply use the results.

We note that many of these PWNs candidates are in regions with LAT-detected pulsars. For these sources, Acero et al. (in prep.) presents the spectral and spatial results both with and without the LAT-detected pulsar in the background model. For the analysis, the spectrum of these sources of these background pulsars was taken from the 2FGL catalog. For simplicity, we present only the analysis with the pulsar included in the background model. This analysis presents our best-guess at the true spectrum of any PWN emission. But we caution that this method could be biased in either oversubtracting or undersubtracting the pulsar depending upon systematics in the fit of the pulsar.

There are three major sources of systematic uncertainty effect the spectrum of these sources. The first systematic is due to uncertainty in our modeling of the Galactic diffuse emission. We estimate this uncertainty following the method of Section 6.2. The second systematic is due to uncertainty in the effective area. We estimated the systematic using the method described in Ackermann et al. (2012). The final systematic is due to our uncertainty in the true morphology of the source. We used as our

Table 8.2. Spatial and spectral results for detected VHE sources

Name	ID	TS _{TeV}	$F_{10\text{ GeV}}^{316\text{ GeV}}$ ($10^{-10}\text{ ph cm}^{-2}\text{ s}^{-1}$)	Γ
HESS J1018–589	O	25	$1.5 \pm 0.5 \pm 0.7$	$2.31 \pm 0.50 \pm 0.49$
HESS J1023–575	PWNc	52	$4.6 \pm 0.9 \pm 1.2$	$1.99 \pm 0.24 \pm 0.32$
HESS J1119–614	PWNc	16	$2.0 \pm 0.6 \pm 0.8$	$1.83 \pm 0.41 \pm 0.36$
HESS J1303–631	PWNc	37	$3.6 \pm 0.9 \pm 2.1$	$1.53 \pm 0.23 \pm 0.37$
HESS J1356–645	PWN	24	$1.1 \pm 0.4 \pm 0.5$	$0.94 \pm 0.40 \pm 0.40$
HESS J1420–607	PWNc	36	$3.4 \pm 0.9 \pm 1.1$	$1.81 \pm 0.29 \pm 0.31$
HESS J1507–622	O	21	$1.5 \pm 0.5 \pm 0.5$	$2.33 \pm 0.48 \pm 0.48$
HESS J1514–591	PWN	156	$6.2 \pm 0.9 \pm 1.3$	$1.72 \pm 0.16 \pm 0.17$
HESS J1616–508	PWNc	75	$9.3 \pm 1.4 \pm 2.3$	$2.18 \pm 0.19 \pm 0.20$
HESS J1632–478	PWNc	137	$11.8 \pm 1.5 \pm 5.3$	$1.82 \pm 0.14 \pm 0.19$
HESS J1634–472	O	33	$5.6 \pm 1.3 \pm 2.5$	$1.96 \pm 0.25 \pm 0.29$
HESS J1640–465	PWNc	47	$5.0 \pm 1.0 \pm 1.7$	$1.95 \pm 0.23 \pm 0.20$
HESS J1708–443	PSR	33	$5.5 \pm 1.3 \pm 3.5$	$2.13 \pm 0.31 \pm 0.33$
HESS J1804–216	O	124	$13.4 \pm 1.6 \pm 3.1$	$2.04 \pm 0.16 \pm 0.24$
HESS J1825–137	PWN	56	$5.6 \pm 1.2 \pm 9.0$	$1.32 \pm 0.20 \pm 0.39$
HESS J1834–087	O	27	$5.5 \pm 1.2 \pm 2.5$	$2.24 \pm 0.34 \pm 0.42$
HESS J1837–069	PWNc	73	$7.5 \pm 1.3 \pm 4.2$	$1.47 \pm 0.18 \pm 0.30$
HESS J1841–055	PWNc	64	$10.9 \pm 0.8 \pm 4.1$	$1.60 \pm 0.27 \pm 0.33$
HESS J1848–018	PWNc	19	$7.4 \pm 1.9 \pm 2.7$	$2.46 \pm 0.50 \pm 0.51$
HESS J1857+026	PWNc	53	$4.2 \pm 0.3 \pm 1.3$	$1.01 \pm 0.24 \pm 0.25$
VER J2016+372	O	31	$1.8 \pm 0.5 \pm 0.8$	$2.45 \pm 0.44 \pm 0.49$

Note. — The results of our analysis search using LAT data for VHE PWNs. Column 2 says the classification of the LAT emission (See Section 8.4). Column 3 is TS_{TeV}, columns 4 is the flux, and 5 is the spectral index all computed assuming a power-law spectral model (See Section 8.3 for a discussion of systematic errors).

systematic error on this the difference in spectrum when the source is fit assuming the published VHE spatial model and when using the spatial model measured by the LAT.

8.4 Sources Detected

We detected 22 sources. For significantly-detected sources, we present the spatial and spectral results for these sources in Table 9.1. Flux upper limits for non-detected sources as well as spectral points in three independent energy bins are can be found in Acero et al. (in prep.).

We attempt to classify the GeV emission into four categories: “PWN” for sources where the GeV emission is clearly identified PWN, “PWNC” for sources where the GeV emission could potentially be due to a PWN, “PSR” for sources where the emission is mostl likely due to pulsed emission inside the pulsar’s magnetosphere, and “O” (for other) when the true nature of emission is uncertain.

We categorize a source as “PWN” or “PWNC” when the emission has a hard spectrum which connects spectrally to the VHE spectrum and when there is some muliwavelenth evidence that the GeV and VHE emission should be due to a pulsar-wind nebula. We put a source in the “PWN” when the VHE emission suggests more strongly that the emission is due to the pulsar wind nebula. We will discuss the muliwavelent evidence for each “PWN” source in Section 8.4.1. We include in Table 9.1 the source classifications.

Mention that Acero et al. (in prep.). has a better discussion of the sources

8.4.1 “PWN” and “PWNC” Sources

In total, we detect fourteen sources which we classify as “PWN” or “PWNC”. Five of these PWN and PWN candidates are first reported in this analysis.

Of these fourteen sources, three are classified as “PWN”. They are HESS J1356–645, MSH 15–52 (HESS J1514–591), and HESS J1825–137. HESS J1356–645 and MSH 15–52 are classified as “PWN” because of the correlation between the X-ray and VHE emission (H.E.S.S. Collaboration et al. 2011d; Aharonian et al. 2005a). HESS J1825–137 is classified as “PWN” because of the energy-dependent morphology observed at VHE energies (Aharonian et al. 2006c). We note that HESS J1356–645 was first presented as an PWN in this work. Once we add the Crab Nebula and Vela–X (not analyzed in this work) to this list, the total number of clearly-identified PWNs detected at GeV energies is five.

In addition, we detect 11 “PWNe”-type sources. Four of these sources are first reported in this work: HESS J1119–614, HESS J1303–631, HESS J1420–607, and HESS J1841–055. These sources are all powered by pulsars energetic enough to power the observed emission (PSR J1119–6127, PSR J1301–6305, PSR J1420–6048, PSR J1838–0537), and they all have a hard spectrum which connects to the spectra observed at VHE energies.

We caution that two of these sources have an associated LAT-detected pulsar and were therefore studied in Chapter 7.

We mention that PSR J1119–6127, PSR J1420–6048, and PSR J1838–0537 are all detected by the LAT and therefore were studied in Section ???. In Section ??, PSR J1119–6127 has $TS = 61.3$ and is classified as a “U”-type source because the spectrum is relatively soft (spectral index ~ 2.2). But in the off-peak analysis, one can observe a faint hard component of the emission at high energy as well as a stronger component at low energy. In the off-peak analysis, we calculated $TS = 13.8$ in the energy range from 10 GeV to 316 GeV when computing an SED with a fixed spectral index of 2. So the off-peak analysis is fairly consistent with this work.

For PSR J1420–6048, ... <http://www.slac.stanford.edu/~lande/pwnpipeline/v37/website/PSRJ1420-6048.html>

The off-peak analysis of PSR J1838–0537 detects a soft and significantly-cutoff spectrum which is <http://www.slac.stanford.edu/~lande/pwnpipeline/v37/website/PSRJ1838-0537.html>. In the residual test-statistic map, there is significant emission towards the position of HESS J1841–055. The off-peak emission is significantly-extended. Therefore our analysis are most-likely compatible.

Mention that Acero et al. (in prep.) has a better discussion of the sources detected for the new PWN

The remaining seven sources have been previously published: HESS J1023–575 (Ackermann et al. 2011a), HESS J1640–465 (Slane et al. 2010), HESS J1616–508 (Lande et al. 2012), HESS J1632–478 (Lande et al. 2012), HESS J1837–069 (Lande et al. 2012), HESS J1848–018 (Tam et al. 2010), and HESS J1857+026 (Rousseau et al. 2012b).

Finally, note that we have add HESS J1848–018 to the list of “PWNe” sources

even though the GeV emission has a soft spectrum because of the careful multiwavelength considerations presented in Lemoine-Goumard et al. (2011).

8.4.2 “O” Sources

The third category is “O”...

8.4.3 “PSR” Sources

The final category is “PSR”...

Forward reference population study in .

Bibliography

- Abdo, A., Ackermann, M., Ajello, M., et al. 2010, *Astrophys.J.*, 722, 1303
- Abdo, A. A., Allen, B., Berley, D., et al. 2007, *ApJ*, 664, L91
- Abdo, A. A., Ackermann, M., Ajello, M., et al. 2009a, *ApJ*, 706, L1
- . 2009b, *ApJS*, 183, 46
- Abdo, A. A., Allen, B. T., Aune, T., et al. 2009c, *ApJ*, 700, L127
- Abdo, A. A., Ackermann, M., Ajello, M., et al. 2009d, *Astroparticle Physics*, 32, 193
- . 2010a, *ApJ*, 714, 927
- Abdo, A. A., Ackermann, M., Ajello, M., et al. 2010b, *A&A*, 523, A46
- Abdo, A. A., Ackermann, M., Ajello, M., et al. 2010c, *Science*, 328, 725
- . 2010d, *ApJS*, 188, 405
- . 2010e, *ApJ*, 708, 1254
- . 2010f, *ApJ*, 718, 348
- Abdo, A. A., Ackermann, M., Ajello, M., et al. 2010, *The Astrophysical Journal*, 713, 146
- Abdo, A. A., Ackermann, M., Ajello, M., et al. 2010a, *Science*, 327, 1103
- . 2010b, *ApJ*, 712, 459

- Abdo, A. A., Ackermann, M., Ajello, M., et al. 2010c, *A&A*, 512, A7
- Abdo, A. A., Ackermann, M., Ajello, M., et al. 2010d, *Physical Review Letters*, 104, 101101
- . 2010e, *ApJS*, 187, 460
- . 2011, *ApJ*, 734, 28
- Acciari, V. A., Aliu, E., Arlen, T., et al. 2009, *ApJ*, 698, L133
- . 2010, *ApJ*, 719, L69
- . 2011, *ApJ*, 738, 3
- Acerro, F., Ackermann, M., Ajello, M., & et al. in prep., *ApJ*
- Acerro, F., Djannati-Ataï, A., Förster, A., et al. 2012, *ArXiv e-prints*
- Ackermann, M., Ajello, M., Allafort, A., et al. in prep, *ApJ*
- Ackermann, M., Ajello, M., Baldini, L., et al. 2011a, *ApJ*, 726, 35
- Ackermann, M., Ajello, M., Allafort, A., et al. 2011b, *ApJ*, 743, 171
- Ackermann, M., Ajello, M., Albert, A., et al. 2012, *ApJS*, 203, 4
- Aharonian, F., Akhperjanian, A. G., Aye, K.-M., et al. 2005a, *A&A*, 435, L17
- . 2005b, *A&A*, 439, 1013
- Aharonian, F., Akhperjanian, A. G., Bazer-Bachi, A. R., et al. 2006a, *A&A*, 460, 743
- . 2006b, *A&A*, 456, 245
- . 2006c, *A&A*, 460, 365
- . 2006d, *A&A*, 448, L43
- . 2006e, *ApJ*, 636, 777

—. 2007a, *A&A*, 472, 489

—. 2007b, *ApJ*, 661, 236

—. 2007c, *A&A*, 464, 235

—. 2008a, *A&A*, 481, 401

Aharonian, F., Akhperjanian, A. G., Barres de Almeida, U., et al. 2008b, *A&A*, 484, 435

—. 2008c, *A&A*, 477, 353

Aharonian, F., Akhperjanian, A. G., Anton, G., et al. 2009, *A&A*, 499, 723

Aharonian, F. A., & Bogovalov, S. V. 2003, *New A*, 8, 85

Aharonian, F. A., Coppi, P. S., & Voelk, H. J. 1994, *ApJ*, 423, L5

Aharonian, F. A., Akhperjanian, A. G., Bazer-Bachi, A. R., et al. 2007d, *A&A*, 469, L1

Ajello, M., Allafort, A., Baldini, L., et al. 2012, *ApJ*, 744, 80

Akaike, H. 1974, *IEEE Transactions on Automatic Control*, 19, 716

Albert, J., Aliu, E., Anderhub, H., et al. 2006, *Science*, 312, 1771

Aliu, E. 2011, in *International Cosmic Ray Conference*, Vol. 7, *International Cosmic Ray Conference*, 227

Arnold, J. R., Metzger, A. E., Anderson, E. C., & van Dilla, M. A. 1962, *J. Geophys. Res.*, 67, 4878

Arons, J. 1996, *Space Sci. Rev.*, 75, 235

Ashworth, William B., J. 1981, *Proceedings of the American Philosophical Society*, 125, pp. 52

- Atwood, W. B., Abdo, A. A., Ackermann, M., et al. 2009, *ApJ*, 697, 1071
- Balbo, M., Saouter, P., Walter, R., et al. 2010, *A&A*, 520, A111
- Baltz, E. A., Berenji, B., Bertone, G., et al. 2008, *J. Cosmology Astropart. Phys.*, 7, 13
- Bamba, A., Ueno, M., Koyama, K., & Yamauchi, S. 2003, *The Astrophysical Journal*, 589, 253
- Bartoli, B., Bernardini, P., Bi, X. J., et al. 2012, *ApJ*, 745, L22
- Baum, W. A., Johnson, F. S., Oberly, J. J., et al. 1946, *Phys. Rev.*, 70, 781
- Bertsch, D. L., Brazier, K. T. S., Fichtel, C. E., et al. 1992, *Nature*, 357, 306
- Bignami, G. F., Boella, G., Burger, J. J., et al. 1975, *Space Science Instrumentation*, 1, 245
- Blandford, R. D., & Romani, R. W. 1988, *MNRAS*, 234, 57P
- Blumenthal, G. R., & Gould, R. J. 1970, *Rev. Mod. Phys.*, 42, 237
- Bogovalov, S. V., & Aharonian, F. A. 2000, *MNRAS*, 313, 504
- Bradt, H., Rappaport, S., & Mayer, W. 1969, *Nature*, 222, 728
- Browning, R., Ramsden, D., & Wright, P. J. 1971, *Nature Physical Science*, 232, 99
- Burnham, K. P., & Anderson, D. R. 2002, *Model selection and multimodel inference: a practical information-theoretic approach*, 2nd edn. (Springer)
- Burnight, T. 1949, *Phys. Rev*, 76, 19
- Caballero, I., & Wilms, J. 2012, *Mem. Soc. Astron. Italiana*, 83, 230
- Carroll, B. W., & Ostlie, D. A. 2006, *An Introduction to Modern Astrophysics*, 2nd edn. (Benjamin Cummings)

- Cash, W. 1979, *ApJ*, 228, 939
- Castelletti, G., Dubner, G., Golap, K., & Goss, W. M. 2006, *A&A*, 459, 535
- Chandrasekhar, S. 1931, *ApJ*, 74, 81
- Chaves, R. C. G., Renaud, M., Lemoine-Goumard, M., & Goret, P. 2008, in *American Institute of Physics Conference Series*, Vol. 1085, American Institute of Physics Conference Series, ed. F. A. Aharonian, W. Hofmann, & F. Rieger, 372–375
- Chen, A. W., Piano, G., Tavani, M., et al. 2011, *A&A*, 525, A33
- Cocke, W. J., Disney, M. J., & Taylor, D. J. 1969, *Nature*, 221, 525
- Critchfield, C. L., Ney, E. P., & Oleksa, S. 1952, *Physical Review*, 85, 461
- Dame, T. M. 2011, *ArXiv e-prints*
- de Geus, E. J., Bronfman, L., & Thaddeus, P. 1990, *A&A*, 231, 137
- de los Reyes, R., Zajczyk, A., Chaves, R. C. G., & for the H. E. S. S. collaboration. 2012, *ArXiv e-prints*
- de Naurois, M., & H.E.S.S. Collaboration. 2013, *Advances in Space Research*, 51, 258
- Demorest, P. B., Pennucci, T., Ransom, S. M., Roberts, M. S. E., & Hessels, J. W. T. 2010, *Nature*, 467, 1081
- Djannati-Ataï, A., de Jager, O. C., Terrier, R., & et al. 2008, in *International Cosmic Ray Conference*, Vol. 2, International Cosmic Ray Conference, 823–826
- Eadie, W. T., Drijard, D., & James, F. E. 1971, *Statistical methods in experimental physics* (North-Holland Pub. Co.)
- Espinoza, C. M., Lyne, A. G., Kramer, M., Manchester, R. N., & Kaspi, V. M. 2011, *ApJ*, 741, L13
- Esposito, J. A., Bertsch, D. L., Chen, A. W., et al. 1999, *ApJS*, 123, 203

- Falanga, M., Kuiper, L., Poutanen, J., et al. 2005, *A&A*, 444, 15
- Feenberg, E., & Primakoff, H. 1948, *Phys. Rev.*, 73, 449
- Fichtel, C. E., Hartman, R. C., Kniffen, D. A., et al. 1975, *ApJ*, 198, 163
- Fisher, R. A. 1925, *Statistical Methods for Research Workers* (Edinburgh: Oliver and Boyd)
- Fritz, G., Henry, R. C., Meekins, J. F., Chubb, T. A., & Friedman, H. 1969, *Science*, 164, 709
- Funk, S., Hinton, J. A., Pühlhofer, G., et al. 2007, *ApJ*, 662, 517
- Gaensler, B. M., Schulz, N. S., Kaspi, V. M., Pivovarov, M. J., & Becker, W. E. 2003, *ApJ*, 588, 441
- Gaensler, B. M., & Slane, P. O. 2006, *ARA&A*, 44, 17
- Gaisser, T. K., Protheroe, R. J., & Stanev, T. 1998, *ApJ*, 492, 219
- Gelfand, J. D., Slane, P. O., & Zhang, W. 2009, *ApJ*, 703, 2051
- Giordano, F., & Fermi LAT Collaboration. 2011, in *High-Energy Emission from Pulsars and their Systems*, ed. D. F. Torres & N. Rea, 69
- Gold, T. 1968, *Nature*, 218, 731
- Goldreich, P., & Julian, W. H. 1969, *ApJ*, 157, 869
- Górski, K. M., Hivon, E., Banday, A. J., et al. 2005, *ApJ*, 622, 759
- Gotthelf, E. V., & Halpern, J. P. 2008, *ApJ*, 681, 515
- Grenier, I. A., Casandjian, J.-M., & Terrier, R. 2005, *Science*, 307, 1292
- Grondin, M.-H., Funk, S., Lemoine-Goumard, M., et al. 2011, *ApJ*, 738, 42
- Gunn, J. E., & Ostriker, J. P. 1969, *Nature*, 221, 454

- H. E. S. S. Collaboration, Abramowski, A., Acero, F., et al. 2012, *A&A*, 541, A5
- Hall, T. A., Bond, I. H., Bradbury, S. M., et al. 2003, *ApJ*, 583, 853
- Hartman, R. C., Bertsch, D. L., Bloom, S. D., et al. 1999, *ApJS*, 123, 79
- Haug, E. 1975, *Zeitschrift Naturforschung Teil A*, 30, 1099
- Hayakawa, S. 1952, *Progress of Theoretical Physics*, 8, 571
- Herschel, W. 1800, *Philosophical Transactions of the Royal Society of London*, 90, pp. 284
- H.E.S.S. Collaboration, Abramowski, A., Acero, F., et al. 2011a, *A&A*, 531, A81
- . 2011b, *A&A*, 528, A143
- H.E.S.S. Collaboration, Acero, F., Aharonian, F., et al. 2011c, *A&A*, 525, A45
- H.E.S.S. Collaboration, Abramowski, A., Acero, F., et al. 2011d, *A&A*, 533, A103
- . 2011e, *A&A*, 525, A46
- Hewish, A., Bell, S. J., Pilkington, J. D. H., Scott, P. F., & Collins, R. A. 1968, *Nature*, 217, 709
- Hewitt, J., Grondin, M.-H., Lemoine-Goumard, M., et al. 2012
- Hofverberg, P. 2011, in *International Cosmic Ray Conference*, Vol. 7, International Cosmic Ray Conference, 247
- Hoppe, S. 2008, in *International Cosmic Ray Conference*, Vol. 2, International Cosmic Ray Conference, 579–582
- Houck, J. C., & Allen, G. E. 2006, *ApJS*, 167, 26
- Hulsizer, R. I., & Rossi, B. 1948, *Phys. Rev.*, 73, 1402
- Hutchinson, G. 1952, *Philosophical Magazine Series 7*, 43, 847

- Hwang, U., Petre, R., & Flanagan, K. A. 2008, *ApJ*, 676, 378
- James, F., & Roos, M. 1975, *Computer Physics Communications*, 10, 343
- Jansky, K. 1933, *Proceedings of the Institute of Radio Engineers*, 21, 1387
- Kargaltsev, O., Pavlov, G. G., & Wong, J. A. 2008, *ArXiv e-prints*
- Kaspi, V. M., & Helfand, D. J. 2002, in *Astronomical Society of the Pacific Conference Series*, Vol. 271, *Neutron Stars in Supernova Remnants*, ed. P. O. Slane & B. M. Gaensler, 3
- Katagiri, H., Tibaldo, L., Ballet, J., et al. 2011, *ApJ*, 741, 44
- Katsuta, J., Uchiyama, Y., Tanaka, T., et al. 2012
- Kennel, C. F., & Coroniti, F. V. 1984, *ApJ*, 283, 710
- Kerr, M. 2010, PhD thesis, University of Washington
- King, I. 1962, *AJ*, 67, 471
- Klein, O., & Nishina, T. 1929, *Zeitschrift fur Physik*, 52, 853
- Kniffen, D. A., & Fichtel, C. E. 1970, *ApJ*, 161, L157
- Krause, J., Carmona, E., Reichardt, I., & for the MAGIC Collaboration. 2011, *ArXiv e-prints*
- Kraushaar, W., Clark, G. W., Garmire, G., et al. 1965, *ApJ*, 141, 845
- Kraushaar, W. L., Clark, G. W., Garmire, G. P., et al. 1972, *ApJ*, 177, 341
- Lande, J., Ackermann, M., Allafort, A., et al. 2012
- Landi, R., de Rosa, A., Dean, A. J., et al. 2007a, *MNRAS*, 380, 926
- Landi, R., Masetti, N., Bassani, L., et al. 2007b, *The Astronomer's Telegram*, 1047,

- Large, M. I., Vaughan, A. E., & Mills, B. Y. 1968, *Nature*, 220, 340
- Lemoine-Goumard, M., Ferrara, E., Grondin, M.-H., Martin, P., & Renaud, M. 2011, *Mem. Soc. Astron. Italiana*, 82, 739
- Li, T.-P., & Ma, Y.-Q. 1983, *ApJ*, 272, 317
- Longair, M. S. 2011, *High Energy Astrophysics*, 3rd edn. (The Edinburgh Building, Cambridge CB2 8RU, UK: Cambridge University Press)
- Markwardt, C. B., & Ogelman, H. 1995, *Nature*, 375, 40
- Matsumoto, H., Ueno, M., Bamba, A., et al. 2007, *PASJ*, 59, 199
- Matsumoto, H., Uchiyama, H., Sawada, M., et al. 2008, *PASJ*, 60, 163
- Mattox, J. R., Bertsch, D. L., Fichtel, C. E., et al. 1992, *ApJ*, 401, L23
- Mattox, J. R., Bertsch, D. L., Chiang, J., et al. 1996, *ApJ*, 461, 396
- Mayer-Hasselwander, H. A., Kanbach, G., Bennett, K., et al. 1982, *A&A*, 105, 164
- McArthur, S. 2011, *ArXiv e-prints*
- McK Mahille, J., Schild, R., Wendorf, F., & Brenner, R. 2007, *African Skies*, 11, 2
- Mizukami, T., Kubo, H., Yoshida, T., et al. 2011, *ApJ*, 740, 78
- Morrison, P. 1958, *Il Nuovo Cimento*, 7, 858
- Murphy, T., Mauch, T., Green, A., et al. 2007, *MNRAS*, 382, 382
- Neronov, A., Semikoz, D. V., Tinyakov, P. G., & Tkachev, I. I. 2011, *A&A*, 526, A90
- Nolan, P. L., Arzoumanian, Z., Bertsch, D. L., et al. 1993, *ApJ*, 409, 697
- Nolan, P. L., Fierro, J. M., Lin, Y. C., et al. 1996, *A&AS*, 120, C61
- Nolan, P. L., Abdo, A. A., Ackermann, M., et al. 2012, *ApJS*, 199, 31

Pacini, F. 1967, *Nature*, 216, 567

—. 1968, *Nature*, 219, 145

Pacini, F., & Salvati, M. 1973, *ApJ*, 186, 249

Paron, S., Dubner, G., Reynoso, E., & Rubio, M. 2008, *A&A*, 480, 439

Petre, R., Becker, C. M., & Winkler, P. F. 1996, *The Astrophysical Journal Letters*, 465, L43

Pollock, A. M. T. 1985, *A&A*, 150, 339

Protasov, R., van Dyk, D. A., Connors, A., Kashyap, V. L., & Siemiginowska, A. 2002, *ApJ*, 571, 545

Rea, N., & Esposito, P. 2011, in *High-Energy Emission from Pulsars and their Systems*, ed. D. F. Torres & N. Rea, 247

Rees, M. J., & Gunn, J. E. 1974, *MNRAS*, 167, 1

Renaud, M., Goret, P., & Chaves, R. C. G. 2008, in *American Institute of Physics Conference Series*, Vol. 1085, *American Institute of Physics Conference Series*, ed. F. A. Aharonian, W. Hofmann, & F. Rieger, 281–284

Reynoso, E. M., Dubner, G. M., Goss, W. M., & Arnal, E. M. 1995, *AJ*, 110, 318

Reynoso, E. M., Green, A. J., Johnston, S., et al. 2003, *MNRAS*, 345, 671

Richards, D. W., & Comella, J. M. 1969, *Nature*, 222, 551

Rodriguez, J., Tomsick, J. A., Foschini, L., et al. 2003, *A&A*, 407, L41

Rousseau, R., Grondin, M.-H., Van Etten, A., et al. 2012a, *A&A*, 544, A3

—. 2012b, *A&A*, 544, A3

- Rowell, G., Horns, D., Fukui, Y., & Moriguchi, Y. 2008, in American Institute of Physics Conference Series, Vol. 1085, American Institute of Physics Conference Series, ed. F. A. Aharonian, W. Hofmann, & F. Rieger, 241–244
- Rybicki, G. B., & Lightman, A. P. 1979, Radiative processes in astrophysics (New York: John Wiley & Sons Ltd)
- Schwarz, G. 1978, The Annals of Statistics, 6, pp. 461
- Sheidaei, F. 2011, in International Cosmic Ray Conference, Vol. 7, International Cosmic Ray Conference, 243
- Slane, P., Castro, D., Funk, S., et al. 2010, The Astrophysical Journal, 720, 266
- Slane, P., Castro, D., Funk, S., et al. 2010, ApJ, 720, 266
- Sreekumar, P., Bertsch, D. L., Hartman, R. C., Nolan, P. L., & Thompson, D. J. 1999, Astroparticle Physics, 11, 221
- Sreekumar, P., Bertsch, D. L., Dingus, B. L., et al. 1992, ApJ, 400, L67
- Sreekumar, P., Bertsch, D. L., Dingus, B. L., et al. 1998, The Astrophysical Journal, 494, 523
- Staelin, D. H., & Reifstein, III, E. C. 1968, Science, 162, 1481
- Strong, A. W., & Moskalenko, I. V. 1998, ApJ, 509, 212
- Sugizaki, M., Mitsuda, K., Kaneda, H., et al. 2001, ApJS, 134, 77
- Swanenburg, B. N., Hermsen, W., Bennett, K., et al. 1978, Nature, 275, 298
- Swanenburg, B. N., Bennett, K., Bignami, G. F., et al. 1981, ApJ, 243, L69
- Tam, P. H. T., Wagner, S. J., Tibolla, O., & Chaves, R. C. G. 2010, A&A, 518, A8
- Tanaka, T., Allafort, A., Ballet, J., et al. 2011, ApJ, 740, L51
- Taylor, A. R., Gibson, S. J., Peracaula, M., et al. 2003, AJ, 125, 3145

- Terrier, R., Mattana, F., Djannati-Atai, A., et al. 2008, in American Institute of Physics Conference Series, Vol. 1085, American Institute of Physics Conference Series, ed. F. A. Aharonian, W. Hofmann, & F. Rieger, 312–315
- Thompson, D. J. 2008, Reports on Progress in Physics, 71, 116901
- Thompson, D. J., Fichtel, C. E., Hartman, R. C., Kniffen, D. A., & Lamb, R. C. 1977a, ApJ, 213, 252
- Thompson, D. J., Fichtel, C. E., Kniffen, D. A., & Ogelman, H. B. 1977b, ApJ, 214, L17
- Thompson, D. J., Bertsch, D. L., Fichtel, C. E., et al. 1993, ApJS, 86, 629
- Tomsick, J. A., Lingenfelter, R., Walter, R., et al. 2003, IAU Circ., 8076, 1
- Torii, K., Kinugasa, K., Toneri, T., et al. 1998, ApJ, 494, L207
- van der Swaluw, E., & Wu, Y. 2001, ApJ, 555, L49
- Vladimirov, A. E., Digel, S. W., Jóhannesson, G., et al. 2011, Computer Physics Communications, 182, 1156
- Weekes, T. C., Cawley, M. F., Fegan, D. J., et al. 1989, ApJ, 342, 379
- Weinstein, A., & for the VERITAS Collaboration. 2009, ArXiv e-prints
- Young, E. T., Lada, C. J., & Wilking, B. A. 1986, ApJ, 304, L45