

Search for Spatially Extended *Fermi*-LAT Sources Using Two Years of Data

J. Lande^{1,2}, M. Ackermann^{3,4}, A. Allafort¹, J. Ballet⁵, K. Bechtol¹, T. H. Burnett⁶,
J. Cohen-Tanugi⁷, A. Drlica-Wagner¹, S. Funk^{1,8}, F. Giordano^{9,10}, M.-H. Grondin^{11,12},
M. Kerr¹, M. Lemoine-Goumard^{13,14}

ABSTRACT

Spatial extension is an important characteristic for correctly associating γ -ray-emitting sources with their counterparts at other wavelengths and for obtaining an unbiased model of their spectra. We present a new method for quantifying the spatial extension of sources detected by the Large Area Telescope (LAT), the primary science instrument on the *Fermi Gamma-ray Space Telescope (Fermi)*. We perform a series of Monte Carlo simulations to validate this tool and calculate

¹W. W. Hansen Experimental Physics Laboratory, Kavli Institute for Particle Astrophysics and Cosmology, Department of Physics and SLAC National Accelerator Laboratory, Stanford University, Stanford, CA 94305, USA

²email: joshualande@gmail.com

³Deutsches Elektronen Synchrotron DESY, D-15738 Zeuthen, Germany

⁴email: markus.ackermann@desy.de

⁵Laboratoire AIM, CEA-IRFU/CNRS/Université Paris Diderot, Service d'Astrophysique, CEA Saclay, 91191 Gif sur Yvette, France

⁶Department of Physics, University of Washington, Seattle, WA 98195-1560, USA

⁷Laboratoire Univers et Particules de Montpellier, Université Montpellier 2, CNRS/IN2P3, Montpellier, France

⁸email: funk@slac.stanford.edu

⁹Dipartimento di Fisica “M. Merlin” dell’Università e del Politecnico di Bari, I-70126 Bari, Italy

¹⁰Istituto Nazionale di Fisica Nucleare, Sezione di Bari, 70126 Bari, Italy

¹¹Max-Planck-Institut für Kernphysik, D-69029 Heidelberg, Germany

¹²Landessternwarte, Universität Heidelberg, Königstuhl, D 69117 Heidelberg, Germany

¹³Université Bordeaux 1, CNRS/IN2p3, Centre d’Études Nucléaires de Bordeaux Gradignan, 33175 Gradignan, France

¹⁴Funded by contract ERC-StG-259391 from the European Community

the LAT threshold for detecting the spatial extension of sources. We then test all sources in the second *Fermi*-LAT catalog (2FGL) for extension. We report the detection of seven new spatially extended sources.

Subject headings: Catalogs; Fermi Gamma-ray Space Telescope; Gamma rays; observations; ISM: supernova remnants; Methods: statistical; pulsar wind nebula

1. Introduction

A number of astrophysical source classes including supernova remnants (SNRs), pulsar wind nebulae (PWNe), molecular clouds, normal galaxies, and galaxy clusters are expected to be spatially resolvable by the Large Area Telescope (LAT), the primary instrument on the *Fermi Gamma-ray Space Telescope* (*Fermi*). Additionally, dark matter satellites are also hypothesized to be spatially extended. See Atwood et al. (2009a) for pre-launch predictions. The LAT has detected seven SNRs which are significantly extended at GeV energies: W51C, W30, IC 443, W28, W44, RX J1713.7–3946, and the Cygnus Loop (Abdo et al. 2009a, 2011a, 2010i,e,h, 2011c; Katagiri et al. 2011). In addition, three extended PWN have been detected by the LAT: MSH 15–52, Vela X, and HESS J1825–137 (Abdo et al. 2010a,f; Grondin et al. 2011). Two nearby galaxies, the Large and Small Magellanic Clouds, and the lobes of one radio galaxy, Centaurus A, were spatially resolved at GeV energies (Abdo et al. 2010j,b,c). A number of additional sources detected at GeV energies are positionally coincident with sources that exhibit large enough extension at other wavelengths to be spatially resolvable by the LAT at GeV energies. In particular, there are 59 GeV sources in the second Fermi Source Catalog (2FGL) that might be associated with extended SNRs (2FGL, Abdo et al. 2011b). Previous analyses of extended LAT sources were performed as dedicated studies of individual sources so we expect that a systematic scan of all LAT detected sources could uncover additional spatially extended sources.

The current generation of air Cherenkov detectors have made it apparent that many sources can be spatially resolved at even higher energies. Most prominent was a survey of the Galactic plane using the High Energy Stereoscopic System (H.E.S.S) which reported 14 spatially extended sources with extensions varying from $\sim 0^\circ.1$ to $\sim 0^\circ.25$ (Aharonian et al. 2006). In fact, within our Galaxy only very few sources detected at TeV energies, most notably the γ -ray binaries LS 5039 (Aharonian et al. 2006a), LS I+61–303 (Albert et al. 2006; Acciari et al. 2011), HESS J0632+057 (Aharonian et al. 2007c), and the Crab nebula (Weekes et al. 1989), have no detectable extension. High-energy γ -rays from TeV sources are produced by the decay of π^0 s produced by hadronic interactions with interstellar matter and by relativistic electrons due to Inverse Compton (IC) scattering and bremsstrahlung

radiation. It is plausible that the GeV and TeV emission from these sources originates from the same population of high-energy particles and so at least some of these sources should be detectable at GeV energies. Studying these TeV sources at GeV energies would help to determine the emission mechanisms producing these high energy photons.

The LAT is a pair conversion telescope that has been surveying the γ -ray sky since 2008 August. The LAT has broad energy coverage (20 MeV to > 300 GeV), wide field of view (~ 2.4 sr), and large effective area (~ 8000 cm² at > 1 GeV) Additional information about the performance of the LAT can be found in Atwood et al. (2009b).

Using 2 years of all-sky survey data, the LAT Collaboration published 2FGL (2FGL, Abdo et al. 2011b). The possible counterparts of many of these sources can be spatially resolved when observed at other frequencies but detecting the spatial extension of these sources at GeV energies is difficult because the size of the point-spread function (PSF) of the LAT is comparable to the typical size of many of these sources.

The capability to spatially resolve GeV γ -ray sources is important for several reasons. Finding a coherent source extension across different energy bands can help to associate a LAT source to an otherwise confused counterpart. Furthermore, γ -rays emission from dark matter annihilation has been predicted to be detectable by the LAT. Some of the dark matter substructure in our Galaxy could be spatially resolvable by the LAT (Baltz et al. 2008). Characterization of spatial extension could help to identify this substructure. Also, due to the strong energy dependence of the LAT PSF, the spatial and spectral characterization of a source cannot be decoupled. An inaccurate spatial model will bias the spectral model of the source and vice versa. Specifically, modeling a spatially extended source as point-like will systematically soften measured spectra. Furthermore, correctly modeling source extension is important for understanding an entire region of the sky. For example, an imperfect model of the spatially extended LMC introduced significant residuals in the surrounding region (Abdo et al. 2010d, 2011b). Such residuals can bias the significance and measured spectra of neighboring sources in the densely populated Galactic plane.

For these reasons, in Section 2 we present a new systematic method for analyzing spatially extended LAT sources. **In Section 3.1, we demonstrate that this method can be used to test the statistical significance of the extension of a LAT source and in Section 3.2 we asses the bias that is introduced by using an incorrect spatial model. In Section 4 we calculate the LAT detection threshold to resolve the extension of a source.** In Section 5, we study the ability of the LAT to distinguish between a single extended source and unresolved closely-spaced point-like sources In Section 6, we further demonstrate that our detection method does not misidentify point-like sources as being extended by testing the extension of active Galactic nuclei (AGN) believed to be

unresolvable. In Section 7, we systematically reanalyze the twelve extended sources included in the 2FGL catalog and in Section 8 we describe a way to estimate systematic errors on the measured extension of a source. In Section 9, we describe a search for new spatially extended LAT sources. Finally, in Section 10 we present the detection of the extension of nine spatially extended sources that were reported in the 2FGL catalog but treated as point-like in the analysis. Two of these sources have been previously analyzed in dedicated publications.

2. Analysis Methods

Morphological studies of sources using the LAT are challenging because of the strongly energy-dependent PSF that is comparable in size to the extension of many sources expected to be detected at GeV energies. Additional complications arise for sources along the Galactic plane due to systematic uncertainties in the model for Galactic diffuse emission.

For energies below ~ 300 MeV, the angular resolution is limited by multiple scattering in the silicon strip tracking section of the detector and is several degrees at 100 MeV. The PSF improves with energy approaching a 68% containment radius of $\sim 0.2^\circ$ at the highest energies (when averaged over the acceptance of the LAT) and is limited by the ratio of the strip pitch to the height of the tracker (Atwood et al. 2009b; Abdo et al. 2009d, 2012).¹ However, since most high energy astrophysical sources have spectra that decrease rapidly with increasing energy, there are typically fewer higher energy photons with improved angular resolution. Therefore sophisticated analysis techniques are required to maximize the sensitivity of the LAT to extended sources.

2.1. Modeling Extended Sources in the pointlike Package

A new maximum-likelihood analysis tool has been developed to address the unique requirements for studying spatially extended sources with the LAT. It works by maximizing the Poisson likelihood to detect the observed distributions of γ -rays (referred to as counts) given a parametrized spatial and spectral model of the sky. The data are binned spatially, using a HEALPix pixellization, and spectrally (Górski et al. 2005) and the likelihood is maximized over all bins in a region. The extension of a source can be modeled by a geometric shape (e.g. a disk or a two-dimensional Gaussian) and the position, extension, and spectrum

¹More information about the performance of the LAT can be found at the *Fermi* Science Support Center (FSSC, <http://fermi.gsfc.nasa.gov>).

of the source can be simultaneously fit.

This type of analysis is not feasible using the standard LAT likelihood analysis tool `gtlike`² because it can only fit the spectral parameters of the model unless a more sophisticated iterative procedure is used. We note that `gtlike` has been used in the past in several studies of source extension in the LAT Collaboration (Abdo et al. 2010j,b,e, 2009a). In these studies, a set of `gtlike` maximum likelihood fits at fixed extensions was used to build a profile of the likelihood as a function of extension. The `gtlike` likelihood profile approach has been shown to correctly reproduce the extension of simulated extended sources assuming that the true position is known (Giordano 2011). But it is not optimal because the position, extension, and spectrum of the source must be simultaneously fit to find the best fit parameters and to maximize the statistical significance of the detection. Furthermore, because the `gtlike` approach is computationally intensive, no large-scale Monte Carlo simulations have been run to calculate its false detection rate.

The approach presented here is based on a second maximum likelihood fitting package developed in the LAT Collaboration called `pointlike` (Abdo et al. 2010d; Kerr 2011). The choice to base the spatial extension fitting on `pointlike` rather than `gtlike` was made due to considerations of computing time. The `pointlike` algorithm was optimized for speed to handle larger numbers of sources efficiently, which is important for our catalog scan and for being able to perform large-scale Monte Carlo simulations to validate the analysis. Details on the `pointlike` package can be found in Kerr (2011). We extended the code to allow a simultaneous fit of the source extension together with the position and the spectral parameters.

2.2. Extension Fitting

In `pointlike`, one can fit the position and extension of a source under the assumption that the source model can be factorized: $M(x, y, E) = S(x, y) \times X(E)$, where $S(x, y)$ is the spatial distribution and $X(E)$ is the spectral distribution. To fit an extended source, `pointlike` convolves the extended source shape with the PSF (as a function of energy) and uses the `minuit` library (James & Roos 1975) to maximize the likelihood by simultaneously varying the position, extension, and spectrum of the source. As will be described in Section 3.1, simultaneously fitting the position, extension, and spectrum is important to maximize the statistical significance of the detection of the extension of a source. To avoid projection effects, the longitude and latitude of the source are not directly fit but instead

²`gtlike` is distributed publicly by the FSSC.

135 the displacement of the source in a reference frame centered on the source.

The significance of the extension of a source can be calculated from the likelihood ratio test of a hypothesis with a spatially extended source and a hypothesis with a point-like source. The test statistic (TS) for this procedure is defined as

$$\text{TS}_{\text{ext}} = 2 \log(\mathcal{L}_{\text{ext}}/\mathcal{L}_{\text{ps}}) \quad (1)$$

136 where \mathcal{L} is the Poisson likelihood. `pointlike` calculates TS_{ext} by fitting a source first with
137 a spatially extended model and then as a point-like source. The interpretation of TS_{ext} in
138 terms of a statistical significance is discussed in Section 3.1.

For extended sources with an assumed radially-symmetric shape, we optimized the calculation by performing one of the integrals analytically. The expected photon distribution can be written as

$$\text{PDF}(\vec{r}) = \int \text{PSF}(|\vec{r} - \vec{r}'|) I_{\text{src}}(\vec{r}') r' dr' d\phi' \quad (2)$$

where \vec{r} represents the position in the sky and $I_{\text{src}}(\vec{r})$ is the spatial distribution of the source. The PSF of the LAT is currently parameterized in the Pass 7_V6 (P7_V6) Source Instrument Response Function (IRFs, Abdo et al. 2012) by a King function (King 1962):

$$\text{PSF}(r) = \frac{1}{2\pi\sigma^2} \left(1 - \frac{1}{\gamma}\right) \left(1 + \frac{u}{\gamma}\right)^{-\gamma}, \quad (3)$$

where $u = (r/\sigma)^2/2$ and σ and γ are free parameters (Kerr 2011). For radially-symmetric extended sources, the angular part of the integral can be evaluated analytically

$$\begin{aligned} \text{PDF}(u) &= \int_0^\infty r' dr' I_{\text{src}}(v) \int_0^{2\pi} d\phi' \text{PSF}(\sqrt{2\sigma^2(u+v-2\sqrt{uv}\cos(\phi-\phi'))}) \\ &= \int_0^\infty dv I_{\text{src}}(v) \left(\frac{\gamma-1}{\gamma}\right) \left(\frac{\gamma}{\gamma+u+v}\right)^\gamma \times {}_2F_1\left(\gamma/2, \frac{1+\gamma}{2}, 1, \frac{4uv}{(\gamma+u+v)^2}\right), \end{aligned} \quad (4)$$

139 where $v = (r'/\sigma)^2/2$ and ${}_2F_1$ is the Gaussian hypergeometric function. This convolution
140 formula reduces the expected photon distribution to a single numerical integral.

141 There will always be a small numerical discrepancy between the expected photon dis-
142 tribution derived from a true point-like source and a very small extended source due to
143 numerical error in the convolution. In most situations, this error is insignificant. But in
144 particular for very bright sources, this numerical error has the potential to bias the TS for
145 the extension test. Therefore, when calculating TS_{ext} , we compare the likelihood fitting the
146 source with an extended spatial model to the likelihood when the extension is fixed to a very
147 small value (10^{-10} degrees in radius for a uniform disk model).

We estimate the error on the extension of a source by fixing the position of the source and varying the extension until the log of the likelihood has decreased by $1/2$, corresponding to a 1σ error (Eadie et al. 1971). Figure 1 demonstrates this method by showing the change in the log of the likelihood when varying the modeled extension of the SNR IC 443. The localization error is calculated by fixing the extension and spectrum of the source to their best fit values and then fitting the log of the likelihood to a 2D Gaussian as a function of position. This localization error algorithm is further described in Abdo et al. (2011b).

2.3. gtlike Analysis Validation

`pointlike` is important for LAT analyses that require many iterations such as source localization and extension fitting. On the other hand, because `gtlike` makes fewer approximations in calculating the likelihood we expect the spectral parameters found with `gtlike` to be slightly more accurate. Furthermore, because `gtlike` is the standard likelihood analysis package for LAT data, it has been more extensively validated for spectral analysis. For those reasons, in the following analysis we used `pointlike` to determine the position and extension of a source and subsequently derived the spectrum using `gtlike`. Both `gtlike` and `pointlike` can be used to estimate the statistical significance of the extension of a source and we required that both methods agree for a source to be considered extended. There was good agreement between the two methods. Unless explicitly mentioned, all TS, TS_{ext} , and spectral parameters were calculated using `gtlike` with the best-fit positions and extension found by `pointlike`.

2.4. Comparing Source Sizes

We considered two models for the surface brightness profile for extended sources: a 2D Gaussian model

$$I_{\text{Gaussian}}(x, y) = \frac{1}{2\pi\sigma^2} \exp(-(x^2 + y^2)/2\sigma^2) \quad (6)$$

or a uniform disk model

$$I_{\text{disk}}(x, y) = \begin{cases} \frac{1}{\pi\sigma^2} & x^2 + y^2 \leq \sigma^2 \\ 0 & x^2 + y^2 > \sigma^2. \end{cases} \quad (7)$$

Although these shapes are significantly different, Figure 2 shows that, after convolution with the LAT PSF, their PDFs are similar for a source that has a 0.5° radius typical of LAT detected extended sources. To allow a valid comparison between the Gaussian and the uniform disk models, we define the source size as the radius containing 68% of the intensity

(r_{68}). **By direct integration, we find**

$$r_{68,\text{Gaussian}} = 1.51\sigma, \quad (8)$$

$$r_{68,\text{disk}} = 0.82\sigma, \quad (9)$$

where σ is defined in Equation 6 and Equation 7 respectively. For the example above, $r_{68} = 0.5$ so $\sigma_{\text{disk}} = 0.61^\circ$ and $\sigma_{\text{Gaussian}} = 0.33^\circ$.

For sources that are comparable in size to the PSF, the differences in the PDF for different spatial models are lost in the noise and the LAT is not sensitive to the detailed spatial structure of these sources. **In section 3.2, we perform a dedicated Monte Carlo simulation to asses possible bias due to incorrectly modeling the spatial structure of an extended source. In our search for extended sources we use only a radially-symmetric uniform disk spatial model. Unless otherwise noted, we quote the radius to the edge (σ) as the size of the source.**

3. Validation of the Analysis Method

3.1. Statistical Significance Testing

We evaluated the false detection probability for the measurement of source extent by fitting the extensions of point-like sources. Mattox et al. (1996) discuss that the TS distribution for a likelihood ratio test on the existence of a source at a given position is

$$P(\text{TS}) = \frac{1}{2}(\chi_1^2(\text{TS}) + \delta(\text{TS})), \quad (10)$$

where $P(\text{TS})$ is the probability of getting a particular value of TS, χ_1^2 is the chi-squared distribution with one degree of freedom, and δ is the Dirac delta function. The particular form of Equation 10 is due to the null hypothesis (source flux $\Phi = 0$) residing on the edge of parameter space and the model hypothesis adding a single degree of freedom. It leads to the often quoted result $\sqrt{TS} = \sigma$, where σ here refers to the significance of the detection. It is plausible to expect a similar distribution for the TS in the test for source extension since the same conditions apply (with the source flux Φ replaced by the source radius r and $r < 0$ being unphysical). To verify Equation 10, we evaluated the empirical distribution function of TS_{ext} computed from simulated sources.

We simulated point-like sources with various spectral forms using the LAT on-orbit simulation tool `gtobssim`³ and fit the sources with `pointlike` using both point-like and ex-

³`gtobssim` is distributed publicly by the FSSC.

tended source hypotheses. These point-like sources were simulated with a power-law spectral model with integrated fluxes above 100 MeV ranging from 3×10^{-9} to 1×10^{-5} $\text{ph cm}^{-2} \text{s}^{-1}$ and spectral indices ranging from 1.5 to 3. These values were picked to represent typical parameters of LAT-detected sources. The point-like sources were simulated on top of an isotropic background with a power-law spectral model with integrated flux above 100 MeV of 1.5×10^{-5} $\text{ph cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$ and spectral index 2.1. This was taken to be the same as the isotropic spectrum measured by EGRET (Sreekumar et al. 1998). This spectrum is comparable to the high-latitude background intensity seen by the LAT. The Monte Carlo simulation was performed over a one-year observation period using a representative spacecraft orbit and livetime. The reconstruction was performed using the P7_V6 Source class event selection and IRFs (Abdo et al. 2012). For each significantly detected point-like source ($\text{TS} \geq 25$), we used `pointlike` to fit the source as an extended source and calculate TS_{ext} . This entire procedure was performed twice, once fitting in the 1 GeV to 100 GeV energy range and once fitting in the 10 GeV to 100 GeV energy range.

For each set of spectral parameters, $\sim 20,000$ statistically independent simulations were performed. For dimmer spectral models, many of the simulations left the source insignificant ($\text{TS} < 25$) and were discarded. Table 1 shows the different spectral models used in our study as well as the number of simulations and the average point-like source significance. The cumulative density of TS_{ext} is plotted in Figures 3 and 4 and compared to the $\chi^2_1/2$ distribution of Equation 10.

Our study shows broad agreement between simulations and Equation 10. To the extent that there is a discrepancy, the simulations tended to produce smaller than expected values of TS_{ext} which would make the formal significance conservative. Considering the distribution in Figures 3 and 4, the choice of a threshold TS_{ext} set to 16 (corresponding to a formal 4σ significance) is reasonable.

These simulations spanned the range of detectable fluxes over two different energy ranges, but only assuming an isotropic background. We performed a second simulation to assess any effects that could be introduced by performing this test in the presence of a highly structured background. Our simulation setup was the same as above except that the sources were simulated on top of and analyzed assuming the presence of the standard Galactic and isotropic diffuse background model used in 2FGL.

In our simulation, we selected our sources to have a random position on the sky such that the absolute value of the galactic latitude of the source is less than 5° , probing the most structured areas of the galaxy.

To limit the number of tests, we selected only one flux level for each of the four spectral indices and we perform this test only in the 1 GeV to 100 GeV energy range. The source flux was selected so that $\text{TS} \sim 50$. This is around the significance level for which one might expect to be able to spatially resolve the source. We do not test brighter sources since the structured background should produce the most significant effect for very bright sources.

Because the Galactic diffuse emission is highly structured, the point-source sensitivity can vary significantly across the galactic plane. To account for this, we scaled the flux (for a given spectral index) so that the source always has approximately the same signal-to-noise ratio:

$$F(\vec{x}) = F(\text{GC}) \times \left(\frac{B(\vec{x})}{B(\text{GC})} \right)^{1/2}. \quad (11)$$

Here, \vec{x} is the position of the source, F is the integral flux of the simulated source from 100 MeV to 100 GeV, $F(\text{GC})$ is the same quantity if the source were at the galactic center, B is the integral of the Galactic diffuse and isotropic diffuse emission model from 1 GeV to 100 GeV and $B(\text{GC})$ is the same quantity if the source were at the galactic center. Table 1 tabulates $F(\text{GC})$ and the average TS for the four spectral models. For each spectral index, we performed $\sim 90,000$ simulations.

Figure 5 shows the cumulative density of TS_{ext} compared to the $\chi^2_1/2$ distribution of Equation 10. For small values of TS_{ext} , there is reasonable agreement between the Monte Carlo simulation and theory. At the highest obtained values of TS_{ext} , there is possibly some discrepancy. But in this range there are only limited statistics because the curve is determined by only a few points.

3.2. Assessing the Bias Due to Incorrect Spatial Models

We performed a Monte Carlo study to assess the biases introduced by incorrectly modeling the spatial structure of an extended source. This is important because any assumed spatial model is just an approximation to the true photon distribution of the source.

To assess this systematic, we simulated the spatially extended ring-type SNR W44. W44 was selected because it is the most significant extended source detected by the LAT that has been reported as having a non-radially symmetric photon-distribution Abdo et al. (2010h).

W44 was simulated with a power-law spectral model. The flux and spectral index of this source were taken from our re-analysis in this paper which is described in Section 7 and summarized in Table 3. In particular W44 was simulated with a flux of $7.12 \times 10^{-8} \text{ ph cm}^{-2} \text{ s}^{-1}$ in the energy range from 1 GeV to 100 GeV with a spectral index of 2.66.

W44 was simulated with an elliptical ring spatial model described in Abdo et al. (2010h). Because that publication does not provide the numbers describing the parameters of the ellipse, we summarize them here for completeness. The elliptical spatial model has a major axis of $0^\circ 3$ and a minor axis of $0^\circ 19$. The ellipse predicts emission only for the outer 0.75% of the ellipse and the ellipse has an angle of 147° measured east of celestial north.

We used a simulation setup similar to that described in Section 3.1, but the source was simulated on top of and analyzed assuming the presence of the standard galactic and isotropic diffuse background model used in 2FGL. This simulation was in the 1 GeV to 100 GeV energy range and the sources were simulated over the two-year time range included in the 2FGL catalog. We performed 985 identical simulations with different seeds to see the statistical spread in the fit values. In the simulations of W44, we disabled the inherent energy dispersion of photons by the LAT. The bias caused by ignoring energy dispersion will be discussed in an upcoming paper by the LAT collaboration (Abdo et al. 2012).

After simulating the ring-like source, we fit the spectral and spatial parameters of the ring-like extended source with successively more complicated spatial models. We obtained the best fit spatial parameters using pointlike and, using these best fit spatial parameters, obtained the best fit spectral parameters using gtlike. In total, we performed the fit using a point-like spatial model, using a radially-symmetric uniform Gaussian and disk spatial model, an Elliptical disk spatial model, and finally the Monte Carlo true Elliptical ring spatial model.

Figure 6 shows the distribution of TS values for each of the simulations when fitting the simulated sources with the different spatial models. Figure 6a shows the distribution of TS values when fitting the simulated Elliptical ring extended source as as a point-like source. the source is very significant as a point source ($\text{TS} \sim 3500$). Figure 6b shows the distirbution of TS_{ext} values when fitting the source as a radially-symmetric uniform disk or Gaussian source. The source is very significantly extended ($\text{TS}_{\text{ext}} \sim 250$). On average TS_{ext} is slightly larger when fitting the source with the Disk spatial model compared to the Gaussian

spatial model. Considering that the simulated source’s true spatial model has a sharp edge, it makes sense that the Disk would provide a better fit.

Figure 6c then shows the change in TS (or twice the improvement in log likelihood) when fitting the simulated source as an elliptical disk compared to fitting the source as a disk. Since the change in TS is ~ 30 , we see that our simulated source is significantly better described with an elliptical spatial model. Finally, Figure 6d shows the change in TS when fitting the source as an elliptical ring compared to fitting it as an elliptical disk. Since the change is ~ 9 , we see that the source is somewhat better described by the true spatial model, but not at a high statistical significance.

Figure 7 shows the distribution of best fit parameters for these simulations. Figure 7a and Figure 7b show the distributions of best fit flux and spectral indices. They show that when fitting this simulated extended source with a point-like spatial model, there is a $\sim 10\%$ bias in the fit flux and $\sim 5\%$ bias in the fit index. But when fitting the sources with any of the extended spatial models, there is no bias in the best fit spectral parameters (within the precision of this Monte Carlo study. The reason for this is because within the finite angular precision and limited statistics of the LAT, all of the extended spatial models can be adjusted to look almost the same. For brighter or larger sources, this could potentially be a larger effect.

Figure 7c plots the best fit 68% containment radius (r_{68}). For the elliptical ring spatial model, as expected there is no significant bias in the best fit size. What is surprising is that for none of the tested spatial models was the bias greater than $\sim 10\%$. This shows that for whatever spatial model is used, the best fit r_{68} is a reasonable measure of the true 68% containment radius.

For completeness, for the elliptical disk and ring spatial models, Figure 7c plots the distribution of best fit eccentricity of the ellipse. It is defined as $\sqrt{1 - b^2/a^2}$ where a is the length of the semi-major axis and b is the length of the semi-minor axis. Figure 7c plots the distribution of best fit orientation angle of the ellipse. For the elliptical spatial models, there is no systematic bias in the eccentricities or orientation angles. For the true spatial model, there is less spread in the distributions of the fit parameters. Finally, for completeness Figure 7d shows that the distribution of best fit ring fractions is unbiased when the source is fit with the for the elliptical ring spatial model.

4. Extended Source Detection Threshold

We calculated the LAT flux threshold to detect spatial extent. We define the detection threshold as the flux at which the value of TS_{ext} averaged over many statistical realizations is $\langle \text{TS}_{\text{ext}} \rangle = 16$ (corresponding to a formal 4σ significance) for a source of a given extension.

We used a simulation setup similar to that described in Section 3.1, but instead of point-like sources we simulated extended sources with radially-symmetric uniform disk spatial models. Additionally, we simulated our sources over the two-year time range included in the 2FGL catalog. For each extension and spectral index, we selected a flux range which bracketed $\text{TS}_{\text{ext}} = 16$ and performed an extension test for > 100 independent realizations of ten fluxes in the range. We calculated $\langle \text{TS}_{\text{ext}} \rangle = 16$ by fitting a line to the flux and TS_{ext} values in the narrow range.

Figure 8 shows the threshold for sources of four spectral indices from 1.5 to 3 and extensions varying from $\sigma = 0.1$ to 2.0 . The threshold is high for small extensions when the source is small compared to the size of the PSF. It drops quickly with increasing source size and reaches a minimum around 0.5 . The threshold increases for large extended sources because the source becomes increasingly diluted by the background. Figure 8 shows the threshold using photons with energies between 100 MeV and 100 GeV and also using only photons with energies between 1 GeV and 100 GeV. Except for very large or very soft sources, the threshold is not substantially improved by including photons with energies between 100 MeV and 1 GeV. This is also demonstrated in Figure 1 which shows TS_{ext} for the SNR IC 443 computed independently in twelve energy bins between 100 MeV and 100 GeV. For IC 443, which has a spectral index ~ 2.4 and an extension ~ 0.35 , almost the entire increase in likelihood from optimizing the source extent in the model comes from energies above 1 GeV. Furthermore, other systematic errors become increasingly large at low energy. For our extension search (Section 9), we therefore used only photons with energies above 1 GeV.

Figure 9 shows the flux threshold as a function of source extension for different background levels ($1\times$, $10\times$, and $100\times$ the nominal background), different spectral indices, and two different energy ranges (1 GeV to 100 GeV and 10 GeV to 100 GeV). The detection threshold is higher for sources in regions of higher background. When studying sources only at energies above 1 GeV, the LAT detection threshold (defined as the 1 GeV to 100 GeV flux at which $\langle \text{TS}_{\text{ext}} \rangle = 16$) depends less strongly on the spectral index of the source. The index dependence of the detection threshold is even weaker when considering only photons with energies above 10 GeV because the PSF changes little from 10 GeV to 100 GeV. Overlaid on Figure 9 are the LAT detected extended sources that will be discussed in Sections 7 and 10. The extension thresholds are tabulated in Table 2.

Finally, Figure 10 shows the projected detection threshold of the LAT to extension with a 10 year exposure against 10 times the isotropic background measured by EGRET. This background is representative of the background near the Galactic plane. For small extended sources, the threshold improves by a factor larger than the square root of the relative exposures because the LAT is signal-limited at high energies where the present analysis is most sensitive. For large extended sources, the relevant background is over a larger spatial range and so the improvement is closer to a factor corresponding to the square root of the relative exposures that is caused by Poisson fluctuations in the background.

5. Testing Against Source Confusion

It is impossible to discriminate using only LAT data between a spatially extended source and multiple point-like sources separated by angular distances comparable to or smaller than the size of the LAT PSF. To assess the plausibility of source confusion for situations in which $\text{TS}_{\text{ext}} \geq 16$, we developed an algorithm in `pointlike` to simultaneously fit the positions and spectra of two point-like sources.

After simultaneously fitting the two positions and two spectra, we define $\text{TS}_{2\text{pts}}$ as twice the increase in the log of the likelihood fitting the region with two point-like sources compared to fitting the region with one point-like source:

$$\text{TS}_{2\text{pts}} = 2 \log(\mathcal{L}_{2\text{pts}}/\mathcal{L}_{\text{ps}}). \quad (12)$$

For the following analysis of LAT data, $\text{TS}_{2\text{pts}}$ was computed by fitting the spectra of the two point-like sources in `gtlike` using the best fit positions of the sources found by `pointlike`.

$\text{TS}_{2\text{pts}}$ cannot be quantitatively compared to TS_{ext} using a simple likelihood ratio test to evaluate which model is significantly better because the models are not nested (Protassov et al. 2002). Even though the comparison of TS_{ext} with $\text{TS}_{2\text{pts}}$ is not a calibrated test, $\text{TS}_{\text{ext}} > \text{TS}_{2\text{pts}}$ indicates that the likelihood for the extended source hypothesis is higher than for two point-like sources and we only consider a source to be extended if $\text{TS}_{\text{ext}} > \text{TS}_{2\text{pts}}$.

It might be desirable to formulate a Bayesian test comparing an extended source to two point-like sources. We considered the Bayesian information criterion (BIC, Schwarz 1978). We found it difficult to apply to LAT data because it required a term including the number of data points and for LAT events, there are many photons included in the ROI that contribute only marginally when considering different models. We also considered the Akaike information

criterion (AIC, Akaike 1974). The AIC is defines as

$$\text{AIC} = 2k - 2 \log \mathcal{L} \quad (13)$$

where k is the number of parameters in the model. The first term in the AIC penalyze models with additonal parameters. In this formulation, the best model is consided to be the one that minimzies the AIC. Because the two point-like source hypotheses has three more parameters compared to the one extended source hypothesis (two spatial parameters and two spectral parameters compared to one extnesion parameter), we note that the $\text{TS}_{\text{ext}} > \text{TS}_{2\text{pts}}$ is mathematically equivalent to the test that

$$\text{AIC}_{\text{ext}} < \text{AIC}_{2\text{pts}} - 6, \quad (14)$$

which gives our original test an equivalent Bayesian formulation by requiring AIC_{ext} to be sufficently smaller than $\text{AIC}_{2\text{pts}}$.

We assessed the power of the $\text{TS}_{\text{ext}} > \text{TS}_{2\text{pts}}$ test with a Monte Carlo study. We simulated one spatially extended source and fit it as both an extended source and as two point-like sources using `pointlike`. We then simulated two point-like sources and fit them with the same two hypotheses. By comparing the distribution of $\text{TS}_{2\text{pts}}$ and TS_{ext} computed by `pointlike` for the two cases, we evaluated how effective the $\text{TS}_{\text{ext}} > \text{TS}_{2\text{pts}}$ test is at rejecting cases of source confusion as well as how likely it is to incorrectly reject that an extended source is spatially extended. All sources were simulated using the same time range as in Section 4 against a background 10 times the isotropic background measured by EGRET, representative of the background near the Galactic plane.

We did this study first in the energy range from 1 GeV to 100 GeV by simulating extended sources of flux $4 \times 10^{-9} \text{ ph cm}^{-2} \text{ s}^{-1}$ integrated from 1 GeV to 100 GeV and a power-law spectral model with spectral index 2. This spectrum was picked to be representative of the new extended sources that were discovered in the following analysis when looking in the 1 GeV to 100 GeV energy range (see Section 10). We simulated these sources using uniform disk spatial models with extensions varying up to 1° . Figure 11a shows the distribution of TS_{ext} and $\text{TS}_{2\text{pts}}$ and Figure 11c shows the distribution of $\text{TS}_{\text{ext}} - \text{TS}_{2\text{pts}}$ as a function of the simulated extension of the source for 200 statistically independent simulations.

Figure 12a shows the same plot but when fitting two simulated point-like sources each with half of the flux of the spatially extended source and with the same spectral index as the extended source. Finally, Figure 12c shows the same plot with each point-like source having the same flux but different spectral indices. One point-like source had a spectral index of 1.5 and the other an index of 2.5. These indices are representative of the range of indices of LAT-detected sources.

The same four plots are shown in Figure 11b, 11d, 12b, and 12d but this time when analyzing a source of flux $10^{-9} \text{ ph cm}^{-2} \text{ s}^{-1}$ (integrated from 10 GeV to 100 GeV) only in the 10 GeV to 100 GeV energy range. This flux is typical of the new extended sources discovered using only photons with energies between 10 GeV and 100 GeV (see Section 10).

Several interesting conclusions can be made from this study. As one would expect, $\text{TS}_{\text{ext}} - \text{TS}_{2\text{pts}}$ is mostly positive when fitting the simulated extended sources. In the 1 GeV to 100 GeV analysis, only 11 of the 200 simulated extended sources had $\text{TS}_{\text{ext}} > 16$ but were incorrectly rejected due to $\text{TS}_{2\text{pts}}$ being greater than TS_{ext} . In the 10 GeV to 100 GeV analysis, only 7 of the 200 sources were incorrectly rejected. From this, we conclude that this test is unlikely to incorrectly reject truly spatially extended sources.

On the other hand, it is often the case that $\text{TS}_{\text{ext}} > 16$ when testing the two simulated point-like sources for extension. This is especially the case when the two sources had the same spectral index. Forty out of 200 sources in the 1 GeV to 100 GeV energy range and 43 out of 200 sources in the 10 GeV to 100 GeV energy range had $\text{TS}_{\text{ext}} > 16$. But in these cases, we always found the single extended source fit to be worse than the two point-like source fit. From this, we conclude that the $\text{TS}_{\text{ext}} > \text{TS}_{2\text{pts}}$ test is powerful at discarding cases in which the true emission comes from two point-like sources.

The other interesting feature in Figure 11a and 11b is that for simulated extended sources with typical sizes ($\sigma \sim 0.5^\circ$), one can often obtain almost as large an increase in likelihood fitting the source as two point-like sources ($\text{TS}_{2\text{pts}} \sim \text{TS}_{\text{ext}}$). This is because although the two point-like sources represent an incorrect spatial model, the second source has four additional degrees of freedom (two spatial and two spectral parameters) and can therefore easily model much of the extended source and statistical fluctuations in the data. This effect is most pronounced when using photons with energies between 1 GeV and 100 GeV where the PSF is broader.

From this Monte Carlo study, we can see the limits of a LAT analysis of spatially extended sources. Section 3.1 showed that we have a statistical test that finds when a LAT source is not well described by the PSF. But this test does not uniquely prove that the emission originates from spatially extended emission instead of from multiple unresolved sources. Demanding that $\text{TS}_{\text{ext}} > \text{TS}_{2\text{pts}}$ is a powerful second test to avoid cases of simple confusion of two point-like sources. But it could always be the case that an extended source is actually the superposition of multiple point-like or extended sources that could be resolved with deeper observations of the region. There is nothing about this conclusion unique to LAT analysis, but the broad PSF of the LAT and the density of sources expected to be GeV emitters in the Galactic plane makes this issue more significant for LAT analysis. When possible, multiwavelength information should be used to help select the best model of the

440 sky.

441 6. Test of 2LAC Sources

442 For all following LAT analyses, we used the same two-year dataset that was used in the
 443 2FGL catalog spanning from 2008 August 4 to 2010 August 1. We applied the same accep-
 444 tance cuts and we used the same P7_V6 Source class event selection and IRFs (Abdo et al.
 445 2012). When analyzing sources in `pointlike`, we used a circular 10° region of interest (ROI)
 446 centered on our source and eight energy bins per logarithmic decade in energy. When refit-
 447 ting the region in `gtlike` using the best fit spatial and spectral models from `pointlike`, we
 448 used the ‘binned likelihood’ mode of `gtlike` on a $14^\circ \times 14^\circ$ ROI with a pixel size of $0^\circ.03$.

449 Unless explicitly mentioned, we used the same background model as 2FGL to represent
 450 the Galactic diffuse, isotropic, and Earth limb emission. To compensate for possible residuals
 451 in the diffuse emission model, the Galactic emission was scaled by a power-law and the
 452 normalization of the isotropic component was left free. Unless explicitly mentioned, we used
 453 all 2FGL sources within 15° of our source as our list of background sources and we refit the
 454 spectral parameters of all sources within 2° of the source.

455 To validate our method, we tested LAT sources associated with AGN for extension.
 456 GeV emission from AGN is believed to originate from collimated jets. Therefore AGN are
 457 not expected to be spatially resolvable by the LAT and provide a good calibration source
 458 to demonstrate that our extension detection method does not misidentify point-like sources
 459 as being extended. We note that megaparsec-scale γ -ray halos around AGNs have been
 460 hypothesized to be resolvable by the LAT (Aharonian et al. 1994). However, no such halo
 461 has been discovered in the LAT data so far (Neronov et al. 2011).

462 Following 2FGL, the LAT Collaboration published the Second LAT AGN Catalog
 463 (2LAC), a list of high latitude ($|b| > 10^\circ$) sources that had a high probability associa-
 464 tion with AGN (Ackermann et al. 2011). 2LAC associated 1016 2FGL sources with AGN.
 465 To avoid systematic problems with AGN classification, we selected only the 885 AGN which
 466 made it into the clean AGN sub-sample defined in the 2LAC paper. An AGN association
 467 is considered clean only if it has a high probability of association $P \geq 80\%$, if it is the only
 468 AGN associated with the catalog source, and if no analysis flags have been set for the source
 469 in the 2FGL catalog. These last two conditions are important for our analysis. Source con-
 470 fusion may look like a spatially extended source and flagged catalog sources may correlate
 471 with unmodeled structure in the diffuse emission.

472 Of the 885 clean AGN, we selected the 733 of these 2FGL sources which were significantly

detected above 1 GeV and fit each of them for extension. The cumulative density of TS_{ext} for these AGN is compared to the $\chi^2_1/2$ distribution of Equation 10 in Figure 13. The TS_{ext} distribution for the AGN shows reasonable agreement with the theoretical distribution and no AGN was found to be significantly extended ($TS_{\text{ext}} > 16$). The observed discrepancy from the theoretical distribution is likely due to small systematics in our model of the LAT PSF and the Galactic diffuse emission (see Section 8). The discrepancy could also in a few cases be due to confusion with a nearby undetected source. We note that the Monte Carlo study of section 3.1 effectively used perfect IRFs and a perfect model of the sky. The overall agreement with the expected distribution demonstrates that we can use TS_{ext} as a measure of the statistical significance of the detection of the extension of a source.

We note that the LAT PSF used in this study was determined empirically by fitting the distributions of gamma rays around bright AGN (see Section 8). Finding that the AGN we test are not extended is not surprising. This validation analysis is not suitable to reject any hypotheses about the existence of megaparsec-scale halos around AGN.

7. Analysis of Extended Sources Identified in the 2FGL Catalog

As further validation of our method for studying spatially extended sources, we re-analyzed the twelve spatially extended sources which were included in the 2FGL catalog (Abdo et al. 2011b). Even though these sources had all been the subjects of dedicated analyses and separate publications, and had been fit with a variety of spatial models, it is valuable to show that these sources are significantly extended using our systematic method assuming radially-symmetric uniform disk spatial models. On the other hand, for some of these sources a uniform disk spatial model does not well describe the observed extended emission and so the dedicated LAT publications provide better models of these sources.

Six extended SNRs were included in the 2FGL catalog: W51C, IC 443, W28, W44, the Cygnus Loop, and W30 (Abdo et al. 2009a, 2010i,e,h; Katagiri et al. 2011; Abdo et al. 2011a). Using photons with energies between 1 GeV and 100 GeV, our analysis significantly detected that these six SNRs are spatially extended.

Two nearby satellite galaxies of the Milky Way the Large Magellanic Cloud (LMC) and the Small Magellanic Cloud (SMC) were included in the 2FGL catalog as spatially extended sources (Abdo et al. 2010j,b). Their extensions were significantly detected using photons with energies between 1 GeV and 100 GeV. Our fit extensions are comparable to the published result, but we note that the previous LAT Collaboration publication on the LMC used a more complicated two 2D Gaussian surface brightness profile when fitting it

(Abdo et al. 2010j).

Three PWNe, MSH 15–52, Vela X, and HESS J1825–137, were fit as extended sources in the 2FGL analysis (Abdo et al. 2010a,f; Grondin et al. 2011). In the present analysis, HESS J1825–137 was significantly detected using photons with energies between 10 GeV and 100 GeV. To avoid confusion with the nearby bright pulsar PSR J1509–5850, MSH 15–52 had to be analyzed at high energies. Using photons with energies above 10 GeV, we fit the extension of MSH 15–52 to be consistent with the published size but with $TS_{\text{ext}}=6.6$.

Our analysis was unable to resolve Vela X which would have required first removing the pulsed photons from the Vela pulsar which was beyond the scope of this paper. Our analysis also failed to detect a significant extension for the Centaurus A Lobes because the shape of the source is significantly different from a uniform radially-symmetric disk (Abdo et al. 2010c).

Our analysis of these sources is summarized in Table 3. This table includes the best fit positions and extensions of these sources when fitting them with a radially-symmetric uniform disk model. It also includes the best fit spectral parameters for each source. The positions and extensions of Vela X and the Centaurus A Lobes were taken from Abdo et al. (2010f,c) and are included in this table for completeness.

8. Systematic Errors on Extension

We developed two criteria for estimating systematic errors on the extensions of the sources. First, we estimated a systematic error due to uncertainty in our knowledge of the LAT PSF. Before launch, the LAT PSF was determined by detector simulations which were verified in accelerator beam tests (Atwood et al. 2009b). However, in-flight data revealed a discrepancy above 3 GeV in the PSF compared to the angular distribution of photons from bright AGN (Abdo et al. 2012). Subsequently, the PSF was fit empirically to bright AGN and this empirical parameterization is used in the P7_V6 IRFs. To account for the uncertainty in our knowledge of the PSF, we refit our extended source candidates using the pre-flight Monte Carlo representation of the PSF and consider the difference in extension found using the two PSFs as a systematic error on the extension of a source. The same approach was used in Abdo et al. (2010i). We believe that our parameterization of the PSF from bright AGN is substantially better than the Monte Carlo representation of the PSF so this systematic error is conservative.

We estimated a second systematic error on the extension of a source due to uncertainty in our model of the Galactic diffuse emission by using an alternative approach to modeling

the diffuse emission which takes as input templates calculated by GALPROP⁴ but then fits each template locally in the surrounding region. The particular GALPROP model that was used as input is described in the LAT analysis of the isotropic diffuse emission (Abdo et al. 2010k). The intensities of various components of the Galactic diffuse emission were fitted individually using a spatial distribution predicted by the model. We distinguished contributions from cosmic-ray interactions with the molecular hydrogen, the atomic and ionized hydrogen, residual gas traced by dust (Grenier et al. 2005), and the interstellar radiation field. We further split the contributions from interactions with molecular and atomic hydrogen to the Galactic diffuse emission according to the distance from the Galactic center in which they are produced. Hence, we replaced the standard diffuse emission model by 18 individually fitted templates to describe individual components of the diffuse emission. A similar crosscheck was used in an analysis of RX J1713.7–3946 by the LAT Collaboration (Abdo et al. 2011c).

It is not expected that this diffuse model is superior to the standard LAT model obtained through an all-sky fit. However, adding degrees of freedom to the background model can remove likely spurious sources that correlate with features in the Galactic diffuse emission. Therefore, this tests systematics that may be due to imperfect modeling of the diffuse emission in the region. Nevertheless, this alternative approach to modeling the diffuse emission does not test all systematics related to the diffuse emission model. In particular, because the alternative approach uses the same underlying gas maps, it is unable to be used to assess systematics due to insufficient resolution of the underlying maps. Structure in the diffuse emission that is not correlated with these maps will also not be assessed by this test.

We do not expect the systematic error due to uncertainties in the PSF to be correlated with the systematic error due to uncertainty in the Galactic diffuse emission. Therefore, the total systematic error on the extension of a source was obtained by adding the two errors in quadrature.

There is another systematic error on the size of a source due to issues modeling nearby sources in crowded regions of the sky. It is beyond the scope of this paper to address this systematic error. Therefore, for sources in crowded regions the systematic errors quoted in this paper may not be the full systematics of this analysis.

⁴GALPROP is a software package for calculating the Galactic γ -ray emission based on a model of cosmic-ray propagation in the Galaxy and maps of the distributions of the components of the interstellar medium (Strong & Moskalenko 1998; Vladimirov et al. 2011). See also <http://galprop.stanford.edu/> for details.

9. Extended Source Search Method

Having demonstrated that we understand the statistical issues associated with analyzing spatially extended sources (Section 3.1 and 6) and that our method can correctly analyze the extended sources included in 2FGL (Section 7), we applied this method to search for new spatially extended GeV sources. The data and general analysis setting is as described in Section 6.

Ideally, we would apply a completely blind and uniform search that tests the extension of each catalog source in the presence of all other 2FGL sources to find a complete list of all spatially extended sources. As our test of AGN in Section 6 showed, at high Galactic latitude where the source density is not as large and the diffuse emission is less structured, this method works well.

But this is infeasible in the Galactic plane where we most likely expect to discover new spatially extended sources. In the Galactic plane, this analysis is challenged by our imperfect model of the diffuse emission and by an imperfect model of nearby sources. The Monte Carlo study in Section 5 showed that the overall likelihood would greatly increase by fitting a spatially extended source as two point-like sources so we expect that spatially extended sources would be modeled in the 2FGL catalog as multiple point-like sources. Furthermore, the positions of other nearby sources in the region close to an extended source could be biased by not correctly modeling the extension of the source. Since 2FGL produced a list of sources significant at energies above 100 MeV whereas we are most sensitive to spatial extension at higher energies, we also expect that at higher energies our analysis would be complicated by 2FGL sources no longer significant and by 2FGL sources whose positions were biased by diffuse emission at lower energies.

To account for these issues, we first produced a large list of possibly extended sources employing very liberal search criteria and then refined the analysis of the promising candidates on a case by case basis. Our strategy was to test all point-like 2FGL sources for extension assuming they had a uniform radially-symmetric disk spatial model and a power-law spectral model. Although not all extended sources are expected to have a shape very similar to a uniform disk, Section 2.4 showed that for many spatially extended sources the wide PSF of the LAT and limited statistics makes this a reasonable approximation. On the other hand, choosing this spatial model biases us against finding extended sources that are not well described by a uniform disk model such as shell-type SNRs.

Before testing for extension, we automatically removed from the background model all other 2FGL sources within 0.5° of the source. This distance is somewhat arbitrary, but was picked in hopes of finding extended sources with sizes on the order of $\sim 1^\circ$ or smaller. On the

other hand, by removing these nearby background sources we expect to also incorrectly add to our list of extended source candidates point-like sources that are confused with nearby sources. To screen out obvious cases of source confusion, we performed the dual localization procedure described in Section 5 to compare the extended source hypothesis to the hypothesis of two independent point-like sources.

As was shown in Section 4, little sensitivity is gained by using photons with energies below 1 GeV. On the other hand, the broad PSF at low energy makes the analysis more susceptible to systematic errors arising from source confusion due to nearby soft point-like sources and by uncertainties in our modeling of the Galactic diffuse emission. For that reason, we performed our search using only photons with energies between 1 GeV and 100 GeV.

We also performed a second search for extended sources using only photons with energies between 10 GeV and 100 GeV. Although this approach tests the same sources, it is complementary because the Galactic diffuse emission is even less dominant above 10 GeV and because source confusion is less of an issue. A similar procedure was used to detect the spatial extensions of MSH 15–52 and HESS J1825–137 with the LAT (Abdo et al. 2010a; Grondin et al. 2011).

When we applied this test to the 1861 point-like sources in the 2FGL catalog, our search found 117 extended source candidates in the 1 GeV to 100 GeV energy range and 11 extended source candidates in the 10 GeV to 100 GeV energy range. Most of the extended sources found above 10 GeV were also found above 1 GeV and in many cases multiple nearby point-like sources were found to be extended even though they fit the same emission region. For example, the sources 2FGL J1630.2–4752, 2FGL J1632.4–4753c, 2FGL J1634.4–4743c, and 2FGL J1636.3–4740c were all found to be spatially extended in the 10 GeV to 100 GeV energy range even though they all fit to similar positions and sizes. For these situations, we manually discarded all but one of the 2FGL sources.

Similarly, many of these sources were confused with nearby point-like sources or influenced by large-scale residuals in the diffuse emission. To help determine which of these fits found truly extended sources and when the extension was influenced by source confusion and diffuse emission, we generated a series of diagnostic plots. For each candidate, we generated a map of the residual TS by adding a new point-like source of spectral index 2 into the region at each position and finding the increase in likelihood when fitting its flux. Figure 14 shows this map around the most significantly extended source IC 443 when it is modeled both as a point-like source and as an extended source. The residual TS map indicates that the spatially extended model for IC 443 is a significantly better description of the observed photons and that there is no significant residual in the region after modeling the source as

being spatially extended. We also generated plots of the sum of all counts within a given distance of the source and compared them to the model predictions assuming the emission originated from a point-like source. An example radial integral plot is shown for the extended source IC 443 in Figure 1. For each source, we also made diffuse-emission-subtracted smoothed counts maps (shown for IC 443 in Figure 1).

We found by visual inspection that in many cases our results were strongly influenced by large-scale residuals in the diffuse emission and hence the extension measure was unreliable. This was especially true in our analysis of sources in the 1 GeV to 100 GeV energy range. An example of such a case is 2FGL J1856.2+0450c analyzed in the 1 GeV to 100 GeV energy range. Figure 15 shows a diffuse-emission-subtracted smoothed counts map for this source with the best fit extension of the source overlaid. There appear to be large-scale residuals in the diffuse emission in this region along the Galactic plane. As a result, 2FGL J1856.2+0450c is fit to an extension of $\sim 2^\circ$ and the result is statistically significant with $\text{TS}_{\text{ext}} = 45.4$. However, by looking at the residuals it is clear that this complicated region is not well modeled. We manually discard sources like this.

We only selected extended source candidates in regions that did not appear dominated by these issues and where there was a multiwavelength counterpart. Because of these systematic issues, this search can not be expected to be complete and it is likely that there are other spatially extended sources that this method missed.

For each candidate that was not biased by neighboring point-like sources or by large-scale residuals in the diffuse emission model, we improved the model of the region by deciding on a case by case basis which background point-like sources should be kept. We kept in our model the sources that we believed represented physically distinct sources and we removed sources that we believed were included in the 2FGL catalog to compensate for residuals induced by not modeling the extension of the source. Soft nearby point-like 2FGL sources that were not significant at higher energies were frozen to the spectras predicted by 2FGL. When deciding which background sources to keep and which to remove, we used as much as possible multiwavelength information about possibly extended source counterparts to help guide our choice. For each extended source presented in Section 10, we describe any modifications from 2FGL of the background model that were performed. **In Table 6, we will summarize the sources that were included in 2FGL to account for residuals induced by not modeling the extension of a nearby source.**

The best fit positions of nearby point-like sources can be influenced by the extended source and vice versa. Similarly, the best fit positions of nearby point-like sources in the 2FGL catalog can be biased by systematic issues at lower energies. Therefore, after selecting the list of background sources, we iteratively refit the positions and spectra of nearby background

sources as well as the positions and extensions of the analyzed spatially extended sources until the overall fit converged globally. For each extended source, we will describe the positions of any relocalized background sources.

After obtaining the overall best fit positions and extensions of all of the sources in the region using `pointlike`, we refit the spectral parameters of the region using `gtlike`. With `gtlike`, we obtained a second measure of TS_{ext} . We only consider a source to be extended when both `pointlike` and `gtlike` agree that $\text{TS}_{\text{ext}} > 16$. We further required that $\text{TS}_{\text{ext}} > 16$ using the alternative approach to modeling the diffuse emission presented in Section 8. We then replaced the spatially extended source with two point-like sources and refit the positions and spectra of the two point-like sources to calculate $\text{TS}_{2\text{pts}}$. We only consider a source to be spatially extended, instead of being the result of confusion of two point-like sources, if $\text{TS}_{\text{ext}} > \text{TS}_{2\text{pts}}$. As was shown in Section 5, this test is fairly powerful at removing situations in which the emission actually originates from two distinct point-like sources instead of one spatially extended source. On the other hand, it is still possible that longer observations could resolve additional structure or new sources that the analysis cannot currently detect. Considering the very complicated morphologies of extended sources observed at other wavelengths and the high density of possible sources that are expected to emit at GeV energies, it is likely that in some of these regions further observations will reveal that the emission is significantly more complicated than the simple radially-symmetric uniform disk model that we assume.

10. New Extended Sources

Nine extended sources not included in the 2FGL catalog were found by our extended source search. Two of these have been previously studied in dedicated publications: RX J1713.7–3946 and Vela Jr. (Abdo et al. 2011c; Tanaka et al. 2011). Two of these sources were found when using photons with energies between 1 GeV and 100 GeV and seven were found when using photons with energies between 10 GeV and 100 GeV. For the sources found at energies above 10 GeV, we restrict our analysis to higher energies because of the issues of source confusion and diffuse emission modeling described in Section 9. The spectral and spatial properties of these nine sources are summarized in Table 4 and the results of our investigation of systematic errors are presented in Table 5. Table 5 also compares the likelihood assuming the source is spatially extended to the likelihood assuming that the emission originates from two independent point-like sources. For these new extended sources, $\text{TS}_{\text{ext}} > \text{TS}_{2\text{pts}}$ so we conclude that the GeV emission does not originate from two physically distinct point-like sources (see Section 5). Table 5 also includes the results of the extension fits using variations of the PSF

and the Galactic diffuse model described in Section 8. There is good agreement between TS_{ext} and the fit size using the standard analysis, the alternative approach to modeling the diffuse emission, and the alternative PSF. This suggests that the sources are robust against mis-modeled features in the diffuse emission model and uncertainties in the PSF. **Finally, in Table 6 we summarize the sources included in 2FGL to account for residuals induced by not modeling the extension of a nearby source.**

10.1. 2FGL J0823.0–4246

2FGL J0823.0–4246 was found by our search to be an extended source candidate in the 1 GeV to 100 GeV energy range and is spatially coincident with the SNR Puppis A. Figure 16 shows a counts map of this source. There are two nearby catalog sources 2FGL J0823.4–4305 and 2FGL J0821.0–4254 that are also coincident with the SNR but that do not appear to represent physically distinct sources. We conclude that these nearby point-like sources were included in the 2FGL catalog to compensate for residuals induced by not modeling the extension of this source and removed them from our model of the sky. After removing these sources, 2FGL J0823.0–4246 was found to have an extension $\sigma = 0.37 \pm 0.03_{\text{stat}} \pm 0.02_{\text{sys}}$ with $\text{TS}_{\text{ext}} = 48.0$. Figure 17 shows the spectrum of this source.

Puppis A has been studied in detail in radio (Castelletti et al. 2006), and X-ray (Petre et al. 1996; Hwang et al. 2008). The fit extension of 2FGL J0823.0–4246 matches well the size of Puppis A in X-ray. The distance of Puppis A was estimated at 2.2 kpc (Reynoso et al. 1995, 2003) and leads to a 1 GeV to 100 GeV luminosity of $\sim 3 \times 10^{34} \text{ erg s}^{-1}$. No molecular clouds have been observed directly adjacent to Puppis A (Paron et al. 2008), similar to the LAT detected Cygnus Loop SNR (Katagiri et al. 2011). The luminosity of Puppis A is also smaller than that of other SNRs believed to interact with molecular clouds (Abdo et al. 2009a, 2010i,h,e,g).

10.2. 2FGL J0851.7–4635

2FGL J0851.7–4635 was found by our search to be an extended source candidate in the 10 GeV to 100 GeV energy range and is spatially coincident with the SNR Vela Jr. This source was recently studied by the LAT Collaboration in Tanaka et al. (2011). Figure 19 shows a counts map of the source. Overlaid on Figure 19 are TeV contours of Vela Jr. (Aharonian et al. 2007a). There are three point-like catalog sources 2FGL J0848.5–4535, 2FGL J0853.5–4711, and 2FGL J0855.4–4625 which correlate with the multiwavelength

emission of this SNR but do not appear to be physically distinct sources. They were most likely included in the 2FGL catalog to compensate for residuals induced by not modeling the extension of Vela Jr. and were removed from our model of the sky.

With this model of the background, 2FGL J0851.7–4635 was found to have an extension of $\sigma = 1^\circ 15 \pm 0^\circ 08_{\text{stat}} \pm 0^\circ 02_{\text{sys}}$ with $\text{TS}_{\text{ext}} = 86.8$. The LAT size matches well the TeV morphology of Vela Jr. While fitting the extension of 2FGL J0851.7–4635, we iteratively relocalized the positions of the nearby point-like catalog source 2FGL J0854.7–4501 to $(l, b) = (266^\circ 24, 0^\circ 49)$ to better fit its position at high energies.

10.3. 2FGL J1615.0–5051

2FGL J1615.0–5051 and 2FGL J1615.2–5138 were both found to be extended source candidates in the 10 GeV to 100 GeV energy range. Because they are less than 1° away from each other, they needed to be analyzed simultaneously. 2FGL J1615.0–5051 is spatially coincident with the extended TeV source HESS J1616–508 and 2FGL J1615.2–5138 is coincident with the extended TeV source HESS J1614–518. Figure 20 shows a counts map of these sources and overlays the TeV contours of HESS J1616–508 and HESS J1614–518 (Aharonian et al. 2006). The figure shows that the 2FGL source 2FGL J1614.9–5212 is very close to 2FGL J1615.2–5138 and correlates with the same extended TeV source as 2FGL J1615.2–5138. We concluded that this source was included in the 2FGL catalog to compensate for residuals induced by not modeling the extension of 2FGL J1615.2–5138 and removed it from our model of the sky.

With this model of the sky, we iteratively fit the extensions of 2FGL J1615.0–5051 and 2FGL J1615.2–5138. 2FGL J1615.0–5051 was found to have an extension $\sigma = 0^\circ 32 \pm 0^\circ 04_{\text{stat}} \pm 0^\circ 01_{\text{sys}}$ and $\text{TS}_{\text{ext}} = 16.7$.

The TeV counterpart of 2FGL J1615.0–5051 was fit with a radially-symmetric Gaussian surface brightness profile with $\sigma = 0^\circ 136 \pm 0^\circ 008$. This TeV size corresponds to a 68% containment radius of $r_{68} = 0^\circ 21 \pm 0^\circ 01$, comparable to the LAT size $r_{68} = 0^\circ 26 \pm 0^\circ 03$. Figure 21 shows that the spectrum of 2FGL J1615.0–5051 at GeV energies connects to the spectrum of HESS J1616–508 at TeV energies.

HESS J1616–508 is located in the region of two SNRs RCW103 (G332.4-04) and Kes 32 (G332.4+0.1) but is not spatially coincident with either of them (Aharonian et al. 2006). HESS J1616–508 is near three pulsars PSR J1614–5048, PSR J1616–5109, and PSR J1617–5055. (Torii et al. 1998; Landi et al. 2007a). Only PSR J1617–5055 is energetically capable of powering the TeV emission and Aharonian et al. (2006) speculated that HESS J1616–508

could be a PWN powered by this young pulsar. Because HESS J1616–508 is 9′ away from PSR J1617–5055, this would require an asymmetric X-ray PWNe to power the TeV emission. *Chandra* ACIS observations revealed an underluminous PWN of size $\sim 1'$ around the pulsar that was not oriented towards the TeV emission, rendering this association uncertain (Kargaltsev et al. 2009). No other promising counterparts were observed at X-ray and soft gamma-ray energies by *Suzaku* (Matsumoto et al. 2007), *Swift*/XRT, IBIS/ISGRBI, BeppoSAX and *XMM-Newton* (Landi et al. 2007a). Kargaltsev et al. (2009) discovered additional diffuse emission towards the center of HESS J1616–508 using archival radio and infrared observations. Deeper observations will likely be necessary to understand this γ -ray source.

10.4. 2FGL J1615.2–5138

2FGL J1615.2–5138 was found to have an extension $\sigma = 0^\circ.42 \pm 0^\circ.04_{\text{stat}} \pm 0.02_{\text{sys}}$ with $\text{TS}_{\text{ext}} = 46.5$. To test for the possibility that 2FGL J1615.2–5138 is not spatially extended but instead composed of two point-like sources (one of them represented in the 2FGL catalog by 2FGL J1614.9–5212), we refit 2FGL J1615.2–5138 as two point-like sources. **Because $\text{TS}_{2\text{pts}} = 35.1$ is less than $\text{TS}_{\text{ext}} = 46.5$, we conclude that this emission does not originate from two nearby point-like sources.**

2FGL J1615.2–5138 is spatially coincident with the extended TeV source HESS J1614–518. H.E.S.S. measured a 2D Gaussian extension of $\sigma = 0^\circ.23 \pm 0^\circ.02$ and $\sigma = 0^\circ.15 \pm 0^\circ.02$ in the semi-major and semi-minor axis. This corresponds to a 68% containment size of $r_{68} = 0^\circ.35 \pm 0^\circ.03$ and $0^\circ.23 \pm 0^\circ.03$, consistent with the LAT size $r_{68} = 0^\circ.34 \pm 0^\circ.03$. Figure 21 shows that the spectrum of 2FGL J1615.2–5138 at GeV energies connects to the spectrum of HESS J1614–518 at TeV energies. Further data collected by H.E.S.S. in 2007 resolve a double peaked structure at TeV energies but no spectral variation across this source, suggesting that the emission is not the confusion of physically separate sources (Rowell et al. 2008). This double peaked structure is also hinted at in the LAT counts map in Figure 20 but is not very significant. The TeV source was also detected by CANGAROO-III (Mizukami et al. 2011).

There are five nearby pulsars, but none are luminous enough to provide the energy output required to power the γ -ray emission (Rowell et al. 2008). HESS J1614–518 is spatially coincident with a young open cluster Pismis 22 (Landi et al. 2007b; Rowell et al. 2008). *Suzaku* detected two promising X-ray candidates. Source A is an extended source consistent with the peak of HESS J1614–518 and source B coincident with Pismis 22 and towards the center but in a relatively dim region of HESS J1614–518 (Matsumoto et al. 2008). Three

hypotheses have been presented to explain this emission: either source A is an SNR powering the γ -ray emission; source A is a PWN powered by an undiscovered pulsar in either source A or B; and finally that the emission may arise from hadronic acceleration in the stellar winds of Pismis 22 (Mizukami et al. 2011).

10.5. 2FGL J1627.0–2425c

2FGL J1627.0–2425c was found by our search to have an extension $\sigma = 0^\circ.42 \pm 0^\circ.05_{\text{stat}} \pm 0^\circ.16_{\text{sys}}$ with $\text{TS}_{\text{ext}} = 32.4$ using photons with energies between 1 GeV and 100 GeV. Figure 18 shows a counts map of this source.

This source is in a region of remarkably complicated diffuse emission. Even though it is 16° from the Galactic plane, this source is on top of the core of the Ophiuchus molecular cloud which contains massive star-forming regions that are bright in infrared. The region also has abundant molecular and atomic gas traced by CO and H I and significant dark gas found only by its association with dust emission (Grenier et al. 2005). Embedded star-forming regions make it even more challenging to measure the column density of dust. Infrared and CO ($J = 1 \rightarrow 0$) contours are overlaid on Figure 18 and show good spatial correlation with the GeV emission (Young et al. 1986; de Geus et al. 1990). This source might represent γ -ray emission from the interactions of cosmic rays with interstellar gas which has not been accounted for in the LAT diffuse emission model.

10.6. 2FGL J1632.4–4753c

2FGL J1632.4–4753c was found by our search to be an extended source candidate in the 10 GeV to 100 GeV energy range but is in a crowded region of the sky. It is spatially coincident with the TeV source HESS J1632–478. Figure 22a shows a counts map of this source and overlays TeV contours of HESS J1632–478 (Aharonian et al. 2006). There are six nearby point-like 2FGL sources that appear to represent physically distinct sources and were included in our background model: 2FGL J1630.2–4752, 2FGL J1631.7–4720c, 2FGL J1632.4–4820c, 2FGL J1635.4–4717c, 2FGL J1636.3–4740c, and 2FGL J1638.0–4703c. On the other hand, one point-like 2FGL source 2FGL J1634.4–4743c correlates with the extended TeV source and at GeV energies does not appear physically separate. **It is very close to the position of 2FGL J1632.4–4753c and does not show in the observed photon distribution separated emission.** We therefore removed this source from our model of the background. Figure 22b shows the same region with the back-

ground sources subtracted. With this model, 2FGL J1632.4–4753c was found to have an extension $\sigma = 0^\circ.35 \pm 0^\circ.04_{\text{stat}} \pm 0^\circ.02_{\text{sys}}$ with $\text{TS}_{\text{ext}} = 26.9$. While fitting the extension of 2FGL J1632.4–4753c, we iteratively relocalized 2FGL J1635.4–4717c to $(l, b) = (337^\circ.23, 0^\circ.35)$ and 2FGL J1636.3–4740c to $(l, b) = (336^\circ.97, -0^\circ.07)$.

H.E.S.S. measured an extension of $\sigma = 0^\circ.21 \pm 0^\circ.05$ and $0^\circ.06 \pm 0^\circ.04$ along the semi-major and semi-minor axes when fitting HESS J1632–478 with an elliptical 2D Gaussian surface brightness profile. This corresponds to a 68% containment size $r_{68} = 0^\circ.31 \pm 0^\circ.08$ and $0^\circ.09 \pm 0^\circ.06$ along the semi-major and semi-minor axis, consistent with the LAT size $r_{68} = 0^\circ.29 \pm 0^\circ.04$. Figure 21 shows that the spectrum of 2FGL J1632.4–4753c at GeV energies connects to the spectrum of HESS J1632–478 at TeV energies.

Aharonian et al. (2006) argued that HESS J1632–478 is positionally coincident with the hard X-ray source IGR J1632–4751 observed by *ASCA*, *INTEGRAL*, and *XMM-Newton* (Sugizaki et al. 2001; Tomsick et al. 2003; Rodriguez et al. 2003), but this source is suspected to be a Galactic X-Ray Binary so the γ -ray extension disfavors the association. Further observations by *XMM-Newton* discovered point-like emission coincident with the peak of the H.E.S.S. source surrounded by extended emission of size $\sim 32'' \times 15''$ (Balbo et al. 2010). They found in archival MGPS-2 data a spatially coincident extended radio source (Murphy et al. 2007) and argued for a single synchrotron and inverse Compton process producing the radio, X-ray, and TeV emission, likely due to a PWN. The increased size at TeV energies compared to X-ray energies has previously been observed in several aging PWNe including HESS J1825–137 (Gaensler et al. 2003; Aharonian et al. 2006b), HESS J1640–465 (Aharonian et al. 2006; Funk et al. 2007), and Vela X (Markwardt & Ogelman 1995; Aharonian et al. 2006c) and can be explained by different synchrotron cooling times for the electrons that produce X-rays and γ -rays.

10.7. 2FGL J1712.4–3941

2FGL J1712.4–3941 was found by our search to be spatially extended using photons with energies between 1 GeV and 100 GeV. This source is spatially coincident with the SNR RX J1713.7–3946 and was recently studied by the LAT Collaboration in Abdo et al. (2011c). To avoid issues related to uncertainties in the nearby Galactic diffuse emission at lower energy, we restricted our analysis only to energies above 10 GeV. Figure 23 shows a smoothed counts map of the source. Above 10 GeV, the GeV emission nicely correlates with the TeV contours of RX J1713.7–3946 (Aharonian et al. 2007b) and 2FGL J1712.4–3941 fit to an extension $\sigma = 0^\circ.56 \pm 0^\circ.04_{\text{stat}} \pm 0^\circ.02_{\text{sys}}$ with $\text{TS}_{\text{ext}} = 38.5$.

10.8. 2FGL J1837.3–0700c

2FGL J1837.3–0700c was found by our search to be an extended source candidate in the 10 GeV to 100 GeV energy range and is spatially coincident with the TeV source HESS J1837–069. This source is in a complicated region. Figure 24a shows a smoothed counts map of the region and overlays the TeV contours of HESS J1837–069 (Aharonian et al. 2006). There are two very nearby point-like 2FGL sources, 2FGL J1836.8–0623c and 2FGL J1839.3–0558c, that clearly represent distinct sources. On the other hand, there is another source 2FGL J1835.5–0649 located between the three sources that appears to correlate with the TeV morphology of HESS J1837–069 but at GeV energies does not appear to represent a physically distinct source. We concluded that this source was included in the 2FGL catalog to compensate for residuals induced by not modeling the extension of this source and removed it from our model of the sky. Figure 24b shows a counts map of this region after subtracting these background sources. After removing 2FGL J1835.5–0649, we tested for source confusion by fitting 2FGL J1837.3–0700c instead as two point-like sources. **Because $\text{TS}_{2\text{pts}} = 10.8$ is less than $\text{TS}_{\text{ext}} = 18.5$, we conclude that this emission does not originate from two nearby point-like sources.**

With this model, 2FGL J1837.3–0700c was found to have an extension $\sigma = 0^\circ.33 \pm 0^\circ.07_{\text{stat}} \pm 0^\circ.05_{\text{sys}}$. While fitting the extension of 2FGL J1837.3–0700c, we iteratively re-localized the two closest background sources along with the extension of 2FGL J1837.3–0700c but their positions did not significantly change. 2FGL J1834.7–0705c moved to $(l, b) = (24^\circ.77, 0^\circ.50)$, 2FGL J1836.8–0623c moved to $(l, b) = (25^\circ.57, 0^\circ.32)$.

H.E.S.S. measured an extension of $\sigma = 0^\circ.12 \pm 0^\circ.02$ and $0^\circ.05 \pm 0^\circ.02$ of the coincident TeV source HESS J1837–069 along the semi-major and semi-minor axis when fitting this source with an elliptical 2D Gaussian surface brightness profile. This corresponds to a 68% containment radius of $r_{68} = 0^\circ.18 \pm 0^\circ.03$ and $0^\circ.08 \pm 0^\circ.03$ along the semi-major and semi-minor axis. The size is not significantly different from the LAT 68% containment radius of $r_{68} = 0^\circ.27 \pm 0^\circ.07$ (less than 2σ). Figure 21 shows that the spectrum of 2FGL J1837.3–0700c at GeV energies connects to the spectrum of HESS J1837–069 at TeV energies.

HESS J1837–069 is coincident with the hard and steady X-ray source AX J1838.0–0655 (Bamba et al. 2003). This source was discovered by RXTE to be a pulsar (PSR J1838–0655) sufficiently luminous to power the TeV emission and was resolved by *Chandra* to be a bright point-like source surrounded by a $\sim 2'$ nebula (Gotthelf & Halpern 2008). The γ -ray emission may be powered by this pulsar. The hard spectral index and spatial extension of 2FGL J1837.3–0700c disfavor a pulsar origin of the LAT emission and suggest instead that the GeV and TeV emission both originate from the pulsar’s wind. There is another X-ray point-like source AX J1837.3–0652 near HESS J1837–069 (Bamba et al. 2003) that was also

resolved into a point-like and diffuse component (Gotthelf & Halpern 2008). Although no pulsations have been detected from it, it could also be a pulsar powering some of the γ -ray emission.

10.9. 2FGL J2021.5+4026

The source 2FGL J2021.5+4026 is associated with the γ -Cygni SNR and has been speculated to originate from the interaction of accelerated particles in the SNR with dense molecular clouds (Pollock 1985; Gaisser et al. 1998). This association was disfavored when the GeV emission from this source was detected to be pulsed (PSR J2021+4026, Abdo et al. 2010l). This pulsar was also observed by AGILE (Chen et al. 2011).

Looking at the same region at energies above 10 GeV, the pulsar is no longer significant but we instead found in our search an extended source candidate. Figure 25 shows a counts map of this source and overlays radio contours of γ -Cygni from the Canadian Galactic Plane Survey (Taylor et al. 2003). There is good spatial overlap between the SNR and the GeV emission.

There is a nearby source 2FGL J2019.1+4040 that correlates with the radio emission of γ -Cygni and at GeV energies does not appear to represent a physically distinct source. We concluded that it was included in the 2FGL catalog to compensate for residuals induced by not modeling the extension of γ -Cygni and removed it from our model of the sky. With this model, 2FGL J2021.5+4026 was found to have an extension $\sigma = 0.63 \pm 0.05_{\text{stat}} \pm 0.04_{\text{sys}}$ with $\text{TS}_{\text{ext}} = 128.9$. Figure 17 shows its spectrum. The inferred size of this source at GeV energies well matches the radio size of γ -Cygni. Milagro detected a 4.2σ excess at energies ~ 30 TeV from this location (Abdo et al. 2009b,c). VERITAS also detected an extended source VER J2019+407 coincident with the SNR above 200 GeV and suggested that the TeV emission could be a shock-cloud interaction in γ -Cygni (Weinstein 2009).

11. Discussion

Twelve extended sources were included in the 2FGL catalog and two additional extended sources were studied in dedicated publications. Using 2 years of LAT data and a new analysis method, we presented the detection of seven additional extended sources. We also reanalyzed the spatial extents of the twelve extended sources in the 2FGL catalog and the two additional sources. The 21 extended LAT sources are located primarily along the Galactic plane and their locations are shown in Figure 26. Most of the LAT detected extended sources are

expected to be of Galactic origin as the distances of extragalactic sources (with the exception of the local group Galaxies) are typically too large to be able to resolve them at γ -ray energies.

For the LAT extended sources also seen at TeV energies, Figure 27 shows that there is a good correlation between the sizes of the sources at GeV and TeV energies. Even so, the sizes of PWNe are expected to vary across the GeV and TeV energy range and the size of HESS J1825–137 is significantly larger at GeV than TeV energies (Grondin et al. 2011). It is interesting to compare the sizes of other PWN candidates at GeV and TeV energies, but definitively measuring a difference in size would require a more in-depth analysis of the LAT data using the same elliptical Gaussian spatial model.

Figure 28 compares the sizes of the 21 extended LAT sources to the 42 extended H.E.S.S. sources.⁵ Because of the large field of view and all-sky coverage, the LAT can more easily measure larger sources. On the other hand, the better angular resolution of air Cherenkov detectors allows them to measure a population of extended sources below the resolution limit of the LAT (currently about $\sim 0.2^\circ$). *Fermi* has a 5 year nominal mission lifetime with a goal of 10 years of operation. As Figure 10 shows, the low background of the LAT at high energies allows its sensitivity to these smaller sources to improve by a factor greater than the square root of the relative exposures. With increasing exposure, the LAT will likely begin to detect and resolve some of these smaller TeV sources.

Figure 29 compares the spectral indices of LAT detected extended sources and of all sources in the 2FGL catalog. This, and Tables 3 and 4, show that the LAT observes a population of hard extended sources at energies above 10 GeV. Figure 21 shows that the spectra of four of these sources (2FGL J1615.0–5051, 2FGL J1615.2–5138, 2FGL J1632.4–4753c, and 2FGL J1837.3–0700c) at GeV energies connects to the spectra of their H.E.S.S. counterparts at TeV energies. This is also true of Vela Jr., HESS J1825–137 (Grondin et al. 2011), and RX J1713.7–3946 (Abdo et al. 2011c). It is likely that the GeV and TeV emission from these sources originates from the same population of high-energy particles.

Many of the TeV detected extended sources now seen at GeV energies are currently unidentified and further multiwavelength follow-up observations will be necessary to understand these particle accelerators. But extending the spectra of these TeV sources towards lower energies with LAT observations may help to determine the origin and nature of the high-energy emission.

The *Fermi* LAT Collaboration acknowledges generous ongoing support from a number

⁵The TeV extension of the 42 extended H.E.S.S. sources comes from the H.E.S.S. Source Catalog <http://www.mpi-hd.mpg.de/hfm/HESS/pages/home/sources/>.

of agencies and institutes that have supported both the development and the operation of the LAT as well as scientific data analysis. These include the National Aeronautics and Space Administration and the Department of Energy in the United States, the Commissariat à l’Energie Atomique and the Centre National de la Recherche Scientifique / Institut National de Physique Nucléaire et de Physique des Particules in France, the Agenzia Spaziale Italiana and the Istituto Nazionale di Fisica Nucleare in Italy, the Ministry of Education, Culture, Sports, Science and Technology (MEXT), High Energy Accelerator Research Organization (KEK) and Japan Aerospace Exploration Agency (JAXA) in Japan, and the K. A. Wallenberg Foundation, the Swedish Research Council and the Swedish National Space Board in Sweden.

Additional support for science analysis during the operations phase is gratefully acknowledged from the Istituto Nazionale di Astrofisica in Italy and the Centre National d’Études Spatiales in France.

This research has made use of `pywcsgrid2`, an open-source plotting package for Python⁶. The authors acknowledge the use of `HEALPix`⁷ (Górski et al. 2005).

REFERENCES

- Abdo, A. A., et al. 2009a, *ApJ*, 706, L1
- . 2009b, *ApJS*, 183, 46
- . 2009c, *ApJ*, 700, L127
- . 2009d, *Astroparticle Physics*, 32, 193
- . 2010a, *ApJ*, 714, 927
- . 2010b, *A&A*, 523, A46+
- . 2010c, *Science*, 328, 725
- . 2010d, *ApJS*, 188, 405
- . 2010e, *ApJ*, 718, 348

⁶<http://leejjoon.github.com/pywcsgrid2/>

⁷<http://healpix.jpl.nasa.gov/>

- 996 —. 2010f, ArXiv e-prints
- 997 —. 2010g, ArXiv e-prints
- 998 —. 2010h, Science, 327, 1103
- 999 —. 2010i, ApJ, 712, 459
- 1000 —. 2010j, A&A
- 1001 —. 2010k, Physical Review Letters, 104, 101101
- 1002 —. 2010l, ApJS, 187, 460
- 1003 —. 2011a, ArXiv e-prints
- 1004 —. 2011b, ArXiv e-prints
- 1005 —. 2011c, ApJ, 734, 28
- 1006 —. 2012, ApJS, in preparation
- 1007 Acciari, V. A., et al. 2009, ApJ, 698, L133
- 1008 —. 2011, ArXiv e-prints
- 1009 Ackermann, M., et al. 2011, ArXiv e-prints
- 1010 Aharonian, F., et al. 2005, A&A, 435, L17
- 1011 —. 2006a, A&A, 460, 743
- 1012 —. 2006b, A&A, 460, 365
- 1013 —. 2006c, A&A, 448, L43
- 1014 Aharonian, F., et al. 2006, ApJ, 636, 777
- 1015 Aharonian, F., et al. 2007a, ApJ, 661, 236
- 1016 —. 2007b, A&A, 464, 235
- 1017 —. 2008, A&A, 481, 401
- 1018 Aharonian, F. A., Coppi, P. S., & Voelk, H. J. 1994, ApJ, 423, L5
- 1019 Aharonian, F. A., et al. 2007c, A&A, 469, L1

- 1020 Akaike, H. 1974, IEEE Transactions on Automatic Control, 19, 716
- 1021 Albert, J., et al. 2006, Science, 312, 1771
- 1022 Atwood, W. B., et al. 2009a, ApJ, 697, 1071
- 1023 —. 2009b, ApJ, 697, 1071
- 1024 Balbo, M., Saouter, P., Walter, R., Pavan, L., Tramacere, A., Pohl, M., & Zurita-Heras,
1025 J.-A. 2010, A&A, 520, A111+
- 1026 Baltz, E. A., et al. 2008, J. Cosmology Astropart. Phys., 7, 13
- 1027 Bamba, A., Ueno, M., Koyama, K., & Yamauchi, S. 2003, ApJ, 589, 253
- 1028 Castelletti, G., Dubner, G., Golap, K., & Goss, W. M. 2006, A&A, 459, 535
- 1029 Chen, A. W., et al. 2011, A&A, 525, A33+
- 1030 Dame, T. M. 2011, ArXiv e-prints
- 1031 de Geus, E. J., Bronfman, L., & Thaddeus, P. 1990, A&A, 231, 137
- 1032 Eadie, W. T., Drijard, D., & James, F. E. 1971, Statistical methods in experimental physics,
1033 ed. Eadie, W. T., Drijard, D., & James, F. E.
- 1034 Funk, S., Hinton, J. A., Pühlhofer, G., Aharonian, F. A., Hofmann, W., Reimer, O., &
1035 Wagner, S. 2007, ApJ, 662, 517
- 1036 Gaensler, B. M., Schulz, N. S., Kaspi, V. M., Pivovarov, M. J., & Becker, W. E. 2003, ApJ,
1037 588, 441
- 1038 Gaisser, T. K., Protheroe, R. J., & Stanev, T. 1998, ApJ, 492, 219
- 1039 Giordano, F. 2011, in High-Energy Emission from Pulsars and their Systems, ed. N. Rea &
1040 D. F. Torres (Springer), 69–73
- 1041 Górski, K. M., Hivon, E., Banday, A. J., Wandelt, B. D., Hansen, F. K., Reinecke, M., &
1042 Bartelmann, M. 2005, ApJ, 622, 759
- 1043 Gotthelf, E. V., & Halpern, J. P. 2008, ApJ, 681, 515
- 1044 Grenier, I. A., Casandjian, J.-M., & Terrier, R. 2005, Science, 307, 1292
- 1045 Grondin, M.-H., et al. 2011, ApJ, 738, 42

- 1046 Hwang, U., Petre, R., & Flanagan, K. A. 2008, *ApJ*, 676, 378
- 1047 James, F., & Roos, M. 1975, *Comput. Phys. Commun.*, 10, 343
- 1048 Kargaltsev, O., Pavlov, G. G., & Wong, J. A. 2009, *The Astrophysical Journal*, 690, 891
- 1049 Katagiri, H., et al. 2011, *ApJ*, 741, 44
- 1050 Kerr, M. 2011, PhD in physics, University of Washington
- 1051 King, I. 1962, *AJ*, 67, 471
- 1052 Krause, J., Carmona, E., & Reichardt, I. 2011, in (Presented at the 2011 Fermi Symposium,
1053 Rome Italy)
- 1054 Landi, R., de Rosa, A., Dean, A. J., Bassani, L., Ubertini, P., & Bird, A. J. 2007a, *MNRAS*,
1055 380, 926
- 1056 Landi, R., Masetti, N., Bassani, L., Cellone, S. A., Romero, G. E., Ubertini, P., & Dean,
1057 A. J. 2007b, *The Astronomer’s Telegram*, 1047, 1
- 1058 Markwardt, C. B., & Ogelman, H. 1995, *Nature*, 375, 40
- 1059 Matsumoto, H., et al. 2007, *PASJ*, 59, 199
- 1060 —. 2008, *PASJ*, 60, 163
- 1061 Mattox, J. R., et al. 1996, *ApJ*, 461, 396
- 1062 Mizukami, T., et al. 2011, *ApJ*, 740, 78
- 1063 Murphy, T., Mauch, T., Green, A., Hunstead, R. W., Piestrzynska, B., Kels, A. P., &
1064 Sztajer, P. 2007, *MNRAS*, 382, 382
- 1065 Neronov, A., Semikoz, D. V., Tinyakov, P. G., & Tkachev, I. I. 2011, *A&A*, 526, A90
- 1066 Paron, S., Dubner, G., Reynoso, E., & Rubio, M. 2008, *A&A*, 480, 439
- 1067 Petre, R., Becker, C. M., & Winkler, P. F. 1996, *ApJ*, 465, L43+
- 1068 Pollock, A. M. T. 1985, *A&A*, 150, 339
- 1069 Protassov, R., van Dyk, D. A., Connors, A., Kashyap, V. L., & Siemiginowska, A. 2002,
1070 *ApJ*, 571, 545
- 1071 Reynoso, E. M., Dubner, G. M., Goss, W. M., & Arnal, E. M. 1995, *AJ*, 110, 318

- 1072 Reynoso, E. M., Green, A. J., Johnston, S., Dubner, G. M., Giacani, E. B., & Goss, W. M.
1073 2003, MNRAS, 345, 671
- 1074 Rodriguez, J., Tomsick, J. A., Foschini, L., Walter, R., Goldwurm, A., Corbel, S., & Kaaret,
1075 P. 2003, A&A, 407, L41
- 1076 Rowell, G., Horns, D., Fukui, Y., & Moriguchi, Y. 2008, in American Institute of Physics
1077 Conference Series, Vol. 1085, American Institute of Physics Conference Series, ed.
1078 F. A. Aharonian, W. Hofmann, & F. Rieger, 241–244
- 1079 Schwarz, G. 1978, The Annals of Statistics, 6, pp. 461
- 1080 Sreekumar, P., et al. 1998, The Astrophysical Journal, 494, 523
- 1081 Strong, A. W., & Moskalenko, I. V. 1998, ApJ, 509, 212
- 1082 Sugizaki, M., Mitsuda, K., Kaneda, H., Matsuzaki, K., Yamauchi, S., & Koyama, K. 2001,
1083 ApJS, 134, 77
- 1084 Tanaka, T., et al. 2011, ApJ, 740, L51
- 1085 Taylor, A. R., et al. 2003, AJ, 125, 3145
- 1086 Tomsick, J. A., Lingenfelter, R., Walter, R., Rodriguez, J., Goldwurm, A., Corbel, S., &
1087 Kaaret, P. 2003, IAU Circ., 8076, 1
- 1088 Torii, K., et al. 1998, ApJ, 494, L207+
- 1089 Vladimirov, A. E., et al. 2011, Computer Physics Communications, 182, 1156
- 1090 Weekes, T. C., et al. 1989, ApJ, 342, 379
- 1091 Weinstein, A. 2009, ArXiv e-prints
- 1092 Young, E. T., Lada, C. J., & Wilking, B. A. 1986, ApJ, 304, L45




Fig. 1.— Counts maps and TS profiles for the SNR IC 443. (a) TS vs. extension of the source. (b) TS_{ext} for individual energy bands. (c) observed radial profile of counts in comparison to the expected profiles for a spatially extended source (solid and colored red in the online version) and for a point-like source (dashed and colored blue in the online version). (d) smoothed counts map after subtraction of the diffuse emission compared to the smoothed LAT PSF (inset). Both were smoothed by a 0.1 2D Gaussian kernel. Plots (a), (c), and (d) use only photons with energies between 1 GeV and 100 GeV. Plots (c) and (d) use only photons which converted in the front part of the tracker and have an improved angular resolution.




Fig. 2.— A comparison of a 2D Gaussian and uniform disk spatial model of extended sources before and after convolving with the PSF for two energy ranges. The solid black line is the PSF that would be observed for a power-law source of spectral index 2. The dashed line and the dash-dotted lines are the brightness profile of a Gaussian with $r_{68} = 0.5$ and the convolution of this profile with the LAT PSF respectively (colored red in the online version). The dash-dot-dotted and the dot-dotted lines are the brightness profile of a uniform disk with $r_{68} = 0.5$ and the convolution of this profile with the LAT PSF respectively (colored blue in the online version).

Table 1. Monte Carlo Spectral Parameters

Spectral Index	Flux ^(a) (ph cm ⁻² s ⁻¹)	$N_{1-100\text{GeV}}$	$\langle\text{TS}\rangle_{1-100\text{GeV}}$	$N_{10-100\text{GeV}}$	$\langle\text{TS}\rangle_{10-100\text{GeV}}$
Isotropic Background					
1.5	3×10^{-7}	18938	22233	18938	8084
	10^{-7}	19079	5827	19079	2258
	3×10^{-8}	19303	1276	19303	541
	10^{-8}	19385	303	19381	142
	3×10^{-9}	18694	62	12442	43
2	10^{-6}	18760	22101	18760	3033
	3×10^{-7}	18775	4913	18775	730
	10^{-7}	18804	1170	18803	192
	3×10^{-8}	18836	224	15256	50
	10^{-8}	17060	50
2.5	3×10^{-6}	18597	19036	18597	786
	10^{-6}	18609	4738	18608	208
	3×10^{-7}	18613	954	15958	53
	10^{-7}	18658	203
	3×10^{-8}	14072	41
3	10^{-5}	18354	19466	18354	215
	3×10^{-6}	18381	4205	15973	54
	10^{-6}	18449	966
	3×10^{-7}	18517	174
	10^{-7}	13714	41
Galactic and Isotropic Diffuse Background ^(b)					
1.5	2.3×10^{-8}	90741	63
2	1.2×10^{-7}	92161	60
2.5	4.5×10^{-7}	86226	47
3	2.0×10^{-6}	94412	61

^(a)Integral 100 MeV to 100 GeV flux.

^(b)**For the Galactic simulations, the quoted fluxes are if it was placed in the galactic center and is scaled by Equation 11.**

Note. — A list of the spectral models of the simulated point-like sources which were tested for extension. For each model, the number of statistically independent simulations and the

average value of TS is also tabulated. **The top rows are the sources simulated on top of an isotropic background and the bottom rows are the simulations on top of the Galactic and isotropic background.**




Fig. 3.— Cumulative distribution of the TS for the extension test when fitting simulated point-like sources in the 1 GeV to 100 GeV energy range. The four plots represent simulated sources of different spectral indices and the different lines (colored in the online version) represent point-like sources with different 100 MeV to 100 GeV integral fluxes. The dashed line (colored red) is the cumulative density function of Equation 10.




Fig. 4.— The same plot as Figure 4 but fitting in the 10 GeV to 100 GeV energy range.




Fig. 5.— The same plot as Figure 3, but fitting sources simulated on top of the Galactic diffuse and isotropic diffuse emission.




Fig. 6.— **The distribution of TS values when fitting statistically independent simulations of the elliptical ring shaped SNR W44. (a) is the distribution of TS values when fitting W44 as a point-like source and (b) is the distribution of TS_{ext} when fitting the SNR with a Gaussian and uniform disk spatial model. (c) is the distribution of the change in TS when fitting the source with an elliptical disk spatial model compared to fitting it with a radially symmetric spatial model and (d) when fitting the source with a elliptical ring spatial model compared to an elliptical disk spatial model.**




Fig. 7.— the distribution of fit parameters for the simulation of Monte Carlo simulations of W44. The plots show the distribution of (a) (a) best fit flux (b) spectral index (c) 68% containment radius (d) eccentricity (e) elliptical orientation angle (f) inner ring fraction.




Fig. 8.— The detection threshold to resolve an extended source with a uniform disk model for a two-year exposure. All sources have an assumed power-law spectrum and the different line styles (colors in the electronic version) correspond to different simulated spectral indices. The lines with no markers correspond to the detection threshold using photons with energies between 100 MeV and 100 GeV while the lines with star-shaped markers correspond to the threshold using photons with energies between 1 GeV and 100 GeV.




Fig. 9.— The LAT detection threshold for four spectral indices and three backgrounds (1×, 10×, and 100× the Sreekumar-like isotropic background) for a two-year exposure. The left-hand plots are the detection threshold when using photons with energies between 1 GeV and 100 GeV and the right-hand plots are the detection threshold when using photons with energies between 10 GeV and 100 GeV. The flux is integrated only in the selected energy range. Overlaid on this plot are the LAT detected extended sources placed by the magnitude of the nearby Galactic diffuse emission and the energy range they were analyzed with. The star-shaped markers (colored red in the electronic version) are sources with a spectral index closer to 1.5, the triangular markers (colored blue) an index closer to 2, and the circular markers (colored green) an index closer to 2.5. The triangular marker in plot (d) below the sensitivity line is MSH 15–52.

Table 2. Extension Detection Threshold

γ	BG	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1.0	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2.0
E>1 GeV																					
1.5	1×	148.1	23.3	11.3	8.0	7.2	6.9	6.7	6.8	7.1	7.4	7.6	7.9	8.1	8.5	9.2	9.9	9.1	9.2	9.0	10.3
	10×	148.4	29.0	18.7	15.2	15.4	15.0	16.1	16.0	16.8	17.7	18.2	19.3	20.9	22.5	23.8	24.8	21.3	22.8	23.4	23.7
	100×	186.8	55.0	43.4	40.7	41.0	41.8	40.9	40.9	42.7	43.6	38.4	39.9	40.6	38.4	36.9	36.3	37.1	38.8	37.2	37.6
2	1×	328.4	43.4	18.9	13.4	11.2	10.4	10.2	10.2	10.2	10.4	10.7	10.9	11.2	11.5	12.4	12.6	13.0	13.4	14.0	14.4
	10×	341.0	55.9	32.3	27.6	26.5	25.4	25.6	25.9	27.4	26.8	27.8	28.7	29.8	30.1	31.0	31.5	31.7	34.0	34.3	35.9
	100×	420.5	128.3	90.2	77.3	73.3	70.8	67.5	64.3	64.2	64.1	62.8	63.6	61.7	61.9	58.4	59.0	61.4	63.3	60.1	58.1
2.5	1×	627.1	75.6	29.8	19.3	15.5	13.5	12.8	12.6	12.5	12.5	12.6	12.9	12.9	13.1	13.5	13.7	14.3	14.8	15.2	15.8
	10×	638.9	99.1	52.1	39.1	34.6	33.0	32.5	32.5	32.8	33.2	34.1	34.3	34.5	35.1	36.6	36.9	35.5	36.0	36.5	37.3
	100×	795.0	262.1	140.9	104.3	90.4	81.2	77.2	75.1	69.7	70.9	66.5	65.6	64.9	64.0	58.9	58.1	60.2	58.4	57.5	55.8
3	1×	841.5	110.6	43.2	25.5	18.7	16.1	14.4	13.6	13.3	13.2	13.1	13.1	13.4	13.6	13.5	13.8	14.2	14.4	14.8	15.4
	10×	921.6	151.3	69.1	47.8	40.7	37.1	35.5	34.5	35.1	35.5	35.3	35.3	35.4	35.5	36.8	37.6	35.3	35.4	36.3	36.6
	100×	1124.1	282.9	181.1	119.8	100.7	91.1	84.3	77.9	73.3	71.8	67.6	66.4	65.5	63.9	59.0	58.6	58.8	57.5	55.4	54.4
E>10 GeV																					
1.5	1×	44.6	8.0	4.3	3.2	2.7	2.6	2.5	2.5	2.4	2.5	2.5	2.6	2.7	2.8	2.9	2.9	3.1	3.2	3.3	3.4
	10×	45.2	9.2	5.8	5.0	4.9	4.9	5.0	5.2	5.3	5.7	5.9	6.3	6.6	6.5	6.8	7.6	7.8	8.2	8.5	8.7
	100×	47.3	13.4	11.6	10.6	10.8	10.8	12.0	12.7	13.2	13.7	15.3	16.1	17.2	18.2	18.9	19.5	20.4	21.0	21.7	22.9
2	1×	49.7	8.4	4.4	3.3	2.8	2.6	2.6	2.6	2.6	2.6	2.7	2.7	2.8	2.9	3.0	3.2	3.2	3.4	3.5	3.5
	10×	48.6	9.5	6.0	5.2	5.0	5.2	5.2	5.3	5.4	5.8	6.4	6.6	7.0	7.1	7.5	8.0	8.3	8.6	9.0	9.2
	100×	51.8	14.7	11.8	11.5	11.5	11.9	13.2	14.0	14.3	15.3	16.2	16.9	18.4	19.2	19.8	21.0	22.0	22.8	23.2	24.3
2.5	1×	53.1	9.1	4.5	3.3	2.8	2.7	2.6	2.5	2.5	2.6	2.7	2.7	2.8	2.8	2.9	3.1	3.2	3.3	3.5	3.6
	10×	53.7	10.5	6.3	5.4	5.1	5.1	5.3	5.4	5.7	6.0	6.3	6.6	6.8	6.9	7.5	8.1	8.3	8.6	8.9	9.2
	100×	57.0	15.6	12.7	11.9	11.8	12.2	13.1	14.3	14.6	15.2	16.3	17.0	18.8	19.2	19.9	21.0	21.9	22.3	23.3	23.7
3	1×	55.5	9.4	4.8	3.4	2.9	2.7	2.6	2.5	2.5	2.5	2.6	2.7	2.7	2.8	2.9	3.0	3.1	3.2	3.4	3.4
	10×	56.0	10.5	6.2	5.3	5.1	5.1	5.1	5.3	5.5	5.7	5.9	6.4	6.4	6.6	7.0	7.8	8.0	8.3	8.6	8.9
	100×	60.3	16.2	12.7	11.7	11.8	12.2	12.6	13.8	14.2	14.6	15.8	16.5	17.6	18.5	19.4	19.8	20.7	21.0	21.8	22.5

Note. — The detection threshold to resolve spatially extended sources with a uniform disk spatial model for a two-year exposure. The threshold is calculated for sources of varying energy ranges, spectral indices, and background levels. The sensitivity was calculated against a Sreekumar-like isotropic background and the second column is the factor that the simulated background was scaled by. The remaining columns are varying sizes of the source. The table quotes integral fluxes in the analyzed energy range (1 GeV to 100 GeV or 10 GeV to 100 GeV) in units of $10^{-10} \text{ ph cm}^{-2} \text{ s}^{-1}$.




Fig. 10.— The projected detection threshold of the LAT to extension after 10 years for a power-law source of spectral index 2 against 10 times the isotropic background in the energy range from 1 GeV to 100 GeV (solid line colored red in the electronic version) and 10 GeV to 100 GeV (dashed line colored blue). The shaded gray regions represent the detection threshold assuming the sensitivity improves from 2 to 10 years by the square root of the exposure (top edge) and linearly with exposure (bottom edge). The lower plot shows the factor increase in sensitivity. For small extended sources, the detection threshold of the LAT to the extension of a source will improve by a factor larger than the square root of the exposure.




Fig. 11.— (a) and (b) are the distribution of TS_{ext} and of $TS_{2\text{pts}}$ when fitting simulated spatially extended sources of varying sizes as both an extended source and as two point-like sources. (c) and (d) are the distribution of $TS_{\text{ext}} - TS_{2\text{pts}}$ for the same simulated sources. (a) and (c) represent sources fit in the 1 GeV to 100 GeV energy range and (b) and (d) represent sources fit in the 10 GeV to 100 GeV energy range. In (c) and (d), the plus-shaped markers (colored red in the electronic version) are fits where $TS_{\text{ext}} \geq 16$.




Fig. 12.— The distribution of $TS_{\text{ext}} - TS_{2\text{pts}}$ when fitting two simulated point-like sources of varying separations as both an extended source and as two point-like sources. (a), and (b) represent simulations of two point-like sources with the same spectral index and (c) and (d) represent simulations of two point-like sources with different spectral indices. (a) and (c) fit the simulated sources in the 1 GeV to 100 GeV energy range and (b) and (d) fit in the 10 GeV to 100 GeV energy range. The plus-shaped markers (colored red in the electronic version) are fits where $TS_{\text{ext}} \geq 16$.




Fig. 13.— The cumulative density of TS_{ext} for the 733 clean AGN in 2LAC that were significant above 1 GeV calculated with `pointlike` (dashed line colored blue in the electronic version) and with `gtlike` (solid line colored black). AGN are too far and too small to be resolved by the LAT. Therefore, the cumulative density of TS_{ext} is expected to follow a $\chi^2_1/2$ distribution (Equation 10, the dash-dotted line colored red).




Fig. 14.— A TS map generated for the region around the SNR IC 443 using photons with energies between 1 GeV and 100 GeV. (a) TS map after subtracting IC 443 modeled as a point-like source. (b) same as (a), but IC 443 modeled as an extended source. The cross represents the best fit position of IC 443.




Fig. 15.— A diffuse-emission-subtracted 1 GeV to 100 GeV counts map of the region around 2FGL J1856.2+0450c smoothed by a 0.1° 2D Gaussian kernel. The plus-shaped marker and circle (colored red in the online version) represent the center and size of the source fit with a radially-symmetric uniform disk spatial model. The black crosses represent the positions of other 2FGL sources. The extension is statistically significant, but the extension encompasses many 2FGL sources and the emission does not look to be uniform. Although the fit is statistically significant, it likely corresponds to residual features of inaccurately modeled diffuse emission picked up by the fit.

Table 3. Analysis of the twelve extended sources included in the 2FGL catalog

Name	GLON (deg.)	GLAT (deg.)	σ (deg.)	TS	TS _{ext}	Pos Err (deg.)	Flux ^(a)	Index
E>1 GeV								
SMC	302.59	−44.42	$1.32 \pm 0.15 \pm 0.31$	95.0	52.9	0.14	2.7 ± 0.3	2.48 ± 0.19
LMC	279.26	−32.31	$1.37 \pm 0.04 \pm 0.11$	1127.9	909.9	0.04	13.6 ± 0.6	2.43 ± 0.06
IC 443	189.05	3.04	$0.35 \pm 0.01 \pm 0.04$	10692.9	554.4	0.01	62.4 ± 1.1	2.22 ± 0.02
Vela X	263.34	−3.11	0.88					
Centaurus A	309.52	19.42	~ 10					
W28	6.50	−0.27	$0.42 \pm 0.02 \pm 0.05$	1330.8	163.8	0.01	56.5 ± 1.8	2.60 ± 0.03
W30	8.61	−0.20	$0.34 \pm 0.02 \pm 0.02$	464.8	76.0	0.02	29.1 ± 1.5	2.56 ± 0.05
W44	34.69	−0.39	$0.35 \pm 0.02 \pm 0.02$	1917.0	224.8	0.01	71.2 ± 0.5	2.66 ± 0.00
W51 C	49.12	−0.45	$0.27 \pm 0.02 \pm 0.04$	1823.4	118.9	0.01	37.2 ± 1.3	2.34 ± 0.03
Cygnus Loop	74.21	−8.48	$1.71 \pm 0.05 \pm 0.06$	357.9	246.0	0.06	11.4 ± 0.7	2.50 ± 0.10
E>10 GeV								
MSH15−52 ^(b)	320.39	−1.22	$0.21 \pm 0.04 \pm 0.04$	76.3	6.6	0.03	0.6 ± 0.1	2.20 ± 0.22
HESS J1825−137 ^(b)	17.56	−0.47	$0.65 \pm 0.04 \pm 0.02$	59.7	33.8	0.05	1.6 ± 0.2	1.63 ± 0.22

^(a)Integral Flux in units of 10^{-9} ph cm $^{-2}$ s $^{-1}$ and integrated in the fit energy range (either 1 GeV to 100 GeV or 10 GeV to 100 GeV).

^(b)The discrepancy in the best fit spectra of MSH15−52 and HESS J1825−137 compared to Abdo et al. (2010a) and Grondin et al. (2011) is due to fitting over a different energy range.

Note. — All sources were fit using a radially-symmetric uniform disk spatial model. GLON and GLAT are Galactic longitude and latitude of the best fit extended source respectively. The first error on σ is statistical and the second is systematic (see Section 8). The errors on the integral fluxes and the spectral indices are statistical

only. Pos Err is the error on the position of the source. Vela X and the Centaurus A Lobes were not fit in our analysis but are included for completeness.

Table 4. Extension fit for the nine additional extended sources

Name	GLON (deg.)	GLAT (deg.)	σ (deg.)	TS	TS _{ext}	Pos Err (deg.)	Flux ^(a)	Index	Counterpart
E>1 GeV									
2FGL J0823.0–4246	260.32	–3.28	$0.37 \pm 0.03 \pm 0.02$	322.2	48.0	0.02	8.4 ± 0.6	2.21 ± 0.09	Puppis A
2FGL J1627.0–2425c	353.07	16.80	$0.42 \pm 0.05 \pm 0.16$	139.9	32.4	0.04	6.3 ± 0.6	2.50 ± 0.14	Ophiuchus
E>10 GeV									
2FGL J0851.7–4635	266.31	–1.43	$1.15 \pm 0.08 \pm 0.02$	116.6	86.8	0.07	1.3 ± 0.2	1.74 ± 0.21	Vela Jr.
2FGL J1615.0–5051	332.37	–0.13	$0.32 \pm 0.04 \pm 0.01$	50.4	16.7	0.04	1.0 ± 0.2	2.19 ± 0.28	HESS J1616–508
2FGL J1615.2–5138	331.66	–0.66	$0.42 \pm 0.04 \pm 0.02$	76.1	46.5	0.04	1.1 ± 0.2	1.79 ± 0.26	HESS J1614–518
2FGL J1632.4–4753c	336.52	0.12	$0.35 \pm 0.04 \pm 0.02$	64.4	26.9	0.04	1.4 ± 0.2	2.66 ± 0.30	HESS J1632–478
2FGL J1712.4–3941 ^(b)	347.26	–0.53	$0.56 \pm 0.04 \pm 0.02$	59.4	38.5	0.05	1.2 ± 0.2	1.87 ± 0.22	RX J1713.7–3946
2FGL J1837.3–0700c	25.08	0.13	$0.33 \pm 0.07 \pm 0.05$	47.0	18.5	0.07	1.0 ± 0.2	1.65 ± 0.29	HESS J1837–069
2FGL J2021.5+4026	78.24	2.20	$0.63 \pm 0.05 \pm 0.04$	237.2	128.9	0.05	2.0 ± 0.2	2.42 ± 0.19	γ -Cygni

^(a)Integral Flux in units of 10^{-9} ph cm^{–2} s^{–1} and integrated in the fit energy range (either 1 GeV to 100 GeV or 10 GeV to 100 GeV).

^(b)The discrepancy in the best fit spectra of 2FGL J1712.4–3941 compared to Abdo et al. (2011c) is due to fitting over a different energy range.

Note. — The columns in this table have the same meaning as those in Table 3. RX J1713.7–3946 and Vela Jr. were previously studied in dedicated publications (Abdo et al. 2011c; Tanaka et al. 2011).

Table 5. Dual localization, alternative PSF, and alternative approach to modeling the diffuse emission

Name	TS _{pointlike}	TS _{gtlike}	TS _{alt,diff}	TS _{extpointlike}	TS _{extgtlike}	TS _{extalt,diff}	σ (deg.)	$\sigma_{\text{alt,diff}}$ (deg.)	$\sigma_{\text{alt,psf}}$ (deg.)	TS _{2pts}
E>1 GeV										
2FGL J0823.0–4246	331.9	322.2	356.0	60.0	48.0	56.0	0.37	0.39	0.39	23.0
2FGL J1627.0–2425c	154.8	139.9	105.7	39.4	32.4	24.8	0.42	0.40	0.58	24.5
E>10 GeV										
2FGL J0851.7–4635	115.2	116.6	123.1	83.9	86.8	89.8	1.15	1.16	1.17	15.5
2FGL J1615.0–5051 ^(a)	48.2	50.4	56.6	15.2	16.7	17.8	0.32	0.33	0.32	13.1
2FGL J1615.2–5138	75.0	76.1	83.8	42.9	46.5	54.1	0.42	0.43	0.43	35.1
2FGL J1632.4–4753c	64.5	64.4	66.8	23.0	26.9	25.5	0.35	0.36	0.37	10.9
2FGL J1712.4–3941	59.8	59.4	39.9	38.4	38.5	30.7	0.56	0.55	0.53	2.7
2FGL J1837.3–0700c	44.5	47.0	39.2	17.6	18.5	16.1	0.33	0.32	0.38	10.8
2FGL J2021.5+4026	239.1	237.2	255.8	139.1	128.9	138.0	0.63	0.65	0.59	37.3

^(a)Using `pointlike`, TS_{ext} for 2FGL J1615.0–5051 was slightly below 16 when the source was fit in the 10 GeV to 100 GeV energy range. To confirm the extension measure, the extension was refit in `pointlike` using a slightly lower energy. In the 5.6 GeV to 100 GeV energy range, we obtained a consistent extension and TS_{ext} = 28.0. In the rest of this paper, we quote the $E > 10\text{GeV}$ results for consistency with the other sources.

Note. — TS_{pointlike}, TS_{gtlike}, and TS_{alt,diff} are the test statistic values from `pointlike`, `gtlike`, and `gtlike` with the alternative

approach to modeling the diffuse emission respectively. $\text{TS}_{\text{extpointlike}}$, $\text{TS}_{\text{extgtlike}}$, and $\text{TS}_{\text{extalt,diff}}$ are the TS values from **pointlike**, **gtlike**, and **gtlike** with the alternative approach to modeling the diffuse emission respectively. σ , $\sigma_{\text{alt,diff}}$, and $\sigma_{\text{alt,psf}}$ are the fit sizes assuming a radially-symmetric uniform disk model with the standard analysis, the alternative approach to modeling the diffuse emission, and the alternative PSF respectively.

Table 6. Nearby Residual Sources

Extended Source	Nearby Residual Sources
2FGL J0823.0–4246	2FGL J0821.0–4254, 2FGL J0823.4–4305
2FGL J1627.0–2425c	...
2FGL J0851.7–4635	2FGL J0848.5–4535, 2FGL J0853.5–4711, 2FGL J0855.4–4625
2FGL J1615.0–5051	...
2FGL J1615.2–5138	2FGL J1614.9–5212
2FGL J1632.4–4753c	2FGL J1634.4–4743c
2FGL J1712.4–3941	...
2FGL J1837.3–0700c	2FGL J1835.5–0649
2FGL J2021.5+4026	2FGL J2019.1+4040

Note. — **For each of the new extended sources, we list any catalog source that was included in 2FGL to account for residuals induced by not modeling the extension of a nearby source. These sources were removed from our model of the sky when studying the extended source.**




Fig. 16.— A diffuse-emission-subtracted 1 GeV to 100 GeV counts map of 2FGL J0823.0–4246 smoothed by a 0.1° 2D Gaussian kernel. The triangular marker (colored red in the online version) represents the 2FGL position of this source. The plus-shaped marker and the circle (colored red) represent the best fit position and extension of this source assuming a radially-symmetric uniform disk model. The two star-shaped markers (colored green) represent 2FGL sources that were removed from the background model. **From left to right, these sources are 2FGL J0823.4–4305 and 2FGL J0821.0–4254.** The lower right inset is the model predicted emission from a point-like source with the same spectrum as 2FGL J0823.4–4305 smoothed by the same kernel. This source is spatially coincident with the Puppis A SNR. The light blue contours correspond to the X-ray image of Puppis A observed by *ROSAT* (Petre et al. 1996).




Fig. 17.— The spectral energy distribution of the extended sources Puppis A (2FGL J0823.0–4246) and γ -Cygni (2FGL J2021.5+4026). The lines (colored red in the online version) are the best fit power-law spectral models of these sources. **Puppis A has a spectral index of 2.21 ± 0.09 and γ -Cygni has a spectral index of 2.42 ± 0.19 .** The spectral errors are statistical only. The upper limit is at the 95% confidence level.




Fig. 18.— A diffuse-emission-subtracted 1 GeV to 100 GeV counts map of (a) the region around 2FGL J1627.0–2425 smoothed by a $0^\circ.1$ 2D Gaussian kernel (b) also the emission from 2FGL J1625.7–2526 subtracted. The triangular marker (colored red in the online version) represents the 2FGL position of this source. The plus-shaped marker and the circle (colored red) represent the best fit position and extension of this source assuming a radially-symmetric uniform disk model and the black cross represents a background 2FGL source. The contours in (a) correspond to the $100\ \mu\text{m}$ image observed by IRAS (Young et al. 1986). The contours in (b) correspond to CO ($J = 1 \rightarrow 0$) emission integrated from $-8\ \text{km s}^{-1}$ to $20\ \text{km s}^{-1}$. They are from de Geus et al. (1990), were cleaned using the moment-masking technique (Dame 2011), and have been smoothed by a $0^\circ.25$ 2D Gaussian kernel.




Fig. 19.— A diffuse-emission-subtracted 10 GeV to 100 GeV counts map of 2FGL J0851.7–4635 smoothed by a 0.25° 2D Gaussian kernel. The triangular marker (colored red in the electronic version) represents the 2FGL position of this source. The plus-shaped marker and the circle (colored red) are the best fit position and extension of this source assuming a radially-symmetric uniform disk model. **The three black crosses represent background 2FGL sources. The three star-shaped markers (colored green) represent other 2FGL sources that were removed from the background model. They are (from left to right) 2FGL J0853.5–4711, 2FGL J0855.4–4625, and 2FGL J0848.5–4535.** The circular and square-shaped marker (colored blue) represents the 2FGL and relocalized position of another 2FGL source. This extended source is spatially coincident with the Vela Jr. SNR. The contours (colored light blue) correspond to the TeV image of Vela Jr. (Aharonian et al. 2007a).




Fig. 20.— A diffuse-emission-subtracted 10 GeV to 100 GeV counts map of 2FGL J1615.0–5051 (upper left) and 2FGL J1615.2–5138 (lower right) smoothed by a $0^\circ.1$ 2D Gaussian kernel. The triangular markers (colored red in the electronic version) represent the 2FGL positions of these sources. The cross-shaped markers and the circles (colored red) represent the best fit positions and extensions of these sources assuming a radially symmetric uniform disk model. **The two black crosses represent background 2FGL sources and the star-shaped marker (colored green) represents 2FGL J1614.9-5212, another 2FGL source that was removed from the background model.** The contours (colored light blue) correspond to the TeV image of HESS J1616–508 (left) and HESS J1614–518 (right) (Aharonian et al. 2006).




Fig. 21.— The spectral energy distribution of four extended sources associated with unidentified extended TeV sources. The black points with circular markers are obtained by the LAT. The points with plus-shaped markers (colored red in the electronic version) are for the associated H.E.S.S. sources. (a) the LAT SED of 2FGL J1615.0–5051 together with the H.E.S.S. SED of HESS J1616–508. (b) 2FGL J1615.2–5138 and HESS J1614–518. (c) 2FGL J1632.4–4753c and HESS J1632–478. (d) 2FGL J1837.3–0700c and HESS J1837–069. The H.E.S.S. data points are from (Aharonian et al. 2006). Both LAT and H.E.S.S. spectral errors are statistical only.




Fig. 22.— A diffuse-emission-subtracted 10 GeV to 100 GeV counts map of 2FGL J1632.4–4753c (a) smoothed by a 0.1° 2D Gaussian kernel and (b) with the emission from the background sources subtracted. The triangular marker (colored red in the electronic version) represents the 2FGL position of this source. The plus-shaped marker and the circle (colored red) are the best fit position and extension of 2FGL J1632.4–4753c assuming a radially-symmetric uniform disk model. The four black crosses represent background 2FGL sources subtracted in (b). The circular and square-shaped markers (colored blue) represent the 2FGL and relocalized positions respectively of two additional background 2FGL sources subtracted in (b). **The star-shaped marker (colored green) represents 2FGL J1634.4–4743c, another 2FGL source that was removed from the background model.** The contours (colored light blue) correspond to the TeV image of HESS J1632–478 (Aharonian et al. 2006).




Fig. 23.— A diffuse-emission-subtracted 10 GeV to 100 GeV counts map of 2FGL J1712.4–3941 (a) smoothed by a 0.15° 2D Gaussian kernel and (b) with the emission from the background sources subtracted. This source is spatially coincident with RX J1713.7–3946 and was recently studied in Abdo et al. (2011c). The triangular marker (colored red in the online version) represents the 2FGL position of this source. The plus-shaped marker and the circle (colored red) are the best fit position and extension of this source assuming a radially symmetric uniform disk model. The two black crosses represent background 2FGL sources subtracted in (b). The contours (colored light blue) correspond to the TeV image (Aharonian et al. 2007b).




Fig. 24.— A diffuse-emission-subtracted 10 GeV to 100 GeV counts map of the region around 2FGL J1837.3–0700c (a) smoothed by a 0.15° 2D Gaussian kernel and (b) with the emission from the background sources subtracted. The triangular marker (colored red in the online version) represents the 2FGL position of this source. The plus-shaped marker and the circle (colored red) represent the best fit position and extension of 2FGL J1837.3–0700c assuming a radially-symmetric uniform disk model. The circular and square-shaped markers (colored blue) represent the 2FGL and the relocalized positions respectively of two background 2FGL sources subtracted in (b). **The star-shaped marker (colored green) represents 2FGL J1835.5–0649, another 2FGL source that was removed from the background model.** The contours (colored light blue) correspond to the TeV image of HESS J1837–069 (Aharonian et al. 2006). The diamond-shaped marker (colored orange) represents the position of PSR J1838–0655 and the hexagonal-shaped marker (colored purple) represents the position AX J1837.3–0652 (Gotthelf & Halpern 2008).




Fig. 25.— A diffuse-emission-subtracted 10 GeV to 100 GeV counts map of the region around 2FGL J2021.5+4026 smoothed by a 0.1° 2D Gaussian kernel. The triangular marker (colored red in the online version) represents the 2FGL position of this source. The plus-shaped marker and the circle (colored red) represent the best fit position and extension of 2FGL J2021.5+4026 assuming a radially-symmetric uniform disk model. **The star-shaped marker (colored green) represents 2FGL J2019.1+4040, a 2FGL source that was removed from the background model.** 2FGL J2021.5+4026 is spatially coincident with the γ -Cygni SNR. The contours (colored light blue) correspond to the 408MHz image of γ -Cygni observed by the Canadian Galactic Plane Survey (Taylor et al. 2003).




Fig. 26.— The 21 spatially extended sources detected by the LAT at GeV energies with 2 years of data. The twelve extended sources included in 2FGL are represented by the circular markers (colored red in the online version). The nine new extended sources are represented by the triangular markers (colored orange). The source positions are overlaid on a 100 MeV to 100 GeV Aitoff projection sky map of the LAT data in Galactic coordinates.




Fig. 27.— A comparison of the sizes of extended sources detected at both GeV and TeV energies. The TeV sizes of W30, 2FGL J1837.3–0700c, 2FGL J1632.4–4753c, 2FGL J1615.0–5051, and 2FGL J1615.2–5138 are from Aharonian et al. (2006). The TeV sizes of MSH 15–52, HESS J1825–137, Vela X, Vela Jr., RX J1713.7–3946 and W28 are from Aharonian et al. (2005, 2006b,c, 2007a,b, 2008). The TeV size of IC 443 is from Acciari et al. (2009) and W51C is from Krause et al. (2011). The TeV sizes of MSH 15–52, HESS J1614–518, HESS J1632–478, and HESS J1837–069 have only been reported with an elliptical 2D Gaussian fit and so the plotted sizes are the geometric mean of the semi-major and semi-minor axis. The LAT extension of Vela X is from Abdo et al. (2010f). The TeV sources were fit assuming a 2D Gaussian surface brightness profile so the plotted GeV and TeV extensions were first converted to r_{68} (see Section 2.4). Because of their large sizes, the shape of RX J1713.7–3946 and Vela Jr. were not directly fit at TeV energies and so are not included in this comparison. On the other hand, dedicated LAT publications on these sources showed that their morphologies are consistent (Abdo et al. 2011c; Tanaka et al. 2011). The LAT extension errors are the statistical and systematic errors added in quadrature.




Fig. 28.— The distributions of the sizes of 18 extended LAT sources at GeV energies (colored blue in the electronic version) and the sizes of the 40 extended H.E.S.S. sources at TeV energies (colored red). The H.E.S.S. sources were fit with a 2D Gaussian surface brightness profile so the LAT and H.E.S.S. sizes were first converted to r_{68} . The GeV size of Vela X is taken from Abdo et al. (2010f). Because of their large sizes, the shape of RX J1713.7–3946 and Vela Jr. were not directly fit at TeV energies and are not included in this comparison. Centaurus A is not included because of its large size.




Fig. 29.— The distribution of spectral indices of the 1873 2FGL sources (colored red in the electronic version) and the 21 spatially extended sources (colored blue). The index of Centaurus A is taken from Abdo et al. (2011b) and the index of Vela X is taken from Abdo et al. (2010f).