

## Cluster Difference Imaging Photometric Survey (CDIPS) II: A Jupiter-Sized Planet in IC 2602

L. G. BOUMA,<sup>1</sup> R. BRAHM,<sup>2,3,4</sup> J. D. HARTMAN,<sup>1</sup> P. EVANS,<sup>5</sup> G. ZHOU,<sup>6</sup> J. DE LEON,<sup>7</sup> C. BERGMANN,<sup>8</sup> K. G. STASSUN,<sup>9,10</sup>  
K. A. COLLINS,<sup>6</sup> S. N. QUINN,<sup>6</sup> C. G. TINNEY,<sup>8,11</sup> C. ZIEGLER,<sup>12</sup> C. BRICEÑO,<sup>13</sup> N. LAW,<sup>14</sup> A. W. MANN,<sup>14</sup> J. LIVINGSTON,<sup>7</sup>  
T. HENNING,<sup>15</sup> A. JORDÁN,<sup>16,4</sup> N. ESPINOZA,<sup>17</sup> W. BHATTI,<sup>1</sup> J. N. WINN,<sup>1</sup> G. Á. BAKOS,<sup>1</sup> G. R. RICKER,<sup>18</sup> R. VANDERSPEK,<sup>18</sup>  
D. W. LATHAM,<sup>6</sup> S. SEAGER,<sup>19</sup> J. M. JENKINS,<sup>20</sup> AND TSO/SPOC/POC REPRESENTATIVES

<sup>1</sup>*Department of Astrophysical Sciences, Princeton University, 4 Ivy Lane, Princeton, NJ 08540, USA*

<sup>2</sup>*Center of Astro-Engineering UC, Pontificia Universidad Católica de Chile, Av. Vicuña Mackenna 4860, 782-0436 Macul, Santiago, Chile*

<sup>3</sup>*Instituto de Astrofísica, Facultad de Física, Pontificia Universidad Católica de Chile, Av. Vicuña Mackenna 4860, 782-0436 Macul, Santiago, Chile*

<sup>4</sup>*Millennium Institute of Astrophysics, Av. Vicuña Mackenna 4860, 782-0436 Macul, Santiago, Chile*

<sup>5</sup>*El Sauce Observatory, Coquimbo Province, Chile*

<sup>6</sup>*Center for Astrophysics | Harvard & Smithsonian, 60 Garden St, Cambridge, MA 02138, USA*

<sup>7</sup>*Department of Astronomy, University of Tokyo, 7-3-1 Hongo, Bunkyo-ky, Tokyo 113-0033, Japan*

<sup>8</sup>*Exoplanetary Science at UNSW, School of Physics, UNSW Sydney, NSW 2052, Australia*

<sup>9</sup>*Vanderbilt University, Department of Physics & Astronomy, 6301 Stevenson Center Lane, Nashville, TN 37235, USA*

<sup>10</sup>*Fisk University, Department of Physics, 1000 17th Avenue N., Nashville, TN 37208, USA*

<sup>11</sup>*Australian Centre for Astrobiology, UNSW Sydney, NSW 2052, Australia*

<sup>12</sup>*Dunlap Institute for Astronomy and Astrophysics, University of Toronto, 50 St. George Street, Toronto, Ontario M5S 3H4, Canada*

<sup>13</sup>*Cerro Tololo Inter-American Observatory, Casilla 603, La Serena, Chile*

<sup>14</sup>*Department of Physics and Astronomy, The University of North Carolina at Chapel Hill, Chapel Hill, NC 27599-3255, USA*

<sup>15</sup>*Max-Planck-Institut für Astronomie, Königstuhl 17, 69117 Heidelberg, Germany*

<sup>16</sup>*Facultad de Ingeniería y Ciencias, Universidad Adolfo Ibáñez, Av. Diagonal las Torres 2640, Peñalolén, Santiago, Chile*

<sup>17</sup>*Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA*

<sup>18</sup>*Department of Physics and Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, Cambridge, MA 02139, USA*

<sup>19</sup>*Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA 02139, USA*

<sup>20</sup>*NASA Ames Research Center, Moffett Field, CA 94035, USA*

(Received May 28, 2020; Revised —; Accepted —)

Submitted to AAS journals.

### ABSTRACT

We report the validation of the transiting planet TOI 837.01, a warm Jupiter-sized planet ( $R_p = 10.XX R_\oplus$ ,  $P = 8.3Y$  d), with mass less than  $X.XX M_{\text{Jup}}$  ( $3\sigma$ ). TOI 837.01 orbits a  $V=10.6$ ,  $T=9.9$  G-dwarf in IC 2602 (the “Southern Pleiades”). It is therefore 50 million years old. We characterize the system using data from the NASA *Transiting Exoplanet Survey Satellite* (TESS), the ESA Gaia mission, ground-based photometry, and spectroscopy from CTIO1.5/CHIRON, XXX/FEROS, and AAT/Veloce. TOI 837.01 joins the growing ranks of adolescent exoplanets orbiting bright host stars, and is amenable for studies of planetary atmospheric and orbital evolution.

**Keywords:** Exoplanets (XXX), Exoplanet evolution (491), Stellar ages (1581), Young star clusters (XXX)

### 1. INTRODUCTION

Recently, young planets have been hype.


TESS (Ricker et al. 2015).

Section 2 Section 3 Section 4 Section 5 Section 6.

### 2. THE DATA

#### 2.1. TESS Photometry

TOI 837 (GAIA DR2 5251470948229949568) was observed by the TESS spacecraft from X to Y in two-minute cadence mode. The star was designated TIC 460205581 in the TESS Input Catalog (Stassun et al. 2018, 2019). The pixel data for an  $11 \times 11$  array surrounding TOI 837 were averaged into 2-minute stacks by the onboard computer. Each  $2048 \times 2048$  image from the CCD was also averaged into 30-minute stacks, and saved as a “full frame image” (FFI).



**Figure 1. TESS lightcurve of TOI 837 (Sectors 10 and 11).** *Top:* PDCSAP mean-subtracted relative flux at 2-minute sampling. Spot-induced stellar variability is the dominant signal. Dashed lines show the five transits observed by TESS. *Bottom:* Zoomed windows of individual transits. Red points show manually identified stellar flares.

The pixel data for an The 2-minute stacks for TOI 837 were reduced to lightcurves by the Science Processing Operations Center (SPOC) at NASA Ames (Jenkins et al. 2016). We mainly used the Presearch Data Conditioning (PDC) lightcurve. The PDC lightcurve aperture used pixels chosen to maximize the SNR of the total flux of the target (?). Non-astrophysical variability was removed by fitting out trends common to many stars (??).

As an independent check on the 2-minute SPOC lightcurve, we examined the lightcurve based upon 30-minute image stacks which was produced as part of the Cluster Difference Imaging Photometric Survey (CDIPS; Bouma et al. 2019). Our CDIPS lightcurve of choice used a circular aperture with radius 1 pixel.

Figure 10.

## 2.2. Ground-based Time-Series Photometry

### 2.3. Imaging

### 2.4. Spectroscopy

#### 2.4.1. CHIRON

9 spectra with CTIO1.5/CHIRON, from January 28 through March 14, 2020. 6 were usable.

#### 2.4.2. FEROS

N spectra with XXX/FEROS

#### 2.4.3. Veloce

M spectra with AAT/Veloce.

## 2.5. Astrometry

## 3. THE STAR

### 3.1. The Cluster

IC 2602 canonically has an age of  $30 \pm 20$  Myr (CITE: Van Leeuwen 2009). Or logage between 7.533 and 7.563 (Bossini et al. 2019).


Or Li age of 40-50 Myr (David+19, comparison with other Li stars).

Reported mean metallicity values  $[Fe/H]$  for the cluster range between slightly super-solar ( $0.04 \pm 0.01$ , Baratella et al. 2020) and slightly sub-solar ( $-0.02 \pm 0.02$ , Netopil et al. 2016).

IC 2602 is supervirial, in the sense that the observed stellar velocity dispersion is larger than the value expected if it were in virial equilibrium by about a factor of two (Bravi et al. 2018).

""""Although IC 2602 does not belong to Sco OB2, it is spatially and dynamically a close relative, and for this reason we include it in this work, if even marginally"" (Damiani+2019).

""""We also find that the open cluster IC 2602 has a similar dynamics to Sco OB2, and its PMS members are currently evaporating and forming a diffuse (size 10deg) halo around its double-peaked core""""



**Figure 2. Phase-folded transit of TOI 837.01.** Gray points are 2-minute PDCSAP flux measurements, detrended with a 0.3-day robust Huber spline (see Section 2.1). Black points are binned to approximately 1 point per 6 minutes. The blue line shows the transit model corresponding to the median transit parameters (Table 2).

### 3.2. Cluster Membership

TOI 837 has been reported as a member of IC 2602 by multiple independent investigators (Kharchenko et al. 2013; Cantat-Gaudin et al. 2018).

The Gaia kinematics are good; gamma velocity is correct; Li (FEROS + Veloce) is strong; the rotation period and vsini agree with sub-Pleiades age expected for an IC 2602 member.

#### 3.2.1. HR Diagram

Figure 3 shows a Hertzsprung-Russell diagram of TOI 837 in reference to the broader IC 2602 cluster, as well as its neighborhood. Cluster members were identified by Cantat-Gaudin et al. (2018) using Gaia-DR2 positions, proper motions, and parallaxes. We included candidate members with formal membership probability exceeding 10%. Save for a few low-mass outliers, most members appear to be young and coeval.

TOI 837 is in its expected position relative to the other members along the cluster isochrone. This photometrically limits the presence of binary companions in the TOI 837 system, to a magnitude difference of perhaps half the brightness of the target star.

Figure 3 also suggests that the membership census of IC 2602 is incomplete. We defined the reference “neighborhood” as the group of at most  $10^4$  randomly selected non-member stars within 5 standard deviations of the mean IC 2602 right ascension, declination, and parallax. We


queried Gaia DR2 for these stars using `astroquery` (Ginsburg et al. 2018). Many low-mass stars appear above the main sequence, even though they were not identified as 5-dimensional kinematic members through the unsupervised ? membership assignment process.

#### 3.2.2. Position and Kinematics

We examine the six-dimensional positions and kinematics in greater depth in Figure 4. The “neighborhood” is defined here identically to as in Figure 3. The axes limits for the right ascension, declination, and parallax dimensions are as in the Gaia archive query. The axes limits for the proper motion and radial velocity dimensions are set at the 25<sup>th</sup> and 75<sup>th</sup> percentiles, in order to give a sense of the population’s distribution, while excluding outlying points. The radial velocities suffer the greatest incompleteness due to the current  $G \approx 12$  magnitude limit of the Gaia-DR2 data processing.

Figure 4 provides strong evidence (i) for the existence of the IC 2602 open cluster and (ii) for the membership of TOI 837 within IC 2602. The only dimension that could perhaps lead to some worry is the parallax, as TOI 837 is one of the closest IC 2602 members reported by Cantat-Gaudin et al. (2018). Fortunately, there are independent means of verifying stellar youth.

#### 3.2.3. Rotation



**Figure 3. Hertzsprung-Russell diagram of TOI 837 and members of IC 2602.** Members (black circles) were identified by Cantat-Gaudin et al. (2018). Gray circles are non-member stars within 5 standard deviations of the mean IC 2602 right ascension, declination, and parallax.  $G$  denotes Gaia broadband magnitudes,  $Bp$  Gaia blue,  $Rp$  Gaia red, and  $\omega_{as}$  the parallax in arcseconds.

TESS photometric rotation period = 3.5days ish.  $v \sin i = 17.48 \pm 0.15$  (CHIRON). The two agree.

For comparison, gyrochrones would “predict” X.XX.

This implies sub-Pleiades age expected for an IC 2602 member.

Figure 5 compares the stellar rotation period of TOI 837 to periods that have been measured in a number of well-studied open clusters. TOI 837 seems to be gyrochronologically coeval with the Pleiades sequence.

We note that this is still in the “sub 500 Myr” range, when classical Skumanich spin-down is not expected to be happening (why is this the number?).

This agrees with recent empirical results from Rebull et al 2020 and perhaps also 2018(?): they observed the Taurus foreground (age 30 Myr) and saw that it was roughly consistent with the Pleiades. Upper Sco had a consistent mean, but a much larger scatter.

#### 3.2.4. Lithium

Rafael’s co-added FEROS spectra gave a Li doublet equivalent width of 154 milliångström.

Lithium depletion for a star of TOI-837’s spectral type happens over timescales  $>100$  Myr (Soderblom+2013). This is because the convective envelopes are very shallow, so they

don’t cycle material down to the  $>3e6$  K core until much later.


Nonetheless, comparison of early-G field stars to e.g., 600 Myr old Hyads has shown that the depletion does happen over gigayear timescales (Berger+2018, fig 7).

Figure 6 compares our measured TOI 837 Li EW to stars in the field, and to young open cluster members. The field star measurements were collected by Berger et al. (2018). We have shown the points with lithium detections at  $SNR > 3$ . TOI-837 is consistent with being younger than almost all CKS planet-hosting field stars.

The young open cluster members were selected for the presence of lithium, as described by Randich et al. (2018). The TOI-837 Li EW measured here (150 milliångström) is consistent with what’s seen for stars of similar colors in  $<100$  Myr old moving groups. (Elliott+2016, Fig 11 – attached). This is sensible, because IC 2602 is supposed to be 50 Myr old.

#### 3.2.5. Literature Membership

The membership of TOI 837 in IC 2602 was noted by Oh et al. (2017), in what they dubbed “Group 5”. Analyzing the clusters of Oh et al. (2017), ? fitted the Gaia, 2MASS, and WISE photometry with MIST isochrones, and reported stel-



**Figure 4.** Positions and kinematics of TOI 837 (star), IC 2602 members (black circles), and stars in the neighborhood (gray circles). Members were identified by [Cantat-Gaudin et al. \(2018\)](#).  $\alpha$  denotes right ascension,  $\delta$  declination,  $\pi$  parallax,  $\mu_{\delta}$  and  $\mu_{\alpha}$  proper motion in each equatorial direction, and RV radial velocity reported by Gaia-DR2. The RVs are for unblended spectra of bright stars ( $G \lesssim 12$ ). The proper motion projection ( $\mu_{\delta}$  vs.  $\mu_{\alpha} \cos \delta$ ) highlights the membership selection function.

lar masses, radii, metallicities, ages, distances, and extinction for 9754 stars, including TOI 837.

TOI 837 was also listed as a member of “Theia 92” by ?. These authors identified 999 candidate cluster members, and reported a cluster isochrone age  $\log t = 7.55$  with uncertainty  $\approx 0.15$  dex. They reported an average extinction of  $A_V = 0.243$  mag.

### 3.3. Stellar Parameters

#### 3.3.1. Physical

( $L_{\text{star}}$ ,  $R_{\text{star}}$ ,  $T_{\text{eff}}$ , Age and  $M_{\text{star}}$ )


From CHIRON spectra, my current best guess spectroscopic parameters are as follows.  $T_{\text{eff}} = 5946 \pm 39$   $\log g = 4.48 \pm 0.03$   $\text{feh} = -0.065 \pm 0.035$   $v \sin i = 17.48 \pm 0.15$

Figure 7. The SED fit, looking really good with  $\text{rchi2} = 1.6$ :


The fit parameters are:  $A_V = 0.20 \pm 0.03$   $F_{\text{bol}} = 1.967 \pm 0.046 \text{ e-9 erg/s/cm}^2$   $R_{\text{star}} = 1.049 \pm 0.019 R_{\text{sun}}$

From the spectroscopic  $\log g$  and the above  $R_{\text{star}}$ , we get an independent mass estimate of  $M_{\text{star}} = 1.21 \pm 0.10 M_{\text{sun}}$ . At face value, the mass seems a bit high for the radius, but I’m guessing that just means the spectroscopic  $\log g$  is a bit too high.





**Figure 5. Rotation periods for TOI 837 and selected open clusters.** The Pleiades (120Myr), Praesepe (670Myr), and NGC6811 (1000Myr) are shown. Their rotation periods were measured by [Rebull et al. \(2016\)](#); [Douglas et al. \(2017, 2019\)](#), and [Curtis et al. \(2019\)](#), respectively.




**Figure 6. Lithium 6708Å equivalent widths for TOI 837, field stars, and young open clusters.** The field star sample is drawn from Kepler planet hosts, and was measured by [Berger et al. \(2018\)](#) using Keck-HIRES. The young open clusters members were surveyed by [Randich et al. \(2018\)](#) using the UVES and GIRAFFE spectrographs at the ESO VLT. [Randich et al. \(2018\)](#) found lithium depletion boundary ages for these clusters of  $37.7^{+5.7}_{-4.8}$  Myr (NGC 2547) and  $43.7^{+4.3}_{-3.9}$  Myr (IC 2602).

*Astrometric information*—The RUWE (CITE, CITE) for TOI 837 is 1.02, indicative that there are no obviously present astrometric companions.

### 3.3.2. Rotation

Stellar vsini Stellar rotation

### 3.4. RVs



**Figure 7. Spectral energy distribution of TOI 837 from archival photometry.** Keivan: please describe, and if possible, send me the data (photometry + the atmosphere model) used to generate the plot so I can regenerate it in a consistent style with the rest of the paper

I don't strongly expect the Veloce RVs or the FEROS ones to lead to a mass measurement. The reason is that with  $v \sin i = 17 \text{ km/s}$ , and 2% rotation amplitude signal, we expect an RV RMS of 300 m/s at the  $P_{\text{rot}} = 3.5$  day rotation period ( $v \sin i \times \text{rotation amplitude}$ ). This is probably larger than the expected planet signal (100 m/s, 8 days).

## 4. THE PLANET

### 4.1. Transit Fitting

Simultaneous dip plus rotation period (GP) fit. Use celerite plus PyMC3/exoplanet. Mention FPP?


### 4.2. Validation of TOI 837.01

""Validation is the process of statistically arguing that a transit-like signal is due to a planet rather than an astrophysical false positive. "" (CITE blender, Torres+04,05,10; vespa Morton 12, 15; pastis Diaz+14 Santerne+15; triceratops Gialalone & Dressing 20). In this study, we provide such statistical arguments.

Exclude possibilities: "The transits are blended from a background, unassociated eclipsing binary system or transiting exoplanet: We take the same approach outlined in Vanderburg et al. (2019). In short, if the transits are a blend from the background system, the true radius ratio is constrained by the ratio of the ingress time (T12) to the..."

Given its galactic latitude of only  $-6^\circ$ , it is not surprising that the field of TOI 837 is crowded. A number of possible false-positive explanations, including the usual neighboring, background, and hierarchical eclipsing binary scenarios, must therefore be considered.

For these scenarios to be viable, the flux from TOI 837 and the true eclipsing binary host would blend together, reducing the "true" eclipse depth  $\delta_{\text{true}}$  to produce the observed depth




**Figure 8. Scene used for blend analysis.** *Top:* Mean TESS image of TOI 837 over Sector 10, with a logarithmic grayscale. The yellow star is the position of TOI 837. Orange circles are neighboring stars with  $T < 16$ , scaled such that brighter stars are larger. The X and / hatches show the apertures used to measure the background and target star flux, respectively. Dashed lines of constant declination and right ascension are shown. *Bottom:* Digitized Sky Survey  $R$ -band image of the same field, with a linear grayscale. The circles show apertures of radii 1, 1.5, and 2.25 pixels used in our blend analysis (Section 4.2). Two stars of interest are “Star A” and “Star B”, which were eventually excluded as being possible sources of the transits.

$\delta_{\text{obs}}$ :

$$\delta_{\text{obs}} = \delta_{\text{true}} \frac{F_{\text{neighbor}}}{F_{\text{total}}}, \quad (1)$$

where the total flux and the flux from the neighbor are labelled as such. The requirement that the true eclipse depth



**Figure 9. SOAR HRCam contrast limits derived from point-source injection-recovery experiments.** Star A ( $\Delta T = 4.7$ ,  $\rho = 2.3''$  west) is detected in the autocorrelation function, in addition to being a resolved Gaia source. It is co-moving with TOI 837, and its parallax and on-sky position imply that it is physically separated from TOI 837 by  $6.6 \pm 0.1$  pc.

is less than one translates to a bound on the faintest possible blended companion stars:

$$\Delta m < -\frac{5}{2} \log_{10} \delta_{\text{obs}}. \quad (2)$$

For TOI 837 ( $T = 9.93$ ), this implies that any star invoked to explain the transit depth must be brighter than  $T = 15.92$ .

There’s also

$$\Delta m < -\frac{5}{2} \log_{10} (\delta_{\text{obs}} t_{12}^2 / t_{13}^2). \quad (3)$$

#### 4.2.1. Visual Binarity


Figure 8 shows the scene. In the upper panels, the pixels used to measure the background level in the SPOC lightcurve are indicated with ‘X’ hatching, and the pixels used in the final lightcurve aperture are shown with ‘/’ hatching.

Stars brighter than  $T = 16$ , as queried from the Gaia DR2 source catalog, are shown with orange circles. The relevant stars for a blend analysis are as follows.

- TOI 837 = TIC 460205581 ( $T=9.9$ ). The target star.
- Star A = TIC 847769574 ( $T=14.6$ ).  $2.3''$  west. Proper motions and parallax imply it is comoving with TOI 837, though with a physical separation of  $6.6 \pm 0.1$  pc, it may not be bound.
- Star B = TIC 460205587 ( $T=13.1$ ).  $5.4''$  north. This is a giant background star.

An additional source, TIC 847769581, is  $4.9''$  from the target, but too faint ( $T=18.8$ ) to be the source of the observed transit signal.

The Gaia DR2 data for Star A seems somewhat poorly behaved. While Star A has  $G = 15.1$ , and  $Bp = 14.9$ , no  $Rp$



**Figure 10. Astrophysical false positive scenarios.** *Top:* for bound companions, *Bottom:* for unassociated companions along the same line of sight.

magnitude is reported. Correspondingly, no RUWE value is available (CITE). The astrometric reduced  $\chi^2$  ( $\chi^2/(N-5)$ , for  $N$  the number of good astrometric observations) seems rather poor, at 8.6. We suspect that the photometric failure to produce an  $R_p$  magnitude, as well as the poor astrometric fit, are likely due to blending with TOI 837.

At the angular resolution of the TESS data, if either Star A or Star B were eclipsing binaries, they could be the sources of the transit signal. Coincidence stellar sources below the resolution limit of Gaia are also a concern (CITE). To rule out these possibilities, we took a number of approaches.

#### 4.2.2. High-resolution imaging



We imaged the system using SOAR-HRCam. **Carl: please describe.** Figure 9. Ziegler et al. (2020)

#### 4.2.3. TESS analysis

We examined the CDIPS FFI lightcurves of the target, which are available on MAST (Bouma et al. 2019). Three lightcurves are available, based on photometric apertures with a radius of 1, 1.5, or 2.5 pixels. In the raw difference-image light-curves, as well as the PCA-detrended light-curves, dips of depth  $\approx 0.35\%$  are visible, and their properties do not significantly vary with aperture size. Within  $\approx 20''$ , the dips are consistent with originating from the target star.

#### 4.2.4. Seeing-limited time-series photometry

**left off here** Co-author P. Evans obtained a total transit that showed that, within  $1.5''$ , the transit comes from TOI 837. We can be even more certain with a pixel-level analysis. This 2.1 arcsec neighbor is also a mid to late M dwarf, and bound given the Gaia parallax + proper motion. The transit duration is rather long for the signal to come from the M dwarf, rather than the G dwarf. I think the state of photometric followup could be improved (there is only one good total transit), but I think over this month and next we can pull another few partials.

A total eclipse of Ziegler’s faint companion would produce a 1.4% dip. A 30% eclipse, more plausible for a NEB scenario, would produce dips at exactly the right depth (0.4%).

$\text{dmag} = 4.7$ , transit depth is 0.4%. This  $\text{dmag}$  converts to a flux fraction of 0.0132. A 30% eclipse on T1 could therefore nominally explain the TESS data.

However in the aperture from this green  $r=2\text{px}$  green LC, say 90% of the flux from T1 is excluded. The eclipse on T1 would therefore need to be much deeper – in fact, over 100%. This argument holds as long as at least 70% of the flux from T1 is excluded, so I think we’re likely in the clear.

#### 4.2.5. Summary

Possible false-positive scenarios are as follows.

- *Neighboring blends.*
- *Hierarchical blends.* Limited by i) FEROS + Veloce RVs, and ii) the star being nicely on the cluster HR diagram. Any extra hidden close-in binary stars would need to be very low-mass M dwarfs, and then the transit duration (and lack of chromaticity???) again helps.
- *Background blends.* An argument on the a priori probability of background blends follows from counting statistics. Within 360 arcseconds, there are 298  $T < 16$  stars in TIC8 (i.e., Gaia DR2). At 2.0 arcseconds, SOAR archives  $\text{dmag}(I)$  of 6. Inside of 2 arcseconds, we expect  $\pi r^2$  times ( $N_{\text{star}}$  per sq arcsec)  $= 9e-3$   $T < 16$  stars. Any with  $T \gtrsim 15$  outside of 0.3 arcseconds would have been detected by SOAR, which brings down the expected number of possible hidden sources by a factor of a few. Within 0.3 arcseconds, where SOAR gets

worse than  $\text{dmag}(I)$  of 3, we expect  $2e-4$   $T < 16$  stars. So (order of magnitude), background blends would be a 0.1% coincidence.

Archival SERC-J and AAO-SES plates are available for the TOI 837 field<sup>1</sup>. These plates were acquired in 1982 and 1992, respectively. For high proper motion stars archival imagery can be used to detect slowly moving background stars that might be an astrophysical false-positive source (e.g., Huang et al. 2018). However TOI 837 has only moved  $\approx 0.7''$  between 1982 and present, in comparison to the  $\approx 2.0''$  FWHM of the target on the plates. We therefore cannot resolve it with great confidence from background sources not already resolved through more modern imaging.

#### 4.3. Additional Companions

None of the extra dips in the PDCSAP light-curve (e.g., “up-down” spikes at BTJD 1572 and 1601) seem likely to be planetary. We checked that i) they were not present in the SAPFLUX light-curves, and ii) that they were not present in the CDIPS light-curves (either raw, or PCA-detrended).

To make this quantitative, we did injection-recovery.

### 5. DISCUSSION

In context,...

### 6. CONCLUSIONS

The authors thank... We also thank the Heising-Simons Foundation for their generous support of this work. The Digitized Sky Survey was produced at the Space Telescope Science Institute under U.S. Government grant NAG W-2166. Figure 8 is based on photographic data obtained using the Oschin Schmidt Telescope on Palomar Mountain. This research made use of the Exoplanet Follow-up Observation Program website, which is operated by the California Institute of Technology, under contract with the National Aeronautics and Space Administration under the Exoplanet Exploration Program.

**Software:** *astrobase* (Bhatti et al. 2018), *astropy* (Astropy Collaboration et al. 2018), *astroquery* (Ginsburg et al. 2018), *cdips-pipeline* (Bhatti et al. 2019), *corner* (Foreman-Mackey 2016), *exoplanet* (Agol et al. 2019), *exoplanet* (Foreman-Mackey et al. 2020), and its dependencies (Agol et al. 2019; Kipping 2013; Luger et al. 2019; Theano Development Team 2016). IPython (Pérez & Granger 2007), *lightkurve* (Lightkurve Collaboration et al. 2018), *matplotlib* (Hunter 2007), *MESA* (Paxton et al. 2011, 2013, 2015), *numpy* (Walt et al. 2011), *pandas* (McKinney 2010), *pyGAM* (Servén et al. 2018), *PyMC3* (Salvatier et al. 2016), *scipy* (Jones et al. 2001), *tesscut* (Brasseur et al. 2019), *webplotdigitizer* (?), *wotan* (Hippke et al. 2019).

<sup>1</sup> [https://archive.stsci.edu/cgi-bin/dss\\_form](https://archive.stsci.edu/cgi-bin/dss_form)

*Facilities:* *Astrometry:* Gaia ([Gaia Collaboration et al. 2016, 2018](#)). *Imaging:* Second Generation Digitized Sky Survey, Keck:II (NIRC2; [www2.keck.hawaii.edu/inst/nirc2](http://www2.keck.hawaii.edu/inst/nirc2)). *Spectroscopy:* Keck:I (HIRES; [Vogt et al. 1994](#)). **VLT (number)**, **UVES and GIRAFFE** (CITE: Pasquini et al 2002) *Photometry:* TESS ([Ricker et al. 2015](#)).

**Table 1.** Model Comparison.

| Description    | $N_s$ | $N_\ell$ | $N_{\text{data}}$ | $N_{\text{param}}$ | $\chi^2$ | $\chi^2_{\text{red}}$ | BIC    | $\Delta\text{BIC}$ |
|----------------|-------|----------|-------------------|--------------------|----------|-----------------------|--------|--------------------|
| Favored        | 2     | 2        | 2585              | 17                 | 3230.2   | 1.258                 | 3363.7 | 0.0                |
| Weakly favored | 3     | 3        | 2585              | 21                 | 3203.4   | 1.249                 | 3368.4 | 4.7                |
| —              | 3     | 2        | 2585              | 19                 | 3222.9   | 1.256                 | 3372.2 | 8.4                |
| Disfavored     | 2     | 3        | 2585              | 19                 | 3244.9   | 1.265                 | 3394.2 | 30.4               |
| —              | 2     | 1        | 2585              | 15                 | 3410.6   | 1.327                 | 3528.5 | 164.7              |
| —              | 3     | 1        | 2585              | 17                 | 3396.4   | 1.323                 | 3530.0 | 166.3              |
| —              | 1     | 2        | 2585              | 15                 | 4158.6   | 1.618                 | 4276.4 | 912.7              |
| —              | 1     | 3        | 2585              | 17                 | 4147.4   | 1.615                 | 4281.0 | 917.2              |
| —              | 1     | 1        | 2585              | 13                 | 4313.5   | 1.677                 | 4415.6 | 1051.9             |

NOTE—  $N_s$  and  $N_\ell$  are the number of harmonics at the short and long periods, respectively.  $N_{\text{data}}$  is the number of fitted flux measurements.  $N_{\text{param}}$  is the number of free parameters in the model. The Bayesian information criterion (BIC) and the difference from the maximum  $\Delta\text{BIC}$  are also listed.

**Table 2.** Best-fit model priors and posteriors.

| Param.         | Unit               | Prior                              | Median       | Mean         | Std. Dev. | 3%           | 97%          |
|----------------|--------------------|------------------------------------|--------------|--------------|-----------|--------------|--------------|
| $P_s$          | d                  | $\mathcal{N}(8.3247; 0.0050)$      | 8.3248321    | 8.3248295    | 0.0003356 | 8.3242288    | 8.3254817    |
| $t_s^{(1)}$    | d                  | $\mathcal{N}(1574.273800; 0.0050)$ | 1574.2727299 | 1574.2727323 | 0.0010770 | 1574.2707084 | 1574.2747189 |
| $\log R_p/R_*$ | —                  | $\log \mathcal{U}(0.01; 1.00)$     | -2.43072     | -2.236       | 0.55106   | -2.74890     | -0.99823     |
| $b$            | —                  | $\mathcal{U}(0; 1 + R_p/R_*)$      | 0.9729       | 1.0221       | 0.1533    | 0.9165       | 1.2972       |
| $u_1$          | —                  | $\mathcal{U}(0.186; 0.486)^{(2)}$  | 0.335        | 0.335        | 0.088     | 0.187        | 0.469        |
| $u_2$          | —                  | $\mathcal{U}(0.075; 0.375)^{(2)}$  | 0.229        | 0.227        | 0.086     | 0.093        | 0.374        |
| Mean           | —                  | $\mathcal{U}(0.99; 1.01)$          | 1.000028     | 1.000028     | 0.000006  | 1.000016     | 1.000039     |
| $R_*$          | $R_\odot$          | $\mathcal{T}(1.05; 0.06)$          | 1.05         | 1.05         | 0.06      | 0.94         | 1.16         |
| $\log g$       | cgs                | $\mathcal{N}(4.48; 0.09)$          | 4.46         | 4.46         | 0.08      | 4.32         | 4.61         |
| $R_p/R_*$      | —                  | —                                  | 0.09         | 0.13         | 0.14      | 0.06         | 0.37         |
| $\rho_*$       | $\text{g cm}^{-3}$ | —                                  | 1.42         | 1.44         | 0.28      | 0.97         | 1.97         |
| $R_p$          | $R_{\text{Jup}}$   | —                                  | 0.90         | 1.38         | 1.44      | 0.58         | 3.87         |
| $a/R_*$        | —                  | —                                  | 17.33        | 17.35        | 1.11      | 15.28        | 19.33        |
| $\cos i$       | —                  | —                                  | 0.06         | 0.06         | 0.01      | 0.05         | 0.08         |
| $T_{14}$       | hr                 | —                                  | 1.79         | 1.79         | 0.05      | 1.68         | 1.87         |
| $T_{13}$       | hr                 | —                                  | 0.27         | 0.28         | 0.12      | 0.01         | NaN          |

Note:  $T_{13}$  is ill-defined for a grazing transit.

(1) To convert mean TESS mid-transit time from BTJD to  $\text{BJD}_{\text{TDB}}$ , add 2,457,000.

(2) Assuming an informative quadratic limb-darkening prior with uniform values about those given for the appropriate  $T_{\text{eff}}$  and  $\log g$  from [Claret \(2017\)](#).

## REFERENCES

- Agol, E., Luger, R., & Foreman-Mackey, D. 2019, [arXiv e-prints](#), [1908.03222](#)
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, *AJ*, **156**, 123
- Baratella, M., D’Orazi, V., Carraro, G., et al. 2020, *Astronomy and Astrophysics*, **634**, A34
- Berger, T. A., Howard, A. W., & Boesgaard, A. M. 2018, *The Astrophysical Journal*, **855**, 115
- Bhatti, W., Bouma, L., & Yee, S. 2019, `cdips-pipeline` v0.1.0, <https://doi.org/10.5281/zenodo.3370324>
- Bhatti, W., Bouma, L. G., & Wallace, J. 2018, `astrobase`, <https://doi.org/10.5281/zenodo.1469822>
- Bossini, D., Vallenari, A., Bragaglia, A., et al. 2019, *Astronomy and Astrophysics*, **623**, A108
- Bouma, L. G., Hartman, J. D., Bhatti, W., Winn, J. N., & Bakos, G. Á. 2019, *ApJS*, **245**, 13
- Brasseur, C. E., Phillip, C., Fleming, S. W., Mullally, S. E., & White, R. L. 2019, *Astrophysics Source Code Library*, [ascl:1905.007](#)
- Bravi, L., Zari, E., Sacco, G. G., et al. 2018, *Astronomy and Astrophysics*, **615**, A37
- Cantat-Gaudin, T., Jordi, C., Vallenari, A., et al. 2018, *Astronomy & Astrophysics*, **618**, A93
- Claret, A. 2017, *Astronomy & Astrophysics*, **600**, A30, [arXiv:1804.10295](#)
- Curtis, J. L., Agüeros, M. A., Douglas, S. T., & Meibom, S. 2019, *The Astrophysical Journal*, **879**, 49
- Douglas, S. T., Agüeros, M. A., Covey, K. R., & Kraus, A. 2017, *The Astrophysical Journal*, **842**, 83
- Douglas, S. T., Curtis, J. L., Agüeros, M. A., et al. 2019, *The Astrophysical Journal*, **879**, 100
- Foreman-Mackey, D. 2016, *The Journal of Open Source Software*, **24**
- Foreman-Mackey, D., Czekala, I., Luger, R., et al. 2020, `exoplanet-dev/exoplanet` v0.2.6
- Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016, *A&A*, **595**, A1
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, *A&A*, **616**, A1
- Ginsburg, A., Sipocz, B., Madhura Parikh, et al. 2018, `Astropy/Astroquery: V0.3.7 Release`
- Hippke, M., David, T. J., Mulders, G. D., & Heller, R. 2019, [arXiv:1906.00966 \[astro-ph\]](#), [arXiv:1906.00966](#)
- Huang, C. X., Burt, J., Vanderburg, A., et al. 2018, *ApJ*, **868**, L39
- Hunter, J. D. 2007, *Computing in Science & Engineering*, **9**, 90
- Jenkins, J. M., Twicken, J. D., McCaulliff, S., et al. 2016, *Software and Cyberinfrastructure for Astronomy IV*, **9913**, 99133E
- Jones, E., Oliphant, T., Peterson, P., et al. 2001, *Open source scientific tools for Python*
- Kharchenko, N. V., Piskunov, A. E., Schilbach, E., Röser, S., & Scholz, R.-D. 2013, *Astronomy and Astrophysics*, **558**, A53
- Kipping, D. M. 2013, *MNRAS*, **435**, 2152
- Lightkurve Collaboration, Cardoso, J. V. d. M., Hedges, C., et al. 2018, *Lightkurve: Kepler and TESS time series analysis in Python*, *Astrophysics Source Code Library*, [ascl:1812.013](#)
- Luger, R., Agol, E., Foreman-Mackey, D., et al. 2019, *AJ*, **157**, 64
- McKinney, W. 2010, in *Proceedings of the 9th Python in Science Conference*, ed. S. van der Walt & J. Millman, 51
- Netopil, M., Paunzen, E., Heiter, U., & Soubiran, C. 2016, *Astronomy and Astrophysics*, **585**, A150
- Oh, S., Price-Whelan, A. M., Hogg, D. W., Morton, T. D., & Spergel, D. N. 2017, *The Astronomical Journal*, **153**, 257
- Paxton, B., Bildsten, L., Dotter, A., et al. 2011, *ApJS*, **192**, 3
- Paxton, B., Cantiello, M., Arras, P., et al. 2013, *ApJS*, **208**, 4
- Paxton, B., Marchant, P., Schwab, J., et al. 2015, *ApJS*, **220**, 15
- Pérez, F., & Granger, B. E. 2007, *Computing in Science and Engineering*, **9**, 21
- Randich, S., Tognelli, E., Jackson, R., et al. 2018, *Astronomy & Astrophysics*, **612**, A99
- Rebull, L. M., Stauffer, J. R., Bouvier, J., et al. 2016, *The Astronomical Journal*, **152**, 113
- Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2015, *Journal of Astronomical Telescopes, Instruments, and Systems*, **1**, 014003
- Salvatier, J., Wiecki, T. V., & Fonnesbeck, C. 2016, *PyMC3: Python probabilistic programming framework*
- Servén, D., Brummitt, C., & Abedi, H. 2018, `dswah/pyGAM: v0.8.0`
- Stassun, K. G., Oelkers, R. J., Pepper, J., et al. 2018, *AJ*, **156**, 102
- Stassun, K. G., Oelkers, R. J., Paegert, M., et al. 2019, [arXiv:1905.10694 \[astro-ph\]](#), [arXiv:1905.10694](#)
- Theano Development Team. 2016, [arXiv e-prints](#), [abs/1605.02688](#)
- Vogt, S. S., Allen, S. L., Bigelow, B. C., et al. 1994, *SPIE Conference Series*, ed. D. L. Crawford & E. R. Craine, Vol. 2198
- Walt, S. v. d., Colbert, S. C., & Varoquaux, G. 2011, *Computing in Science & Engineering*, **13**, 22
- Ziegler, C., Tokovinin, A., Briceño, C., et al. 2020, *AJ*, **159**, 19