

= Cretaceous ? Paleogene extinction event =

The Cretaceous ? Paleogene (K ? Pg) extinction event , also known as the Cretaceous ? Tertiary (K ? T) extinction , was a mass extinction of some three @-@ quarters of the plant and animal species on Earth ? including all non @-@ avian dinosaurs ? that occurred over a geologically short period of time approximately 66 million years ago . With the exception of some ectothermic species in aquatic ecosystems like the leatherback sea turtle and crocodiles , no tetrapods weighing more than 55 pounds (25 kilos) survived . It marked the end of the Cretaceous period and with it , the entire Mesozoic Era , opening the Cenozoic Era that continues today .

In the geologic record , the K ? Pg event is marked by a thin layer of sediment called the K ? Pg boundary , which can be found throughout the world in marine and terrestrial rocks . The boundary clay shows high levels of the metal iridium , which is rare in the Earth 's crust but abundant in asteroids .

As originally proposed in 1980 by a team of scientists led by Luis Alvarez , it is now generally thought that the K ? Pg extinction was triggered by a massive comet or asteroid impact 66 million years ago and its catastrophic effects on the global environment , including a lingering impact winter that made it impossible for plants and plankton to carry out photosynthesis . The impact hypothesis , also known as the Alvarez hypothesis , was bolstered by the discovery of the 180 @-@ kilometre @-@ wide (112 mi) Chicxulub crater in the Gulf of Mexico in the early 1990s , which provided conclusive evidence that the K ? Pg boundary clay represented debris from an asteroid impact . The fact that the extinctions occurred at the same time as the impact provides strong situational evidence that the K ? Pg extinction was caused by the asteroid . It was possibly accelerated by the creation of the Deccan Traps . However , some scientists maintain the extinction was caused or exacerbated by other factors , such as volcanic eruptions , climate change , or sea level change , separately or together .

A wide range of species perished in the K ? Pg extinction . The most well @-@ known victims are the non @-@ avian dinosaurs . However , the extinction also destroyed a plethora of other terrestrial organisms , including certain mammals , pterosaurs , birds , lizards , insects , and plants . In the oceans , the K ? Pg extinction killed off plesiosaurs and the giant marine lizards (Mosasauridae) and devastated fish , sharks , mollusks (especially ammonites , which became extinct) and many species of plankton . It is estimated that 75 % or more of all species on Earth vanished . Yet the devastation caused by the extinction also provided evolutionary opportunities . In the wake of the extinction , many groups underwent remarkable adaptive radiations ? a sudden and prolific divergence into new forms and species within the disrupted and emptied ecological niches resulting from the event . Mammals in particular diversified in the Paleogene , producing new forms such as horses , whales , bats , and primates . Birds , fish and perhaps lizards also radiated .

= = Microbiota = =

The K ? Pg boundary represents one of the most dramatic turnovers in the fossil record for various calcareous nanoplankton that formed the calcium deposits that gave the Cretaceous its name . The turnover in this group is clearly marked at the species level . Statistical analysis of marine losses at this time suggests that the decrease in diversity was caused more by a sharp increase in extinctions than by a decrease in speciation . The K ? Pg boundary record of dinoflagellates is not as well @-@ understood , mainly because only microbial cysts provide a fossil record , and not all dinoflagellate species have cyst @-@ forming stages , thereby likely causing diversity to be underestimated . Recent studies indicate that there were no major shifts in dinoflagellates through the boundary layer .

= = Extinction patterns = =

The K ? Pg extinction event was severe , global , rapid , and selective . In terms of severity , the event eliminated a vast number of species . Based on marine fossils , it is estimated that 75 % or

more of all species were made extinct by the K ? Pg extinction event .

The event appears to have affected all continents at the same time . Non @-@ avian dinosaurs , for example , are known from the Maastrichtian of North America , Europe , Asia , Africa , South America and Antarctica , but are unknown from the Cenozoic anywhere in the world . Similarly , fossil pollen shows devastation of the plant communities in areas as far apart as New Mexico , Alaska , China , and New Zealand .

Even though the boundary event was severe , there was significant variability in the rate of extinction between and within different clades . Species that depended on photosynthesis declined or became extinct as atmospheric particles blocked sunlight and reduced the solar energy reaching the Earth 's surface . This plant extinction caused a major reshuffling of the dominant plant groups . Omnivores , insectivores and carrion @-@ eaters survived the extinction event , perhaps because of the increased availability of their food sources . No purely herbivorous or carnivorous mammals seem to have survived . Rather , the surviving mammals and birds fed on insects , worms , and snails , which in turn fed on dead plant and animal matter . Scientists hypothesize that these organisms survived the collapse of plant @-@ based food chains because they fed on detritus (non @-@ living organic material) .

In stream communities , few animal groups became extinct because such communities rely less directly on food from living plants and more on detritus that washes in from the land , buffering them from extinction . Similar , but more complex patterns have been found in the oceans . Extinction was more severe among animals living in the water column than among animals living on or in the sea floor . Animals in the water column are almost entirely dependent on primary production from living phytoplankton , while animals living on or in the ocean floor feed on detritus or can switch to detritus feeding . Coccolithophorids and mollusks (including ammonites , rudists , freshwater snails and mussels) , and those organisms whose food chain included these shell builders , became extinct or suffered heavy losses . For example , it is thought that ammonites were the principal food of mosasaurs , a group of giant marine reptiles that became extinct at the boundary . The largest air @-@ breathing survivors of the event , crocodyliforms and champsosaurs , were semi @-@ aquatic and had access to detritus . Modern crocodilians can live as scavengers and can survive for months without food , and their young are small , grow slowly , and feed largely on invertebrates and dead organisms or fragments of organisms for their first few years . These characteristics have been linked to crocodilian survival at the end of the Cretaceous .

After the K ? Pg extinction event , biodiversity required substantial time to recover , despite the existence of abundant vacant ecological niches .

Radiolaria have left a geological record since at least the Ordovician times , and their mineral fossil skeletons can be tracked across the K ? Pg boundary . There is no evidence of mass extinction of these organisms , and there is support for high productivity of these species in southern high latitudes as a result of cooling temperatures in the early Paleocene . Approximately 46 % of diatom species survived the transition from the Cretaceous to the Upper Paleocene . This suggests a significant turnover in species , but not a catastrophic extinction of diatoms , across the K ? Pg boundary .

The occurrence of planktonic foraminifera across the K ? Pg boundary has been studied since the 1930s . Research spurred by the possibility of an impact event at the K ? Pg boundary resulted in numerous publications detailing planktonic foraminiferal extinction at the boundary . However , there is debate ongoing between groups that think the evidence indicates substantial extinction of these species at the K ? Pg boundary , and those who think the evidence supports multiple extinctions and expansions through the boundary .

Numerous species of benthic foraminifera became extinct during the K ? Pg extinction event , presumably because they depend on organic debris for nutrients , since the biomass in the ocean is thought to have decreased . However , as the marine microbiota recovered , it is thought that increased speciation of benthic foraminifera resulted from the increase in food sources . Phytoplankton recovery in the early Paleocene provided the food source to support large benthic foraminiferal assemblages , which are mainly detritus @-@ feeding . Ultimate recovery of the benthic populations occurred over several stages lasting several hundred thousand years into the

early Paleocene .

== Marine invertebrates ==

There is variability in the fossil record as to the extinction rate of marine invertebrates across the K ? Pg boundary . The apparent rate is influenced by the lack of fossil records rather than actual extinction .

Ostracods , a class of small crustaceans that were prevalent in the upper Maastrichtian , left fossil deposits in a variety of locations . A review of these fossils shows that ostracod diversity was lower in the Paleocene than any other time in the Cenozoic . However , current research cannot ascertain whether the extinctions occurred prior to or during the boundary interval itself .

Approximately 60 % of late @-@ Cretaceous Scleractinia coral genera failed to cross the K ? Pg boundary into the Paleocene . Further analysis of the coral extinctions shows that approximately 98 % of colonial species , ones that inhabit warm , shallow tropical waters , became extinct . The solitary corals , which generally do not form reefs and inhabit colder and deeper (below the photic zone) areas of the ocean were less impacted by the K ? Pg boundary . Colonial coral species rely upon symbiosis with photosynthetic algae , which collapsed due to the events surrounding the K ? Pg boundary . However , the use of data from coral fossils to support K ? Pg extinction and subsequent Paleocene recovery must be weighed against the changes that occurred in coral ecosystems through the K ? Pg boundary .

The numbers of cephalopod , echinoderm , and bivalve genera exhibited significant diminution after the K ? Pg boundary . Most species of brachiopods , a small phylum of marine invertebrates , survived the K ? Pg extinction event and diversified during the early Paleocene .

Except for nautiloids (represented by the modern order Nautilida) and coleoids (which had already diverged into modern octopodes , squids , and cuttlefish) all other species of the molluscan class Cephalopoda became extinct at the K ? Pg boundary . These included the ecologically significant belemnoids , as well as the ammonoids , a group of highly diverse , numerous , and widely distributed shelled cephalopods . Researchers have pointed out that the reproductive strategy of the surviving nautiloids , which rely upon few and larger eggs , played a role in outsurviving their ammonoid counterparts through the extinction event . The ammonoids utilized a planktonic strategy of reproduction (numerous eggs and planktonic larvae) , which would have been devastated by the K ? Pg extinction event . Additional research has shown that subsequent to this elimination of ammonoids from the global biota , nautiloids began an evolutionary radiation into shell shapes and complexities theretofore known only from ammonoids .

Approximately 35 % of echinoderm genera became extinct at the K ? Pg boundary , although taxa that thrived in low @-@ latitude , shallow @-@ water environments during the late Cretaceous had the highest extinction rate . Mid @-@ latitude , deep @-@ water echinoderms were much less affected at the K ? Pg boundary . The pattern of extinction points to habitat loss , specifically the drowning of carbonate platforms , the shallow @-@ water reefs in existence at that time , by the extinction event .

Other invertebrate groups , including rudists (reef @-@ building clams) and inoceramids (giant relatives of modern scallops) , also became extinct at the K ? Pg boundary .

== Fish ==

There are substantial fossil records of jawed fishes across the K ? Pg boundary , which provides good evidence of extinction patterns of these classes of marine vertebrates . Within cartilaginous fish , approximately 7 out of the 41 families of Neoselachian , modern sharks , suffered during this event and Batoids , skates and rays , lost nearly all the identifiable species , while more than 90 % of teleost fish (bony fish) families survived .

While the deep sea realm was able to remain seemingly unaffected , there was an equal loss between the open marine apex predators and the durophagous demersal feeders on the continental shelf . The loss and origination of sharks and batoids at family level are more pronounced . Sharks

and Batoids first appeared in the Danian Age 66 to 61 @.@ 6 million years ago , starting with two shark families (Carcharhinidae , Isuridae) and a single batoid family (Torpedinidae) , resulting in a total origination percentage of only 8 % of marine animals . Only 25 shark families and nine batoid families survived the K @-@ T boundary event . In the late cretaceous period , the Maastrichtian age , 28 shark families and 13 batoid families thrived , before the event occurred . Forty @-@ seven of all marine genera cross the K / T boundary , 85 % being sharks . Batoids display with 15 % a comparably low survival rate .

There is evidence of a mass kill of bony fishes at a fossil site immediately above the K ? Pg boundary layer on Seymour Island near Antarctica , apparently precipitated by the K ? Pg extinction event . However , the marine and freshwater environments of fishes mitigated environmental effects of the extinction event .

= = = Terrestrial invertebrates = = =

Insect damage to the fossilized leaves of flowering plants from fourteen sites in North America were used as a proxy for insect diversity across the K ? Pg boundary and analyzed to determine the rate of extinction . Researchers found that Cretaceous sites , prior to the extinction event , had rich plant and insect @-@ feeding diversity . However , during the early Paleocene , flora were relatively diverse with little predation from insects , even 1 @.@ 7 million years after the extinction event .

= = = Terrestrial plants = = =

There is overwhelming evidence of global disruption of plant communities at the K ? Pg boundary . Extinctions are seen both in studies of fossil pollen , and fossil leaves . In North America , the data suggests massive devastation and mass extinction of plants at the K ? Pg boundary sections , although there were substantial megafloral changes before the boundary . In North America , approximately 57 % of plant species became extinct . In high southern hemisphere latitudes , such as New Zealand and Antarctica , the mass die @-@ off of flora caused no significant turnover in species , but dramatic and short @-@ term changes in the relative abundance of plant groups . In some regions , the Paleocene recovery of plants began with recolonizations by fern species , represented as a fern spike in the geologic record ; this same pattern of fern recolonization was observed after the 1980 Mount St. Helens eruption .

Due to the wholesale destruction of plants at the K ? Pg boundary , there was a proliferation of saprotrophic organisms , such as fungi , that do not require photosynthesis and use nutrients from decaying vegetation . The dominance of fungal species lasted only a few years while the atmosphere cleared and there was plenty of organic matter to feed on . Once the atmosphere cleared , photosynthetic organisms , like ferns and other plants , returned . Polyploidy appears to have enhanced the ability of flowering plants to survive the extinction , probably because the additional copies of the genome such plants possessed allowed them to more readily adapt to the rapidly changing environmental conditions that followed the impact .

= = = Amphibians = = =

There is limited evidence for extinction of amphibians at the K ? Pg boundary . A study of fossil vertebrates across the K ? Pg boundary in Montana concluded that no species of amphibian became extinct . Yet there are several species of Maastrichtian amphibian , not included as part of this study , which are unknown from the Paleocene . These include the frog Theatonius lancensis and the albanerpetontid Albanerpeton galaktion ; therefore some amphibians do seem to have become extinct at the boundary . The relatively low levels of extinction seen among amphibians probably reflect the low extinction rates seen in freshwater animals .

= = = Non @-@ archosaur reptiles = = =

The two living non @-@ archosaurian reptile taxa , testudines (turtles) and lepidosaurs (lizards and tuataras) , along with choristoderes (semi @-@ aquatic archosauromorphs that would die out in the early Miocene) , survived across the K ? Pg boundary . Over 80 % of Cretaceous turtle species passed through the K ? Pg boundary . Additionally , all six turtle families in existence at the end of the Cretaceous survived into the Paleogene and are represented by living species . Living lepidosaurs include the tuataras (the only living rhynchocephalians) and the squamates . The rhynchocephalians were a widespread and relatively successful group of lepidosaurs during the early Mesozoic , but began to decline by the mid @-@ Cretaceous , though they were very successful in the Late Cretaceous of South America . They are represented today by a single genus located exclusively in New Zealand .

The order Squamata , which is represented today by lizards , including snakes and amphisbaenians (worm lizards) , radiated into various ecological niches during the Jurassic and was successful throughout the Cretaceous . They survived through the K ? Pg boundary and are currently the most successful and diverse group of living reptiles with more than 6 @,@ 000 extant species . Many families of terrestrial squamates became extinct at the boundary , such as monstrosaurians and polyglyphanodonts , and fossil evidence indicates they suffered very heavy losses in the KT event , only recovering 10 million years after it . Giant non @-@ archosaurian aquatic reptiles such as mosasaurs and plesiosaurs , which were the top marine predators of their time , became extinct by the end of the Cretaceous . The ichthyosaurs had already disappeared before the mass extinction occurred .

= = = Archosaurs = = =

The archosaur clade includes two surviving groups , crocodilians and birds , along with the various extinct groups of non @-@ avian dinosaurs and pterosaurs .

= = = = Crocodyliforms = = = =

Ten families of crocodilians or their close relatives are represented in the Maastrichtian fossil records , of which five died out prior to the K ? Pg boundary . Five families have both Maastrichtian and Paleocene fossil representatives . All of the surviving families of crocodyliforms inhabited freshwater and terrestrial environments ? except for the Dyrosauridae , which lived in freshwater and marine locations . Approximately 50 % of crocodyliform representatives survived across the K ? Pg boundary , the only apparent trend being that no large crocodiles survived . Crocodyliform survivability across the boundary may have resulted from their aquatic niche and ability to burrow , which reduced susceptibility to negative environmental effects at the boundary . Jouve and colleagues suggested in 2008 that juvenile marine crocodyliforms lived in freshwater environments like modern marine crocodile juveniles , which would have helped them survive where other marine reptiles became extinct ; freshwater environments were not as strongly affected by the K ? Pg extinction event as marine environments .

The Choristodera , a generally crocodile @-@ like group of uncertain phylogeny (possibly archosaurian) also survived the event , only to become extinct in the Miocene .

= = = = Pterosaurs = = = =

One family of pterosaurs , Azhdarchidae , was definitely present in the Maastrichtian , and it likely became extinct at the K ? Pg boundary . These large pterosaurs were the last representatives of a declining group that contained 10 families during the mid @-@ Cretaceous . Several other pterosaur lineages may have been present during the Maastrichtian , such as the ornithocheirids , pteranodontids and / or nyctosaurids , as well as a possible tapejarid , though they are represented by fragmentary remains that are difficult to assign to any given group . While this was occurring , modern birds were undergoing diversification ; traditionally it was thought that they replaced archaic birds and pterosaur groups , possibly due to direct competition , or they simply filled empty niches ,

but there is no correlation between pterosaur and avian diversities that are conclusive to a competition hypothesis , and small pterosaurs were present in the Late Cretaceous .

===== Birds =====

Most paleontologists regard birds as the only surviving dinosaurs (see Origin of birds) . It is thought that all non @-@ avian theropods became extinct , including then @-@ flourishing groups like enantiornithines and hesperornithiforms . Several analyses of bird fossils show divergence of species prior to the K ? Pg boundary , and that duck , chicken and ratite bird relatives coexisted with non @-@ avian dinosaurs . Large collections of bird fossils representing a range of different species provides definitive evidence for the persistence of archaic birds to within 300 @,@ 000 years of the K ? Pg boundary . The absence of these birds in the Paleogene is evidence that a mass extinction of archaic birds took place there . A small fraction of the Cretaceous bird species survived the impact , giving rise to today 's birds . The only bird group known for certain to have survived the K ? Pg boundary is the Aves . Avians may have been able to survive the extinction as a result of their abilities to dive , swim , or seek shelter in water and marshlands . Many species of avians can build burrows , or nest in tree holes or termite nests , all of which provided shelter from the environmental effects at the K ? Pg boundary . Long @-@ term survival past the boundary was assured as a result of filling ecological niches left empty by extinction of non @-@ avian dinosaurs .

===== Non @-@ avian dinosaurs =====

Excluding a few controversial claims , scientists agree that all non @-@ avian dinosaurs became extinct at the K ? Pg boundary . The dinosaur fossil record has been interpreted to show both a decline in diversity and no decline in diversity during the last few million years of the Cretaceous , and it may be that the quality of the dinosaur fossil record is simply not good enough to permit researchers to distinguish between the options . Since there is no evidence that late Maastrichtian non @-@ avian dinosaurs could burrow , swim or dive , they were unable to shelter themselves from the worst parts of any environmental stress that occurred at the K ? Pg boundary . It is possible that small dinosaurs (other than birds) did survive , but they would have been deprived of food , as herbivorous dinosaurs would have found plant material scarce and carnivores would have quickly found prey in short supply .

The growing consensus about the endothermy of dinosaurs (see dinosaur physiology) helps to understand their full extinction in contrast with their close relatives , the crocodilians . Ectothermic (" cold @-@ blooded ") crocodiles have very limited needs for food (they can survive several months without eating) while endothermic (" warm @-@ blooded ") animals of similar size need much more food to sustain their faster metabolism . Thus , under the circumstances of food chain disruption previously mentioned , non @-@ avian dinosaurs died , while some crocodiles survived . In this context , the survival of other endothermic animals , such as some birds and mammals , could be due , among other reasons , to their smaller needs for food , related to their small size at the extinction epoch .

Whether the extinction occurred gradually or suddenly has been debated , as both views have support from the fossil record . A study of 29 fossil sites in Catalan Pyrenees of Europe in 2010 supports the view that dinosaurs there had great diversity until the asteroid impact , with over 100 living species . However , more recent research indicates that this figure is obscured by taphonomical biases and the sparsity of the continental fossil record . The results of this study , which were based on estimated real global biodiversity , showed that between 628 and 1078 non @-@ avian dinosaur species were alive at the end of the Cretaceous and underwent sudden extinction after the Cretaceous ? Paleogene extinction event . Alternatively , interpretation based on the fossil @-@ bearing rocks along the Red Deer River in Alberta supports the gradual extinction of non @-@ avian dinosaurs ; during the last 10 million years of the Cretaceous layers there , the number of dinosaur species seems to have decreased from about 45 to about 12 . Other scientists have pointed out the same .

Several researchers support the existence of Paleocene dinosaurs . Evidence of this existence is based on the discovery of dinosaur remains in the Hell Creek Formation up to 1 @. @ 3 m (4 @. @ 3 ft) above and 40 thousand years later than the K ? Pg boundary . Pollen samples recovered near a fossilized hadrosaur femur recovered in the Ojo Alamo Sandstone at the San Juan River indicate that the animal lived during the Cenozoic , approximately 64 @. @ 5 Ma (about 1 million years after the K ? Pg extinction event) . If their existence past the K ? Pg boundary can be confirmed , these hadrosaurids would be considered a dead clade walking . Scientific consensus is that these fossils were eroded from their original locations and then re @-@ buried in much later sediments (also known as reworked fossils) .

= = = Mammals = = =

All major Cretaceous mammalian lineages , including monotremes (egg @-@ laying mammals) , multituberculates , marsupials and placentals , dryolestoids , and gondwanatheres survived the K ? Pg extinction event , although they suffered losses . In particular , marsupials largely disappeared from North America , and the Asian deltatheroidans , primitive relatives of extant marsupials , became extinct . In the Hell Creek beds of North America , at least half of the ten known multituberculate species and all eleven marsupial species are not found above the boundary . Multituberculates in Europe and North America survived relatively unscathed and quickly bounced back in the Palaeocene , but Asian forms were decimated , never again to represent a significant component on mammalian faunas .

Mammalian species began diversifying approximately 30 million years prior to the K ? Pg boundary . Diversification of mammals stalled across the boundary . Current research indicates that mammals did not explosively diversify across the K ? Pg boundary , despite the environment niches made available by the extinction of dinosaurs . Several mammalian orders have been interpreted as diversifying immediately after the K ? Pg boundary , including Chiroptera (bats) and Cetartiodactyla (a diverse group that today includes whales and dolphins and even @-@ toed ungulates) , although recent research concludes that only marsupial orders diversified after the K ? Pg boundary .

K ? Pg boundary mammalian species were generally small , comparable in size to rats ; this small size would have helped them find shelter in protected environments . In addition , it is postulated that some early monotremes , marsupials , and placentals were semiaquatic or burrowing , as there are multiple mammalian lineages with such habits today . Any burrowing or semiaquatic mammal would have had additional protection from K ? Pg boundary environmental stresses .

= = Evidence = =

= = = North American fossils = = =

In North American terrestrial sequences , the extinction event is best represented by the marked discrepancy between the rich and relatively abundant late @-@ Maastrichtian palynomorph record and the post @-@ boundary fern spike .

At present the most informative sequence of dinosaur @-@ bearing rocks in the world from the K ? Pg boundary is found in western North America , particularly the late Maastrichtian @-@ age Hell Creek Formation of Montana . This formation , when compared with the older (approximately 75 Ma) Judith River / Dinosaur Park Formations (from Montana and Alberta respectively) provides information on the changes in dinosaur populations over the last 10 million years of the Cretaceous . These fossil beds are geographically limited , covering only part of one continent .

The middle ? late Campanian formations show a greater diversity of dinosaurs than any other single group of rocks . The late Maastrichtian rocks contain the largest members of several major clades : Tyrannosaurus , Ankylosaurus , Pachycephalosaurus , Triceratops and Torosaurus , which suggests food was plentiful immediately prior to the extinction .

In addition to rich dinosaur fossils , there are also plant fossils that illustrate the reduction in plant species across the K ? Pg boundary . In the sediments below the K ? Pg boundary the dominant plant remains are angiosperm pollen grains , but the actual boundary layer contains little pollen and is dominated by fern spores . More usual pollen levels gradually resume above the boundary layer . This is reminiscent of areas blighted by modern volcanic eruptions , where the recovery is led by ferns , which are later replaced by larger angiosperm plants .

== Marine fossils ==

The mass extinction of marine plankton appears to have been abrupt and right at the K ? Pg boundary . Ammonite genera became extinct at or near the K ? Pg boundary ; however , there was a smaller and slower extinction of ammonite genera prior to the boundary that was associated with a late Cretaceous marine regression . The gradual extinction of most inoceramid bivalves began well before the K ? Pg boundary , and a small , gradual reduction in ammonite diversity occurred throughout the very late Cretaceous . Further analysis shows that several processes were in progress in the late Cretaceous seas and partially overlapped in time , then ended with the abrupt mass extinction . The diversity of marine life decreased when the climate near the K @-@ T boundary increased in temperature . The temperature increased about three to four degrees very rapidly between 65 @.@ 4 and 65 @.@ 2 million years ago , which is around the time of the extinction event . Not only did the climate temperature increase , but the water temperate decreased causing a drastic decrease in marine diversity .

== Megatsunamis ==

The scientific consensus is that the asteroid impact at the K ? Pg boundary left tsunami deposits and sediments around the area of the Caribbean Sea and Gulf of Mexico . These deposits have been identified in the La Popa basin in northeastern Mexico , platform carbonates in northeastern Brazil , and Atlantic deep @-@ sea sediments .

== Duration ==

The length of time taken for the extinction to occur is a controversial issue , because some theories about the extinction 's causes require a rapid extinction over a relatively short period (from a few years to a few thousand years) while others require longer periods . The issue is difficult to resolve because of the Signor ? Lipps effect ; that is , the fossil record is so incomplete that most extinct species probably died out long after the most recent fossil that has been found . Scientists have also found very few continuous beds of fossil @-@ bearing rock which cover a time range from several million years before the K ? Pg extinction to a few million years after it . The sedimentation rate and thickness of K @-@ Pg clay from three sites suggest short duration of event , perhaps less than ten thousand years .

== Chicxulub asteroid impact ==

== Evidence for impact ==

In 1980 , a team of researchers consisting of Nobel prize @-@ winning physicist Luis Alvarez , his son geologist Walter Alvarez , and chemists Frank Asaro and Helen Michel discovered that sedimentary layers found all over the world at the Cretaceous ? Paleogene boundary contain a concentration of iridium many times greater than normal (30 , 160 and 20 times in three sections originally studied) . Iridium is extremely rare in Earth 's crust because it is a siderophile element , and therefore most of it traveled with the iron as it sank into Earth 's core during planetary differentiation . As iridium remains abundant in most asteroids and comets , the Alvarez team

suggested that an asteroid struck the Earth at the time of the K ? Pg boundary . There were earlier speculations on the possibility of an impact event , but this was the first hard evidence of an impact .

This hypothesis was viewed as radical when first proposed , but additional evidence soon emerged . The boundary clay was found to be full of minute spherules of rock , crystallized from droplets of molten rock formed by the impact . Shocked quartz and other minerals were also identified in the K ? Pg boundary . Shocked minerals have their internal structure deformed , and are created by intense pressures such as those associated with nuclear blasts or meteorite impacts . The identification of giant tsunami beds along the Gulf Coast and the Caribbean also provided evidence for impact , and suggested that the impact may have occurred nearby ? as did the discovery that the K ? Pg boundary became thicker in the southern United States , with meter @-@ thick beds of debris occurring in northern New Mexico .

Further research identified the giant Chicxulub crater , buried under Chicxulub on the coast of Yucatán , as the source of the K ? Pg boundary clay . Identified in 1990 based on work by geophysicist Glen Penfield in 1978 , the crater is oval , with an average diameter of roughly 180 kilometres (110 mi) , about the size calculated by the Alvarez team . The discovery of the crater ? a necessary prediction of the impact hypothesis ? provided conclusive evidence for a K ? Pg impact , and strengthened the hypothesis that the extinction was caused by an impact .

In 2007 , a hypothesis was put forth that argued the impactor that killed the dinosaurs belonged to the Baptistina family of asteroids . Concerns have been raised regarding the reputed link , in part because very few solid observational constraints exist of the asteroid or family . Indeed , it was recently discovered that 298 Baptistina does not share the same chemical signature as the source of the K ? Pg impact . Although this finding may make the link between the Baptistina family and K ? Pg impactor more difficult to substantiate , it does not preclude the possibility . A 2011 WISE study of reflected light from the asteroids of the family estimated the break @-@ up at 80 Ma , giving it insufficient time to shift orbits and impact the Earth by 66 Ma .

In a 2013 paper , Paul Renne of the Berkeley Geochronology Center reported that the date of the asteroid event is 66 ± 0.011 million years ago , based on argon ? argon dating . He further posits that the mass extinction occurred within 32 @, @ 000 years of this date .

== Effects of impact ==

In March 2010 , an international panel of scientists endorsed the asteroid hypothesis , specifically the Chicxulub impact , as being the cause of the extinction . A team of 41 scientists reviewed 20 years of scientific literature and in so doing also ruled out other theories such as massive volcanism . They had determined that a 10 @-@ to @-@ 15 @-@ kilometre (6 @-@ 2 to 9 @-@ 3 mi) space rock hurtled into Earth at Chicxulub on Mexico 's Yucatán Peninsula . The collision would have released the same energy as 100 teratonnes of TNT (420 ZJ) , over a billion times the energy of the atomic bombings of Hiroshima and Nagasaki .

The consequences of the Chicxulub impact were of global extent . Some of these phenomena were brief occurrences that immediately followed the impact , but there were also long @-@ term geochemical and climatic disruptions that were catastrophic to the ecology .

The reentry of ejecta into Earth 's atmosphere would include a brief (hours long) but intense pulse of infrared radiation , killing exposed organisms . Global firestorms likely resulted from the heat pulse . Recent research indicates that the global debris layer deposited by the impact contained enough soot to suggest that the entire terrestrial biosphere had burned .

The impact would have inhibited photosynthesis by creating a dust cloud that blocked sunlight for up to a year . Further , the asteroid struck a region of sulfur @-@ rich carbonate rock , much of which was vaporized , thereby injecting sulfuric acid aerosols into the stratosphere , which might have reduced sunlight reaching the Earth 's surface by more than 50 % , and would have caused rain and ocean water to become acidic . The acidification of the oceans would kill many organisms that build shells from calcium carbonate . At Brazos section , the paleo @-@ sea surface temperature dropped as much as 7 ? for decades after the impact . It would take at least ten years

for such aerosols to dissipate , and would account for the extinction of plants and phytoplankton , and of organisms dependent on them (including predatory animals as well as herbivores) . Some creatures whose food chains were based on detritus would have a reasonable chance of survival .

If widespread fires occurred , they would have increased the CO

2 content of the atmosphere and caused a temporary greenhouse effect once the dust clouds and aerosol settled , and this would have exterminated the most vulnerable organisms that survived the period immediately after the impact .

Most paleontologists now agree that an asteroid did hit the Earth at approximately the end of the Cretaceous , but there is an ongoing dispute whether the impact was the sole cause of the extinctions .

= = Alternative hypotheses = =

The fact that the extinctions occur at the same time as the Chicxulub asteroid impact strongly supports the impact hypothesis of extinction . However , some scientists continue to dispute the role of the Chicxulub impact in driving the extinction , and to suggest that other events may have contributed to the end @-@ Cretaceous mass extinction . In particular , volcanic eruptions , climate change , sea level change , and other impact events have been suggested to play a role in driving the K ? Pg extinction .

= = = Deccan Traps = = =

Before 2000 , arguments that the Deccan Traps flood basalts caused the extinction were usually linked to the view that the extinction was gradual , as the flood basalt events were thought to have started around 68 Mya and lasted more than 2 million years . The most recent evidence shows that the traps erupted over a period of 800 @,@ 000 years spanning the K ? Pg boundary , and therefore may be responsible for the extinction and the delayed biotic recovery thereafter .

The Deccan Traps could have caused extinction through several mechanisms , including the release of dust and sulfuric aerosols into the air , which might have blocked sunlight and thereby reduced photosynthesis in plants . In addition , Deccan Trap volcanism might have resulted in carbon dioxide emissions that increased the greenhouse effect when the dust and aerosols cleared from the atmosphere .

In the years when the Deccan Traps hypothesis was linked to a slower extinction , Luis Alvarez (who died in 1988) replied that paleontologists were being misled by sparse data . While his assertion was not initially well @-@ received , later intensive field studies of fossil beds lent weight to his claim . Eventually , most paleontologists began to accept the idea that the mass extinctions at the end of the Cretaceous were largely or at least partly due to a massive Earth impact . However , even Walter Alvarez has acknowledged that there were other major changes on Earth even before the impact , such as a drop in sea level and massive volcanic eruptions that produced the Indian Deccan Traps , and these may have contributed to the extinctions . The duration of event was less than 10 ky , and the time span is too short to be explained by Deccan volcanism . Geophysical models and high @-@ precision radiometric dating suggest that the Chicxulub impact could have triggered some of the largest Deccan eruptions , and potentially could have triggered eruptions at active volcanoes anywhere on Earth .

= = = Multiple impact event = = =

One other crater also appears to have been formed at about the time of the K ? Pg boundary . Other crater @-@ like topographic features have also been proposed as impact craters formed in connection with Cretaceous @-@ Paleogene extinction . This suggests to some the possibility of near @-@ simultaneous multiple impacts , perhaps from a fragmented asteroidal object , similar to the Shoemaker ? Levy 9 impact with Jupiter . In addition to the 180 km (110 mi) Chicxulub Crater , there is the 24 km (15 mi) Boltysh crater in Ukraine (65 @.@ 17 ± 0 @.@ 64 Ma) , the 20 km (12

mi) Silverpit crater in the North Sea (59 @. @ 5 ± 14 @. @ 5 Ma) possibly formed by bolide impact , and the controversial and much larger 600 km (370 mi) Shiva crater . Any other craters that might have formed in the Tethys Ocean would have been obscured by tectonic events like the northward drift of Africa and India .

= = = Maastrichtian sea @-@ level regression = = =

There is clear evidence that sea levels fell in the final stage of the Cretaceous by more than at any other time in the Mesozoic era . In some Maastrichtian stage rock layers from various parts of the world , the later layers are terrestrial ; earlier layers represent shorelines and the earliest layers represent seabeds . These layers do not show the tilting and distortion associated with mountain building , therefore , the likeliest explanation is a " regression " , that is , a drop in sea level . There is no direct evidence for the cause of the regression , but the explanation currently accepted as most likely is that the mid @-@ ocean ridges became less active and therefore sank under their own weight .

A severe regression would have greatly reduced the continental shelf area , which is the most species @-@ rich part of the sea , and therefore could have been enough to cause a marine mass extinction . However research concludes that this change would have been insufficient to cause the observed level of ammonite extinction . The regression would also have caused climate changes , partly by disrupting winds and ocean currents and partly by reducing the Earth 's albedo and therefore increasing global temperatures .

Marine regression also resulted in the loss of epeiric seas , such as the Western Interior Seaway of North America . The loss of these seas greatly altered habitats , removing coastal plains that ten million years before had been host to diverse communities such as are found in rocks of the Dinosaur Park Formation . Another consequence was an expansion of freshwater environments , since continental runoff now had longer distances to travel before reaching oceans . While this change was favorable to freshwater vertebrates , those that prefer marine environments , such as sharks , suffered .

= = = Multiple causes = = =

In a review article , J. David Archibald and David E. Fastovsky discussed a scenario combining three major postulated causes : volcanism , marine regression , and extraterrestrial impact . In this scenario , terrestrial and marine communities were stressed by the changes in and loss of habitats . Dinosaurs , as the largest vertebrates , were the first affected by environmental changes , and their diversity declined . At the same time , particulate materials from volcanism cooled and dried areas of the globe . Then , an impact event occurred , causing collapses in photosynthesis @-@ based food chains , both in the already @-@ stressed terrestrial food chains and in the marine food chains . The major difference between this hypothesis and the single @-@ cause hypotheses is that its proponents view the suggested single causes as either not sufficient in strength to cause the extinctions or not likely to produce the taxonomic pattern of the extinction .

= = Recovery and radiation = =

The K ? Pg extinction had a profound effect on the evolution of life on Earth . The elimination of dominant Cretaceous groups allowed other organisms to take their place , spurring a remarkable series of adaptive radiations in the Paleogene . The most striking example is the replacement of dinosaurs by mammals . After the K ? Pg extinction , mammals evolved rapidly to fill the niches left vacant by the dinosaurs . Within mammalian genera , new species were approximately 9 @. @ 1 % larger after the K ? Pg boundary .

Other groups also underwent major radiations . Based on molecular sequencing and fossil dating , Neoaves appeared to radiate after the K ? Pg boundary . They even produced giant , flightless forms , such as the herbivorous *Gastornis* and *Dromornithidae* , and the predatory *Phorusrhacidae* .

The extinction of Cretaceous lizards and snakes may have led to the radiation of modern groups such as iguanas , monitor lizards , and boas . On land , giant boid and enormous madtsoiid snakes appeared , and in the seas , giant sea snakes radiated . Teleost fish diversified explosively , filling the niches left vacant by the extinction . Groups appearing in the Paleocene and Eocene include billfish , tunas , eels , and flatfish . Major changes are also seen in Paleogene insect communities . Many groups of ants were present in the Cretaceous , but in the Eocene ants became dominant and diverse , with larger colonies . Butterflies diversified as well , perhaps to take the place of leaf @-@ eating insects wiped out by the extinction . The advanced mound @-@ building termites , Termitidae , also appear to have risen in importance .