

= Fleiss ' kappa =

Fleiss ' kappa ( named after Joseph L. Fleiss ) is a statistical measure for assessing the reliability of agreement between a fixed number of raters when assigning categorical ratings to a number of items or classifying items . This contrasts with other kappas such as Cohen 's kappa , which only work when assessing the agreement between two raters . The measure calculates the degree of agreement in classification over that which would be expected by chance . There is no generally agreed @-@ upon measure of significance , although guidelines have been given .

Fleiss ' kappa can be used only with binary or nominal @-@ scale ratings . No version is available for ordered @-@ categorical ratings .

= = Introduction = =

Fleiss ' kappa is a generalisation of Scott 's pi statistic , [ 1 ] a statistical measure of inter @-@ rater reliability . [ 2 ] It is also related to Cohen 's kappa statistic and Youden 's J statistic which may be more appropriate in certain instances [ 3 ] [ 4 ] . Whereas Scott 's pi and Cohen 's kappa work for only two raters , Fleiss ' kappa works for any number of raters giving categorical ratings , to a fixed number of items . It can be interpreted as expressing the extent to which the observed amount of agreement among raters exceeds what would be expected if all raters made their ratings completely randomly . It is important to note that whereas Cohen 's kappa assumes the same two raters have rated a set of items , Fleiss ' kappa specifically allows that although there are a fixed number of raters ( e.g. , three ) , different items may be rated by different individuals ( Fleiss , 1971 , p.378 ) . That is , Item 1 is rated by Raters A , B , and C ; but Item 2 could be rated by Raters D , E , and F .

Agreement can be thought of as follows , if a fixed number of people assign numerical ratings to a number of items then the kappa will give a measure for how consistent the ratings are . The kappa , <formula> , can be defined as ,

( 1 )

<formula>

The factor <formula> gives the degree of agreement that is attainable above chance , and , <formula> gives the degree of agreement actually achieved above chance . If the raters are in complete agreement then <formula> . If there is no agreement among the raters ( other than what would be expected by chance ) then <formula> .

An example of the use of Fleiss ' kappa may be the following : Consider fourteen psychiatrists are asked to look at ten patients . Each psychiatrist gives one of possibly five diagnoses to each patient . These are compiled into a matrix , and Fleiss ' kappa can be computed from this matrix ( see example below ) to show the degree of agreement between the psychiatrists above the level of agreement expected by chance .

= = Equations = =