

= Oort cloud =

The Oort cloud (/ ɔːr t / or / ɔːr t / , named after the astronomer Jan Oort) , sometimes called the Öpik - Oort cloud , is a theoretical cloud of predominantly icy planetesimals believed to surround the Sun to as far as somewhere between 50 000 and 200 000 AU (0.8 and 3.2 ly) . It is divided into two regions : a disc - shaped inner Oort cloud (or Hills cloud) and a spherical outer Oort cloud . Both regions lie beyond the heliosphere and in interstellar space . The Kuiper belt and the scattered disc , the other two reservoirs of trans - Neptunian objects , are less than one thousandth as far from the Sun as the Oort cloud .

The outer limit of the Oort cloud defines the cosmographical boundary of the Solar System and the extent of the Sun 's Hill sphere . The outer Oort cloud is only loosely bound to the Solar System , and thus is easily affected by the gravitational pull both of passing stars and of the Milky Way itself . These forces occasionally dislodge comets from their orbits within the cloud and send them toward the inner Solar System . Based on their orbits , most of the short - period comets may come from the scattered disc , but some may still have originated from the Oort cloud .

Astronomers conjecture that the matter composing the Oort cloud formed closer to the Sun and was scattered far into space by the gravitational effects of the giant planets early in the Solar System 's evolution . Although no confirmed direct observations of the Oort cloud have been made , it may be the source of all long - period and Halley - type comets entering the inner Solar System , and many of the centaurs and Jupiter - family comets as well .

= = Hypothesis = =

In 1932 , the Estonian astronomer Ernst Öpik postulated that long - period comets originated in an orbiting cloud at the outermost edge of the Solar System . The idea was independently revived by Dutch astronomer Jan Oort as a means to resolve a paradox . Over the course of the Solar System 's existence the orbits of comets are unstable and eventually dynamics dictate that a comet must either collide with the Sun or a planet or else be ejected from the Solar System by planetary perturbations . Moreover , their volatile composition means that as they repeatedly approach the Sun , radiation gradually boils the volatiles off until the comet splits or develops an insulating crust that prevents further outgassing . Thus , Oort reasoned , a comet could not have formed while in its current orbit and must have been held in an outer reservoir for almost all of its existence .

There are two main classes of comet , short - period comets (also called ecliptic comets) and long - period comets (also called nearly isotropic comets) . Ecliptic comets have relatively small orbits , below 10 AU , and follow the ecliptic plane , the same plane in which the planets lie . All long - period comets have very large orbits , on the order of thousands of AU , and appear from every direction in the sky . Oort noted that there was a peak in numbers of long - period comets with aphelia (their farthest distance from the Sun) of roughly 20 000 AU , which suggested a reservoir at that distance with a spherical , isotropic distribution . Those relatively rare comets with orbits of about 10 000 AU have probably gone through one or more orbits through the Solar System and have had their orbits drawn inward by the gravity of the planets .

= = Structure and composition = =

The Oort cloud is thought to occupy a vast space from somewhere between 2 000 and 5 000 AU (0.03 and 0.08 ly) to as far as 50 000 AU (0.79 ly) from the Sun . Some estimates place the outer edge at between 100 000 and 200 000 AU (1.58 and 3.16 ly) . The region can be subdivided into a spherical outer Oort cloud of 20 000 - 50 000 AU (0.32 - 0.79 ly) , and a torus - shaped inner Oort cloud of 2 000 - 20 000 AU (0.03 - 0.3 ly) . The outer cloud is only weakly bound to the Sun and supplies the long - period (and possibly Halley - type) comets to inside the orbit of Neptune . The inner Oort cloud is also known as the Hills cloud , named after Jack G. Hills , who proposed its existence in 1981 . Models predict that the inner cloud should have tens

or hundreds of times as many cometary nuclei as the outer halo ; it is seen as a possible source of new comets to resupply the tenuous outer cloud as the latter 's numbers are gradually depleted . The Hills cloud explains the continued existence of the Oort cloud after billions of years .

The outer Oort cloud may have trillions of objects larger than 1 km (0 @. @ 62 mi) , and billions with absolute magnitudes brighter than 11 (corresponding to approximately 20 @- @ kilometre (12 mi) diameter) , with neighboring objects tens of millions of kilometres apart . Its total mass is not known , but , assuming that Halley 's Comet is a suitable prototype for comets within the outer Oort cloud , roughly the combined mass is 3×10^{25} kilograms (6 @. @ 6×10^{25} lb) , or five times that of Earth . Earlier it was thought to be more massive (up to 380 Earth masses) , but improved knowledge of the size distribution of long @- @ period comets led to lower estimates . The mass of the inner Oort cloud has not been characterized .

If analyses of comets are representative of the whole , the vast majority of Oort @- @ cloud objects consist of ices such as water , methane , ethane , carbon monoxide and hydrogen cyanide . However , the discovery of the object 1996 PW , an object whose appearance was consistent with a D @- @ type asteroid in an orbit typical of a long @- @ period comet , prompted theoretical research that suggests that the Oort cloud population consists of roughly one to two percent asteroids . Analysis of the carbon and nitrogen isotope ratios in both the long @- @ period and Jupiter @- @ family comets shows little difference between the two , despite their presumably vastly separate regions of origin . This suggests that both originated from the original protosolar cloud , a conclusion also supported by studies of granular size in Oort @- @ cloud comets and by the recent impact study of Jupiter @- @ family comet Tempel 1 .

= = Origin = =

The Oort cloud is thought to be a remnant of the original protoplanetary disc that formed around the Sun approximately 4 @. @ 6 billion years ago . The most widely accepted hypothesis is that the Oort cloud 's objects initially coalesced much closer to the Sun as part of the same process that formed the planets and minor planets , but that gravitational interaction with young gas giants such as Jupiter ejected the objects into extremely long elliptic or parabolic orbits . Recent research has been cited by NASA hypothesizing that a large number of Oort cloud objects are the product of an exchange of materials between the Sun and its sibling stars as they formed and drifted apart , and it is suggested that many ? possibly the majority of ? Oort cloud objects did not form in close proximity to the Sun . Simulations of the evolution of the Oort cloud from the beginnings of the Solar System to the present suggest that the cloud 's mass peaked around 800 million years after formation , as the pace of accretion and collision slowed and depletion began to overtake supply .

Models by Julio Ángel Fernández suggest that the scattered disc , which is the main source for periodic comets in the Solar System , might also be the primary source for Oort cloud objects . According to the models , about half of the objects scattered travel outward toward the Oort cloud , whereas a quarter are shifted inward to Jupiter 's orbit , and a quarter are ejected on hyperbolic orbits . The scattered disc might still be supplying the Oort cloud with material . A third of the scattered disc 's population is likely to end up in the Oort cloud after 2 @. @ 5 billion years .

Computer models suggest that collisions of cometary debris during the formation period play a far greater role than was previously thought . According to these models , the number of collisions early in the Solar System 's history was so great that most comets were destroyed before they reached the Oort cloud . Therefore , the current cumulative mass of the Oort cloud is far less than was once suspected . The estimated mass of the cloud is only a small part of the 50 ? 100 Earth masses of ejected material .

Gravitational interaction with nearby stars and galactic tides modified cometary orbits to make them more circular . This explains the nearly spherical shape of the outer Oort cloud . On the other hand , the Hills cloud , which is bound more strongly to the Sun , has not acquired a spherical shape . Recent studies have shown that the formation of the Oort cloud is broadly compatible with the hypothesis that the Solar System formed as part of an embedded cluster of 200 ? 400 stars . These early stars likely played a role in the cloud 's formation , since the number of close stellar passages

within the cluster was much higher than today , leading to far more frequent perturbations .

In June 2010 Harold F. Levison and others suggested on the basis of enhanced computer simulations that the Sun " captured comets from other stars while it was in its birth cluster " . Their results imply that " a substantial fraction of the Oort cloud comets , perhaps exceeding 90 % , are from the protoplanetary discs of other stars " .

= = Comets = =

Comets are thought to have two separate points of origin in the Solar System . Short @-@ period comets (those with orbits of up to 200 years) are generally accepted to have emerged from either the Kuiper belt or the scattered disc , which are two linked flat discs of icy debris beyond Neptune 's orbit at 30 AU and jointly extending out beyond 100 AU from the Sun . Long @-@ period comets , such as comet Hale ? Bopp , whose orbits last for thousands of years , are thought to originate in the Oort cloud . The orbits within the Kuiper belt are relatively stable , and so very few comets are thought to originate there . The scattered disc , however , is dynamically active , and is far more likely to be the place of origin for comets . Comets pass from the scattered disc into the realm of the outer planets , becoming what are known as centaurs . These centaurs are then sent farther inward to become the short @-@ period comets .

There are two main varieties of short @-@ period comet : Jupiter @-@ family comets (those with semi @-@ major axes of less than 5 AU) and Halley @-@ family comets . Halley @-@ family comets , named for their prototype , Halley 's Comet , are unusual in that although they are short @-@ period comets , it is hypothesized that their ultimate origin lies in the Oort cloud , not in the scattered disc . Based on their orbits , it is suggested they were long @-@ period comets that were captured by the gravity of the giant planets and sent into the inner Solar System . This process may have also created the present orbits of a significant fraction of the Jupiter @-@ family comets , although the majority of such comets are thought to have originated in the scattered disc .

Oort noted that the number of returning comets was far less than his model predicted , and this issue , known as " cometary fading " , has yet to be resolved . No known dynamical process can explain this undercount of observed comets . Hypotheses for this discrepancy include the destruction of comets due to tidal stresses , impact or heating ; the loss of all volatiles , rendering some comets invisible , or the formation of a non @-@ volatile crust on the surface . Dynamical studies of Oort cloud comets have shown that their occurrence in the outer @-@ planet region is several times higher than in the inner @-@ planet region . This discrepancy may be due to the gravitational attraction of Jupiter , which acts as a kind of barrier , trapping incoming comets and causing them to collide with it , just as it did with Comet Shoemaker ? Levy 9 in 1994 .

= = Tidal effects = =

Most of the comets seen close to the Sun seem to have reached their current positions through gravitational perturbation of the Oort cloud by the tidal force exerted by the Milky Way . Just as the Moon 's tidal force deforms Earth 's oceans , causing the tides to rise and fall , the galactic tide also distorts the orbits of bodies in the outer Solar System . In the charted regions of the Solar System , these effects are negligible compared to the gravity of the Sun , but in the outer reaches of the system , the Sun 's gravity is weaker and the gradient of the Milky Way 's gravitational field has substantial effects . Galactic tidal forces stretch the cloud along an axis directed toward the galactic centre and compress it along the other two axes ; these small perturbations can shift orbits in the Oort cloud to bring objects close to the Sun . The point at which the Sun 's gravity concedes its influence to the galactic tide is called the tidal truncation radius . It lies at a radius of 100 @,@ 000 to 200 @,@ 000 AU , and marks the outer boundary of the Oort cloud .

Some scholars theorise that the galactic tide may have contributed to the formation of the Oort cloud by increasing the perihelia (smallest distances to the Sun) of planetesimals with large aphelia (largest distances to the Sun) . The effects of the galactic tide are quite complex , and depend heavily on the behaviour of individual objects within a planetary system . Cumulatively , however ,

the effect can be quite significant : up to 90 % of all comets originating from the Oort cloud may be the result of the galactic tide . Statistical models of the observed orbits of long @-@ period comets argue that the galactic tide is the principal means by which their orbits are perturbed toward the inner Solar System .

= = Stellar perturbations and stellar companion hypotheses = =

Besides the galactic tide , the main trigger for sending comets into the inner Solar System is thought to be interaction between the Sun 's Oort cloud and the gravitational fields of nearby stars or giant molecular clouds . The orbit of the Sun through the plane of the Milky Way sometimes brings it in relatively close proximity to other stellar systems . For example , 70 thousand years ago , Scholz 's star passed through the outer Oort cloud (although its low mass and high relative velocity limited its effect) . During the next 10 million years the known star with the greatest possibility of perturbing the Oort cloud is Gliese 710 . This process also scatters Oort cloud objects out of the ecliptic plane , potentially also explaining its spherical distribution .

In 1984 , Physicist Richard A. Muller postulated that the Sun has a heretofore undetected companion , either a brown dwarf or a red dwarf , in an elliptical orbit within the Oort cloud . This object , known as Nemesis , was hypothesized to pass through a portion of the Oort cloud approximately every 26 million years , bombarding the inner Solar System with comets . However , to date no evidence of Nemesis has been found , and many lines of evidence (such as crater counts) , have thrown its existence into doubt . Recent scientific analysis no longer supports the idea that extinctions on Earth happen at regular , repeating intervals . Thus , the Nemesis hypothesis is no longer needed .

A somewhat similar hypothesis was advanced by astronomer John J. Matese of the University of Louisiana at Lafayette in 2002 . He contends that more comets are arriving in the inner Solar System from a particular region of the Oort cloud than can be explained by the galactic tide or stellar perturbations alone , and that the most likely cause is a Jupiter @-@ mass object in a distant orbit . This hypothetical gas giant was nicknamed Tyche . The WISE mission , an all @-@ sky survey using parallax measurements in order to clarify local star distances , was capable of proving or disproving the Tyche hypothesis . In 2014 , NASA announced that the WISE survey had ruled out any object as they had defined it .

= = Modified Newtonian dynamics within the Oort cloud = =

Modified Newtonian dynamics (MOND) suggests that at their distances from the Sun , the objects comprising the Oort cloud should experience accelerations of the order of 10^{-10} m / s² , and thus should be within the realms at which deviations from Newtonian predictions come into effect . According to this hypothesis , which was proposed to account for the discrepancies in the galaxy rotation curve , which are more commonly attributed to dark matter , acceleration ceases to be linearly proportional to force at very low accelerations . If correct , this would have significant implications regarding the formation and structure of the Oort cloud . However , the majority of cosmologists do not consider MOND a valid hypothesis because it is unable to explain the movement of galactic clusters or account accurately for the cosmic microwave background .

= = Future exploration = =

Space probes have yet to reach the area of the Oort cloud . Voyager 1 , the fastest and farthest of the interplanetary space probes currently leaving the Solar System , will reach the Oort cloud in about 300 years and would take about 30 @,@ 000 years to pass through it . However , around 2025 , the radioisotope thermoelectric generators on Voyager 1 will no longer supply enough power to operate any of its scientific instruments , preventing any exploration by Voyager 1 . The other four probes currently escaping the Solar System either are already or are predicted to be non @-@ functional when they reach the Oort cloud ; however , it may be possible to find an object from the

cloud that has been knocked into the inner Solar System .

In the 1980s there was a concept for a probe to reach 1 @, @ 000 AU in 50 years called TAU ; among its missions would be to look for the Oort cloud .

In the 2014 Announcement of Opportunity for the Discovery program , an observatory to detect the objects in the Oort cloud (and Kuiper belt) called the " The Whipple Mission " was proposed . It would monitor distant stars with a photometer , looking for transits up to 10 thousand AU away . The observatory was proposed for halo orbiting around L2 with a suggested 5 @-@ year mission . It has been suggested that the Kepler observatory may also be able to detect objects in the Oort cloud .