

# Numerical Linear Algebra Vector Spaces

Franck K. Mutombo  
[franckm@aims.ac.za](mailto:franckm@aims.ac.za)

Professor of Mathematics  
[University of Lubumbashi](#)

use this email to contact me:  
[franck.mutombo@aims.ac.rw](mailto:franck.mutombo@aims.ac.rw)

October 2024

# Linear Algebra I

# Outline

Vector spaces

Euclidean spaces

Linear maps

Metric spaces

Normed spaces

Inner product space

# Vector spaces

## Vector spaces

### Note:

We focus on the **field of real** numbers ( $\mathbb{R}$ ) but most of the results can be **generalized** to the **field of complex** numbers ( $\mathbb{C}$ ) in a straightforward fashion.

## Vector spaces

A vector space or *linear space* (over the field  $\mathbb{R}$ ) consists of

- (a) a **set** of vectors  $\mathcal{V}$
- (b) an **addition** operation:  $\mathcal{V} \times \mathcal{V} \rightarrow \mathcal{V}$
- (c) a **scalar multiplication** operation:  $\mathbb{R} \times \mathcal{V} \rightarrow \mathcal{V}$
- (d) a **distinguished** element  $\mathbf{0} \in \mathcal{V}$

## Vector spaces

A vector space or *linear space* (over the field  $\mathbb{R}$ ) consists of

- (a) a **set** of vectors  $\mathcal{V}$
- (b) an **addition** operation:  $\mathcal{V} \times \mathcal{V} \rightarrow \mathcal{V}$
- (c) a **scalar multiplication** operation:  $\mathbb{R} \times \mathcal{V} \rightarrow \mathcal{V}$
- (d) a **distinguished** element  $\mathbf{0} \in \mathcal{V}$

and satisfies the following properties:

1.  $\mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x}$ ,  $\forall \mathbf{x}, \mathbf{y} \in \mathcal{V}$  (**commutative under addition**)
2.  $(\mathbf{x} + \mathbf{y}) + \mathbf{z} = \mathbf{x} + (\mathbf{y} + \mathbf{z})$ ,  $\forall \mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathcal{V}$  (**associative under addition**)
3.  $\mathbf{0} + \mathbf{x} = \mathbf{x}$ ,  $\forall \mathbf{x} \in \mathcal{V}$  (**0 being additive identity**)
4.  $\forall \mathbf{x} \in \mathcal{V} \exists (-\mathbf{x})$  such that  $\mathbf{x} + (-\mathbf{x}) = \mathbf{0}$  ( **$-\mathbf{x}$  being additive inverse**)
5.  $(\alpha\beta)\mathbf{x} = \alpha(\beta\mathbf{x})$ ,  $\forall \alpha, \beta \in \mathbb{R} \quad \forall \mathbf{x} \in \mathcal{V}$  (**associative under scalar mult.**)
6.  $\alpha(\mathbf{x} + \mathbf{y}) = \alpha\mathbf{x} + \alpha\mathbf{y}$ ,  $\forall \alpha \in \mathbb{R} \quad \forall \mathbf{x}, \mathbf{y} \in \mathcal{V}$  (**distributive**)
7.  $1\mathbf{x} = \mathbf{x}$ ,  $\forall \mathbf{x} \in \mathcal{V}$  (**1 being multiplicative identity**)

## Vector spaces contd.

### Example (Vector space)

- ▶  $\mathcal{V}_1 = \{\mathbf{0}\}$  for  $\mathbf{0} \in \mathbb{R}^n$
- ▶  $\mathcal{V}_2 = \mathbb{R}^n$
- ▶  $\mathcal{V}_3 = \sum_{i=1}^k \alpha_i \mathbf{x}_i$  for  $\alpha_i \in \mathbb{R}$ ,  $k < n$ , and  $\mathbf{x}_i \in \mathbb{R}^n$



## Vector spaces contd.

### Example (Vector space)

- ▶  $\mathcal{V}_1 = \{\mathbf{0}\}$  for  $\mathbf{0} \in \mathbb{R}^n$
- ▶  $\mathcal{V}_2 = \mathbb{R}^n$
- ▶  $\mathcal{V}_3 = \sum_{i=1}^k \alpha_i \mathbf{x}_i$  for  $\alpha_i \in \mathbb{R}$ ,  $k < n$ , and  $\mathbf{x}_i \in \mathbb{R}^n$

It is straight forward to show that  $\mathcal{V}_1$ ,  $\mathcal{V}_2$ , and  $\mathcal{V}_3$  satisfy properties 1–7 above. However, using a [subspace](#) argument is much simpler.

## Vector spaces contd.

### Example (Vector space)

- ▶  $\mathcal{V}_1 = \{\mathbf{0}\}$  for  $\mathbf{0} \in \mathbb{R}^n$
- ▶  $\mathcal{V}_2 = \mathbb{R}^n$
- ▶  $\mathcal{V}_3 = \sum_{i=1}^k \alpha_i \mathbf{x}_i$  for  $\alpha_i \in \mathbb{R}$ ,  $k < n$ , and  $\mathbf{x}_i \in \mathbb{R}^n$

It is straight forward to show that  $\mathcal{V}_1$ ,  $\mathcal{V}_2$ , and  $\mathcal{V}_3$  satisfy properties 1–7 above. However, using a **subspace** argument is much simpler.

### Definition (Subspace)

A **subspace** is a vector space that is a *subset* of another vector space.

## Vector spaces contd.

### Example (Vector space)

- ▶  $\mathcal{V}_1 = \{\mathbf{0}\}$  for  $\mathbf{0} \in \mathbb{R}^n$
- ▶  $\mathcal{V}_2 = \mathbb{R}^n$
- ▶  $\mathcal{V}_3 = \sum_{i=1}^k \alpha_i \mathbf{x}_i$  for  $\alpha_i \in \mathbb{R}$ ,  $k < n$ , and  $\mathbf{x}_i \in \mathbb{R}^n$

It is straight forward to show that  $\mathcal{V}_1$ ,  $\mathcal{V}_2$ , and  $\mathcal{V}_3$  satisfy properties 1–7 above. However, using a **subspace** argument is much simpler.

### Definition (Subspace)

A **subspace** is a vector space that is a *subset* of another vector space.

### Example (Subspace)

$\mathcal{V}_3$  (and actually  $\mathcal{V}_1$  as well as  $\mathcal{V}_2$ ) in the example above is subspace of  $\mathbb{R}^n$ .

## Vector spaces contd.

### Definition (Linear independence)

A set of vectors,  $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n\}$ , is **linearly independent** if

$$\alpha_1 \mathbf{x}_1 + \alpha_2 \mathbf{x}_2 + \dots + \alpha_n \mathbf{x}_n = \mathbf{0} \Rightarrow \alpha_1 = \alpha_2 = \dots = \alpha_n = 0.$$

## Vector spaces contd.

### Definition (Linear independence)

A set of vectors,  $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n\}$ , is **linearly independent** if

$$\alpha_1 \mathbf{x}_1 + \alpha_2 \mathbf{x}_2 + \dots + \alpha_n \mathbf{x}_n = \mathbf{0} \Rightarrow \alpha_1 = \alpha_2 = \dots = \alpha_n = 0.$$

### Definition (Span)

The **span** of  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in \mathcal{V}$  is the set of all vectors that are linear combination of them, i.e.

$$\text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_n\} = \{\mathbf{v} \in \mathcal{V} : \exists \alpha_1, \dots, \alpha_n, \text{ such that } \mathbf{v} = \alpha_1 \mathbf{v}_1, \dots, \alpha_n \mathbf{v}_n\}$$

## Vector spaces contd.

### Definition (Linear independence)

A set of vectors,  $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n\}$ , is **linearly independent** if

$$\alpha_1 \mathbf{x}_1 + \alpha_2 \mathbf{x}_2 + \dots + \alpha_n \mathbf{x}_n = \mathbf{0} \Rightarrow \alpha_1 = \alpha_2 = \dots = \alpha_n = 0.$$

### Definition (Span)

The **span** of  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in \mathcal{V}$  is the set of all vectors that are linear combination of them, i.e.

$$\text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_n\} = \{\mathbf{v} \in \mathcal{V} : \exists \alpha_1, \dots, \alpha_n, \text{ such that } \mathbf{v} = \alpha_1 \mathbf{v}_1, \dots, \alpha_n \mathbf{v}_n\}$$

### Definition (Basis)

A set of vectors that is **linearly independent** and its **spans** is the whole of  $\mathcal{V}$ , is said to be a **basis** of  $\mathcal{V}$ .

## Vector spaces contd.

### Definition (Dimension)

The **dimension** of a vector space,  $\mathcal{V}$ , (denoted  $\dim(\mathcal{V})$ ) is the number of vectors in the basis of  $\mathcal{V}$ .

If dimension of a vector space is finite, it is said to be **finite-dimensional**. Otherwise it is **infinite-dimensional**.

## Vector spaces contd.

### Example

Let  $\mathbb{R}_n[x]$  denote the vector space of polynomial of degree at most  $n$  over the field of real numbers.

1. Show that it is indeed a vector space!
2. Since every degree  $n$  polynomial can be written as  $a_0x^0 + a_1x^1 + \cdots + a_nx^n$ , what is the dimension of  $\mathbb{R}_n[x]$ ?
3. Show that the family  $\{x^i\}_{i=0}^n$  is linearly independent.
4. Hence is the family  $\{x^i\}_{i=0}^n$  a basis?



## Euclidean spaces

## Euclidean spaces

- ▶ The Euclidean spaces is when  $\mathcal{V} = \mathbb{R}^n$  (most popular vector space)

## Euclidean spaces

- ▶ The Euclidean spaces is when  $\mathcal{V} = \mathbb{R}^n$  (most popular vector space)
- ▶ The vectors in this space consist of  $n$ -tuples of real numbers, i.e.

$$\mathbf{x} = (x_1, x_2, \dots, x_n)$$

## Euclidean spaces

- ▶ The Euclidean spaces is when  $\mathcal{V} = \mathbb{R}^n$  (most popular vector space)
- ▶ The vectors in this space consist of  $n$ -tuples of real numbers, i.e.

$$\mathbf{x} = (x_1, x_2, \dots, x_n)$$

- ▶ For this course and typically in Data Science, vectors will be column vectors, i.e.

$$\mathbf{x} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

## Euclidean spaces

- ▶ The **Euclidean spaces** is when  $\mathcal{V} = \mathbb{R}^n$  (most popular vector space)
- ▶ The vectors in this space consist of  **$n$ -tuples** of real numbers, i.e.

$$\mathbf{x} = (x_1, x_2, \dots, x_n)$$

- ▶ For this course and typically in Data Science, vectors will be **column vectors**, i.e.

$$\mathbf{x} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

- ▶ Addition and scalar multiplication are defined **component-wise** on vectors in  $\mathbb{R}^n$ , i.e.

$$\mathbf{x} + \mathbf{y} = \begin{bmatrix} x_1 + y_1 \\ \vdots \\ x_n + y_n \end{bmatrix} \quad \text{and} \quad \alpha \mathbf{x} = \begin{bmatrix} \alpha x_1 \\ \vdots \\ \alpha x_n \end{bmatrix}$$

## Euclidean spaces

- ▶ The **Euclidean spaces** is when  $\mathcal{V} = \mathbb{R}^n$  (most popular vector space)
- ▶ The vectors in this space consist of  **$n$ -tuples** of real numbers, i.e.

$$\mathbf{x} = (x_1, x_2, \dots, x_n)$$

- ▶ For this course and typically in Data Science, vectors will be **column vectors**, i.e.

$$\mathbf{x} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

- ▶ Addition and scalar multiplication are defined **component-wise** on vectors in  $\mathbb{R}^n$ , i.e.

$$\mathbf{x} + \mathbf{y} = \begin{bmatrix} x_1 + y_1 \\ \vdots \\ x_n + y_n \end{bmatrix} \quad \text{and} \quad \alpha \mathbf{x} = \begin{bmatrix} \alpha x_1 \\ \vdots \\ \alpha x_n \end{bmatrix}$$

- ▶ Euclidean space is used to mathematically represent **physical space**, with notions such as **distance**, **length**, and **angles**.

# Linear maps

## Linear maps

### Definition (Linear map)

A **linear map** is a function  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{U}$ , where  $\mathcal{V}$  and  $\mathcal{U}$  are vector spaces, that satisfies

1.  $\mathcal{T}(\mathbf{x} + \mathbf{y}) = \mathcal{T}(\mathbf{x}) + \mathcal{T}(\mathbf{y})$  for all  $\mathbf{x}, \mathbf{y} \in \mathcal{V}$
2.  $\mathcal{T}(\alpha \mathbf{x}) = \alpha \mathcal{T}(\mathbf{x})$  for all  $\mathbf{x} \in \mathcal{V}$ ,  $\alpha \in \mathbb{R}$



## Linear maps

### Definition (Linear map)

A **linear map** is a function  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{U}$ , where  $\mathcal{V}$  and  $\mathcal{U}$  are vector spaces, that satisfies

1.  $\mathcal{T}(\mathbf{x} + \mathbf{y}) = \mathcal{T}(\mathbf{x}) + \mathcal{T}(\mathbf{y})$  for all  $\mathbf{x}, \mathbf{y} \in \mathcal{V}$
2.  $\mathcal{T}(\alpha \mathbf{x}) = \alpha \mathcal{T}(\mathbf{x})$  for all  $\mathbf{x} \in \mathcal{V}$ ,  $\alpha \in \mathbb{R}$

- ▶ A **composition** of linear maps,  $\mathcal{T}$  and  $\mathcal{S}$  will be denoted as  $\mathcal{T} \circ \mathcal{S}$
- ▶ A linear map from  $\mathcal{V}$  to itself is called a **linear operator**

## Linear maps

### Definition (Linear map)

A **linear map** is a function  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{U}$ , where  $\mathcal{V}$  and  $\mathcal{U}$  are vector spaces, that satisfies

1.  $\mathcal{T}(\mathbf{x} + \mathbf{y}) = \mathcal{T}(\mathbf{x}) + \mathcal{T}(\mathbf{y})$  for all  $\mathbf{x}, \mathbf{y} \in \mathcal{V}$
2.  $\mathcal{T}(\alpha \mathbf{x}) = \alpha \mathcal{T}(\mathbf{x})$  for all  $\mathbf{x} \in \mathcal{V}$ ,  $\alpha \in \mathbb{R}$

- ▶ A **composition** of linear maps,  $\mathcal{T}$  and  $\mathcal{S}$  will be denoted as  $\mathcal{T} \circ \mathcal{S}$
- ▶ A linear map from  $\mathcal{V}$  to itself is called a **linear operator**
- ▶ A linear map **preserves** addition and scalar multiplication, hence reflect the structure of vector spaces  $\Rightarrow$  a **homomorphism** of vector spaces
- ▶ An **invertible** homomorphism (where the inverse is also a homomorphism) is called an **isomorphism**

## Linear maps

### Definition (Linear map)

A **linear map** is a function  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{U}$ , where  $\mathcal{V}$  and  $\mathcal{U}$  are vector spaces, that satisfies

1.  $\mathcal{T}(\mathbf{x} + \mathbf{y}) = \mathcal{T}(\mathbf{x}) + \mathcal{T}(\mathbf{y})$  for all  $\mathbf{x}, \mathbf{y} \in \mathcal{V}$
2.  $\mathcal{T}(\alpha \mathbf{x}) = \alpha \mathcal{T}(\mathbf{x})$  for all  $\mathbf{x} \in \mathcal{V}$ ,  $\alpha \in \mathbb{R}$

- ▶ A **composition** of linear maps,  $\mathcal{T}$  and  $\mathcal{S}$  will be denoted as  $\mathcal{T} \circ \mathcal{S}$
- ▶ A linear map from  $\mathcal{V}$  to itself is called a **linear operator**
- ▶ A linear map **preserves** addition and scalar multiplication, hence reflect the structure of vector spaces  $\Rightarrow$  a **homomorphism** of vector spaces
- ▶ An **invertible** homomorphism (where the inverse is also a homomorphism) is called an **isomorphism**
- ▶  $\mathcal{V}$  and  $\mathcal{W}$  are **isomorphic**, when there is an **isomorphism** from  $\mathcal{V}$  to  $\mathcal{W}$
- ▶ Every real  **$n$ -dimensional** vector space is isomorphic to  $\mathbb{R}^n$

## The matrix of a linear map

- ▶ In the finite-dimensional setting, **matrices** are used to represent and manipulate **vectors** and **linear maps** on a computer

## The matrix of a linear map

- In the finite-dimensional setting, **matrices** are used to represent and manipulate **vectors** and **linear maps** on a computer

### Definition (Matrix of a linear map)

Suppose  $\mathbf{v}_1, \dots, \mathbf{v}_n$  and  $\mathbf{u}_1, \dots, \mathbf{u}_m$  are **bases** of  $\mathcal{V}$  and  $\mathcal{U}$  respectively, and  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{U}$  is a linear map. Then the **matrix** of  $\mathcal{T}$ , with entries  $A_{ij}$  for  $i = 1, \dots, m$  and  $j = 1, \dots, n$  is defined by

$$\mathcal{T}(\mathbf{v}_j) = A_{1j}\mathbf{u}_1 + A_{2j}\mathbf{u}_2 + \cdots + A_{mj}\mathbf{u}_m,$$

i.e. the  $j$ th column of  $\mathbf{A}$  is the coordinates of  $\mathcal{T}(\mathbf{v}_j)$  in the chosen basis for  $\mathcal{U}$ .

## The matrix of a linear map

- In the finite-dimensional setting, **matrices** are used to represent and manipulate **vectors** and **linear maps** on a computer

### Definition (Matrix of a linear map)

Suppose  $\mathbf{v}_1, \dots, \mathbf{v}_n$  and  $\mathbf{u}_1, \dots, \mathbf{u}_m$  are **bases** of  $\mathcal{V}$  and  $\mathcal{U}$  respectively, and  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{U}$  is a linear map. Then the **matrix** of  $\mathcal{T}$ , with entries  $A_{ij}$  for  $i = 1, \dots, m$  and  $j = 1, \dots, n$  is defined by

$$\mathcal{T}(\mathbf{v}_j) = A_{1j}\mathbf{u}_1 + A_{2j}\mathbf{u}_2 + \dots + A_{mj}\mathbf{u}_m,$$

i.e. the  $j$ th column of  $\mathbf{A}$  is the coordinates of  $\mathcal{T}(\mathbf{v}_j)$  in the chosen basis for  $\mathcal{U}$ .

- Conversely, every matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$  **induces** a linear map  $\mathcal{T} : \mathbb{R}^n \rightarrow \mathbb{R}^m$  given by

$$\mathcal{T}(\mathbf{x}) = \mathbf{A}\mathbf{x}$$

## The matrix of a linear map

- In the finite-dimensional setting, **matrices** are used to represent and manipulate **vectors** and **linear maps** on a computer

### Definition (Matrix of a linear map)

Suppose  $\mathbf{v}_1, \dots, \mathbf{v}_n$  and  $\mathbf{u}_1, \dots, \mathbf{u}_m$  are **bases** of  $\mathcal{V}$  and  $\mathcal{U}$  respectively, and  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{U}$  is a linear map. Then the **matrix** of  $\mathcal{T}$ , with entries  $A_{ij}$  for  $i = 1, \dots, m$  and  $j = 1, \dots, n$  is defined by

$$\mathcal{T}(\mathbf{v}_j) = A_{1j}\mathbf{u}_1 + A_{2j}\mathbf{u}_2 + \cdots + A_{mj}\mathbf{u}_m,$$

i.e. the  $j$ th column of  $\mathbf{A}$  is the coordinates of  $\mathcal{T}(\mathbf{v}_j)$  in the chosen basis for  $\mathcal{U}$ .

- Conversely, every matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$  **induces** a linear map  $\mathcal{T} : \mathbb{R}^n \rightarrow \mathbb{R}^m$  given by

$$\mathcal{T}(\mathbf{x}) = \mathbf{A}\mathbf{x}$$

- And the matrix of this map with respect to the **standard bases** of  $\mathbb{R}^n$  and  $\mathbb{R}^m$  is simply  $\mathbf{A}$

## The matrix of a linear map

- ▶ If  $\mathbf{A} \in \mathbb{R}^{m \times n}$ , its **transpose**,  $\mathbf{A}^T \in \mathbb{R}^{n \times m}$  is  $(\mathbf{A}^T)_{ij} = A_{ji}$  for each  $(i, j)$
- ▶  $\Rightarrow$  the **columns** of  $\mathbf{A}$  become the **rows** of  $\mathbf{A}^T$ , and the **rows** of  $\mathbf{A}$  become the **columns** of  $\mathbf{A}^T$



## The matrix of a linear map

- ▶ If  $\mathbf{A} \in \mathbb{R}^{m \times n}$ , its **transpose**,  $\mathbf{A}^T \in \mathbb{R}^{n \times m}$  is  $(\mathbf{A}^T)_{ij} = A_{ji}$  for each  $(i, j)$
- ▶  $\Rightarrow$  the **columns** of  $\mathbf{A}$  become the **rows** of  $\mathbf{A}^T$ , and the **rows** of  $\mathbf{A}$  become the **columns** of  $\mathbf{A}^T$
- ▶ Some nice **algebraic** properties of the **transpose**:
  1.  $(\mathbf{A}^T)^T = \mathbf{A}$
  2.  $(\mathbf{A} + \mathbf{B})^T = \mathbf{A}^T + \mathbf{B}^T$
  3.  $(\alpha \mathbf{A})^T = \alpha \mathbf{A}^T$
  4.  $(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$

## Nullspace & range space

- ▶ Some of the most important **subspaces** are those induced by **linear maps**

## Nullspace & range space

- Some of the most important **subspaces** are those induced by **linear maps**

### Definition (Nullspace and range)

If  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{U}$  is a linear map, we define the **nullspace** of  $\mathcal{T}$  as

$$\text{null}(\mathcal{T}) = \{\mathbf{v} \in \mathcal{V} \mid \mathcal{T}(\mathbf{v}) = \mathbf{0}\},$$

and the **range** of  $\mathcal{T}$  as

$$\text{range}(\mathcal{T}) = \{\mathbf{u} \in \mathcal{U} \mid \exists \mathbf{v} \in \mathcal{V} \text{ such that } \mathcal{T}(\mathbf{v}) = \mathbf{u}\}$$

## Nullspace & range space

- Some of the most important **subspaces** are those induced by **linear maps**

### Definition (Nullspace and range)

If  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{U}$  is a linear map, we define the **nullspace** of  $\mathcal{T}$  as

$$\text{null}(\mathcal{T}) = \{\mathbf{v} \in \mathcal{V} \mid \mathcal{T}(\mathbf{v}) = \mathbf{0}\},$$

and the **range** of  $\mathcal{T}$  as

$$\text{range}(\mathcal{T}) = \{\mathbf{u} \in \mathcal{U} \mid \exists \mathbf{v} \in \mathcal{V} \text{ such that } \mathcal{T}(\mathbf{v}) = \mathbf{u}\}$$

- The **nullspace** and **range** of a linear map are always **subspaces** of its **domain** and **codomain**, respectively

## Nullspace & range space

- ▶ Some of the most important **subspaces** are those induced by **linear maps**

### Definition (Nullspace and range)

If  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{U}$  is a linear map, we define the **nullspace** of  $\mathcal{T}$  as

$$\text{null}(\mathcal{T}) = \{\mathbf{v} \in \mathcal{V} \mid \mathcal{T}(\mathbf{v}) = \mathbf{0}\},$$

and the **range** of  $\mathcal{T}$  as

$$\text{range}(\mathcal{T}) = \{\mathbf{u} \in \mathcal{U} \mid \exists \mathbf{v} \in \mathcal{V} \text{ such that } \mathcal{T}(\mathbf{v}) = \mathbf{u}\}$$

- ▶ The **nullspace** and **range** of a linear map are always **subspaces** of its **domain** and **codomain**, respectively
- ▶ The **columnspace** of a matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$  is the **span of its columns** (considered as vectors in  $\mathbb{R}^m$ ), denoted  $\text{range}(\mathbf{A})$
- ▶ The **rowspace** of a matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$  is the **span of its rows** (considered as vectors in  $\mathbb{R}^n$ ), denoted  $\text{range}(\mathbf{A}^T)$

## Nullspace & range space

- Some of the most important **subspaces** are those induced by **linear maps**

### Definition (Nullspace and range)

If  $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{U}$  is a linear map, we define the **nullspace** of  $\mathcal{T}$  as

$$\text{null}(\mathcal{T}) = \{\mathbf{v} \in \mathcal{V} \mid \mathcal{T}(\mathbf{v}) = \mathbf{0}\},$$

and the **range** of  $\mathcal{T}$  as

$$\text{range}(\mathcal{T}) = \{\mathbf{u} \in \mathcal{U} \mid \exists \mathbf{v} \in \mathcal{V} \text{ such that } \mathcal{T}(\mathbf{v}) = \mathbf{u}\}$$

- The **nullspace** and **range** of a linear map are always **subspaces** of its **domain** and **codomain**, respectively
- The **columnspace** of a matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$  is the **span of its columns** (considered as vectors in  $\mathbb{R}^m$ ), denoted  $\text{range}(\mathbf{A})$
- The **rowspace** of a matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$  is the **span of its rows** (considered as vectors in  $\mathbb{R}^n$ ), denoted  $\text{range}(\mathbf{A}^T)$
- The **rank** satisfies:  $\text{rank}(\mathbf{A}) = \dim(\text{range}(\mathbf{A})) = \dim(\text{range}(\mathbf{A}^T))$

# Metric spaces

## Metric spaces

- **Metrics** generalize the notion of **distance** from Euclidean space (although metric spaces need not be vector spaces)



## Metric spaces

- **Metrics** generalize the notion of **distance** from Euclidean space (although metric spaces need not be vector spaces)

### Definition (Metric)

A metric on a set  $\mathcal{X}$  is a function  $d(\cdot, \cdot) : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  that  $\forall \mathbf{x}, \mathbf{y} \in \mathcal{X}$  satisfies

- (a)  $d(\mathbf{x}, \mathbf{y}) \geq 0$  for all  $\mathbf{x}$  and  $\mathbf{y}$  (*nonnegativity*)
- (b)  $d(\mathbf{x}, \mathbf{y}) = 0$  if and only if  $\mathbf{x} = \mathbf{y}$  (*definiteness*)
- (c)  $d(\mathbf{x}, \mathbf{y}) = d(\mathbf{y}, \mathbf{x})$  (*symmetry*)
- (d)  $d(\mathbf{x}, \mathbf{y}) \leq d(\mathbf{x}, \mathbf{z}) + d(\mathbf{z}, \mathbf{y})$  (*triangle inequality*)

## Metric spaces

- **Metrics** generalize the notion of **distance** from Euclidean space (although metric spaces need not be vector spaces)

### Definition (Metric)

A metric on a set  $\mathcal{X}$  is a function  $d(\cdot, \cdot) : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  that  $\forall \mathbf{x}, \mathbf{y} \in \mathcal{X}$  satisfies

- (a)  $d(\mathbf{x}, \mathbf{y}) \geq 0$  for all  $\mathbf{x}$  and  $\mathbf{y}$  (*nonnegativity*)
- (b)  $d(\mathbf{x}, \mathbf{y}) = 0$  if and only if  $\mathbf{x} = \mathbf{y}$  (*definiteness*)
- (c)  $d(\mathbf{x}, \mathbf{y}) = d(\mathbf{y}, \mathbf{x})$  (*symmetry*)
- (d)  $d(\mathbf{x}, \mathbf{y}) \leq d(\mathbf{x}, \mathbf{z}) + d(\mathbf{z}, \mathbf{y})$  (*triangle inequality*)

- A **pseudo-metric** satisfies (a), (c) and (d) but not necessarily (b)

## Metric spaces

- **Metrics** generalize the notion of **distance** from Euclidean space (although metric spaces need not be vector spaces)

### Definition (Metric)

A metric on a set  $\mathcal{X}$  is a function  $d(\cdot, \cdot) : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  that  $\forall \mathbf{x}, \mathbf{y} \in \mathcal{X}$  satisfies

- (a)  $d(\mathbf{x}, \mathbf{y}) \geq 0$  for all  $\mathbf{x}$  and  $\mathbf{y}$  (*nonnegativity*)
- (b)  $d(\mathbf{x}, \mathbf{y}) = 0$  if and only if  $\mathbf{x} = \mathbf{y}$  (*definiteness*)
- (c)  $d(\mathbf{x}, \mathbf{y}) = d(\mathbf{y}, \mathbf{x})$  (*symmetry*)
- (d)  $d(\mathbf{x}, \mathbf{y}) \leq d(\mathbf{x}, \mathbf{z}) + d(\mathbf{z}, \mathbf{y})$  (*triangle inequality*)

- Metrics allow **limits** to be defined for mathematical objects other than reals

## Metric spaces

- **Metrics** generalize the notion of **distance** from Euclidean space (although metric spaces need not be vector spaces)

### Definition (Metric)

A metric on a set  $\mathcal{X}$  is a function  $d(\cdot, \cdot) : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  that  $\forall \mathbf{x}, \mathbf{y} \in \mathcal{X}$  satisfies

- (a)  $d(\mathbf{x}, \mathbf{y}) \geq 0$  for all  $\mathbf{x}$  and  $\mathbf{y}$  (*nonnegativity*)
- (b)  $d(\mathbf{x}, \mathbf{y}) = 0$  if and only if  $\mathbf{x} = \mathbf{y}$  (*definiteness*)
- (c)  $d(\mathbf{x}, \mathbf{y}) = d(\mathbf{y}, \mathbf{x})$  (*symmetry*)
- (d)  $d(\mathbf{x}, \mathbf{y}) \leq d(\mathbf{x}, \mathbf{z}) + d(\mathbf{z}, \mathbf{y})$  (*triangle inequality*)

- Metrics allow **limits** to be defined for mathematical objects other than reals

### Example

1.  $\{x_n\} \subseteq \mathcal{S}$  converges to  $x$  if for any  $\epsilon > 0$ ,  $\exists N \in \mathbb{N} \mid d(x_n, x) < \epsilon \forall n \geq N$
2. Euclidean distance:  $d_E(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\|_2$
3. Loss/cost functions: (a) MSE, (b) MAE, (c) Cross Entropy, etc

## Normed spaces

## Normed spaces

- ▶ Norms generalize the notion of length from Euclidean space

## Normed spaces

- Norms generalize the notion of length from Euclidean space

### Definition (Vector norm)

A norm on a real vector space  $\mathcal{V}$  is a function  $\|\cdot\| : \mathcal{V} \rightarrow \mathbb{R}$  that for all  $\mathbf{x}, \mathbf{y} \in \mathcal{V}$  and  $\lambda \in \mathbb{R}$  satisfies

- (a)  $\|\mathbf{x}\| \geq 0$  *(nonnegativity)*
- (b)  $\|\mathbf{x}\| = 0$  if and only if  $\mathbf{x} = \mathbf{0}$  *(definitiveness)*
- (c)  $\|\lambda\mathbf{x}\| = |\lambda|\|\mathbf{x}\|$  *(homogeneity)*
- (d)  $\|\mathbf{x} + \mathbf{y}\| \leq \|\mathbf{x}\| + \|\mathbf{y}\|$  *(triangle inequality)*

## Normed spaces

- **Norms** generalize the notion of **length** from Euclidean space

### Definition (Vector norm)

A norm on a real vector space  $\mathcal{V}$  is a function  $\| \cdot \| : \mathcal{V} \rightarrow \mathbb{R}$  that for all  $\mathbf{x}, \mathbf{y} \in \mathcal{V}$  and  $\lambda \in \mathbb{R}$  satisfies

- (a)  $\|\mathbf{x}\| \geq 0$  *(nonnegativity)*
- (b)  $\|\mathbf{x}\| = 0$  if and only if  $\mathbf{x} = \mathbf{0}$  *(definitiveness)*
- (c)  $\|\lambda\mathbf{x}\| = |\lambda|\|\mathbf{x}\|$  *(homogeneity)*
- (d)  $\|\mathbf{x} + \mathbf{y}\| \leq \|\mathbf{x}\| + \|\mathbf{y}\|$  *(triangle inequality)*

- A vector space endowed with a **norm** is called a **normed (vector) space**



## Normed spaces

- **Norms** generalize the notion of **length** from Euclidean space

### Definition (Vector norm)

A norm on a real vector space  $\mathcal{V}$  is a function  $\| \cdot \| : \mathcal{V} \rightarrow \mathbb{R}$  that for all  $\mathbf{x}, \mathbf{y} \in \mathcal{V}$  and  $\lambda \in \mathbb{R}$  satisfies

- (a)  $\|\mathbf{x}\| \geq 0$  *(nonnegativity)*
- (b)  $\|\mathbf{x}\| = 0$  if and only if  $\mathbf{x} = \mathbf{0}$  *(definitiveness)*
- (c)  $\|\lambda\mathbf{x}\| = |\lambda|\|\mathbf{x}\|$  *(homogeneity)*
- (d)  $\|\mathbf{x} + \mathbf{y}\| \leq \|\mathbf{x}\| + \|\mathbf{y}\|$  *(triangle inequality)*

- A vector space endowed with a **norm** is called a **normed (vector) space**
- Note that any norm on  $\mathcal{V}$  induces a **distance metric** on  $\mathcal{V}$ :

$$d(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\|$$

## Normed spaces

- **Norms** generalize the notion of **length** from Euclidean space

### Definition (Vector norm)

A norm on a real vector space  $\mathcal{V}$  is a function  $\|\cdot\| : \mathcal{V} \rightarrow \mathbb{R}$  that for all  $\mathbf{x}, \mathbf{y} \in \mathcal{V}$  and  $\lambda \in \mathbb{R}$  satisfies

- (a)  $\|\mathbf{x}\| \geq 0$  *(nonnegativity)*
- (b)  $\|\mathbf{x}\| = 0$  if and only if  $\mathbf{x} = \mathbf{0}$  *(definitiveness)*
- (c)  $\|\lambda\mathbf{x}\| = |\lambda|\|\mathbf{x}\|$  *(homogeneity)*
- (d)  $\|\mathbf{x} + \mathbf{y}\| \leq \|\mathbf{x}\| + \|\mathbf{y}\|$  *(triangle inequality)*

- A vector space endowed with a **norm** is called a **normed (vector) space**
- Note that any norm on  $\mathcal{V}$  induces a **distance metric** on  $\mathcal{V}$ :
 
$$d(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\|$$
- This satisfies axioms for metrics, hence any **normed space is also a metric space** (since axioms for norms  $\Rightarrow$  axioms for metrics)

## Vector Norms

- ▶ This course will focus on a few specific norms on  $\mathbb{R}^n$ , i.e. *p-norms*

## Vector Norms

- ▶ This course will focus on a few specific norms on  $\mathbb{R}^n$ , i.e. *p-norms*
- ▶ For  $\mathbf{x} \in \mathbb{R}^n$ , the *p-norm* is defined as  $\|\mathbf{x}\|_p := \left(\sum_{i=1}^n |x_i|^p\right)^{1/p}$

## Vector Norms

- ▶ This course will focus on a few specific norms on  $\mathbb{R}^n$ , i.e. ***p*-norms**
- ▶ For  $\mathbf{x} \in \mathbb{R}^n$ , the ***p*-norm** is defined as  $\|\mathbf{x}\|_p := \left(\sum_{i=1}^n |x_i|^p\right)^{1/p}$

### Example

- |     |                      |   |                  |
|-----|----------------------|---|------------------|
| (1) | $\ell_2$ -norm:      | $\ \mathbf{x}\ _2 := \sqrt{\sum_{i=1}^p x_i^2}$     | (Euclidean norm) |
| (2) | $\ell_1$ -norm:      | $\ \mathbf{x}\ _1 := \sum_{i=1}^p  x_i $            | (Manhattan norm) |
| (3) | $\ell_\infty$ -norm: | $\ \mathbf{x}\ _\infty := \max_{i=1,\dots,p}  x_i $ | (Chebyshev norm) |

## Vector Norms

- ▶ This course will focus on a few specific norms on  $\mathbb{R}^n$ , i.e. *p-norms*
- ▶ For  $\mathbf{x} \in \mathbb{R}^n$ , the *p-norm* is defined as  $\|\mathbf{x}\|_p := \left( \sum_{i=1}^n |x_i|^p \right)^{1/p}$

### Example

- |     |                      |   |                  |
|-----|----------------------|---|------------------|
| (1) | $\ell_2$ -norm:      | $\ \mathbf{x}\ _2 := \sqrt{\sum_{i=1}^p x_i^2}$       | (Euclidean norm) |
| (2) | $\ell_1$ -norm:      | $\ \mathbf{x}\ _1 := \sum_{i=1}^p  x_i $              | (Manhattan norm) |
| (3) | $\ell_\infty$ -norm: | $\ \mathbf{x}\ _\infty := \max_{i=1, \dots, p}  x_i $ | (Chebyshev norm) |

- ▶  $p \geq 1$  is required for the general definition of the *p-norm* because the *triangle inequality* fails to hold if  $p < 1$

## Vector Norms

- ▶ This course will focus on a few specific norms on  $\mathbb{R}^n$ , i.e. ***p*-norms**
- ▶ For  $\mathbf{x} \in \mathbb{R}^n$ , the ***p*-norm** is defined as  $\|\mathbf{x}\|_p := \left(\sum_{i=1}^n |x_i|^p\right)^{1/p}$

### Example

- (1)  $\ell_2$ -norm:  $\|\mathbf{x}\|_2 := \sqrt{\sum_{i=1}^n x_i^2}$  (Euclidean norm)
- (2)  $\ell_1$ -norm:  $\|\mathbf{x}\|_1 := \sum_{i=1}^n |x_i|$  (Manhattan norm)
- (3)  $\ell_\infty$ -norm:  $\|\mathbf{x}\|_\infty := \max_{i=1,\dots,p} |x_i|$  (Chebyshev norm)

- ▶  $p \geq 1$  is required for the general definition of the *p*-norm because the **triangle inequality** **fails** to hold if  $p < 1$
- ▶ In a finite-dimensional vector space  $\mathcal{V}$ , all norms on  $\mathcal{V}$  are **equivalent**, i.e. for two norms  $\|\cdot\|_A$  and  $\|\cdot\|_B$  there exist constants  $\alpha, \beta > 0$  such that

$$\alpha \|\mathbf{x}\|_A \leq \|\mathbf{x}\|_B \leq \beta \|\mathbf{x}\|_A \quad \text{for all } \mathbf{x} \in \mathcal{V}$$

## Vector norms contd.

### Definition (Quasi-norm)

A **quasi-norm** satisfies all the norm properties except (d) triangle inequality, which is replaced by  $\|\mathbf{x} + \mathbf{y}\| \leq c(\|\mathbf{x}\| + \|\mathbf{y}\|)$  for a constant  $c \geq 1$ .



## Vector norms contd.

### Definition (Quasi-norm)

A **quasi-norm** satisfies all the norm properties except (d) triangle inequality, which is replaced by  $\|\mathbf{x} + \mathbf{y}\| \leq c(\|\mathbf{x}\| + \|\mathbf{y}\|)$  for a constant  $c \geq 1$ .

### Definition (Semi(pseudo)-norm)

A **semi(pseudo)-norm** satisfies all the norm properties except (b) definiteness.

## Vector norms contd.

### Definition (Quasi-norm)

A **quasi-norm** satisfies all the norm properties except (d) triangle inequality, which is replaced by  $\|\mathbf{x} + \mathbf{y}\| \leq c(\|\mathbf{x}\| + \|\mathbf{y}\|)$  for a constant  $c \geq 1$ .

### Definition (Semi(pseudo)-norm)

A **semi(pseudo)-norm** satisfies all the norm properties except (b) definiteness.

### Example

- ▶ The  $\ell_q$ -norm becomes a quasi-norm when  $q \in (0, 1)$  with  $c = 2^{1/q} - 1$
- ▶ The **total variation norm** (TV-norm) defined (in 1D):  
 $\|\mathbf{x}\|_{\text{TV}} := \sum_{i=1}^{p-1} |x_{i+1} - x_i|$  is a **semi-norm** since it fails to satisfy (b);  
 e.g.,  $\mathbf{x} = (1, 1, \dots, 1)^T$  has  $\|\mathbf{x}\|_{\text{TV}} = 0$  even though  $\mathbf{x} \neq \mathbf{0}$ .

## Vector norms contd.

### Definition (“ $\ell_0$ -norm”)

$$\|\mathbf{x}\|_0 = |\{i : x_i \neq 0\}| \quad \equiv \lim_{q \rightarrow 0} \|\mathbf{x}\|_q^q$$

## Vector norms contd.

### Definition (“ $\ell_0$ -norm”)

$$\|\mathbf{x}\|_0 = |\{i : x_i \neq 0\}| \quad \equiv \lim_{q \rightarrow 0} \|\mathbf{x}\|_q^q$$

- The “ $\ell_0$ -norm” counts the non-zero components of  $\mathbf{x}$

## Vector norms contd.

### Definition (“ $\ell_0$ -norm”)

$$\|\mathbf{x}\|_0 = |\{i : x_i \neq 0\}| \quad \equiv \lim_{q \rightarrow 0} \|\mathbf{x}\|_q^q$$

- ▶ The “ $\ell_0$ -norm” counts the non-zero components of  $\mathbf{x}$
- ▶ It is **not** a norm – it does not satisfy norm properties (c) and (d)
- ▶  $\Rightarrow$  it is also neither a **quasi-** nor a **semi-norm**

## Vector norms contd.

### Definition (“ $\ell_0$ -norm”)

$$\|\mathbf{x}\|_0 = |\{i : x_i \neq 0\}| \quad \equiv \lim_{q \rightarrow 0} \|\mathbf{x}\|_q^q$$

- ▶ The “ $\ell_0$ -norm” counts the non-zero components of  $\mathbf{x}$
- ▶ It is **not** a norm – it does not satisfy norm properties (c) and (d)
- ▶  $\Rightarrow$  it is also neither a **quasi-** nor a **semi-norm**

### Example (Compressed sensing)

Find  $\arg \min_{\mathbf{x} \in \mathbb{R}^p} \|\mathbf{x}\|_0$  subject to:  $\mathbf{Ax} = \mathbf{y}$ .

## Vector norms contd.

### Definition (“ $\ell_0$ -norm”)

$$\|\mathbf{x}\|_0 = |\{i : x_i \neq 0\}| \quad \equiv \lim_{q \rightarrow 0} \|\mathbf{x}\|_q^q$$

- ▶ The “ $\ell_0$ -norm” counts the non-zero components of  $\mathbf{x}$
- ▶ It is **not** a norm – it does not satisfy norm properties (c) and (d)
- ▶  $\Rightarrow$  it is also neither a **quasi-** nor a **semi-norm**

### Example (Compressed sensing)

Find  $\arg \min_{\mathbf{x} \in \mathbb{R}^p} \|\mathbf{x}\|_0$  subject to:  $\mathbf{Ax} = \mathbf{y}$ .

### Example (Sparse approximation)

Find  $\arg \min_{\mathbf{x} \in \mathbb{R}^p} \|\mathbf{Ax} - \mathbf{y}\|_2^2$  subject to:  $\|\mathbf{x}\|_0 \leq s$ .

## Vector norms contd.

### Definition (Dual norm)

Let  $\|\cdot\|$  be a norm in  $\mathbb{R}^p$ , then the **dual norm** denoted by  $\|\cdot\|^*$  is defined:

$$\|\mathbf{x}\|^* = \sup_{\|\mathbf{y}\| \leq 1} \mathbf{x}^T \mathbf{y}, \quad \text{for all } \mathbf{x}, \mathbf{y} \in \mathbb{R}^p$$



## Vector norms contd.

### Definition (Dual norm)

Let  $\|\cdot\|$  be a norm in  $\mathbb{R}^p$ , then the **dual norm** denoted by  $\|\cdot\|^*$  is defined:

$$\|\mathbf{x}\|^* = \sup_{\|\mathbf{y}\| \leq 1} \mathbf{x}^T \mathbf{y}, \quad \text{for all } \mathbf{x}, \mathbf{y} \in \mathbb{R}^p$$

- The **dual** of the *dual norm* is the **original (primal) norm**, i.e.,  $\|\mathbf{x}\|^{**} = \|\mathbf{x}\|$

## Vector norms contd.

### Definition (Dual norm)

Let  $\|\cdot\|$  be a norm in  $\mathbb{R}^p$ , then the **dual norm** denoted by  $\|\cdot\|^*$  is defined:

$$\|\mathbf{x}\|^* = \sup_{\|\mathbf{y}\| \leq 1} \mathbf{x}^T \mathbf{y}, \quad \text{for all } \mathbf{x}, \mathbf{y} \in \mathbb{R}^p$$

- ▶ The **dual** of the *dual norm* is the **original (primal) norm**, i.e.,  $\|\mathbf{x}\|^{**} = \|\mathbf{x}\|$
- ▶ Hölder's inequality  $\Rightarrow \|\cdot\|_q$  is a **dual norm** of  $\|\cdot\|_r$  when  $\frac{1}{q} + \frac{1}{r} = 1$ .

## Vector norms contd.

### Example 1

- i)  $\|\cdot\|_2$  is **dual** of  $\|\cdot\|_2$  (i.e.,  $\|\cdot\|_2$  is *self-dual*):  
$$\sup\{\mathbf{z}^T \mathbf{x} \mid \|\mathbf{x}\|_2 \leq 1\} = \|\mathbf{z}\|_2.$$
- ii)  $\|\cdot\|_1$  is **dual** of  $\|\cdot\|_\infty$ , (and *vice versa*):  $\sup\{\mathbf{z}^T \mathbf{x} \mid \|\mathbf{x}\|_\infty \leq 1\} = \|\mathbf{z}\|_1.$

## Vector norms contd.

### Example 1

- i)  $\|\cdot\|_2$  is **dual** of  $\|\cdot\|_2$  (i.e.,  $\|\cdot\|_2$  is *self-dual*):  
$$\sup\{\mathbf{z}^T \mathbf{x} \mid \|\mathbf{x}\|_2 \leq 1\} = \|\mathbf{z}\|_2.$$
- ii)  $\|\cdot\|_1$  is **dual** of  $\|\cdot\|_\infty$ , (and *vice versa*):  $\sup\{\mathbf{z}^T \mathbf{x} \mid \|\mathbf{x}\|_\infty \leq 1\} = \|\mathbf{z}\|_1.$

### Example 2

What is the **dual norm** of  $\|\cdot\|_q$  for  $q = 1 + 1/\log(p)$ ?

## Vector norms contd.

### Example 1

- i)  $\|\cdot\|_2$  is **dual** of  $\|\cdot\|_2$  (i.e.,  $\|\cdot\|_2$  is *self-dual*):  
 $\sup\{\mathbf{z}^T \mathbf{x} \mid \|\mathbf{x}\|_2 \leq 1\} = \|\mathbf{z}\|_2.$
- ii)  $\|\cdot\|_1$  is **dual** of  $\|\cdot\|_\infty$ , (and *vice versa*):  $\sup\{\mathbf{z}^T \mathbf{x} \mid \|\mathbf{x}\|_\infty \leq 1\} = \|\mathbf{z}\|_1.$

### Example 2

What is the **dual norm** of  $\|\cdot\|_q$  for  $q = 1 + 1/\log(p)$ ?

### Solution

By Hölder's inequality,  $\|\cdot\|_r$  is the **dual norm** of  $\|\cdot\|_q$  if  $\frac{1}{q} + \frac{1}{r} = 1$ . Therefore,  $r = 1 + \log(p)$  for  $q = 1 + 1/\log(p)$ .

## Inner product space

## Inner product space

### Definition (Inner product)

An **inner product** on a real vector space  $\mathcal{V}$  is a function  $\langle \cdot, \cdot \rangle : \mathcal{V} \times \mathcal{V} \rightarrow \mathbb{R}$  such that for all  $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathcal{V}$  and  $\alpha, \beta \in \mathbb{R}$  it satisfies

1.  $\langle \mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{y}, \mathbf{x} \rangle$ , (symmetry)
2.  $\langle (\alpha \mathbf{x} + \beta \mathbf{y}), \mathbf{z} \rangle = \alpha \langle \mathbf{x}, \mathbf{z} \rangle + \beta \langle \mathbf{y}, \mathbf{z} \rangle$ , (linearity)
3.  $\langle \mathbf{x}, \mathbf{x} \rangle \geq 0$  (positive definiteness)

## Inner product space

### Definition (Inner product)

An **inner product** on a real vector space  $\mathcal{V}$  is a function  $\langle \cdot, \cdot \rangle : \mathcal{V} \times \mathcal{V} \rightarrow \mathbb{R}$  such that for all  $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathcal{V}$  and  $\alpha, \beta \in \mathbb{R}$  it satisfies

1.  $\langle \mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{y}, \mathbf{x} \rangle$ , (symmetry)
2.  $\langle (\alpha \mathbf{x} + \beta \mathbf{y}), \mathbf{z} \rangle = \alpha \langle \mathbf{x}, \mathbf{z} \rangle + \beta \langle \mathbf{y}, \mathbf{z} \rangle$ , (linearity)
3.  $\langle \mathbf{x}, \mathbf{x} \rangle \geq 0$  (positive definiteness)

- The **inner product** (also **dot product**) of any two vectors  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$  is defined as  $\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^T \mathbf{y} = \sum_i^n x_i y_i$



## Inner product space

### Definition (Inner product)

An **inner product** on a real vector space  $\mathcal{V}$  is a function  $\langle \cdot, \cdot \rangle : \mathcal{V} \times \mathcal{V} \rightarrow \mathbb{R}$  such that for all  $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathcal{V}$  and  $\alpha, \beta \in \mathbb{R}$  it satisfies

1.  $\langle \mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{y}, \mathbf{x} \rangle$ , (symmetry)
2.  $\langle (\alpha \mathbf{x} + \beta \mathbf{y}), \mathbf{z} \rangle = \alpha \langle \mathbf{x}, \mathbf{z} \rangle + \beta \langle \mathbf{y}, \mathbf{z} \rangle$ , (linearity)
3.  $\langle \mathbf{x}, \mathbf{x} \rangle \geq 0$  (positive definiteness)

- The **inner product** (also **dot product**) of any two vectors  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$  is defined as  $\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^T \mathbf{y} = \sum_i^n x_i y_i$
- Important relations involving the inner product:
  - **Hölder's inequality**:  $|\langle \mathbf{x}, \mathbf{y} \rangle| \leq \|\mathbf{x}\|_q \|\mathbf{y}\|_r$ , where  $r > 1$  and  $\frac{1}{q} + \frac{1}{r} = 1$
  - **Cauchy-Schwarz** is a special case of Hölder's inequality ( $q = r = 2$ )